



## 11th meeting of the BRICS Astronomy Working Group

13 to 17 October 2025

Instituto Nacional de Pesquisas Espaciais (INPE)  
São José dos Campos, São Paulo, Brasil

# The Impact of $\beta$ -Dependent Tearing Instability Suppression on Cosmic Ray Acceleration and Multi-Messenger Transients

Gabriel L. Ferreira-Santos

First Name:	Gabriel L.
Last Name:	Ferreira-Santos
Institution/Affiliation:	INPE
Country of Residence:	Brazil
Preferred type of presentation	Oral
Will you attend in person or online?	—
Email	<a href="mailto:gabriel.ferreira@inpe.br">gabriel.ferreira@inpe.br</a>

### Abstract

Magnetic reconnection is a primary mechanism for powering energetic, transient astrophysical phenomena, with profound implications for multi-messenger astronomy. Theoretical estimates for reconnection rates rely on the presence/absence of instabilities, such as the Tearing Instability, whose growth rate in classical magnetohydrodynamic (MHD) theory is independent of the plasma- $\beta$  (the ratio of thermal to magnetic pressure). However, many astrophysical environments are filled with a weakly collisional plasma and are characterized by high- $\beta$  conditions, where the validity of classical MHD is limited. This work investigates the theory of Tearing Instability under these more realistic conditions using a non-ideal gyrotropic-MHD framework. Our analysis revealed a novel and previously unknown scaling relation for the instability's maximum growth rate in high- $\beta$  regimes, which scales as  $\gamma_{\text{max}} \propto \beta^{-1/4}$ . This suppression arises from dynamically generated pressure anisotropy that introduces a stabilizing restoring force, absent in standard MHD. Our results suggest that the onset of plasmoid-mediated fast reconnection and subsequent particle acceleration by first order Fermi process may be significantly inhibited or delayed in high- $\beta$  regions of astrophysical sources. For multi-messenger astronomy, this  $\beta$ -dependent suppression introduces a crucial physical constraint on the production of UHECRs, VHE gamma-rays, and high-energy neutrinos. This challenges the universal applicability of reconnection models that neglect pressure anisotropy and suggests that plasma- $\beta$  could be a key modulating factor in the multi-messenger output of the most energetic cosmic accelerators.