



## 11th meeting of the BRICS Astronomy Working Group

13 to 17 October 2025

Instituto Nacional de Pesquisas Espaciais (INPE)  
São José dos Campos, São Paulo, Brasil

# Compton-induced cascade $\gamma$ -rays in radio galaxy NGC 1275.

Mfuphi Ntshatsha

First Name:	Mfuphi
Last Name:	Ntshatsha
Institution/Affiliation:	University of Johannesburg
Country of Residence:	South Africa
Preferred type of presentation	Oral
Will you attend in person or online?	—
Email	mfuphin95@gmail.com

### Abstract

Among active galactic nuclei (AGN), blazars are the brightest emitters of high- (HE,  $E \geq 100$  MeV) to very-high-energy (VHE,  $E \geq 100$  GeV)  $\gamma$ -rays from their jets. Radio galaxies, being the misaligned parent population of the blazar class, were historically not observed at these frequencies. However, there is a growing number of radio galaxies detected in HE-VHE  $\gamma$ -rays in recent years. In this work, we leverage and refine a Monte-Carlo photon and electron-positron ( $e^\pm$ ) pair tracking code in the AGN environment of the radio galaxy NGC 1275. In the code, we consider the isotropic broad emission line and anisotropic Shakura-Sunyaev (SS) accretion disk radiation fields, with mild magnetic fields in the AGN environment. We find that cascade  $\gamma$ -rays from inverse-Compton scattering by relativistic  $e^\pm$  pairs of these external radiation fields can explain the *Fermi* Large Area Telescope's (LAT) and Major Atmospheric Cherenkov Experiment (MACE) observations from the radio galaxy NGC 1275. We present a set of plausible parameters obtained from the code by fitting the source's spectral energy distribution (SED) during flaring events reported in the period December 2022 to January 2023.