

An efficient real-space density functional algorithm^{1,2}

with S. A. Chin³, S. Janecek^{1,4}, M. Liebrecht¹ and R. Zillich¹

¹Institut für theoretische Physik, Johannes Kepler Universität, Linz, Austria

²Department of Physics, University at Buffalo SUNY, Buffalo NY 14260

³Department of Physics, Texas A&M University, USA

⁴Institut de Ciència de Materials de Barcelona (ICMAB-CSIC), Campus de Bellaterra,
08193 Barcelona, Spain

Oct. 5, 2010



FWF

Der Wissenschaftsfonds.


- 1 First exposures to DFT
 - Surface energy of simple metals
 - Closed-shell atoms
- 2 Density Functional Theory
 - Messing with the Many-Body Problem
 - Operator factorizations
 - High-order propagation
 - Operator factorizations in a magnetic field
 - Gauge invariance
- 3 Some results
 - Magnetic susceptibilities
 - NMR shifts
- 4 Clusters in a quantum matrix
 - Magnesium-Helium Interaction
 - Combination of DFT and PIMC
- 5 Summary

My first exposures to DFT

E. K. and Walter Kohn, **PRL57**, 862 (1986)

Surface energy for simple model systems in:

- LDA (Lang, Kohn, 1971)




My first exposures to DFT

E. K. and Walter Kohn, **PRL57**, 862 (1986)

Surface energy for simple model systems in:

- LDA (Lang, Kohn, 1971)
- JF-variational theory (EK, Qian, Kohn 1985)




My first exposures to DFT

E. K. and Walter Kohn, **PRL**57, 862 (1986)

Surface energy for simple model systems in:

- LDA (Lang, Kohn, 1971)
- JF-variational theory (EK, Qian, Kohn 1985)
- Diffusion Monte Carlo (Li, Needs, Martin, Ceperley 1992)




My first exposures to DFT

E. K. and Walter Kohn, **PRL**57, 862 (1986)

Surface energy for simple model systems in:

- LDA (Lang, Kohn, 1971)
- JF-variational theory (EK, Qian, Kohn 1985)
- Diffusion Monte Carlo (Li, Needs, Martin, Ceperley 1992)



Conclusion

- LDA is significantly off
- LDA must be fixed by ad-hoc corrections (GGA) **BUT**
- LDA is the only choice in complicated geometries

Surface energy of simple metals


Poking into a wasp-nest

- FHNC-EL generalizable to non-uniform Fermi systems
(E. K. and W. Kohn and G.-X. Qian: PR **B32**, 5693 (1985);
E. K. and W. Kohn: PRL **57**, 862 (1986).)

Surface energy of simple metals

Poking into a wasp-nest


- FHNC-EL generalizable to non-uniform Fermi systems (E. K. and W. Kohn and G.-X. Qian: PR **B32**, 5693 (1985); E. K. and W. Kohn: PRL **57**, 862 (1986).)
- Significant difference between LDA and HNC (“the authors should explain why HNC works so poorly”) ☹



Surface energy of simple metals

Poking into a wasp-nest


- FHNC-EL generalizable to non-uniform Fermi systems (E. K. and W. Kohn and G.-X. Qian: PR **B32**, 5693 (1985); E. K. and W. Kohn: PRL **57**, 862 (1986).)
- Significant difference between LDA and HNC (“the authors should explain why HNC works so poorly”) ☹
- Excellent agreement of FHNC-EL for metallic surface energies with GFMC ☺



Analysis of surface energies

... a learning experience in many-body physics


- Try LDA;



Analysis of surface energies

... a learning experience in many-body physics

- Try LDA;
- Try RPA;




Analysis of surface energies

... a learning experience in many-body physics

- Try LDA;
- Try RPA;
- Try RPA with a “LDA screened interaction”


$$V_{ph}(r) = v_c(r)C_{bulk}(r, r_{cm});$$



Analysis of surface energies

... a learning experience in many-body physics


- Try LDA;
- Try RPA;
- Try RPA with a “LDA screened interaction”
$$V_{ph}(r) = v_c(r)C_{bulk}(r, r_{cm});$$
- Try RPA with a non-locally screened interaction $V_{ph}(\mathbf{r}_1, \mathbf{r}_2) = v_c(|\mathbf{r}_1 - \mathbf{r}_2|)C(\mathbf{r}_1 - \mathbf{r}_2, \mathbf{r}_{cm});$



Analysis of surface energies

... a learning experience in many-body physics

- Try LDA;
- Try RPA;
- Try RPA with a “LDA screened interaction”
 $V_{ph}(r) = v_c(r)C_{bulk}(r, r_{cm});$
- Try RPA with a non-locally screened interaction $V_{ph}(\mathbf{r}_1, \mathbf{r}_2) = v_c(|\mathbf{r}_1 - \mathbf{r}_2|)C(\mathbf{r}_1 - \mathbf{r}_2, \mathbf{r}_{cm});$
- **Conclusion:** The local structure is just different !



Closed-shell atoms

Correlation energy – everything beyond Hartree-Fock

	<i>expt.</i>	<i>FHNC</i>	<i>LDA</i>	<i>GGA</i>	<i>GGA'</i>
<i>Be</i>	-0.0944	-0.107	-0.224	-0.099	-0.094
<i>Ne</i> ⁺⁶	-0.18	-0.109	-0.333	-0.026	-0.136
<i>Ne</i>	-0.393	-0.421	-0.74	-0.41	-0.39
<i>Na</i> ⁺¹	-0.396	-0.419			
<i>Mg</i> ⁺²	-0.402	-0.418			
<i>Al</i> ⁺³	-0.409	-0.418			
<i>Si</i> ⁺⁴	-0.417	-0.419			
<i>P</i> ⁺⁵	-0.426	-0.419			
<i>S</i> ⁺⁶	-0.434	-0.421			
...					

“LDA is 99.5 percent accurate” (A well known DFT practitioner)

Density Functional Theory

Messing with the Many-Body Problem

Electronic Hamiltonian in a magnetic field

$$H = \underbrace{\frac{1}{2m} \sum_{i=1}^N [-i\hbar \nabla_i + e\mathbf{A}(\mathbf{r}_i)]^2}_1 + \underbrace{\sum_{i<j}^N \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|}}_2 + \underbrace{\sum_{i=1}^N V_{\text{ext}}(\mathbf{r}_i)}_3$$

Density Functional Theory

Messing with the Many-Body Problem

Electronic Hamiltonian in a magnetic field

$$H = \underbrace{\frac{1}{2m} \sum_{i=1}^N [-i\hbar \nabla_i + e\mathbf{A}(\mathbf{r}_i)]^2}_1 + \underbrace{\sum_{i<j}^N \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|}}_2 + \underbrace{\sum_{i=1}^N V_{\text{ext}}(\mathbf{r}_i)}_3$$

(1) **Kinetic energy.** Vector potential \mathbf{A} .

Density Functional Theory

Messing with the Many-Body Problem

Electronic Hamiltonian in a magnetic field

$$H = \underbrace{\frac{1}{2m} \sum_{i=1}^N [-i\hbar \nabla_i + e\mathbf{A}(\mathbf{r}_i)]^2}_1 + \underbrace{\sum_{i<j}^N \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|}}_2 + \underbrace{\sum_{i=1}^N V_{\text{ext}}(\mathbf{r}_i)}_3$$

(1) Kinetic energy. Vector potential \mathbf{A} .

(2) Electron-electron interaction.

Density Functional Theory

Messing with the Many-Body Problem

Electronic Hamiltonian in a magnetic field

$$H = \underbrace{\frac{1}{2m} \sum_{i=1}^N [-i\hbar \nabla_i + e\mathbf{A}(\mathbf{r}_i)]^2}_1 + \underbrace{\sum_{i<j}^N \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|}}_2 + \underbrace{\sum_{i=1}^N V_{\text{ext}}(\mathbf{r}_i)}_3$$

(1) Kinetic energy. Vector potential \mathbf{A} .

(2) Electron-electron interaction.

(3) Ion core (pseudo-)potentials. A mess....

Kohn-Sham Equation

Mapping the many-body problem onto a one-body problem

Kohn-Sham equations:

Effective Schrödinger (“Kohn-Sham”) equation

$$\left[\frac{1}{2m} (\mathbf{p} + e\mathbf{A}(\mathbf{r}))^2 + \hat{V}_{\text{KS}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r}) \right] \psi_{i,\sigma}(\mathbf{r}) = \epsilon_{i,\sigma} \psi_{i,\sigma}(\mathbf{r})$$

KS-wavefunctions $\psi_{i,\sigma}(\mathbf{r})$

Kohn-Sham Equation

Mapping the many-body problem onto a one-body problem

Kohn-Sham equations:

Effective Schrödinger (“Kohn-Sham”) equation

$$\left[\frac{1}{2m} (\mathbf{p} + e\mathbf{A}(\mathbf{r}))^2 + \hat{V}_{\text{KS}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r}) \right] \psi_{i,\sigma}(\mathbf{r}) = \epsilon_{i,\sigma} \psi_{i,\sigma}(\mathbf{r})$$

Densities

$$\rho_{\sigma}(\mathbf{r}) = \sum_{i=1}^{N_{\sigma}} |\psi_{i,\sigma}(\mathbf{r})|^2, \quad \rho(\mathbf{r}) = \sum_{\sigma} \rho_{\sigma}(\mathbf{r})$$

Kohn-Sham Equation

Mapping the many-body problem onto a one-body problem

Kohn-Sham equations:

Effective Schödinger (“Kohn-Sham”) equation

$$\left[\frac{1}{2m}(\mathbf{p} + e\mathbf{A}(\mathbf{r}))^2 + \hat{V}_{\text{KS}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r}) \right] \psi_{i,\sigma}(\mathbf{r}) = \epsilon_{i,\sigma} \psi_{i,\sigma}(\mathbf{r})$$

Densities

$$\rho_{\sigma}(\mathbf{r}) = \sum_{i=1}^{N_{\sigma}} |\psi_{i,\sigma}(\mathbf{r})|^2, \quad \rho(\mathbf{r}) = \sum_{\sigma} \rho_{\sigma}(\mathbf{r})$$

Effective potential

$$V_{\text{KS}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r}) = \hat{V}_{\text{ext}} + V_{\text{c}}[\rho](\mathbf{r}) + V_{\text{xc}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r})$$

Kohn-Sham Equation

Mapping the many-body problem onto a one-body problem

Kohn-Sham equations:

Effective Schödinger (“Kohn-Sham”) equation

$$\left[\frac{1}{2m}(\mathbf{p} + e\mathbf{A}(\mathbf{r}))^2 + \hat{V}_{\text{KS}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r}) \right] \psi_{i,\sigma}(\mathbf{r}) = \epsilon_{i,\sigma} \psi_{i,\sigma}(\mathbf{r})$$

Densities

$$\rho_{\sigma}(\mathbf{r}) = \sum_{i=1}^{N_{\sigma}} |\psi_{i,\sigma}(\mathbf{r})|^2, \quad \rho(\mathbf{r}) = \sum_{\sigma} \rho_{\sigma}(\mathbf{r})$$

Effective potential

$$V_{\text{KS}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r}) = \hat{V}_{\text{ext}} + V_{\text{c}}[\rho](\mathbf{r}) + V_{\text{xc}}^{\sigma}[\rho^{\uparrow}, \rho^{\downarrow}](\mathbf{r})$$

Self-consistency iterations

Solving Schrödinger-like equations

The imaginary timestep method

Define: *evolution operator* $\mathcal{T}(\epsilon) \equiv e^{-\epsilon \hat{H}}$

Solving Schrödinger-like equations

The imaginary timestep method

Define: *evolution operator* $\mathcal{T}(\epsilon) \equiv e^{-\epsilon \hat{H}}$

Algorithm

Solving Schrödinger-like equations

The imaginary timestep method

Define: *evolution operator* $\mathcal{T}(\epsilon) \equiv e^{-\epsilon \hat{H}}$

Algorithm

- 1 begin with initial set of states $\{\psi_1^{(0)} \dots \psi_n^{(0)}\}$

Solving Schrödinger-like equations

The imaginary timestep method

Define: *evolution operator* $\mathcal{T}(\epsilon) \equiv e^{-\epsilon\hat{H}}$

Algorithm

- 1 begin with initial set of states $\{\psi_1^{(0)} \dots \psi_n^{(0)}\}$
- 2 for $k = 0, \dots$ apply evolution operator:

$$\phi_i^{(k+1)}(\mathbf{r}) = \mathcal{T}(\epsilon)\psi_i^{(k)}$$

Solving Schrödinger-like equations

The imaginary timestep method

Define: *evolution operator* $\mathcal{T}(\epsilon) \equiv e^{-\epsilon \hat{H}}$

Algorithm

- 1 begin with initial set of states $\{\psi_1^{(0)} \dots \psi_n^{(0)}\}$
- 2 for $k = 0, \dots$ apply evolution operator:
$$\phi_i^{(k+1)}(\mathbf{r}) = \mathcal{T}(\epsilon)\psi_i^{(k)}$$
- 3 orthonormalize $\psi_i^{(k+1)} = \sum_j c_{ij} \phi_j^{(k+1)}$, go to (2)

Solving Schrödinger-like equations

The imaginary timestep method

Define: *evolution operator* $\mathcal{T}(\epsilon) \equiv e^{-\epsilon \hat{H}}$

Algorithm

- 1 begin with initial set of states $\{\psi_1^{(0)} \dots \psi_n^{(0)}\}$
- 2 for $k = 0, \dots$ apply evolution operator:
$$\phi_i^{(k+1)}(\mathbf{r}) = \mathcal{T}(\epsilon)\psi_i^{(k)}$$
- 3 orthonormalize $\psi_i^{(k+1)} = \sum_j c_{ij} \phi_j^{(k+1)}$, go to (2)

Then:

- The sequence $\{\psi_i^{(k)}(\mathbf{r})\}$ converges towards the low-lying eigenstates.

Eigenvalue solver – operator factorizations

Calculating $e^{-\epsilon \hat{H}} = e^{-\epsilon(\hat{T} + \hat{V})}$

Eigenvalue solver – operator factorizations

$$\text{Calculating } e^{-\epsilon \hat{H}} = e^{-\epsilon(\hat{T} + \hat{V})}$$

“Trotter Formula” - Second order factorization

$$\mathcal{T}_2(\epsilon) = e^{-\frac{1}{2}\epsilon \hat{V}} e^{-\epsilon \hat{T}} e^{-\frac{1}{2}\epsilon \hat{V}} + \mathcal{O}(\epsilon^3)$$

- Calculate $e^{-\frac{1}{2}\epsilon \hat{V}}$ in real space, $e^{-\epsilon \hat{T}}$ in momentum space.

Eigenvalue solver – operator factorizations

$$\text{Calculating } e^{-\epsilon \hat{H}} = e^{-\epsilon(\hat{T} + \hat{V})}$$

“Trotter Formula” - Second order factorization

$$\mathcal{T}_2(\epsilon) = e^{-\frac{1}{2}\epsilon \hat{V}} e^{-\epsilon \hat{T}} e^{-\frac{1}{2}\epsilon \hat{V}} + \mathcal{O}(\epsilon^3)$$

- Calculate $e^{-\frac{1}{2}\epsilon \hat{V}}$ in real space, $e^{-\epsilon \hat{T}}$ in momentum space.

Convergence rate

- Evolution time τ to obtain a certain accuracy depends on \hat{V} , but not on the exact form of the factorization of $\mathcal{T}(\epsilon)$;
- \Rightarrow Number of iterations needed: $N \approx \tau/\epsilon$

Eigenvalue solver – operator factorizations

$$\text{Calculating } e^{-\epsilon \hat{H}} = e^{-\epsilon(\hat{T} + \hat{V})}$$

“Trotter Formula” - Second order factorization

$$\mathcal{T}_2(\epsilon) = e^{-\frac{1}{2}\epsilon \hat{V}} e^{-\epsilon \hat{T}} e^{-\frac{1}{2}\epsilon \hat{V}} + \mathcal{O}(\epsilon^3)$$

- Calculate $e^{-\frac{1}{2}\epsilon \hat{V}}$ in real space, $e^{-\epsilon \hat{T}}$ in momentum space.

Convergence rate

- Evolution time τ to obtain a certain accuracy depends on \hat{V} , but not on the exact form of the factorization of $\mathcal{T}(\epsilon)$;
- \Rightarrow Number of iterations needed: $N \approx \tau/\epsilon$

Large ϵ
Small ϵ



Fast convergence, poor accuracy
Slow convergence, good accuracy

Fourth order factorization

Suzuki '96, Chin '97, Chin and Forbert '01

$$\mathcal{T}_4(\epsilon) = e^{-\frac{1}{6}\epsilon\hat{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{2}{3}\epsilon\tilde{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{1}{6}\epsilon\hat{V}} + \mathcal{O}(\epsilon^5)$$

$$\tilde{V} = \hat{V} + \frac{\epsilon^2}{48}[\hat{V}, [\hat{T}, \hat{V}]] \Rightarrow \tilde{V}(\mathbf{r}) = V(\mathbf{r}) + \frac{\hbar^2\epsilon^2}{48m}|\nabla V(\mathbf{r})|^2$$

Fourth order factorization

Suzuki '96, Chin '97, Chin and Forbert '01

$$\mathcal{T}_4(\epsilon) = e^{-\frac{1}{6}\epsilon\hat{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{2}{3}\epsilon\tilde{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{1}{6}\epsilon\hat{V}} + \mathcal{O}(\epsilon^5)$$

$$\tilde{V} = \hat{V} + \frac{\epsilon^2}{48}[\hat{V}, [\hat{T}, \hat{V}]] \Rightarrow \tilde{V}(\mathbf{r}) = V(\mathbf{r}) + \frac{\hbar^2\epsilon^2}{48m}|\nabla V(\mathbf{r})|^2$$

- Local potential: \Rightarrow calculate in coordinate space

Fourth order factorization

Suzuki '96, Chin '97, Chin and Forbert '01

$$\mathcal{T}_4(\epsilon) = e^{-\frac{1}{6}\epsilon\hat{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{2}{3}\epsilon\tilde{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{1}{6}\epsilon\hat{V}} + \mathcal{O}(\epsilon^5)$$

$$\tilde{V} = \hat{V} + \frac{\epsilon^2}{48}[\hat{V}, [\hat{T}, \hat{V}]] \Rightarrow \tilde{V}(\mathbf{r}) = V(\mathbf{r}) + \frac{\hbar^2\epsilon^2}{48m}|\nabla V(\mathbf{r})|^2$$

- Local potential: \Rightarrow calculate in coordinate space
- Double commutator term $\propto |\nabla V(\mathbf{r})|^2$: \Rightarrow calculate in coordinate space.

Fourth order factorization

Suzuki '96, Chin '97, Chin and Forbert '01

$$\mathcal{T}_4(\epsilon) = e^{-\frac{1}{6}\epsilon\hat{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{2}{3}\epsilon\tilde{V}} e^{-\frac{1}{2}\epsilon\hat{T}} e^{-\frac{1}{6}\epsilon\hat{V}} + \mathcal{O}(\epsilon^5)$$

$$\tilde{V} = \hat{V} + \frac{\epsilon^2}{48}[\hat{V}, [\hat{T}, \hat{V}]] \Rightarrow \tilde{V}(\mathbf{r}) = V(\mathbf{r}) + \frac{\hbar^2\epsilon^2}{48m}|\nabla V(\mathbf{r})|^2$$


- Local potential: \Rightarrow calculate in coordinate space
- Double commutator term $\propto |\nabla V(\mathbf{r})|^2$: \Rightarrow calculate in coordinate space.
- Kinetic energy: \Rightarrow calculate in momentum space

Plane waves are eigenstates of \hat{T} in any uniform discretization

$$t_n(k) = \left(\frac{2\sin\kappa}{h}\right)^2 \left[1 + \frac{1}{3}\sin^2\kappa + \frac{8}{45}\sin^4\kappa + \frac{4}{35}\sin^6\kappa + \dots\right]$$

$$\kappa = kh/2 \quad h: \text{discretization step size}$$

Convergence Comparison



- typical speed gain (system dependent): 10 - 100
- expectation energy $\langle \psi_i(\epsilon) | \hat{H} | \psi_i(\epsilon) \rangle = e_i(\epsilon) + \mathcal{O}(\epsilon^8)$ gives good convergence tests.

High-order propagation: ...

Can we do even better ?

The “nogo theorem” (Chin, Suzuki)

- There is no factorization of the diffusion operator of the form

$$\mathcal{T}(\epsilon) = e^{-\epsilon(\hat{T} + \hat{V})} = \prod_{i=1}^M e^{-a_i \epsilon \hat{T}} e^{-b_i \epsilon \hat{V}} + \mathcal{O}(\epsilon^N)$$

with **positive coefficients** a_i, b_i

High-order propagation: ...

Can we do even better ?

The “nogo theorem” (Chin, Suzuki)

- There is no factorization of the diffusion operator of the form

$$\mathcal{T}(\epsilon) = e^{-\epsilon(\hat{T} + \hat{V})} = \prod_{i=1}^M e^{-a_i \epsilon \hat{T}} e^{-b_i \epsilon \hat{V}} + \mathcal{O}(\epsilon^N)$$

with **positive coefficients** a_i, b_i

- There is no single-product approximation of order higher than 4.

High-order propagation: ...

Can we do even better ?

The “nogo theorem” (Chin, Suzuki) and the way out

- There is no factorization of the diffusion operator of the form

$$\mathcal{T}(\epsilon) = e^{-\epsilon(\hat{T} + \hat{V})} = \prod_{i=1}^M e^{-a_i \epsilon \hat{T}} e^{-b_i \epsilon \hat{V}} + \mathcal{O}(\epsilon^N)$$

with **positive coefficients** a_i, b_i

- There is no single-product approximation of order higher than 4.
- Multi-Product Expansion are a way out

$$e^{-\epsilon(\hat{T} + \hat{V})} = \sum_{k=1}^n c_k \mathcal{T}_2^k \left(\frac{\epsilon}{k} \right) + \mathcal{O}(\epsilon^{2n+1}) \equiv \mathcal{T}_n(\epsilon) + \mathcal{O}(\epsilon^{2n+1}),$$


coefficients:

$$c_i = \prod_{j=1(\neq i)}^n k_i^2 / (k_i^2 - k_j^2)$$

High-Order propagation

How easy is it to solve the Schrödinger equation in 3D ?


- Higher order methods can converge in very few steps



High-Order propagation

How easy is it to solve the Schrödinger equation in 3D ?


- Higher order methods can converge in very few steps
- Diffusion is typically an order of magnitude faster than Lanczos



High-Order propagation

How easy is it to solve the Schrödinger equation in 3D ?


- Higher order methods can converge in very few steps
- Diffusion is typically an order of magnitude faster than Lanczos
- Higher-order methods are computationally more costly but easy to parallelize



High-Order propagation

How easy is it to solve the Schrödinger equation in 3D ?

- Higher order methods can converge in very few steps
- Diffusion is typically an order of magnitude faster than Lanczos
- Higher-order methods are computationally more costly but easy to parallelize




Solving a local Schrödinger equation in 3D is embarrassingly easy

Program for download:

S. Janecek and E. K., Computer Physics
Communications 178, 835 (2008)

Convergence for realistic systems





Convergence deteriorates very little

- for small systems: CO

Recall $N \propto 1/\epsilon$

Convergence for realistic systems




Convergence deteriorates very little

- for small systems: CO
- for large systems: C_{60}

Recall $N \propto 1/\epsilon$

Convergence for realistic systems





Convergence deteriorates very little

- for small systems: CO
- for large systems: C_{60}
- ...to be quantitative: CO

Recall $N \propto 1/\epsilon$

Convergence for realistic systems



Convergence deteriorates very little

- for small systems: CO
- for large systems: C_{60}
- ...to be quantitative: CO
- ...to impress: C_{60}

Recall $N \propto 1/\epsilon$

Homogenous Magnetic Fields

$$e^{-\frac{1}{2}\epsilon\hat{T}} : \hat{T} = \frac{1}{2m}\mathbf{P}^2 \rightarrow \hat{T} = \frac{1}{2m}\mathbf{\Pi}^2 = \frac{1}{2m}(\Pi_x^2 + \Pi_y^2),$$
$$\mathbf{\Pi} = \mathbf{p} + e\mathbf{A}(\mathbf{r})$$

Homogenous Magnetic Fields

$$e^{-\frac{1}{2}\epsilon\hat{T}} : \hat{T} = \frac{1}{2m}\mathbf{p}^2 \rightarrow \hat{T} = \frac{1}{2m}\mathbf{\Pi}^2 = \frac{1}{2m}(\Pi_x^2 + \Pi_y^2),$$
$$\mathbf{\Pi} = \mathbf{p} + e\mathbf{A}(\mathbf{r})$$

- $e^{-\frac{1}{2}\epsilon\hat{T}}$ not longer diagonal in Fourier space

Homogenous Magnetic Fields

$$e^{-\frac{1}{2}\epsilon\hat{T}} : \hat{T} = \frac{1}{2m}\mathbf{p}^2 \rightarrow \hat{T} = \frac{1}{2m}\mathbf{\Pi}^2 = \frac{1}{2m}(\Pi_x^2 + \Pi_y^2),$$
$$\mathbf{\Pi} = \mathbf{p} + e\mathbf{A}(\mathbf{r})$$

- $e^{-\frac{1}{2}\epsilon\hat{T}}$ not longer diagonal in Fourier space
- free electrons in homogenous magnetic field \leftrightarrow 1D harmonic oscillator (Landau '30)

Homogenous Magnetic Fields

$$e^{-\frac{1}{2}\epsilon\hat{T}} : \hat{T} = \frac{1}{2m}\mathbf{p}^2 \rightarrow \hat{T} = \frac{1}{2m}\mathbf{\Pi}^2 = \frac{1}{2m}(\Pi_x^2 + \Pi_y^2),$$
$$\mathbf{\Pi} = \mathbf{p} + e\mathbf{A}(\mathbf{r})$$

- $e^{-\frac{1}{2}\epsilon\hat{T}}$ not longer diagonal in Fourier space
- free electrons in homogenous magnetic field \leftrightarrow 1D harmonic oscillator (Landau '30)
- density matrix for 1D harmonic oscillator can be calculated exactly (Feynman, Statistical Mechanics)

Homogenous Magnetic Fields

$$e^{-\frac{1}{2}\epsilon\hat{T}} : \hat{T} = \frac{1}{2m}\mathbf{p}^2 \rightarrow \hat{T} = \frac{1}{2m}\mathbf{\Pi}^2 = \frac{1}{2m}(\Pi_x^2 + \Pi_y^2),$$
$$\mathbf{\Pi} = \mathbf{p} + e\mathbf{A}(\mathbf{r})$$

- $e^{-\frac{1}{2}\epsilon\hat{T}}$ not longer diagonal in Fourier space
- free electrons in homogenous magnetic field \leftrightarrow 1D harmonic oscillator (Landau '30)
- density matrix for 1D harmonic oscillator can be calculated exactly (Feynman, Statistical Mechanics)

Exact Factorization

$$e^{-\frac{\epsilon}{2m}(\Pi_x^2 + \Pi_y^2)} = e^{-\frac{\epsilon}{4m}C_E(\xi)\Pi_x^2} e^{-\frac{\epsilon}{2m}C_M(\xi)\Pi_y^2} e^{-\frac{\epsilon}{4m}C_E(\xi)\Pi_x^2}$$

$$\xi = \epsilon \frac{\hbar e B}{m}, \quad C_E(\xi), C_M(\xi) : \text{simple analytic functions}$$

Exact calculation of the kinetic energy: Algorithm

Redo kinetic energy:

$$\underbrace{e^{-\epsilon \hat{T}} \psi(x, y, z)}_{(4)} = \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2}}_{(3)} \underbrace{e^{-\frac{\epsilon}{2m} C_M(\xi) \Pi_y^2}}_{(2)} \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2} e^{-\frac{\epsilon}{2m} p_z^2} \psi(x, y, x)}_{(1)}$$

Exact calculation of the kinetic energy: Algorithm

Redo kinetic energy:

$$\underbrace{e^{-\epsilon \hat{T}} \psi(\mathbf{x}, \mathbf{y}, \mathbf{z})}_{(4)} = \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2}}_{(3)} \underbrace{e^{-\frac{\epsilon}{2m} C_M(\xi) \Pi_y^2}}_{(2)} \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2} e^{-\frac{\epsilon}{2m} p_z^2} \psi(\mathbf{x}, \mathbf{y}, \mathbf{x})}_{(1)}$$

Algorithm: Choose an n -point derivative formula

- (1) For all \mathbf{y} , Fourier transform the \mathbf{x} and the \mathbf{z} -coordinate of each state to (k_x, k_z) -space.

Multiply by $e^{-\frac{\epsilon \hbar^2}{2m} C_E(\xi) t_n(k_x + eB y / \hbar) - \frac{\epsilon \hbar^2}{2m} t_n(k_z)}$.

\Rightarrow N 2d-Fourier transforms.

Counting operations

$$N \times N^2 \ln N^2$$

Exact calculation of the kinetic energy: Algorithm

Redo kinetic energy:

$$\underbrace{e^{-\epsilon \hat{T}} \psi(\mathbf{x}, \mathbf{y}, \mathbf{z})}_{(4)} = \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2}}_{(3)} \underbrace{e^{-\frac{\epsilon}{2m} C_M(\xi) \Pi_y^2}}_{(2)} \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2} e^{-\frac{\epsilon}{2m} p_z^2} \psi(\mathbf{x}, \mathbf{y}, \mathbf{x})}_{(1)}$$

Algorithm: Choose an n -point derivative formula

- (2) For all k_x, k_z , Fourier transform now y to k_y -space.

Multiply by $e^{-\frac{\epsilon \hbar^2 t_n(k_y)}{2m} C_M(\xi)}$.

$\Rightarrow N^2$ 1d-Fourier transforms.

Counting operations

$$N \times N^2 \ln N^2 + N^2 \times N \ln N$$

Exact calculation of the kinetic energy: Algorithm

Redo kinetic energy:

$$\underbrace{e^{-\epsilon \hat{T}} \psi(x, y, z)}_{(4)} = \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2}}_{(3)} \underbrace{e^{-\frac{\epsilon}{2m} C_M(\xi) \Pi_y^2}}_{(2)} \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2} e^{-\frac{\epsilon}{2m} p_z^2} \psi(x, y, x)}_{(1)}$$

Algorithm: Choose an n -point derivative formula

- (3) For all k_x, k_z , do the inverse transformation back to y .
Multiply by $e^{-\frac{\epsilon \hbar^2}{2m} C_E(\xi) t_n(k_x + eBy/\hbar)^2}$.
 $\Rightarrow N^2$ 1d-Fourier transforms.

Counting operations

$$N \times N^2 \ln N^2 + N^2 \times N \ln N + N^2 \times N \ln N$$

Exact calculation of the kinetic energy: Algorithm

Redo kinetic energy:

$$\underbrace{e^{-\epsilon \hat{T}} \psi(\mathbf{x}, \mathbf{y}, \mathbf{z})}_{(4)} = \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2}}_{(3)} \underbrace{e^{-\frac{\epsilon}{2m} C_M(\xi) \Pi_y^2}}_{(2)} \underbrace{e^{-\frac{\epsilon}{4m} C_E(\xi) \Pi_x^2} e^{-\frac{\epsilon}{2m} p_z^2} \psi(\mathbf{x}, \mathbf{y}, \mathbf{x})}_{(1)}$$

Algorithm: Choose an n -point derivative formula

- (4) For all y , Fourier transform now k_x, k_z back to the (x, z) -space.
 \Rightarrow N 2d-Fourier transforms.

Counting operations

$$N \times N^2 \ln N^2 + N^2 \times N \ln N + N^2 \times N \ln N + N \times N^2 \ln N^2 = 2N^3 \ln N^3$$

Operations: \Rightarrow Same as two 3D-FFt's !

Gauge invariance

What we learned in E&M and quantum mechanics

- The gauge transformation

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}'(\mathbf{r}) = \mathbf{A}(\mathbf{r}) + \nabla\chi(\mathbf{r})$$

$$\psi(\mathbf{r}) \rightarrow \psi'(\mathbf{r}) = e^{ie\chi(\mathbf{r})/\hbar}\psi(\mathbf{r})$$

does not change the physics.

Gauge invariance

What we learned in E&M and quantum mechanics

- The gauge transformation

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}'(\mathbf{r}) = \mathbf{A}(\mathbf{r}) + \nabla\chi(\mathbf{r})$$

$$\psi(\mathbf{r}) \rightarrow \psi'(\mathbf{r}) = e^{ie\chi(\mathbf{r})/\hbar}\psi(\mathbf{r})$$

does not change the physics.

- This is true only if we have no discretization errors.
For example: For a discretized derivative operator

$$f'(x) \approx \frac{f(x+h) - f(x-h)}{2h}$$

chain- and product rules are only approximately satisfied.

What we learned in E&M and quantum mechanics

- The gauge transformation

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}'(\mathbf{r}) = \mathbf{A}(\mathbf{r}) + \nabla\chi(\mathbf{r})$$

$$\psi(\mathbf{r}) \rightarrow \psi'(\mathbf{r}) = e^{ie\chi(\mathbf{r})/\hbar}\psi(\mathbf{r})$$

does not change the physics.

- This is true only if we have no discretization errors.
For example: For a discretized derivative operator

$$f'(x) \approx \frac{f(x+h) - f(x-h)}{2h}$$

chain- and product rules are only approximately satisfied.

- However: To be efficient, we want the mesh as coarse as possible.

Discrete gauge invariance

Why is this relevant ?

Suppose we have a uniform field $\mathbf{B} = B\mathbf{e}_z$. Then $\mathbf{A} = -By\mathbf{e}_x$.

Discrete gauge invariance

Why is this relevant ?

Suppose we have a uniform field $\mathbf{B} = B\mathbf{e}_z$. Then $\mathbf{A} = -By\mathbf{e}_x$.

The gauge function $\chi(\mathbf{r}) = -By_0x$ would change

$$\mathbf{A} \rightarrow \mathbf{A}' = -B(y - y_0)\mathbf{e}_x$$

Discrete gauge invariance

Why is this relevant ?

Suppose we have a uniform field $\mathbf{B} = B\mathbf{e}_z$. Then $\mathbf{A} = -By\mathbf{e}_x$.

The gauge function $\chi(\mathbf{r}) = -By_0x$ would change

$$\mathbf{A} \rightarrow \mathbf{A}' = -B(y - y_0)\mathbf{e}_x$$

Use *some discretization* of the derivative operator: *The result would depend on the gauge origin y_0 .*

Discrete gauge invariance

Why is this relevant ?

Suppose we have a uniform field $\mathbf{B} = B\mathbf{e}_z$. Then $\mathbf{A} = -By\mathbf{e}_x$.

The gauge function $\chi(\mathbf{r}) = -By_0x$ would change

$$\mathbf{A} \rightarrow \mathbf{A}' = -B(y - y_0)\mathbf{e}_x$$

Use *some discretization* of the derivative operator: *The result would depend on the gauge origin y_0 .*

The simple reason:

$$\frac{\partial}{\partial x} e^{ie\chi(x)/\hbar} = \frac{ie}{\hbar} \frac{\partial \chi(x)}{\partial x} e^{ie\chi(x)/\hbar}$$

but

$$\frac{e^{ie\chi(x+h)/\hbar} - e^{ie\chi(x-h)/\hbar}}{2h} \neq \frac{ie}{\hbar} \left[\frac{\chi(x+h) - \chi(x-h)}{2h} \right] e^{ie\chi(x)/\hbar}$$

Is there a solution ?

What we find in the literature:

Indeed, within a finite linear variational subspace, gauge-origin invariance can never be obtained exactly, only approximately for small displacements of the gauge origin. In such cases, therefore, the calculated energies and properties will depend on the choice of gauge origin.

T. Helgaker, M. Jazuński, and K. Ruud,
Chem. Rev. **99**, 293 (1999).

The way out

Let's solve the problem anyway !

Schwinger's gauge transport function

Let

$$f_i(\mathbf{r}) = \int^{x_i} A_i(\mathbf{r}) dx_i$$

The way out

Let's solve the problem anyway !

Schwinger's gauge transport function

Let

$$f_i(\mathbf{r}) = \int^{x_i} A_i(\mathbf{r}) dx_i$$

Then

$$\left[-i\hbar \frac{\partial}{\partial x_i} + eA_i(\mathbf{r}) \right]^2 = e^{-ief_i(\mathbf{r})/\hbar} \frac{\partial^2}{\partial x_i^2} e^{-ief_i(\mathbf{r})/\hbar}$$

The way out

Let's solve the problem anyway !

Schwinger's gauge transport function

Let

$$f_i(\mathbf{r}) = \int^{x_i} A_i(\mathbf{r}) dx_i$$

Then

$$\left[-i\hbar \frac{\partial}{\partial x_i} + eA_i(\mathbf{r}) \right]^2 = e^{-ief_i(\mathbf{r})/\hbar} \frac{\partial^2}{\partial x_i^2} e^{-ief_i(\mathbf{r})/\hbar}$$

Then: A gauge transformation

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}'(\mathbf{r}) = \mathbf{A}(\mathbf{r}) + \nabla \chi(\mathbf{r})$$

changes the gauge transport functions

$$f_i(\mathbf{r}) \rightarrow f'_i(\mathbf{r}) = f_i(\mathbf{r}) + \chi(\mathbf{r})$$

The way out

Let's solve the problem anyway !

Schwinger's gauge transport function

Let

$$f_i(\mathbf{r}) = \int^{x_i} A_i(\mathbf{r}) dx_i$$

Then

$$\left[-i\hbar \frac{\partial}{\partial x_i} + eA_i(\mathbf{r}) \right]^2 = e^{-ief_i(\mathbf{r})/\hbar} \frac{\partial^2}{\partial x_i^2} e^{-ief_i(\mathbf{r})/\hbar}$$

Then: A gauge transformation

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}'(\mathbf{r}) = \mathbf{A}(\mathbf{r}) + \nabla \chi(\mathbf{r})$$

changes the gauge transport functions

$$f_i(\mathbf{r}) \rightarrow f'_i(\mathbf{r}) = f_i(\mathbf{r}) + \chi(\mathbf{r})$$

⇒ takes care of the phase factor in the wave function.

Homogeneous fields:

Let $\mathbf{A} = -By\mathbf{e}_x$, $f_x(\mathbf{r}) = -Bxy$.

Represent the wave function in momentum space:

$$\psi(\mathbf{x}) = \sum_k \psi_k e^{ikx}.$$

$$\begin{aligned} \left[-i\hbar \frac{\partial}{\partial x_i} + eA_i(\mathbf{r}) \right]^2 \psi(\mathbf{x}) &= e^{ieBxy/\hbar} \frac{\partial^2}{\partial x^2} e^{-ieBxy/\hbar} \sum_k \psi_k e^{ikx} \\ &= \sum_k \psi_k e^{ieBxy/\hbar} \frac{\partial^2}{\partial x^2} e^{i(k - eBy/\hbar)x} = \sum_k t_n(k - eBy/\hbar) \psi_k e^{ikx} \end{aligned}$$


with – for an n -point second derivative formula

$$t_n(k) = \left(\frac{2\sin\kappa}{h} \right)^2 \left[1 + \frac{1}{3}\sin^2\kappa + \frac{8}{45}\sin^4\kappa + \frac{4}{35}\sin^6\kappa + \dots \right].$$

Induced currents

Current density:

$$\mathbf{j}(\mathbf{r}) = \frac{e}{2m} \sum_i [\psi_i^* \boldsymbol{\Pi} \psi_i + \psi_i (\boldsymbol{\Pi} \psi_i)^*] = \frac{e}{m} \sum_i \Re \{ \psi_i^* \boldsymbol{\Pi} \psi_i \},$$



Induced current density in
a C₆H₆ molecule at 1 Tesla

Featured in Phys. Rev. B
Kaleidoscope

S. Janecek and E.
K., Phys. Rev.
B77, 245115 (2008)


Magnetic susceptibility

Induced magnetic dipole moment

$$\mathbf{m} = \frac{1}{2} \int \mathbf{r} \times \mathbf{j}(\mathbf{r}) d^3\mathbf{r}.$$

Susceptibility:

$$\begin{aligned}\chi_{ij} &= \frac{\partial M_i}{\partial H_j} = \frac{\partial B_j}{\partial H_j} \frac{\partial M_i}{\partial B_j} \\ &= \frac{N\mu_0}{V} \frac{\partial}{\partial B_i} m_j.\end{aligned}$$



NMR shifts: Measure local field strengths

Induced magnetic field:

$$\mathbf{B}^{\text{ind}}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} \times \mathbf{j}(\mathbf{r}')$$

NMR shifts: Measure local field strengths


Induced magnetic field:

$$\mathbf{B}^{\text{ind}}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} \times \mathbf{j}(\mathbf{r}')$$

Chemical shift tensor

$$\sigma(\mathbf{R}) = \frac{\partial \mathbf{B}^{\text{ind}}(\mathbf{R})}{\partial \mathbf{B}^{\text{ext}}}.$$

NMR shift for C atoms in C_6H_6



NMR shifts: Measure local field strengths

Induced magnetic field:

$$\mathbf{B}^{\text{ind}}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} \times \mathbf{j}(\mathbf{r}')$$


Chemical shift tensor

$$\sigma(\mathbf{R}) = \frac{\partial \mathbf{B}^{\text{ind}}(\mathbf{R})}{\partial \mathbf{B}^{\text{ext}}}.$$

No miracles !

Results are just as good as DFT
lets them be.

NMR shift for C atoms in C_6H_6



NMR shifts: Measure local field strengths

Induced magnetic field:

$$\mathbf{B}^{\text{ind}}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} \times \mathbf{j}(\mathbf{r}')$$


Chemical shift tensor

$$\sigma(\mathbf{R}) = \frac{\partial \mathbf{B}^{\text{ind}}(\mathbf{R})}{\partial \mathbf{B}^{\text{ext}}}.$$

No miracles !

Results are just as good as DFT lets them be.

NMR shift for H atoms in C_6H_6



NMR shifts: Measure local field strengths

Induced magnetic field:

$$\mathbf{B}^{\text{ind}}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} \times \mathbf{j}(\mathbf{r}')$$

Chemical shift tensor

$$\sigma(\mathbf{R}) = \frac{\partial \mathbf{B}^{\text{ind}}(\mathbf{R})}{\partial \mathbf{B}^{\text{ext}}}.$$


No miracles !

Results are just as good as DFT lets them be.

Gauge invariance

Results are stable against gauge transformations,

Sensitive against gauge transformations
(if one does it wrong !)



Motivation

First set of applications

Cluster physics is a research field of high interest

- Clusters represent the connection between molecules and solids.
- Preparation of clusters in their perfect ground-state is possible.

Motivation

First set of applications

Cluster physics is a research field of high interest

- Clusters represent the connection between molecules and solids.
- Preparation of clusters in their perfect ground-state is possible.

Clusters are created and embedded in

- ultracold, supersonic helium beams
- superfluid helium nanodroplets
- Question: Does the helium change cluster properties ?

Motivation

First set of applications

Cluster physics is a research field of high interest

- Clusters represent the connection between molecules and solids.
- Preparation of clusters in their perfect ground-state is possible.


Clusters are created and embedded in

- ultracold, supersonic helium beams
- superfluid helium nanodroplets
- Question: Does the helium change cluster properties ?

Model for cluster-helium interaction is required!

Physical Model

Helium atoms: quantum mechanics



Ions: fixed positions,
classical mechanics

Electrons: quantum mechanics

Magnesium-Helium Interaction Model


- electrostatic interactions: Helium atoms are considered as polarizable spheres
- Pauli repulsion: Phenomenological
- long-range part: Non-local dispersion corrections

The new interactions have to be added to the **self-consistency** cycle.


Effective Potential in Helium environment:

$$v_s[\rho](\mathbf{r}) = v(\mathbf{r}) + v_H[\rho](\mathbf{r}) + v_{xc}[\rho](\mathbf{r}) + v_C[\rho, \rho_{He}](\mathbf{r}) + v_P[\rho, \rho_{He}](\mathbf{r})$$


Calculating Cluster-Helium Potentials



calculate reference energy of
undisturbed cluster




calculate energy-response of cluster to
He-atom




move He-atom


Mg-He Potential




Mg₄-He Potential in z-Direction



Mg₁₀-He Potential in x-Direction



Mg₁₀-He Potential in z-Direction



Path Integral Monte Carlo - PIMC

- Quantum Monte Carlo method in the path integral formulation of quantum mechanics
- well suited for condensed systems like liquid Helium
- in every move the total potential energy is needed


Helium Potential caused by one Mg-cluster

$$v_{He}[\rho](\mathbf{r}_{He}) = v_C[\rho](\mathbf{r}_{He}) + v_P[\rho](\mathbf{r}_{He}) + v_{nl}(\mathbf{r}_{He})$$


Combination of DFT and PIMC

Compute $v_{He}[\rho]$


Combination of DFT and PIMC




Combination of DFT and PIMC



Combination of DFT and PIMC



Combination of DFT and PIMC



Mg_nHe_m : clusters: Iso-Surfaces of Helium Density

- A Mg_{07} cluster with 40 ^4He atoms $\text{Mg}_7\text{He}_{40}$
- A Mg_{07} cluster with 80 ^4He atoms $\text{Mg}_7\text{He}_{80}$
- A Mg_{11} cluster with 60 ^4He atoms $\text{Mg}_{11}\text{He}_{60}$

Summary: A solution looking for problems

Designing a DFT cluster code from scratch

- Forward factorization eigenvalue solver
 - Speed gain in eigenvalue solver by a factor of 10-100
 - Intellectual input from Monte Carlo methods
 - Trivial to parallelize

Summary: A solution looking for problems

Designing a DFT cluster code from scratch

- Forward factorization eigenvalue solver
 - Speed gain in eigenvalue solver by a factor of 10-100
 - Intellectual input from Monte Carlo methods
 - Trivial to parallelize
- Response iteration method for density update
 - Intellectual input from Jastrow-Feenberg theory

Summary: A solution looking for problems

Designing a DFT cluster code from scratch

- Forward factorization eigenvalue solver
 - Speed gain in eigenvalue solver by a factor of 10-100
 - Intellectual input from Monte Carlo methods
 - Trivial to parallelize
- Response iteration method for density update
 - Intellectual input from Jastrow-Feenberg theory
- Exact calculation of the exponential of the canonical kinetic energy operator without computational overhead
 - Intellectual input from statistical physics

Summary: A solution looking for problems

Designing a DFT cluster code from scratch

- Forward factorization eigenvalue solver
 - Speed gain in eigenvalue solver by a factor of 10-100
 - Intellectual input from Monte Carlo methods
 - Trivial to parallelize
- Response iteration method for density update
 - Intellectual input from Jastrow-Feenberg theory
- Exact calculation of the exponential of the canonical kinetic energy operator without computational overhead
 - Intellectual input from statistical physics
- Gauge invariant in any discretization
 - Intellectual input from quantum field theory
- Combination of QMC and DFT

Summary: A solution looking for problems

Designing a DFT cluster code from scratch

- Forward factorization eigenvalue solver
 - Speed gain in eigenvalue solver by a factor of 10-100
 - Intellectual input from Monte Carlo methods
 - Trivial to parallelize
- Response iteration method for density update
 - Intellectual input from Jastrow-Feenberg theory
- Exact calculation of the exponential of the canonical kinetic energy operator without computational overhead
 - Intellectual input from statistical physics
- Gauge invariant in any discretization
 - Intellectual input from quantum field theory
- Combination of QMC and DFT

Results for magnetic susceptibilities and NMR shifts:
Just as good as the pseudopotentials and density functionals.