

25.

(a) **Solution:**

Let  $D_{\text{right}}$  denote distance travelled by the car as estimated by the right sum. Then the right sum, in terms of  $t$  and  $v(t)$ , can be calculated using

$$\begin{aligned} D_{\text{right}} &= RIGHT(5) \\ &= \Delta t \sum_{i=1}^5 v(t_i) \end{aligned} \tag{1}$$

From the table, it can be observed that the width of each subinterval is 0.2 h. Otherwise, here's the calculation for completeness:

$$\begin{aligned} \Delta t &= \frac{b-a}{n} \\ &= \frac{1-0}{5} \\ &= 0.2 \text{ h} \end{aligned}$$

Then let's use (1) to calculate the estimated distance:

$$\begin{aligned} D_{\text{right}} &= \Delta t \sum_{i=1}^5 v(t_i) \\ &= 0.2(30 + 30 + 70 + 90 + 90) \\ &= 62 \text{ km} \end{aligned}$$

(b) **Solution:**

Let  $G_{\text{right}}$  denote the amount of gas consumed by the car as estimated by and let  $g(v(t))$  be the fuel gas efficiency of the car given a speed  $v(t)$ .

Unit analysis: We want L from the distances (speed  $\times$  time) in part (a) and the efficiency from the table.

$$\underbrace{\frac{\cancel{\text{h}}}{t}}_{v(t)} \times \underbrace{\frac{\text{km}}{\cancel{\text{h}}}}_{g(v(t))} \times \frac{\text{L}}{100 \cancel{\text{km}}} = \frac{1}{100} \text{L}$$

So then the terms we're summing over are  $f(t_i) = v(t_i)g(v(t_i))$ , which implies that the quantity that we're looking for is

$$\begin{aligned} G_{\text{right}} &= RIGHT(5) \\ &= \Delta t \sum_{i=1}^5 f(t_i) \\ &= \Delta t \sum_{i=1}^5 v(t_i)g(v(t_i)). \end{aligned} \tag{2}$$

With  $\Delta t$  as before, using (2) to calculate the estimated gas consumed yields

$$\begin{aligned}
 G_{\text{right}} &= \Delta t \sum_{i=1}^5 v(t_i)g(v(t_i)) \\
 &= 0.2 \left( 30 \cdot \frac{15}{100} + 30 \cdot \frac{15}{100} + 70 \cdot \frac{7}{100} + 90 \cdot \frac{8}{100} + 90 \cdot \frac{8}{100} \right) \\
 &= 5.66 \text{ L}
 \end{aligned}$$

26.

(a) **Solution:**

The length of each of the four subintervals is

$$\begin{aligned}
 \Delta x &= \frac{b-a}{n} \\
 &= \frac{\frac{\pi}{2} - 0}{4} \\
 &= \frac{\pi}{8}
 \end{aligned}$$

With  $f(x) = \cos(x)$  and  $n = 4$ , the trapezoidal rule gives us

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} f(x) \, dx &\approx TRAP(4) \\
 &= \Delta x \left( \frac{f(x_0) + f(x_4)}{2} + \sum_{i=1}^3 f(x_i) \right) \\
 &= \frac{\pi}{8} \left( \frac{\cos(0) + \cos(\frac{\pi}{2})}{2} + \cos\left(\frac{\pi}{8}\right) + \cos\left(\frac{\pi}{4}\right) + \cos\left(\frac{3\pi}{8}\right) \right) \\
 &\approx 0.987
 \end{aligned}$$

(b) **Solution:**




Figure 1: A sketch graph of the trapezoidal rule for approximating the integral of  $\cos(x)$  from 0 to  $\frac{\pi}{2}$ .

From Figure 1, we can deduce that the estimate from part (a) is an **underestimate** of the exact integral value.