

2026 FAMILY CARE REPORT

When it's no longer safe for a parent to *live alone.*

The early warning signs, risk factors, and decision thresholds most families miss — until a fall, a hospital visit, or an emergency forces action.

Based on 2025 caregiving research and the latest national data on aging, home safety, and care transitions.





01 A note before you begin

A short letter *before you read further.*

If you're reading this, it likely means you've started to notice changes in someone you care about — and you're trying to figure out what to do next.

We know this isn't something people casually search for. That's not always easy to sit with.

One of the biggest challenges families face is that decline doesn't happen all at once. It's gradual. Subtle. Easy to explain away.

Most people don't realize how serious things have become until something forces the issue — a fall, a hospital visit, or a moment that felt too close for comfort. We put this report together to help with that exact gap.

Not to alarm you. Not to push you into a decision. But to give you a clearer way to look at what's happening, understand what to watch for, and feel more confident about what to do next.

Because the goal isn't to overreact. It's to avoid being caught off guard.

As you read, you'll start to see patterns — things that feel familiar, things you may have already noticed, and things you may not have connected yet. That clarity is what matters.

And if nothing else, we hope this helps you feel a little more prepared moving forward.

With care,

M&L IN-HOME CARE

02 *The reason this decision feels so heavy*

Why this decision *is* so hard.

The reality is, the majority of these situations don't feel urgent — until they suddenly are.

1 in 4

Adults over 65 falls each year — and most families don't realize the risk until after it happens.



Decline rarely happens all at once. It shows up in small ways:

- A missed step
- A forgotten detail
- A little more effort to do everyday things

Each one feels minor on its own, so it's easy to move past it and carry on.

It's also easy to normalize what's happening.

So you find yourself saying things like:

- "This is just part of getting older."
- "They've always been a little forgetful."
- "It's probably nothing serious."

And sometimes, that's true. But sometimes, it isn't.

At the same time, your parent can still seem completely fine in other moments. They're often able to hold a conversation, recognize people, and insist they don't need help. That contrast makes it harder to trust your instincts about what's actually changing.



You're also not just dealing with one thing.

There's family, distance, daily responsibilities — and a parent who may not want to talk about it. So even if something feels off, it's easy to push the decision down the list and come back to it later.

And when you do think about taking action, it can feel bigger than it actually is. Getting help might feel like:

- Taking away independence
- Making something "official"
- Changing the family dynamic

Which makes it easier to wait than risk acting too soon.

In the end, most families end up in the same place — somewhere in between. You might be thinking:

"I feel like something's changing. But I'm not sure how serious it is yet. Maybe we just wait and see..."

A COMMON STARTING POINT FOR MOST FAMILIES

This is a very normal place to be. The challenge is that this stage of uncertainty is where most delays happen — not because families don't care, but because they don't have a clear way to evaluate what's actually going on.

And when things aren't clear, waiting feels like the safest option. But in most cases, what actually helps is clarity, and a decision at the right time.



03 From vague worry to identifiable signals

What families *usually* notice *first*.

Most serious issues don't start with a crisis. They start with small, noticeable changes.

According to the National Institute on Aging, early signs of declining independence often show up in everyday behaviors long before a major incident happens. These aren't diagnoses — they're signals. And they're often

the first indication that more support may be needed.

Common early signs families notice

01 Confusion or forgetfulness

Repeating the same questions, forgetting appointments, getting confused about time or place. Early cognitive changes are one of the most common precursors to loss of independence — and they're often dismissed at first.

02 Changes in mobility or balance

Slower walking, hesitation when standing or sitting, needing support to move around safely. According to the CDC, 1 in 4 adults over 65 falls each year, and many of those falls are linked to subtle mobility changes that were already present.

03 Decline in personal hygiene

Wearing the same clothes repeatedly, skipping bathing routines, a general drop in self-care. This is often one of the earlier signs that daily tasks are becoming harder to manage independently.

04 Weight changes or poor nutrition

Noticeable weight loss, uneaten food, difficulty preparing meals. Research shows that poor nutrition in older adults is closely tied to increased health risks, including weakness, falls, and hospitalization.

05 Increased isolation

Fewer social interactions, avoiding activities they used to enjoy, staying home more often. Isolation is more than emotional — it's been linked to faster cognitive and physical decline.

Why these signs matter

Individually, each of these can seem small. Easy to explain. Easy to overlook. But when they start to show up together — or become more frequent — they often point to a bigger shift.

What most families miss

It's not one major event that signals a change. It's a pattern. And that pattern usually appears weeks or months before something serious happens.

If you've started noticing one or more of these signs, you're not overreacting. You're catching something early — and that's exactly when you have the most options.

04 The risks that don't look like risks

The hidden risks families underestimate.

Most of the real risks at home don't look like risks at all. In many cases, the biggest dangers are the result of small changes that quietly increase risk over time.

**01 Fall risk increases gradually**

Falls are rarely random. They're usually the result of changes that build over time — reduced balance, slower reaction time, weaker muscles, subtle mobility issues. According to the CDC, falls are the leading cause of injury-related deaths among adults 65+, and most of those falls are linked to factors that were already present beforehand.

02 Medication becomes harder to manage

Many older adults take multiple medications each day. Over time, this can lead to missed doses, incorrect timing, or accidental double-dosing. Even small medication errors can lead to serious complications — dizziness, confusion, or hospitalization.

03 The home environment becomes less safe

A home that once felt completely safe can slowly become harder to navigate. Common risks: loose rugs or clutter, poor lighting, stairs or uneven surfaces, bathrooms without proper support. None of these are major on their own — but combined with mobility changes, they increase the likelihood of a fall.

04 Vision and balance changes go unnoticed

Vision and balance issues often happen gradually — misjudging distances, difficulty navigating familiar spaces, increased risk of tripping or falling. Because these changes happen slowly, they're often overlooked until something happens.

05 The post-hospital window is high risk

According to the Agency for Healthcare Research and Quality (AHRQ), the transition from hospital to home is one of the highest-risk periods for older adults. Without proper support, the likelihood of complications or readmission increases. Many families assume things are "back to normal" after discharge — but this is often when additional support is needed most.

Why these risks matter together

Individually, each of these risks may seem manageable. But when they start to overlap, the likelihood of something serious increases. It's not just one factor — it's how they combine.

The key insight

Most serious incidents aren't caused by one major issue. They're the result of multiple smaller risks that were already there — just not fully recognized.

If you've noticed even a few of these changes, it doesn't mean something bad is about to happen. But it does mean it's worth taking a closer look. Because the earlier you understand the risk, the more control you have over what happens next.



05 *Why doing nothing isn't neutral*

Why "waiting and watching" can backfire.

Most families don't choose to wait. They wait because things don't feel urgent yet. And on the surface, that makes sense.

The challenge is that risk doesn't stay the same over time. It usually increases quietly.

What the data shows

A single fall doesn't happen in isolation.

- After one fall, the risk of falling again significantly increases.
- Falls are one of the leading causes of injury and hospitalization in adults over 65.
- Many of these incidents are linked to changes that were already present beforehand.

Cognitive decline follows a similar pattern.

- Subtle memory issues often appear long before a diagnosis — typically, a serious fall or hospitalization is what brings those issues to light.
- Studies have shown that a noticeable percentage of older adults receive a dementia diagnosis within a year after a major fall.

There's a critical window after hospital visits.

- The transition from hospital to home is one of the highest-risk periods.
- Without proper support, the likelihood of complications or readmission increases.
- Many families assume things are "back to normal" after discharge — but that's not always the case.

Why this matters

None of these situations start as emergencies. They build over time. And when action is delayed, decisions often have to be made quickly, under pressure.

What most families don't expect

Waiting doesn't just delay the decision. It often reduces your options.

- Fewer choices
- Less time to plan
- More stress when something does happen

Taking action early doesn't mean rushing into care. It means understanding what's happening before things become urgent. So if something changes — you're not reacting. You're prepared.

06 *A simple way to see where you are*

A practical family *risk framework*.

"Okay... I see some of the signs. But how serious is this, really?"

Not every situation requires immediate action. But clarity makes the decision-making process easier.

Most situations fall into one of three categories:

- **Green** — things are stable
- **Yellow** — early warning signs are showing
- **Red** — risk is elevated and action is needed

This isn't a diagnosis. It's a practical way to understand where things stand right now.

Green — Mostly Independent

At this stage, things are generally stable. There may be small changes, but they're not impacting safety in a meaningful way.

YOU MIGHT NOTICE

- Minor forgetfulness that doesn't affect daily life
- Independent with meals, hygiene, and movement
- No recent falls or hospital visits
- Able to manage household tasks and finances independently

WHAT TO DO

- Keep an eye on changes over time
- Start light conversations about future support
- Make small home-safety improvements if needed

Yellow — Early Warning Signs

This is where most families start to feel unsure. Things aren't urgent, but they're no longer fully stable.

YOU MIGHT NOTICE

- Missed medications or confusion around routines
- A minor fall or near fall
- Increased forgetfulness or repeated questions
- Decline in hygiene or daily habits
- Eating less or struggling with meals
- Pulling back from social activity

WHAT TO DO

- Start actively evaluating support options
- Increase check-ins or oversight
- Consider part-time help or a professional assessment

Red — Elevated Risk

At this stage, safety is becoming a real concern. Waiting often leads to decisions being made under pressure.

YOU MIGHT NOTICE

- Recent fall or hospital visit
- Difficulty walking or moving safely
- Confusion that affects daily decisions
- Inability to manage medications properly
- Struggling with basic daily tasks (bathing, dressing, eating)
- Wandering or disorientation

WHAT TO DO

- Take action to put support in place
- Involve professionals to assess needs
- Focus on safety and stability first

How to use this

You don't need to overanalyze it. Just ask yourself: *"Which category feels closest to what I'm seeing right now?"* Even if it's not a perfect fit, you'll usually land somewhere near one of these.

This isn't about labeling the situation. It's about understanding it clearly enough to make a good decision. Once things are clear, the next step

07 *A simple family planning tool*

Five decisions families should make before a crisis.

When something does change — and it usually does — families who've already thought through these questions make better, faster decisions with far less stress.



01 Who manages medication?

Is there a system, or is it memory-based? Have there been missed doses or confusion?

02 Who checks in — and how often?

Who is actually seeing your parent regularly? Would they notice if something changed?

03 What changes at home have already happened?

Have routines shifted? Are certain tasks being avoided? Has the home itself been adjusted?


04 What happens after the next fall or ER visit?

Who steps in immediately? Is there a plan — or would decisions be made in the moment?

05 What level of support would be acceptable if risk increases?

Would part-time help feel okay? Would full-time care ever be considered? What would your parent realistically accept?

You're not deciding now — you're defining boundaries. Even partial answers to these questions can make a big difference when something changes.



08 It's smaller than you think

What taking action early actually looks like.

You've likely started to recognize some of the signs already. The next step isn't to rush into a big decision. It's to get clear on a few smaller ones first.

01 Who is responsible for keeping an eye on things day-to-day?

In many families, this is unclear. Everyone assumes someone is paying attention — but no one is fully accountable.

Food for thought: Who checks in regularly? How often are they actually seeing your parent? Would they notice if something changed?

Why this matters — small changes are easy to miss without consistent oversight.

02 How are medications being managed right now?

Medication issues are one of the most common causes of complications — and they're often invisible until something goes wrong.

Consider: Who ensures medications are taken correctly? Is there a system in place, or is it memory-based? Have there been any missed doses or confusion?

Why this matters — even small errors can lead to serious health issues.

03 What changes have already happened at home?

Most families underestimate how much has already shifted. Take a step back and look at what's different.

Observe: Have routines changed? Are certain tasks being avoided? Has the home been adjusted in any way?

Why this matters — these changes are often early signals of increasing risk.

04 What happens if something does go wrong?

This is the question most families avoid. But it's one of the most important.

Ask yourself: What happens after a fall or ER visit? Who steps in immediately? Is there a plan — or would decisions be made in the moment?

Why this matters — decisions made under pressure are rarely the best ones.

05 What level of support would feel acceptable if things got worse?

This helps remove uncertainty later. You're not deciding now — you're just defining boundaries.

Consider: Would part-time help feel okay? Would full-time care ever be considered? What would your parent realistically accept?

Why this matters — when expectations are unclear, decisions become harder.

You don't need perfect answers.

This isn't about getting everything figured out today. It's about starting the conversation — either with yourself or with your family. Even having partial answers to these questions can make a big difference when something changes.

| *Clarity now creates a lot more control later.*

09 *A clearer picture, a calmer next step*

Where to go *from here*.

By now, you've probably seen one of two things — either things are still mostly stable and you just need to stay aware, or there are signs that it may be time to look more closely at what kind of support could help.

Both are useful to know. Because the goal of this report was never to push you into a decision. It was to help you reduce uncertainty.

That's what most families need first. Not pressure. Not urgency. Just a clearer picture of what's happening and what to do next.

The hardest situations usually aren't the ones where families made the wrong choice. They're the ones where families had to make a choice *too quickly*, with too little clarity, after something had already happened.

That's why earlier awareness matters.

It gives you more time to:

- Notice patterns before they become bigger problems
- Talk through options without feeling rushed
- Make a thoughtful decision based on what's actually needed

And if after reading this, you're still unsure — that's okay too. Uncertainty doesn't mean you're behind. It means you need a clearer outside perspective. That's exactly where a simple conversation can help.

FREE CONSULTATION • NO OBLIGATION

Book your free *Home Care Readiness Session*.

In a short conversation, we'll help you review the current situation, identify the most likely risks, and give you a clear next-step recommendation. No pressure. No obligation. Just clarity.

[Schedule your free session →](#)

MLINHOMECARE.COM • (000) 000-0000

