

## 1 T-duality of Heterotic Strings<sup>1</sup>

We use  $d \leq 10$  to denote the number of noncompact dimensions; the remaining  $m \geq d$  dimensions are compactified. For heterotic strings, the  $I \geq 10$  dimensions of the left-moving sector are already compactified on a lattice  $\Gamma_{16}$  or  $\Gamma_8 \times \Gamma_8$ . Here we use the label  $I$  to index the 16 internal dimensions.

(a) Generally, if we compactify an open string on the  $x^m$  direction:  $x^m \cong x^m + 2\pi R$  with constant backgrounds  $A_m$ , then its zero mode spectrum, *with winding*  $w = 0$ , can be obtained from canonical quantization of the effective point particle action, with an additional gauge action term in the form of a Wilson line<sup>2</sup>:

$$-W_q = -iq \int dx^m A_m \sim -iq \int d\tau A_m \dot{X}^m \quad (1)$$

By imposing that the canonical momentum to be periodic along  $x^m$ , we find that:

$$k_m = \frac{n_m}{R} - qA_m \quad (2)$$

To obtain the winding states, we have to reproduce the above action from the world-sheet description. For heterotic strings with  $m < 10 \leq I \leq 26$ , this can be achieved by adding the following term to the usual world-sheet action<sup>3</sup>:

$$S_A \propto \int d^2\sigma \epsilon^{ab} A_\mu^I \partial_a X^\mu \partial_b X_I \quad (3)$$

With proper normalizations to match the result in (2).

Canonical quantization then produces<sup>4</sup>:

$$k_m = \frac{n_m}{R} \pm \frac{w_m R}{\alpha'} - q_I A_m^I - \frac{w_n R}{2} A_I^n A_m^I, \quad (4)$$

$$k_L^I = \sqrt{\frac{2}{\alpha'}} (q^I + w^m R A_m^I), \quad (5)$$

The “ $\pm$ ” signs in  $k_m$  correspond to the left and right-moving sector, respectively. Only the left-moving sector has an additional 16 dimensional internal torus, therefore  $k^I$  is labeled with an “ $L$ ”.

Note that the charge  $q^I$  now takes value on the  $\Gamma_{16}$  or  $\Gamma_8 \times \Gamma_8$  lattice, and:

$$l \circ l' = \frac{\alpha'}{2} (k_L^I k'_{L,I} + k_L^m k'_{L,m} - k_R^m k'_{R,m}) = q^I q'_I + 2nw \quad (6)$$

We can then see that the new “extended” lattice indeed satisfies the even and self-dual conditions, which follows from the even and self-dual properties of  $\Gamma_{16}$  or  $\Gamma_8 \times \Gamma_8$ .

<sup>1</sup>I would like to thank Lucy Smith for help with this problem.

<sup>2</sup>Reference: *Polchinski*, Chapter 8.

<sup>3</sup>Reference: *Blumenhagen et al*, Basic Concepts of String Theory.

<sup>4</sup>Reference: *Polchinski*, Chapter 11.

(b) With  $m = 9$  and  $G_{dd} = 1$ , we have:

$$W_q = \exp(-iq_I \theta^I), \quad A_9^I = -\frac{\theta^I}{2\pi R} \quad (7)$$

Note that  $W_q$  captures the phase change of the paths that wind around  $x^9$ ; the extra phase from a non-trivial Wilson line might affect the boundary condition of some states while leaving others intact, thus breaking the original gauge symmetry. Our discussions here closely follow *Polchinski*, Chapter 11.

For the  $\text{SO}(32)$  theory with:

$$RA_9^I = \text{diag}\left(\left(\frac{1}{2}\right)^8, 0^8\right) \quad (8)$$

Adjoint states are labeled by a pair of indices valued in  $1, \dots, 32$ ; those with one index from  $1 \leq A \leq 16$  and one from  $17 \leq A \leq 32$  are antiperiodic due to the additional phase  $e^{i\pi} = -1$  from the Wilson line, so the gauge symmetry is reduced to  $\text{SO}(16) \times \text{SO}(16)$ .

Similarly, for the  $E_8 \times E_8$  theory with:

$$R'A_9^I = \text{diag}(1, 0^7, 1, 0^7) \quad (9)$$

Note that  $\Gamma_8$ , the root lattice of  $E_8$ , is basically the root lattice union an additional spinor weight lattice of  $\text{SO}(16)$ . With the above Wilson line, the integer-charged states from the  $\text{SO}(16)$  root lattice in each  $E_8$  remain periodic, while the half-integer charged states from the  $\text{SO}(16)$  spinor lattices become antiperiodic, due to the additional phase  $e^{i\frac{1}{2} \cdot 2\pi} = -1$ . Again the gauge symmetry is broken down to  $\text{SO}(16) \times \text{SO}(16)$ .

In summary, with the above Wilson line, the  $\text{SO}(32)$  and  $E_8 \times E_8$  theory shares a unbroken gauge of  $\text{SO}(16) \times \text{SO}(16)$ . Consider the spectrum of the  $\text{SO}(16) \times \text{SO}(16)$  *neutral states*, i.e. those with internal momentum:

$$k_L^I = \sqrt{\frac{2}{\alpha'}} (q^I + wRA_9^I) = 0 \quad (10)$$

For the  $\text{SO}(32)$  theory, since  $q^I \in \Gamma_{16}$  while  $RA_9^I = \text{diag}\left(\left(\frac{1}{2}\right)^8, 0^8\right)$ , we must have  $w = 2m$  for this to hold. The same goes for the  $E_8 \times E_8$  theory; therefore, we have:

$$k_{L,R} = \frac{\tilde{n}}{R} \pm \frac{2mR}{\alpha'}, \quad k'_{L,R} = \frac{\tilde{n}'}{R'} \pm \frac{2m'R'}{\alpha'}, \quad (11)$$

$$\tilde{n} = n + 2m, \quad \tilde{n}' = n' + 2m' \quad (12)$$

(c) If the two theories are related by T-duality, then we should expect:

$$(k_L, k_R) \longleftrightarrow (k'_L, -k'_R), \quad (13)$$

Under suitable mapping of parameters. Indeed, it is straightforward to verify that  $(\tilde{n}, m) \leftrightarrow (m', \tilde{n}')$  realizes this, along with  $RR' = \alpha'/2$ . The above arguments can then be generalized to higher levels, by acting on fermionized left-moving fields  $\lambda^A$  and carefully organizing representations. We see that the two heterotic string theories are equivalent under T-duality.

## 2 String Junction<sup>5</sup>

For a string junction to be mechanically stable, the tension force exerted on the junction must cancel each other; this is a Newtonian mechanics problem, but with  $(p, q)$ -string tension given by the BPS bound:

$$\tau_{(p,q)} = \frac{\sqrt{p^2 + q^2/g^2}}{2\pi\alpha'} \quad (14)$$

Stability of the system implies that the BPS bound should be saturated.

From Newtonian mechanics, we know that three forces cannot cancel each other unless they are co-planar. Therefore, a 3-string junction must be co-planar in order to be stable. Suppose they lie in the  $(X^1, X^2)$  plane, then the tension exerted on the junction can be expressed as:

$$\vec{T}_i = \tau_{(p_i, q_i)} (\cos \theta_i, \sin \theta_i), \quad i = 1, 2, 3, \quad (15)$$

$\sum_i \vec{T}_i = 0$  gives two equations, and we have two independent unknowns (the angle between two pairs of strings); therefore if a solution exists, it should be unique up to rotations and reflections.

In fact, a solution can be found by simple observations:

$$\cos \theta_i = \frac{p_i}{\sqrt{p^2 + q^2/g^2}}, \quad \sin \theta_i = \frac{q_i/g}{\sqrt{p^2 + q^2/g^2}}, \quad (16)$$

It satisfies  $\sum_i \vec{T}_i = 0$  since that total  $(p, q)$  vanishes at each junction.

To find the remaining supersymmetries of this system, we start from the original supersymmetries of a  $(p, q)$  string (which saturates the BPS bound) extended along the  $\hat{X} = (\cos \theta, \sin \theta)$  direction:

$$\frac{1}{2L\tau_{(p,q)}} \left\{ \begin{bmatrix} Q_\alpha \\ \tilde{Q}_\alpha \end{bmatrix}, \begin{bmatrix} Q_\beta^\dagger & \tilde{Q}_\beta^\dagger \end{bmatrix} \right\} = \delta_{\alpha\beta} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + (\Gamma^0 \Gamma^\theta)_{\alpha\beta} \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix}, \quad (17)$$

$$\Gamma^\theta = \hat{X} \cdot \vec{\Gamma} = \Gamma^1 \cos \theta + \Gamma^2 \sin \theta \quad (18)$$

We see that the algebra depends on  $\theta$ , i.e. it is different for strings in different directions. However, if we can find a (maximal) subalgebra that is independent of  $\theta$ , then we would have found the remaining supersymmetries of the full system<sup>6</sup>.

We begin with diagonalizing the matrix on the RHS with:

$$U(\frac{\theta}{2}) = \begin{bmatrix} \cos \frac{\theta}{2} & -\sin \frac{\theta}{2} \\ \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{bmatrix}, \quad (19)$$

$$\frac{1}{2L\tau_{(p,q)}} \left\{ U^T \begin{bmatrix} Q_\alpha \\ \tilde{Q}_\alpha \end{bmatrix}, \begin{bmatrix} Q_\beta^\dagger & \tilde{Q}_\beta^\dagger \end{bmatrix} U \right\} = \begin{bmatrix} (\mathbb{1} + \Gamma^0 \Gamma^\theta)_{\alpha\beta} & 0 \\ 0 & (\mathbb{1} - \Gamma^0 \Gamma^\theta)_{\alpha\beta} \end{bmatrix}, \quad (20)$$

Note that  $(\mathbb{1} + \Gamma^0 \Gamma^\theta)(\mathbb{1} - \Gamma^0 \Gamma^\theta) = 0$  and  $(\mathbb{1} + \Gamma^0 \Gamma^\theta) + (\mathbb{1} - \Gamma^0 \Gamma^\theta) = 2\mathbb{1}$ , i.e. they are orthogonal to each other; acting  $(\mathbb{1} \pm \Gamma^0 \Gamma^\theta)$  on both sides, we find the following combinations, which gives the 16 SUSYs of a  $(p, q)$  string:

$$(\mathbb{1} - \Gamma^0 \Gamma^\theta) \left( \cos \frac{\theta}{2} Q + \sin \frac{\theta}{2} \tilde{Q} \right)_\alpha = 0 = (\mathbb{1} + \Gamma^0 \Gamma^\theta) \left( -\sin \frac{\theta}{2} Q + \cos \frac{\theta}{2} \tilde{Q} \right)_\beta \quad (21)$$

<sup>5</sup>Reference: [arXiv:0812.4408](https://arxiv.org/abs/0812.4408).

<sup>6</sup>The  $\tau_{(p,q)}$  factor can be absorbed by rescaling generators, hence does not matter in our discussions.

For further simplification, we can isolate the  $\theta$  dependence in  $\Gamma^\theta$  by working in a specific representation of the Clifford algebra, e.g. the Dirac representation given by *Polchinski*. Then  $\alpha$  is given by 10D spinor components:  $\alpha = (s_0, s_1, s_2, s_3, s_4)$ ,  $s_i = \pm$ , with additional chirality constraints from both  $Q$  and  $\tilde{Q}$ :  $\prod_i s_i = +$ . In the end, we have 16 independent components as expected.

Details of the expansion is given in [arXiv:0812.4408](#). When the dust settles, we find that the 16 SUSYs in (21) is given by:

$$\sin \frac{\theta}{2} \left( \cos \frac{\theta}{2} Q + \sin \frac{\theta}{2} \tilde{Q} \right)_{(++s)} + \cos \frac{\theta}{2} \left( \cos \frac{\theta}{2} Q + \sin \frac{\theta}{2} \tilde{Q} \right)_{(--s)}, \quad (22a)$$

$$\sin \frac{\theta}{2} \left( \cos \frac{\theta}{2} Q + \sin \frac{\theta}{2} \tilde{Q} \right)_{(+-s)} - \cos \frac{\theta}{2} \left( \cos \frac{\theta}{2} Q + \sin \frac{\theta}{2} \tilde{Q} \right)_{(-+s)}, \quad (22b)$$

$$\cos \frac{\theta}{2} \left( -\sin \frac{\theta}{2} Q + \cos \frac{\theta}{2} \tilde{Q} \right)_{(++s)} - \sin \frac{\theta}{2} \left( -\sin \frac{\theta}{2} Q + \cos \frac{\theta}{2} \tilde{Q} \right)_{(--s)}, \quad (22c)$$

$$\cos \frac{\theta}{2} \left( -\sin \frac{\theta}{2} Q + \cos \frac{\theta}{2} \tilde{Q} \right)_{(+-s)} + \sin \frac{\theta}{2} \left( -\sin \frac{\theta}{2} Q + \cos \frac{\theta}{2} \tilde{Q} \right)_{(-+s)}, \quad (22d)$$

$$s = (s_2 s_3 s_4), \quad \prod_i s_i = + \quad (23)$$

By trial and error, we can find the 8 linear combinations that are independent of  $\theta$ ; they are:

$$(a) + (c) \implies \tilde{Q}_{(++s)} + Q_{(--s)}, \quad (24)$$

$$(b) + (d) \implies \tilde{Q}_{(+-s)} - Q_{(-+s)}, \quad (25)$$

Therefore, the string junction is  $\frac{8}{32} = \frac{1}{4}$  BPS.

### **3 Two and Three-Point Functions in AdS/CFT**