

**1 Stringy Physics!**

$$T(z) = -\frac{1}{\alpha'} : \partial X^\mu \partial X_\mu : , \quad \tilde{T}(\bar{z}) = -\frac{1}{\alpha'} : \bar{\partial} X^\mu \bar{\partial} X_\mu : , \quad (1)$$

$$V_k = : e^{ik \cdot X(z, \bar{z})} : , \quad G_{e,k} = e_{\mu\nu} : \partial X_z^\mu \bar{\partial} X_{\bar{z}}^\nu e^{ik \cdot X(z, \bar{z})} : , \quad (2)$$

We sometimes use subscripts like  $\partial X_z^\mu$  to denote variable dependence to avoid clutter.

(a) The weight of a primary operator is given by its OPE with  $T$  and  $\tilde{T}$ . For exponential operators, there is a neat formula for cross contractions<sup>1</sup>:

$$\begin{aligned} T(z) V_k(w, \bar{w}) &= \exp \left\{ \int d^2 z' \int d^2 w' \overline{X_{z'}^\mu X_{w'}^\nu} \frac{\delta}{\delta X_{z'}^\mu} \frac{\delta}{\delta X_{w'}^\nu} \right\} : T_z e^{ik \cdot X_w} : \\ &= \exp \left\{ \int d^2 z' \overline{X_{z'}^\mu X_w^\nu} \frac{\delta}{\delta X_{z'}^\mu} ik_\nu \right\} : T_z e^{ik \cdot X_w} : \\ &= : \left\{ \exp \left( ik_\nu \int d^2 z' \overline{X_{z'}^\mu X_w^\nu} \frac{\delta}{\delta X_{z'}^\mu} \right) T_z \right\} e^{ik \cdot X_w} : \\ &\sim -\frac{1}{\alpha'} : \left\{ 2\partial_z (ik_\sigma \overline{X_z^\mu X_w^\sigma}) \partial_z X_\mu + \partial_z (ik_\rho \overline{X_z^\mu X_w^\rho}) \partial_z (ik_\sigma \overline{X_{z,\mu}^\sigma X_w^\sigma}) \right\} e^{ik \cdot X_w} : \\ &\sim -\frac{1}{\alpha'} : \left\{ 2 \left( -\frac{\alpha'}{2} \frac{ik^\mu}{z-w} \right) \partial_z X_\mu + \left( -\frac{\alpha'}{2} \frac{ik^\mu}{z-w} \right) \left( -\frac{\alpha'}{2} \frac{ik_\mu}{z-w} \right) \right\} e^{ik \cdot X_w} : \\ &\sim \frac{\alpha' k^2}{4} \frac{V_k(w, \bar{w})}{(z-w)^2} + \frac{\partial V_k(w, \bar{w})}{z-w} \end{aligned} \quad (3)$$

Here we've used the result that  $ik_\sigma \overline{X_z^\mu X_w^\sigma} = ik^\mu (-\frac{\alpha'}{2}) \ln |z-w|^2$ . We see that  $V_k$  is a primary of weight  $(1, 1)$  iff.  $\frac{\alpha' k^2}{4} = 1$ , or  $m^2 = -k^2 = -\frac{4}{\alpha'}$ . This is the mass shell condition for the closed string tachyon (at level 0). On the other hand,

$$G_{e,k} = e_{\mu\nu} G_k^{\mu\nu}, \quad (4)$$

$$\begin{aligned} T(z) G_k^{\mu\nu}(0) &\sim : T_z \partial X_0^\mu \bar{\partial} X_0^\nu e^{ik \cdot X_0} : + : \cancel{T_z \partial X_0^\mu \bar{\partial} X_0^\nu e^{ik \cdot X_0}} : + : T_z \partial X_0^\mu \bar{\partial} X_0^\nu e^{ik \cdot X_0} : \\ &\quad + : T_z \partial X_0^\mu \bar{\partial} X_0^\nu e^{ik \cdot X_0} : + : T_z \partial X_0^\mu \bar{\partial} X_0^\nu e^{ik \cdot X_0} : \\ &\sim \left( \frac{1}{z^2} G_k^{\mu\nu}(0) + \frac{1}{z} : \partial^2 X_0^\mu \bar{\partial} X_0^\nu e^{ik \cdot X_0} : \right) + \left( \frac{\alpha' k^2}{4} \frac{1}{z^2} G_k^{\mu\nu}(0) + \frac{1}{z} : \partial X_0^\mu \bar{\partial} X_0^\nu \partial e^{ik \cdot X_0} : \right) \\ &\quad - \frac{2}{\alpha'} \left( -\frac{\alpha'}{2} \eta^{\sigma\mu} \frac{1}{z^2} \right) \left( -\frac{\alpha'}{2} \frac{ik_\sigma}{z} \right) : \bar{\partial} X_0^\nu e^{ik \cdot X_0} : \\ &\sim ik^\mu : \bar{\partial} X_0^\nu e^{ik \cdot X_0} : \left( -\frac{\alpha'}{2} \right) \frac{1}{z^3} + \left( 1 + \frac{\alpha' k^2}{4} \right) \frac{G_k^{\mu\nu}(0)}{z^2} + \frac{\partial G_k^{\mu\nu}(0)}{z}, \end{aligned} \quad (5)$$

$$\tilde{T}(\bar{z}) G_k^{\mu\nu}(0) \sim ik^\nu : \partial X_0^\mu e^{ik \cdot X_0} : \left( -\frac{\alpha'}{2} \right) \frac{1}{\bar{z}^3} + \left( 1 + \frac{\alpha' k^2}{4} \right) \frac{G_k^{\mu\nu}(0)}{\bar{z}^2} + \frac{\partial G_k^{\mu\nu}(0)}{\bar{z}}, \quad (6)$$

<sup>1</sup>Reference: Polchinski, and [physics.stackexchange.com/a/389193](https://physics.stackexchange.com/a/389193).

Therefore,  $G_{e,k}$  is a primary of weight  $(1,1)$  iff.  $1 + \frac{\alpha' k^2}{4} = 1$  and  $k^\mu e_{\mu\nu} = 0 = k^\nu e_{\mu\nu}$ . The first equation gives the mass shell condition  $m^2 = -k^2 = 0$  for a massless boson, while the second equation constrains the polarization to be transverse. These are the physical constraints for a massless gauge boson, which is the level 1 excitation for a bosonic closed string.

(b) The form of any primary 3-point function is completely fixed by  $\text{PSL}(2, \mathbb{C})$  invariance<sup>2</sup>. In fact, for any holomorphic  $\phi_i(z_i)$  with weight  $h_i$ , by translational invariance, we have:

$$\langle \phi_1(z_1) \phi_2(z_2) \phi_3(z_3) \rangle = f(z_{12}, z_{23}, z_{31}), \quad z_{ij} = z_i - z_j, \quad (7)$$

Furthermore, scaling invariance requires that  $f$  is quasi-homogeneous:

$$\begin{aligned} z \mapsto z' = \lambda^{-1} z, \quad f &\mapsto \langle \lambda^{h_1} \phi_1(\lambda z_1) \lambda^{h_2} \phi_2(\lambda z_2) \lambda^{h_3} \phi_3(\lambda z_3) \rangle \\ &= \lambda^{h_1+h_2+h_3} f(\lambda z_{12}, \lambda z_{23}, \lambda z_{31}) \\ &= f(z_{12}, z_{23}, z_{31}), \end{aligned} \quad (8)$$

$$f = \sum_{a+b+c=\sum_i h_i} f_{abc} = \sum_{a+b+c=\sum_i h_i} \frac{C_{abc}}{z_{12}^a z_{23}^b z_{31}^c} \quad (9)$$

On the other hand, for special conformal transformation<sup>3</sup>  $\frac{1}{z} \mapsto \frac{1}{z'} = \frac{1}{z} + a$ , we have:

$$z \mapsto z' = \frac{1}{\frac{1}{z} + \bar{a}} = \frac{z}{1 + z\bar{a}} = w(z), \quad \frac{\partial z}{\partial z'} = \frac{1}{(1 - z\bar{a})^2} = \frac{1}{\kappa^2}, \quad z_{ij} = \frac{z'_{ij}}{\kappa_i \kappa_j}, \quad (10)$$

$$f \mapsto f(w^{-1}(z_{12}), w^{-1}(z_{23}), w^{-1}(z_{31})) \frac{1}{\kappa_1^{2h_1} \kappa_2^{2h_2} \kappa_3^{2h_3}} = f(z_{12}, z_{23}, z_{31}), \quad (11)$$

$$f_{abc}(w^{-1}(z_{12}), w^{-1}(z_{23}), w^{-1}(z_{31})) = f_{abc}(z_{12}, z_{23}, z_{31}) \kappa_1^{c+a} \kappa_2^{a+b} \kappa_3^{b+c}, \quad (12)$$

We see that  $f$  is invariant under special conformal transformation iff.  $f = f_{abc}$  where:

$$c + a = 2h_1, \quad a + b = 2h_2, \quad b + c = 2h_3, \quad (13)$$

$$\text{i.e. } a = h_1 + h_2 - h_3, \quad b = h_2 + h_3 - h_1, \quad c = h_3 + h_1 - h_2, \quad (14)$$

In the above discussions we've restricted  $\phi_i$  to be holomorphic; for *spin-less*  $\phi_i = \phi_i(z, \bar{z})$ ,  $h_i = \tilde{h}_i$ ,  $\Delta_i = h_i + \tilde{h}_i$ , the holomorphic and anti-holomorphic contributions can be nicely combined, and we have:

$$f = \frac{C}{|z_{12}|^{2a} |z_{23}|^{2b} |z_{31}|^{2c}}, \quad (15)$$

$$2a = \Delta_1 + \Delta_2 - \Delta_3, \quad 2b = \Delta_2 + \Delta_3 - \Delta_1, \quad 2c = \Delta_3 + \Delta_1 - \Delta_2, \quad (16)$$

$$\langle V_{k_1}(z_1, \bar{z}_1) V_{k_2}(z_2, \bar{z}_2) G_{e,k_3}(z_3, \bar{z}_3) \rangle = \frac{A(k_1, k_2, e)}{|z_{12}|^2 |z_{23}|^2 |z_{31}|^2} \quad (17)$$

<sup>2</sup>Reference: Blumenhagen, *Introduction to CFT*, and also *Di Francesco et al.*

<sup>3</sup>See *Di Francesco et al.* and also [github.com/davidsd/ph229](https://github.com/davidsd/ph229).



$$\begin{aligned}
G_{k_2}^{\rho\sigma}(z_2, \bar{z}_2) G_{k_3}^{\mu\nu}(z_3, \bar{z}_3) \sim \dots + \left(-\frac{\alpha'}{2} \eta^{\rho\mu} \frac{1}{z_{23}^2}\right) \left(-\frac{\alpha'}{2} \eta^{\sigma\nu} \frac{1}{\bar{z}_{23}^2}\right) \times 1 \\
+ \left(-\frac{\alpha'}{2} \eta^{\rho\mu} \frac{1}{z_{23}^2}\right) \left(-\frac{\alpha'}{2} \frac{ik_3^\sigma}{\bar{z}_{23}}\right) \left(-\frac{\alpha'}{2} \frac{ik_2^\nu}{\bar{z}_{32}}\right) + (z \leftrightarrow \bar{z}, \rho \leftrightarrow \sigma, \mu \leftrightarrow \nu) \\
+ \left(-\frac{\alpha'}{2} \frac{ik_2^\rho}{z_{23}}\right) \left(-\frac{\alpha'}{2} \frac{ik_2^\sigma}{\bar{z}_{23}}\right) \left(-\frac{\alpha'}{2} \frac{ik_3^\mu}{z_{32}}\right) \left(-\frac{\alpha'}{2} \frac{ik_3^\nu}{\bar{z}_{32}}\right) + \dots \\
O_{k_1, k_2}(z_2, \bar{z}_2) G_{k_3}^{\mu\nu}(z_3, \bar{z}_3) \sim \dots - k_1^\mu k_1^\nu \left(\frac{\alpha'^2}{4}\right) \frac{1}{|z_{23}|^4} \\
- i^2 (k_1^\mu k_2^\nu + k_1^\nu k_2^\mu) \left(\frac{\alpha'}{2} (k_1 \cdot k_3)\right) \left(\frac{\alpha'^2}{4}\right) \frac{1}{|z_{23}|^4} \\
- i^4 k_3^\mu k_3^\nu \left(\frac{\alpha'}{2} (k_1 \cdot k_2)\right)^2 \left(\frac{\alpha'^2}{4}\right) \frac{1}{|z_{23}|^4} + \dots
\end{aligned} \tag{26}$$

Again, apply the on-shell conditions, and we find that:

$$\frac{\alpha'}{2} k_1 \cdot k_2 = -2, \quad \frac{\alpha'}{2} k_1 \cdot k_3 = -\frac{\alpha'}{2} k_1 \cdot (k_1 + k_2) = -\frac{\alpha'}{2} k_1^2 - \frac{\alpha'}{2} k_1 \cdot k_2 = -2 - (-2) = 0, \tag{27}$$

$$\begin{aligned}
A(k_1, k_2, e) &= -\frac{\alpha'^2}{4} (4e_{\mu\nu} k_3^\mu k_3^\nu + e_{\mu\nu} k_1^\mu k_1^\nu) = -\frac{\alpha'^2}{4} e_{\mu\nu} k_1^\mu k_1^\nu \\
&= -\frac{\alpha'^2}{4} e_{\mu\nu} (k_2 + k_3)^\mu (k_2 + k_3)^\nu = -\frac{\alpha'^2}{4} e_{\mu\nu} k_2^\mu k_2^\nu \\
&= -\frac{\alpha'^2}{8} e_{\mu\nu} (k_1^\mu k_1^\nu + k_2^\mu k_2^\nu) \\
&= -\frac{\alpha'^2}{8} e_{\mu\nu} (k_{12}^\mu k_{12}^\nu + (k_1^\mu k_2^\nu + k_1^\nu k_2^\mu)),
\end{aligned} \tag{28}$$

On the other hand,

$$0 = e_{\mu\nu} k_3^\mu k_3^\nu = e_{\mu\nu} (k_1 + k_2)^\mu (k_1 + k_2)^\nu = e_{\mu\nu} (k_{12}^\mu k_{12}^\nu + 2(k_1^\mu k_2^\nu + k_1^\nu k_2^\mu)) \tag{29}$$

$$A(k_1, k_2, e) = -\frac{\alpha'^2}{8} e_{\mu\nu} k_{12}^\mu k_{12}^\nu \left(1 - \frac{1}{2}\right) = -\frac{\alpha'^2}{16} e_{\mu\nu} k_{12}^\mu k_{12}^\nu \tag{30}$$

## 2 Strings Scattering Off a Heavy Particle:

A heavy particle can be modeled by some D0-brane with Neumann boundary condition in the  $X_0$  direction<sup>4</sup>. The scattering of a closed string tachyon off the heavy particle can then be computed via a disc diagram with two insertions.

(a) The conformal Killing group (CKG) of the disc is  $\text{PSL}(2, \mathbb{R})$ . It is a 3 dimensional  $\mathbb{R}$  Lie group, generated by 3 conformal Killing vectors (CKV's); therefore, it is possible to partially fix the positions of the two insertions  $V_1, V_2$ . On the upper half plane, this can be implemented by putting  $z_1, z_2$  on the imaginary axis, with  $z_2$  fixed and  $z_1$  integrated<sup>5</sup>:

$$\mathcal{A} = g_c^2 e^{-\lambda} \int_0^{z_2} dz_1 \left\langle : c_1^x e^{ik_1 \cdot X_1} : : c_2 \tilde{c}_2 e^{ik_2 \cdot X_2} : \right\rangle, \quad z_2 = i, \quad z_1 = iy, \quad y \in [0, 1] \tag{31}$$

<sup>4</sup>Reference: [arXiv:hep-th/9611214](#), [arXiv:hep-th/9605168](#), and *Polchinski*.

<sup>5</sup>Reference: [arXiv:0812.4408](#). I would like to thank Lucy Smith for pointing this out.

Here  $c^x$  comes from the CKV that brings  $z_1 \rightarrow iy$ . On the disc this can be taken to be a rotation around  $z_2$ ; when mapped to the upper half plane and at around the imaginary axis, this is simply a translation along the  $x = \frac{1}{2}(z + \bar{z})$  direction<sup>6</sup>, i.e.

$$\text{CKV: } \partial_x = \delta_x^a \partial_a \implies \text{Ghost: } c^x, \quad (32)$$

$$c^x \partial_x + c^y \partial_y = c^z \partial_z + c^{\bar{z}} \partial_{\bar{z}}, \quad c^x = \frac{1}{2}(c^z + c^{\bar{z}}) = \frac{1}{2}(c(z) + \tilde{c}(\bar{z})), \quad (33)$$

The ghost contribution is then:

$$\begin{aligned} \langle c_1^x c_2 \tilde{c}_2 \rangle &= \langle c^x(z_1) c(z_2) \tilde{c}(\bar{z}_2) \rangle = \frac{1}{2} \left( \langle c(z_1) c(z_2) \tilde{c}(\bar{z}_2) \rangle + \langle \tilde{c}(z_1) c(z_2) \tilde{c}(\bar{z}_2) \rangle \right) \\ &= \frac{1}{2} \left( \langle c(z_1) c(z_2) c(z'_2) \rangle + \langle c(z'_1) c(z_2) c(z'_2) \rangle \right), \quad z' = \bar{z}, \\ &= \frac{C_{D^2}^g}{2} (z_{12} z_{12'} z_{22'} + z_{1'2} z_{1'2'} z_{22'}), \quad z_1, z_2 \in i\mathbb{R}, \\ &= 2C_{D^2}^g (z_1^2 - z_2^2) z_2 \end{aligned} \quad (34)$$

On the other hand, the  $e^{ik_j \cdot X_j}$  contributions is similar to what we've computed in [1], except that now we should be careful about the boundary conditions of  $X^\mu$ , which affect the  $XX$  contraction in the formulae. For Neumann boundary condition:  $\partial_y X^0 = 0$ , the half-plane propagator from  $z'$  can be constructed with an image at  $\bar{z}'$  with *the same charge*, i.e. we have:

$$\overline{X_1^0 X_2^0} = -\frac{\alpha'}{2} \eta^{00} \ln |z_1 - z_2|^2 - \frac{\alpha'}{2} \eta^{00} \ln |z_1 - \bar{z}_2|^2 \quad (35)$$

While for Dirichlet boundary  $X^i = \text{const}$ , we can always select the origin so that  $X^i = 0$ , and in this case the image should have *the opposite charge*, i.e.

$$\overline{X_1^i X_2^j} = -\frac{\alpha'}{2} \delta^{ij} \ln |z_1 - z_2|^2 + \frac{\alpha'}{2} \delta^{ij} \ln |z_1 - \bar{z}_2|^2, \quad (36)$$

$$\begin{aligned} \implies :e^{ik_1 \cdot X_1} : :e^{ik_2 \cdot X_2} : &= \exp \left( ik_{1,\mu} ik_{2,\nu} \overline{X_1^\mu X_2^\nu} \right) :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \\ &= |z_{12}|^{\alpha' k_1 \cdot k_2} |\bar{z}_{1\bar{2}}|^{\alpha' (-k_1^0 k_2^0 - \mathbf{k}_1 \cdot \mathbf{k}_2)} :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \end{aligned} \quad (37)$$

Before further calculations, we note that the normal ordering defined here on  $D^2$  differs from that on the usual  $\mathbb{C}^2$ ; in fact, there are also self-contractions with image charge<sup>7</sup>:

$$\overline{X^\mu(z, \bar{z}) X^\nu(\bar{z}, z)} = G_r^{\mu\nu}(z, \bar{z}) = \mp \frac{\alpha'}{2} \eta^{\mu\nu} \ln |z - \bar{z}|^2, \quad (38)$$

$$\implies \left\langle :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \right\rangle_{D^2} = \left\langle :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \right\rangle_{\mathbb{C}^2} \exp \left( \frac{1}{2} \sum_n ik_{n,\mu} ik_{n,\nu} \overline{X_n^\mu X_n^\nu} \right), \quad n = 1, 2 \quad (39)$$

The “ $\mp$ ” sign choice depends on the boundary condition.

<sup>6</sup>Reference: *Polchinski*, Chapter 5 & 6.

<sup>7</sup>This is very much similar to the torus situation, where we also have to consider self-contractions with image charges. More rigorous discussion of  $G^r$  is given in *Polchinski*.

Therefore,

$$\begin{aligned}
\left\langle :e^{ik_1 \cdot X_1} : : e^{ik_2 \cdot X_2} : \right\rangle_{D^2} &= \left\langle :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \right\rangle_{\mathbb{C}^2} \exp \left( ik_{1,\mu} ik_{2,\nu} \overline{X_1^\mu} X_2^\nu \right) \exp \left( \frac{1}{2} \sum_n ik_{n,\mu} ik_{n,\nu} \overline{X_n^\mu} X_n^\nu \right) \\
&= \left\langle :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \right\rangle_{\mathbb{C}^2} \exp \left( \frac{1}{2} \sum_{m,n} ik_{m,\mu} ik_{n,\nu} \overline{X_m^\mu} X_n^\nu \right) \\
&= \left\langle :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \right\rangle_{\mathbb{C}^2} |z_{12}|^{\alpha' k_1 \cdot k_2} |z_{1\bar{2}}|^{\alpha' (-k_1^0 k_2^0 - \mathbf{k}_1 \cdot \mathbf{k}_2)} \prod_n |z_{n\bar{n}}|^{\frac{\alpha'}{2} (-(k_n^0)^2 - \mathbf{k}_n^2)}
\end{aligned} \tag{40}$$

Note that  $X^i$  has no zero mode due to the Dirichlet boundary, hence  $\int \mathcal{D}X$  gives a delta function in only the Neumann direction:  $\delta(k_1^0 + k_2^0)$ . Physically, this means that only the energy is conserved; the momentum  $k^i$  is not conserved since the heavy D0-brane does not recoil. It is therefore convenient to define these on shell variables:

$$s = \omega^2 = (k_1^0)^2 = (k_2^0)^2, \quad t = -(\mathbf{k}_1 + \mathbf{k}_2)^2 = -\mathbf{k}_1^2 - \mathbf{k}_2^2 - 2\mathbf{k}_1 \cdot \mathbf{k}_2 = 2 \left( -\omega^2 - \mathbf{k}_1 \cdot \mathbf{k}_2 - \frac{4}{\alpha'} \right), \tag{41}$$

$$\mathbf{k}_1 \cdot \mathbf{k}_2 = -\frac{t}{2} - \omega^2 - \frac{4}{\alpha'}, \quad k_1 \cdot k_2 = -\omega(-\omega) + \mathbf{k}_1 \cdot \mathbf{k}_2 \tag{42}$$

Here we've used the on-shell condition:  $m^2 = -k^2 = \omega^2 - \mathbf{k}^2 = -\frac{4}{\alpha'}$  for tachyons. The previous expressions can then be simplified to:

$$\begin{aligned}
\left\langle :e^{ik_1 \cdot X_1} : : e^{ik_2 \cdot X_2} : \right\rangle_{D^2} &= \left\langle :e^{ik_1 \cdot X_1} e^{ik_2 \cdot X_2} : \right\rangle_{\mathbb{C}^2} |z_{12}|^{-\frac{\alpha'}{2}t-4} |z_{1\bar{2}}|^{+\frac{\alpha'}{2}t+4+2\alpha'\omega^2} \prod_n |2z_n|^{-\alpha'\omega^2-2} \\
&= iC_{D^2}^X 2\pi \delta(k_1^0 + k_2^0) |z_{12}|^{-\frac{\alpha'}{2}t-4} |z_{1\bar{2}}|^{+\frac{\alpha'}{2}t+4+2\alpha'\omega^2} \prod_n |2z_n|^{-\alpha'\omega^2-2} \\
&= iC_{D^2}^X 2\pi \delta(k_1^0 + k_2^0) f(|z_{12}|, |z_{1\bar{2}}|, |z_1|, |z_2|),
\end{aligned} \tag{43}$$

$$\begin{aligned}
\mathcal{A} &= g_c^2 e^{-\lambda} \cdot iC_{D^2}^X 2\pi \delta(k_1^0 + k_2^0) \cdot 2C_{D^2}^g \int_0^{z_2} dz_1 (z_1^2 - z_2^2) z_2 f(|z_{12}|, |z_{1\bar{2}}|, |z_1|, |z_2|) \\
&= g_c^2 C_{D^2} 2\pi \delta(k_1^0 + k_2^0) \cdot 2i \int_0^1 dy ((iy)^2 - i^2) i \cdot f(1-y, 1+y, 2y, 2) \\
&= -ig_c^2 C_{D^2} 2\pi \delta(k_1^0 + k_2^0) \cdot 2 \cdot 2^{-2\alpha'\omega^2-4} \int_0^1 dy (1-y^2) f(1-y, 1+y, y, 1),
\end{aligned} \tag{44}$$

$$\begin{aligned}
\int_0^1 dy (1-y^2) f(1-y, 1+y, y, 1) &= \int_0^1 dy (1-y)^{-\frac{\alpha'}{2}t-4+1} (1+y)^{+\frac{\alpha'}{2}t+4+2\alpha'\omega^2+1} y^{-\alpha'\omega^2-2} \\
&= \int_0^1 dy y^{a-1} (1-y)^{2b-1} (1+y)^{-2a-2b+1}, \quad y' = \frac{1-y}{1+y}, \\
&= -2^{1-2a} \int_0^1 dy' (-y')^{2b-1} (1-y'^2)^{a-1} \\
&= 2^{-2a} \int_0^1 d(y'^2) (y'^2)^{b-1} (1-y'^2)^{a-1} \\
&= 2^{-2a} B \left( a = -\alpha'\omega^2 - 1, b = -\frac{\alpha't}{4} - 1 \right)
\end{aligned} \tag{45}$$

Here  $B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$  is the Euler Beta function.

Putting everything together, we obtain:

$$\begin{aligned}\mathcal{A} &= -ig_c^2 C_{D^2} 2\pi \delta(k_1^0 + k_2^0) \cdot \frac{1}{2} B\left(-\alpha'\omega^2 - 1, -\frac{\alpha't}{4} - 1\right) \\ &= -ig_c^2 C_{D^2} \pi \delta(k_1^0 + k_2^0) B\left(-\alpha'\omega^2 - 1, -\frac{\alpha't}{4} - 1\right)\end{aligned}\quad (46)$$

In fact  $g_c^2 C_{D^2}$  can be further computed by path integral or by comparing physical results. Here we settle for this generic coefficient since it's already enough for our following discussions<sup>8</sup>.

(b) The Regge limit is found by taking the high energy limit while keeping the momentum transfer fixed; in this case it is achieved by:

$$\text{Regge: } s = \omega^2 \rightarrow \infty, \quad t = -(\mathbf{k}_1 + \mathbf{k}_2)^2 \text{ fixed}, \quad (47)$$

$$\begin{aligned}\mathcal{A} \propto B\left(a = -\alpha's - 1, b = -\frac{\alpha't}{4} - 1\right) &= \frac{\Gamma(-\alpha's - 1)}{\Gamma(-\alpha's - \frac{\alpha't}{4} - 2)} \Gamma\left(-\frac{\alpha't}{4} - 1\right) \\ &\sim \left\{e\left(\alpha's + \frac{\alpha't}{4} + 3\right)\right\}^{\frac{\alpha't}{4} + 1} \Gamma\left(-\frac{\alpha't}{4} - 1\right) \\ &\sim (e\alpha'\omega^2)^{\frac{\alpha't}{4} + 1} \Gamma\left(-\frac{\alpha't}{4} - 1\right) \\ &\sim (\omega^2)^{\frac{\alpha't}{4} + 1} \Gamma\left(-\frac{\alpha't}{4} - 1\right)\end{aligned}\quad (48)$$

Here we've used the Stirling's approximation<sup>9</sup>:  $\ln \Gamma(z + 1) = \ln z! \sim z \ln z - z$ . On the other hand, the hard scattering limit is found by keeping the scattering angle fixed, i.e.

$$\text{Hard scattering: } s = \omega^2 \rightarrow \infty, \quad (t/s) \equiv \lambda \text{ fixed}, \quad (49)$$

$$\mathcal{A} \propto B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} \sim \exp\left\{-\alpha'\left(s \ln(\alpha's) + \frac{t}{4} \ln \frac{\alpha't}{4} + \frac{u}{4} \ln \frac{\alpha'u}{4}\right)\right\}, \quad (50)$$

$$s = \omega^2 = (k_1^0)^2 = (k_2^0)^2, \quad t = -(\mathbf{k}_1 + \mathbf{k}_2)^2, \quad u = -(\mathbf{k}_1 - \mathbf{k}_2)^2, \quad (51)$$

$$s + \frac{t}{4} + \frac{u}{4} = -\frac{4}{\alpha'}, \quad (52)$$

Here we've introduced an additional  $u$  variable, and we see that the result is symmetric under  $t \leftrightarrow u$ . We find that under the above limits, the amplitude exhibits similar behaviors as the Veneziano amplitude.

(c) Note that  $\Gamma(z)$  has no zeros on  $\mathbb{C}^2$ , therefore the poles of  $B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$  is given by the poles of  $\Gamma(a)$  and  $\Gamma(b)$ :

$$a = -\alpha's - 1 = 0, -1, -2, \dots, \quad b = -\frac{\alpha't}{4} - 1 = 0, -1, -2, \dots, \quad (53)$$

---

<sup>8</sup>And I have run out of time and energy.

<sup>9</sup>For the validity of Stirling's approximation when  $z \in \mathbb{C}$ ,  $\arg z = \pi - \epsilon$  and  $|z| \rightarrow \infty$ , see [Wikipedia: Stirling's formula for the gamma function](#).

The first set of poles gives:

$$\omega^2 = -\frac{1}{\alpha'}, 0, \frac{1}{\alpha'}, \frac{2}{\alpha'}, \dots \quad (54)$$

This is precisely the open string spectrum in  $D = 26$ . Going back to (43)~(45), we see that they come from the singularities as  $y = |z_1| \rightarrow 0$  or  $|z_{12}| \rightarrow 1$ , i.e. the two insertions are far apart.

This is the  $s$ -channel contribution; physically, it means that the closed string is first absorbed by the D0-brane, then it propagates as an intermediate open string on the brane, and is re-emitted in the end.

The other set of poles gives:

$$-t = (\mathbf{k}_1 + \mathbf{k}_2)^2 = \frac{4}{\alpha'}, 0, -\frac{4}{\alpha'}, -\frac{8}{\alpha'}, \dots \quad (55)$$

Only two poles are realistic due to  $\mathbf{k}^2 \geq 0$ . This is the  $t$ -channel contribution, which corresponds to a closed string exchange between the D0-brane and the incoming particle. The above factorization channels are nicely illustrated in the diagram below, borrowed from [arXiv:hep-th/9611214](https://arxiv.org/abs/hep-th/9611214).



### 3 Compton Scattering Between $U(N)$ Gluons and Adjoint Tachyons:

Such scattering is captured by a disc diagram with  $AATT$  insertions at the boundary; following the same recipe as before, we write down<sup>10</sup>:

$$\begin{aligned} \mathcal{A} = & g_o'^2 g_o^2 e^{-\lambda} \int dy \left\langle : e_\mu \partial_y X^\mu e^{ik \cdot X}(y) : : c_1^y e_{1,\nu} \partial_y X_1^\nu e^{ik_1 \cdot X_1} : : c_2^y e^{ik_2 \cdot X_2} : : c_3^y e^{ik_2 \cdot X_2} : \right\rangle \\ & \times \text{Tr} (\lambda^a \lambda^{a_1} \lambda^{a_2} \lambda^{a_3} : \text{in } (y, y_1, y_2, y_3) \text{ order}) \\ & + (2 \leftrightarrow 3) \end{aligned} \quad (56)$$

Where  $y_i = y_{1,2,3}$  are fixed while  $y_0 = y$  is integrated. Note that for [2] we use  $(x, y)$  to label the points on the upper half plane so that the boundary coordinate is  $x = (x, 0)$ ; here in order to match with *Polchinski*, we label the boundary points with  $y = (y, 0)$  instead.

<sup>10</sup>I would like to thank 谷夏 for help with this problem.



We see that  $\mathcal{A}$  is structurally similar to the 4-point Veneziano amplitude  $\mathcal{A}_{TTTT}$  and also the 3-point amplitude  $\mathcal{A}_{TTA}$ ; by analogy we can evaluate  $\mathcal{A}$  in a similar fashion<sup>11</sup>:

$$\begin{aligned} \mathcal{A} = & g_o'^2 g_o^2 e^{-\lambda} \prod_{i < j} |y_{ij}|^{2\alpha' k_i \cdot k_j + 1} e_\mu e_{1,\nu} \\ & \times \int dy \prod_j |y_{0j}|^{2\alpha' k \cdot k_j} \left\{ -2\alpha' \frac{\eta^{\mu\nu}}{y_{01}^2} + (-2\alpha') \left( \frac{ik_1^\mu}{y_{01}} + \frac{ik_2^\mu}{y_{02}} + \frac{ik_3^\mu}{y_{03}} \right) (-2\alpha') \left( \frac{ik^\nu}{y_{10}} + \frac{ik_2^\nu}{y_{12}} + \frac{ik_3^\nu}{y_{13}} \right) \right\} \\ & \times \text{Tr}(\cdots) iC_{D_2}^X C_{D_2}^g (2\pi)^{26} \delta^{26}(\sum_i k_i) \\ & + (2 \leftrightarrow 3) \end{aligned} \quad (57)$$

Notice the doubling due to the boundary contraction  $\overline{X_1^\mu} X_2^\nu = 2 \times (-\frac{\alpha'}{2}) \ln y_{12}^2 = -2\alpha' \ln y_{12}$ . Further simplifications can be achieved by using the on-shell and physical conditions for open string states:

$$k^2 = k_1^2 = 0, \quad e \cdot k = e_1 \cdot k_1 = 0, \quad k_{2,3}^2 = \frac{1}{\alpha'} = -m^2, \quad (58)$$

$$\begin{aligned} s &= -\alpha' (k + k_1)^2 = -2\alpha' k \cdot k_1, \\ t &= -\alpha' (k + k_2)^2 = -2\alpha' k \cdot k_2 - 1, \\ u &= -\alpha' (k + k_3)^2 = -2\alpha' k \cdot k_3 - 1, \end{aligned} \quad (59)$$

$$s + t + u = -2, \quad (60)$$

$$2\alpha' k_1 \cdot k_2 = \alpha' (k_1 + k_2)^2 - \alpha' k_1^2 - \alpha' k_2^2 = \alpha' (k + k_3)^2 - 1 = -u - 1 = 2\alpha' k \cdot k_3, \quad (61)$$

$$2\alpha' k_1 \cdot k_3 = -t - 1 = 2\alpha' k \cdot k_2, \quad (62)$$

$$2\alpha' k_2 \cdot k_3 = \alpha' (k + k_1)^2 - 2 = -s - 2, \quad (63)$$

$$\begin{aligned} \mathcal{A} = & i g_o'^2 g_o^2 C_{D^2} |y_{12}|^{-u} |y_{13}|^{-t} |y_{23}|^{-s-2} e_\mu e_{1,\nu} \\ & \times \int dy |y_{01}|^{-s} |y_{02}|^{-t-1} |y_{03}|^{-u-1} \left\{ \frac{\eta^{\mu\nu}}{y_{01}^2} - 2\alpha' \left( \frac{ik_1^\mu}{y_{01}} + \frac{ik_2^\mu}{y_{02}} + \frac{ik_3^\mu}{y_{03}} \right) \left( -\frac{ik^\nu}{y_{01}} + \frac{ik_2^\nu}{y_{12}} + \frac{ik_3^\nu}{y_{13}} \right) \right\} \\ & \times \text{Tr}(\cdots) (-2\alpha') (2\pi)^{26} \delta^{26}(\sum_i k_i) \\ & + (2 \leftrightarrow 3) \end{aligned} \quad (64)$$

Fix  $y_1 \rightarrow \infty, y_2 = 0, y_3 = 1$ , and we get:

$$\begin{aligned} \mathcal{A} = & i g_o'^2 g_o^2 C_{D^2} |y_1|^{-u-t} e_\mu e_{1,\nu} \\ & \times \int dy |y_{01}|^{-s} |y_{02}|^{-t-1} |y_{03}|^{-u-1} \left\{ \frac{\eta^{\mu\nu}}{y_{01}^2} - 2\alpha' \left( \frac{ik_1^\mu}{y_{01}} + \frac{ik_2^\mu}{y_{02}} + \frac{ik_3^\mu}{y_{03}} \right) \left( -\frac{ik^\nu}{y_{01}} + \frac{ik_2^\nu}{y_{12}} + \frac{ik_3^\nu}{y_{13}} \right) \right\} \\ & \times \text{Tr}(\cdots) \cdots + \cdots \end{aligned} \quad (65)$$

---

<sup>11</sup>For the  $\partial X$  contribution, see *Polchinski*, [arXiv:1403.4553](#) and [arXiv:1507.02172](#). Note that there are contractions between two  $\partial X$  fields.

$$\begin{aligned}
\mathcal{A} &= i g_o'^2 g_o^2 C_{D^2} \\
&\times \int dy |y_{02}|^{-t-1} |y_{03}|^{-u-1} \left| \frac{y_1^{s+2}}{y_{01}^s} \right| \left\{ \frac{e \cdot e_1}{y_{01}^2} - 2\alpha' e_\mu \left( \frac{i k_1^\mu}{y_{01}} + \frac{i k_2^\mu}{y_{02}} + \frac{i k_3^\mu}{y_{03}} \right) e_{1,\nu} \left( -\frac{i k^\nu}{y_{01}} + \frac{i k_2^\nu}{y_{12}} + \frac{i k_3^\nu}{y_{13}} \right) \right\} \\
&\times \text{Tr}(\dots) \dots + \dots \\
&= i g_o'^2 g_o^2 C_{D^2} \int dy \left\{ (e \cdot e_1) f(t, u) - 2\alpha' g(t, u) \right\} \times \dots + \dots
\end{aligned} \tag{66}$$

Note that the definition of  $f(t, u)$  and  $g(t, u)$  contains the  $\text{Tr}(\dots)$  factor.

The limit  $y_1 \rightarrow \infty$  should be treated with care. In fact, the integral along the boundary splits into three ranges<sup>12</sup>:

$$\int dy = \left\{ \int_{y_1 \rightarrow -\infty}^{y_3=0} + \int_{y_2=0}^{y_3=1} + \int_{y_3=1}^{y_1 \rightarrow +\infty} \right\} dy \tag{67}$$

Notice the difference of  $y_1 \rightarrow \pm\infty$ ; this is due to the  $S^1$  topology of the boundary. Therefore,

$$\int dy f(t, u) = \int dy |y_{02}|^{-t-1} |y_{03}|^{-u-1} \left| \frac{y_1}{y_{01}} \right|^{s+2} \text{Tr}(\dots), \tag{68}$$

$$\text{Tr}(\lambda^a \lambda^{a_1} \lambda^{a_2} \lambda^{a_3}) \equiv T_{0123}, \tag{69}$$

$$\begin{aligned}
\int_{y_1 \rightarrow -\infty}^{y_2=0} dy f(t, u) &= \int_{y_1 \rightarrow -\infty}^0 dy (1-y)^{-t-1} (-y)^{-u-1} \left( \frac{-y_1}{y-y_1} \right)^{s+2} T_{1023} = T_{1023} B(-u, -s-1), \\
\int_{y_3=1}^{y_1 \rightarrow +\infty} dy f(t, u) &= \int_1^{y_1 \rightarrow +\infty} dy (y-1)^{-t-1} y^{-u-1} \left( \frac{y_1}{y_1-y} \right)^{s+2} T_{1230} = T_{1230} B(-t, -s-1), \\
\int_{y_3=0}^{y_2=1} dy f(t, u) &= \int_0^1 dy (1-y)^{-t-1} y^{-u-1} t_{1203} = T_{1203} B(-t, -u),
\end{aligned} \tag{70}$$

$$\int dy f(t, u) = T_{1203} B(-t, -u) + T_{1230} B(-t, -s-1) + T_{1023} B(-u, -s-1) \tag{71}$$

Under  $2 \leftrightarrow 3$ , we have  $t \leftrightarrow u$ ; therefore,

$$\begin{aligned}
\int dy f(t, u) + (2 \leftrightarrow 3) &= (T_{1203} + T_{1302}) B(-t, -u) \\
&\quad + (T_{1230} + T_{1032}) B(-t, -s-1) \\
&\quad + (T_{1023} + T_{1320}) B(-u, -s-1) \\
&\equiv \alpha B(-t, -u) + \beta B(-t, -s-1) + \gamma B(-u, -s-1)
\end{aligned} \tag{72}$$

---

<sup>12</sup>Reference: *Polchinski's* discussion of  $\mathcal{A}_{TTTT}$ .

The  $g(t, u)$  factor can be computed in a similar manner. In the end we have<sup>13</sup>:

$$\begin{aligned}
 \mathcal{A} = & ig_o'^2 g_o^2 C_{D^2} (2\pi)^{26} \delta^{26}(\sum_i k_i) \\
 & \times (-2\alpha') \left\{ (e_0 \cdot e_1) (\alpha B(-t, -u) + \beta B(-t, -s-1) + \gamma B(-u, -s-1)) \right. \\
 & \left. + (-2\alpha')(e_0 \cdot k_2)(e_1 \cdot k_3) (\alpha B(-t, -u) - \beta B(-t, -s-1) - \gamma B(-u, -s-1)) + (2 \leftrightarrow 3) \right\}
 \end{aligned}
 \tag{73}$$

---

<sup>13</sup>Reference: [arXiv:0801.3358](#).