

CMV primary episode analysis

Bryan Mayer

March 6, 2016


Contents

Description	1
Overview of the data	2
General statistics, peak shedding, duration	2
Comparison of oral and plasma shedding	4
Three phase classification	4
Summary of three phases	5
Compare three phase to two phase	5
Plot of features	6
Cubic model of clearance and rebound analysis	7
Correlation analysis	8
Age and features	8
Mother HIV	8
Saving figures and table	8

Description

This is the main analysis of the primary CMV oral shedding episodes. Here, we extract key characteristics of the shedding episodes (e.g., peak load), determine the cutoffs for the three phases, analyze correlation between features, and analyze rebound behavior.

Overview of the data



General statistics, peak shedding, duration


	n	14
total negative swabs		4
median age (range) days		70 (31, 205)
percent mothers hiv +		50
total oral swabs		763
total plasma swabs		52
median duration (range) days		324.5 (244, 756)
median peak shedding duration (range) weeks		20.57 (7, 32.43)
median first positive (range) log10 conc.		2.86 (2.22, 4.12)
median peak shedding (range) log10 conc.		8.34 (6.96, 8.91)

Table 1: Summary statistics


PatientID2	n	days
B	2	10, 111
C	1	503
H	1	220

Table 2: Negative swabs

Red region denotes shedding within 1 log of peak



Comparison of oral and plasma shedding



total oral swabs	763
total plasma swabs	52
total positive plasma swabs	0
percent positive plasma swabs	55.76923
median log10 pls viral load (range)	2.5 (1.78, 3.92)
Pearson correlation with oral	0.22 (95% CI: -0.06, 0.46)
Spearman correlation with oral	0.2 (95% CI: -0.1, 0.47)
Pearson correlation with oral (pos plasma only)	-0.04 (95% CI: -0.4, 0.33)
Spearman correlation with oral (pos plasma only)	-0.13 (95% CI: -0.47, 0.26)
lmm oral count ~ binary pls predictor	0.81 (95% CI: -0.11, 1.73)

Table 3: Plasma load analysis

Three phase classification

Each episode has three characteristic phases:

1. expansion: oral episode start to 1 log before the peak
2. transition: 1 log before the peak until the peak
3. clearance: peak until end of observation

Summary of three phases

	Expansion	Transition	Clearance
Median duration (range) wk.	7 (3.71, 27.71)	6 (2, 17)	32.07 (15.43, 92.14)
Median swab change (range) wk.	0.64 (-1.77, 2.99)	0.09 (-1.03, 1.64)	-0.07 (-3.6, 3.13)
Median slope (range) wk.	0.7 (0.1, 1.62)	0.06 (0, 0.27)	-0.1 (-0.23, -0.02)
Median double/half-time (range) wk.	0.43 (0.19, 3.11)	NA (NA, NA)	3.11 (1.3, 15.3)
Median R-square (range)	0.92 (0.63, 0.98)	0.13 (0, 0.92)	0.7 (0.1, 0.85)

Table 4: Summary of phase features

Infant B had the lowest expansion R2 of 0.6276844 . Infant B had the a low clearance R2 of 0.1646749 . Infant N had the a low clearance R2 of 0.1028881 .

Compare three phase to two phase


The three phase classification comes from an apparent transition phase between the start of the episode and the peak viral load. In the biphasic classification, expansion phase ends at peak (instead of 1 log before peak). The R-square values are generally worse when we make that assumption.

Infant ID	Growth to peak	Growth to 1 log before peak	Triphasic > biphasic
A	0.67	0.96	TRUE
B	0.72	0.63	FALSE
C	0.80	0.84	TRUE
D	0.94	0.95	TRUE
E	0.85	0.89	TRUE
F	0.61	0.95	TRUE
G	0.93	0.85	FALSE
H	0.88	0.81	FALSE
I	0.76	0.94	TRUE
J	0.74	0.93	TRUE
K	0.79	0.87	TRUE
L	0.75	0.94	TRUE
M	0.65	0.98	TRUE
N	0.59	0.92	TRUE


Table 5: R-square comparing growth in two phase vs three phase classification

11 episodes perform better with transition phase

Plot of features



Cubic model of clearance and rebound analysis



Pair1	Pair2	raw_cor	p_value	boot_cor	boot_lower95CI	boot_upper95CI
Age (days)	Peak	-0.396	0.202	-0.381	-0.759	0.169
Age (days)	Expansions days	-0.285	0.369	-0.292	-0.830	0.429
Age (days)	Transition days	-0.156	0.627	-0.161	-0.690	0.513
Age (days)	Clearance days	0.190	0.555	0.193	-0.472	0.846
Peak	Expansions days	-0.095	0.768	-0.103	-0.734	0.584
Peak	Transition days	-0.382	0.221	-0.382	-0.884	0.316
Peak	Clearance days	0.084	0.795	0.086	-0.495	0.648
Expansions days	Transition days	0.247	0.438	0.272	-0.423	0.731
Expansions days	Clearance days	0.161	0.618	0.151	-0.506	0.757
Transition days	Clearance days	-0.284	0.371	-0.303	-0.865	0.445

Table 6: Spearman Correlations - excluded 2 older infants. Bootstrap takes median.

Pair1	Pair2	raw_cor	p_value	boot_cor	boot_lower95CI	boot_upper95CI
Age (days)	Peak	-0.590	0.026	-0.586	-0.846	-0.118
Age (days)	Expansions days	0.192	0.512	0.202	-0.521	0.720
Age (days)	Transition days	-0.308	0.284	-0.302	-0.815	0.312
Age (days)	Clearance days	0.044	0.881	0.049	-0.553	0.673
Peak	Expansions days	-0.385	0.175	-0.393	-0.874	0.299
Peak	Transition days	-0.104	0.725	-0.103	-0.690	0.534
Peak	Clearance days	0.095	0.748	0.093	-0.434	0.599
Expansions days	Transition days	-0.089	0.763	-0.091	-0.560	0.483
Expansions days	Clearance days	0.056	0.848	0.050	-0.532	0.653
Transition days	Clearance days	-0.202	0.489	-0.217	-0.778	0.492

Table 7: Spearman Correlations - all data. Bootstrap takes median.

Correlation analysis

Age and features

Mother HIV

outcome	estimate	p.value
Age (days)	-1.286	0.961
Peak	0.195	0.588
Expansions days	-6.857	0.772
Transition days	3.714	0.817
Clearance days	-97.000	0.219

Table 8: Linear model (unadjusted) with momhiv (reference = neg) as predictor.

Saving figures and table