

### 3.4 Location and scale families

**Fact.** if  $f(x)$  is a valid pdf on  $\mathbb{R}$   
then for any  $\mu \in \mathbb{R}$ ,  $\sigma > 0$   
the function

$$g(x|\mu, \sigma) := \frac{1}{\sigma} f\left(\frac{x-\mu}{\sigma}\right), x \in \mathbb{R}$$

is a valid pdf on  $\mathbb{R}$ .

**Proof**

- clearly,  $g \geq 0$  because  $f \geq 0$ .

- check the integral:

$$\begin{aligned} & \int_{\mathbb{R}} g(x|\mu, \sigma) dx \\ &= \int_{\mathbb{R}} \frac{1}{\sigma} f\left(\frac{x-\mu}{\sigma}\right) dx \\ &= \int \frac{1}{\sigma} f(y) \sigma dy \\ &= \int f(y) dy = 1. \end{aligned}$$

Let  $(x-\mu)/\sigma = y$   
 $\Rightarrow x = \mu + \sigma y$   
 $\rightarrow dx = \sigma dy$

Definition Let  $f(x)$  be any pdf (on  $\mathbb{R}$ ).

The family of pdf  $\left\{ g(x|\mu, \sigma) := \frac{1}{\sigma} f\left(\frac{x-\mu}{\sigma}\right), \mu \in \mathbb{R}, \sigma > 0 \right\}$ ,

is called the location-scale family of distributions

$\begin{cases} \mu : \text{location parameter} \\ \sigma : \text{scale parameter} \end{cases}$

Special cases:


①  $\left\{ g(x|\mu) := f(x-\mu) \mid \mu \in \mathbb{R} \right\}$  is a location family

②  $\left\{ g(x|\sigma) := \frac{1}{\sigma} g\left(\frac{x}{\sigma}\right) \mid \sigma > 0 \right\}$  is a scale family

$$g(x|\mu) = f(x-\mu)$$


↙ ↘  
μ < 0      μ > 0

μ > 0



$$\text{if } X \sim f \quad \text{then} \quad X + \mu \sim g(x) = f(x - \mu)$$

$$\text{if } X \sim f \quad \text{then} \quad \sigma X \sim g(x) = \frac{1}{\sigma} f\left(\frac{x}{\sigma}\right)$$



$$\text{if } X \sim f \quad \text{then} \quad \sigma X + \mu \sim g(x) = \frac{1}{\sigma} f\left(\frac{x - \mu}{\sigma}\right)$$

① Let  $f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2}$  then we obtain l-s family

$$\left\{ g(x|\mu, \sigma) = \frac{1}{\sigma} f\left(\frac{1}{\sigma}(x-\mu)\right) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} \mid \mu \in \mathbb{R}, \sigma > 0 \right\}$$

② Let  $f(x) = \frac{1}{\Gamma(\alpha)} x^{\alpha-1} e^{-x}$ , then we obtain

$$\left\{ g(x|\mu, \sigma) = \frac{1}{\Gamma(\alpha)} \frac{1}{\sigma} \left(\frac{x-\mu}{\sigma}\right)^{\alpha-1} e^{-\frac{x-\mu}{\sigma}} \mid \mu \in \mathbb{R}, \sigma > 0 \right\}$$

Not a gamma family

$$\left\{ g(x|\sigma) = \frac{1}{\Gamma(\alpha)} \frac{1}{\sigma} \left(\frac{x}{\sigma}\right)^{\alpha-1} e^{-\frac{x}{\sigma}} \mid \sigma \in \mathbb{R}_+ \right\}$$

$$\equiv \left\{ \text{Gamma}(\alpha, \sigma) \mid \sigma \in \mathbb{R}_+ \right\}$$

③ Let  $f(x) = e^{-\lambda} \lambda^x / x!$ ,  $x = 0, 1, 2, \dots$

define  $g(x|\sigma) = \frac{1}{\sigma} e^{-\lambda} \frac{\lambda^{x/\sigma}}{(x/\sigma)!}$  for  $x = \sigma y, y \in \mathbb{N}, \sigma > 0$ .

then  $g$  is a valid pmf supported by multiples of  $\sigma$ .

Not a Poisson family but inherits many properties of Poisson dist.

Theorem Suppose  $Y$  is a R.V. with pdf  $f(z)$  and  $EY$  and  $\text{var } Y$  exists  
if  $X$  is a R.V. with pdf  $\frac{1}{\sigma} f\left(\frac{z-\mu}{\sigma}\right)$ , then

$$\begin{cases} EX = \sigma EY + \mu \\ \text{var } X = \sigma^2 \text{var } Y. \end{cases}$$