




Model

Son Nguyen

Linear

- Given the data

The diagram shows a dataset with three columns: x_1 , x_2 , and y . The first two columns are grouped under the label "Input" with a red arrow, and the third column is labeled "target" with a red arrow. The data rows are as follows:

x_1	x_2	y
1	0	-2
2	1	0
3	-2	-1
4	3	1

- How are y and x related?

A review of Linear Model for Regression

- Given the data

x_1	x_2	y
1	0	-2
2	1	0
3	-2	-1
4	3	1

- Linear model predicts y is a linear combination of x_1, x_2

$$\hat{y} = w_0 + \underline{w_1}x_1 + \underline{w_2}x_2$$

Intercept     

slopes

A review of Linear Model for Regression

- Given the data

x_1	x_2	y
1	0	-2
2	1	0
3	-2	-1
4	3	1

- Linear model predicts y is a linear combination of x_1, x_2

$$\hat{y} = w_0 + w_1x_1 + w_2x_2$$

- The goal of linear model is to solve for w_0, w_1 and w_2
- To **train** a linear model is to find w_0, w_1 and w_2

How to find the coefficients?

- **Step 1:** Define the loss function $l(y, \hat{y})$

How to find the coefficients?

- **Step 1:** Define the loss function $l(y, \hat{y})$
- **Step 2:** Find w that minimizes the total loss function.

How to find the coefficients?

- Least Squared Method uses the **square loss**

$$l(\hat{y}, y) = (\hat{y} - y)^2$$

How to find the coefficients?

- Least Squared Method uses the **square loss**

$$l(\hat{y}, y) = (\hat{y} - y)^2$$

- We want to find w_0 , w_1 and w_2 that minimizes a **loss function**.

x_1	x_2	y	$\hat{y} = w_0 + w_1 x_1 + w_2 x_2$	$(\hat{y} - y)^2$
1	0	-2	$w_0 + w_1 \cdot 1 + w_2 \cdot 0$	$(w_0 + w_1 \cdot 1 + w_2 \cdot 0 + 2)^2$
2	1	0	$w_0 + w_1 \cdot 2 + w_2 \cdot 1$	$(w_0 + w_1 \cdot 2 + w_2 \cdot 1 - 0)^2$
3	-2	-1	$w_0 + w_1 \cdot 3 + w_2 \cdot -2$	$(w_0 + w_1 \cdot 3 + w_2 \cdot -2 + 1)^2$
4	3	1	$w_0 + w_1 \cdot 4 + w_2 \cdot 3$	$(w_0 + w_1 \cdot 4 + w_2 \cdot 3 - 1)^2$

How to find the coefficients?

- Least Squared Method uses the **square loss**

$$l(\hat{y}, y) = (\hat{y} - y)^2$$

- We want to find w_0 , w_1 and w_2 that minimizes a **loss function**.

x_1	x_2	y	$\hat{y} = w_0 + w_1 x_1 + w_2 x_2$	$(\hat{y} - y)^2$
1	0	-2	$w_0 + w_1 \cdot 1 + w_2 \cdot 0$	$(w_0 + w_1 \cdot 1 + w_2 \cdot 0 + 2)^2$
2	1	0	$w_0 + w_1 \cdot 2 + w_2 \cdot 1$	$(w_0 + w_1 \cdot 2 + w_2 \cdot 1 - 0)^2$
3	-2	-1	$w_0 + w_1 \cdot 3 + w_2 \cdot -2$	$(w_0 + w_1 \cdot 3 + w_2 \cdot -2 + 1)^2$
4	3	1	$w_0 + w_1 \cdot 4 + w_2 \cdot 3$	$(w_0 + w_1 \cdot 4 + w_2 \cdot 3 - 1)^2$

- The total loss function:

$$L = L(w_0, w_1, w_2) = (w_0 + w_1 + 2)^2 + (w_0 + 2w_1 + w_2)^2 + (w_0 + 3w_1 - 2w_2 + 1)^2 + (w_0 + 4w_1 + 3w_2 - 1)^2$$

need
to
minimize

}

How to find the coefficients?

- Least Squared Method uses the **square loss**

$$l(\hat{y}, y) = (\hat{y} - y)^2$$

- We want to find w_0 , w_1 and w_2 that minimizes a **loss function**.
- The total loss function:

$$L = L(w_0, w_1, w_2) = (w_0 + w_1 + 2)^2 + (w_0 + 2w_1 + w_2)^2 \\ + (w_0 + 3w_1 - 2w_2 + 1)^2 + (w_0 + 4w_1 + 3w_2 - 1)^2$$

- Solve for the partial derivatives equaling 0 to find w_0 , w_1 and w_2 .

$$x^2 + 2x + 3$$

$$f'(x) = 2x + 2 = 0$$

$$x = -1$$

How about other loss functions?

- Absolute loss:

$$L(\hat{y}, y) = |\hat{y} - y|$$

- The total loss function:

$$L = L(w_0, w_1, w_2) = |w_0 + w_1 + 2| + |w_0 + 2w_1 + w_2| \\ + |w_0 + 3w_1 - 2w_2 + 1| + |w_0 + 4w_1 + 3w_2 - 1|$$

- Use Linear Programming to find w_0 , w_1 and w_2 that minimizes the total loss.


- Least absolute deviations regression

square loss = $(y - \hat{y})^2$

↑ true ↑ prediction

abs. loss = $|y - \hat{y}|$


Linear Models



least Squared L.m

How about other loss functions?

Ordinary least squares regression	<u>Least absolute deviations regression</u>
Not very robust	<u>Robust</u>
Stable solution	Unstable solution
Always one solution	Possibly multiple solutions



A general framework

- **Problem:** Given the data of $x_1, x_2, \dots, x_d, y$, establish the best relation between y and $x = [x_1, x_2, \dots, x_d]$.

for example : linear relation

- A solution framework:

- Step 1: Assume the model function $\hat{y} = f(x, w)$, where w is a parameter vector.
- Step 2: Define the loss function $l(y, \hat{y})$
- Step 3: Find w that minimizes the loss function using gradient descent

example : squared loss

LASSO

- Consider a linear model

$$y = 100x_1 + \underline{0.01}x_2 + 50x_3 - \underline{0.002}x_4$$

- x_2 and x_4 are not important because the coefficients are too small.

- We want to get rid of x_2 and x_4

LASSO - Principle

- LASSO forces the sum of the absolute value of the coefficients to be less than a fixed value.
- which forces certain coefficients (slopes) to be set to zero
- effectively making the model simpler

Linear Model vs. LASSO - Principle

- Linear Model minimizes

$$L(w_0, w_1, w_2)$$

loss function

- LASSO minimizes

$$L(w_0, w_1, w_2) + \alpha (|w_1| + |w_2|)$$

- The greater α , the easier w_1 and w_2 will be zeros.

- When $\alpha = 0$, LASSO is the linear model.

loss f.


α is a hyper-parameter of LASSO

notice that the least square linear model does not
have a hyper-parameter

LASSO for Linear Model

$\text{Alpha} = 0$


Linear model $\equiv$ LASSO



alpha $\rightarrow$
slope $\rightarrow$ 0


LASSO for Linear Model

Alpha = 10




LASSO for Linear Model

Alpha = 20




LASSO for Linear Model

Alpha = 30



LASSO for Linear Model

Alpha = 40



the effect of
 x on y is
gone

LASSO for Variables Selection

- Data

x_1	x_2	x_3	x_4	x_5	x_6	y
...	...	...	...	...	...	...
...	...	...	...	...	...	...
...	...	...	...	...	...	...
...	...	...	...	...	...	...

data

- Assume that the truth relation between the input $x_1, x_2, x_3, x_4, x_5, x_6$ and the output y is

$$y = 4x_2 + 3x_4 + 7x_6$$

the truth

- We see that only x_2, x_4 and x_6 impact y

not a model

- LASSO can help to identify variables that have effect on y

LASSO for Variables Selection

- The result when training the linear model and the LASSO

	w_1	w_2	w_3	w_4	w_5	w_6
Truth	0	4	0	<u>3</u>	0	7
Linear Model	<u>-0.244061</u>	<u>3.54013</u>	<u>0.221939</u>	<u>2.6042</u>	<u>0.0982158</u>	<u>6.83617</u>
LASSO	-0	2.65623	0	<u>1.84839</u>	0	<u>5.80624</u>

- In Linear Model, x_1 , x_3 and x_5 have effect on y (which is WRONG!)
- In LASSO, x_1 , x_3 and x_5 have no effect on y (CORRECT!)
- LASSO can also be applied before another model

w_2 : true = 4

Lasso predicts : 2.6

linear model : 3.5

Logistic Regression

x_1	x_2	y
1	0	1
2	1	0
3	-2	0
4	3	1

← binary target

- How are y and x related?

Linear model

Regression

classification (Logistic Regression)

Logistic Regression

x_1	x_2	y
1	0	1
2	1	0
3	-2	0
4	3	1

- Logistic Regression models $P(y = 1|x) = \hat{y}$ as:

$$\hat{y} = \frac{1}{1 + e^{-(w_0 + w_1 \cdot x_1 + w_2 \cdot x_2)}}$$


probability of
being positive

- OR,

$$\log \left(\frac{\hat{y}}{1 - \hat{y}} \right) = w_0 + w_1 \cdot x_1 + w_2 \cdot x_2$$

linear

where $\hat{y}$ is the predicted value of the probability of $y = 1$ given x_1 and x_2 .



Logistic Regression

x_1	x_2	y
1	0	1
2	1	0
3	-2	0
4	3	1

- Logistic Regression models $P(y = 1|x) = \hat{y}$ as:

$$\hat{y} = \frac{1}{1 + e^{-(w_0 + w_1 \cdot x_1 + w_2 \cdot x_2)}}$$

- OR,

$$\log \left(\frac{\hat{y}}{1 - \hat{y}} \right) = w_0 + \underbrace{w_1}_{\text{red circle}} \cdot x_1 + \underbrace{w_2}_{\text{red circle}} \cdot x_2$$

Note: Red handwritten annotations in the original image include a circle around w_1 , a circle around w_2 , and an arrow pointing from the word "bias" to w_0 .

where $\hat{y}$ is the predicted value of the probability of $y = 1$ given x_1 and x_2 .

Logistic Regression

$$\log \left(\frac{\hat{y}}{1 - \hat{y}} \right) = w_0 + w_1 \cdot x_1 + w_2 \cdot x_2$$

- $\left(\frac{\hat{y}}{1 - \hat{y}} \right)$ is also called odd-ratio. 
- Logistic Regression assumes that the log of the odd ratio is linear. 

linear model assumes that y is linear
=

How to find w_0, w_1, w_2 ?

- **Step 1:** Define the loss function $l(\hat{y}, y)$
- **Step 2:** Find w that minimizes the total loss function

Logistic Regression

- Define the loss function: We use the log-loss or cross-entropy loss function

$$l(\hat{y}, y) = -y \log(\hat{y}) - (1 - y) \log(1 - \hat{y})$$


- Total Loss:

$$\begin{aligned} L(w_0, w_1, w_2) = & -\log\left(\frac{1}{1 + e^{-w_0 - w_1}}\right) \\ & -\log\left(1 - \frac{1}{1 + e^{-w_0 - 2w_1 - w_2}}\right) \\ & -\log\left(1 - \frac{1}{1 + e^{-w_0 - 3w_1 + w_2}}\right) \\ & -\log\left(\frac{1}{1 + e^{-w_0 - 4w_1 - 3w_2}}\right) \end{aligned}$$


- We need to find w_0, w_1, w_2 that minimizes the total loss

square loss $(y - \hat{y})^2$


Logistic Regression



Logistic Regression




Logistic Regression




LASSO for Logistic Regression

- The idea is the same as for linear model


LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression



LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression


LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression




LASSO for Logistic Regression



LASSO for Logistic Regression



LASSO for Logistic Regression



LASSO for Logistic Regression

