

Chapter 8

Phenotypic Diversity of Colored Phytochemicals in Sorghum Accessions With Various Pericarp Pigments

Haley Davis^{*}, Xiaoyu Su^{*}, Yanting Shen^{*}, Jingwen Xu^{*}, Donghai Wang[†], J. Scott Smith[‡], Fadi Aramouni[‡] and Weiqun Wang^{*}

^{*}Department of Food Nutrition Dietetics & Health, Kansas State University, Manhattan, KS, United States, [†]Department of Biological & Agricultural Engineering, Kansas State University, Manhattan, KS, United States, [‡]Food Science Institute, Kansas State University, Manhattan, KS, United States

Chapter Outline

1 Introduction	123	6 Health Benefits of Colored Phytochemicals	127
2 Pericarp Pigments	124	7 Discussion	129
3 Phytochemicals	125	8 Conclusion	130
4 Anthocyanins	125	Acknowledgments	130
5 Carotenoids	126	References	130

1 INTRODUCTION

The philosophy that most Americans share is that eating fruits and vegetables of all different colors is important because colorful food is healthy for you. But what is unknown is that there are many other colorful foods, like sorghum, that could be beneficial to the diet as well. Sorghum is the fifth leading whole grain cereal crop in the world after wheat, maize, rice, and barley, primarily grown in tropical, subtropical, and arid regions such as Asia and Africa. In countries like the United States, Australia, and Brazil, this cereal is mainly used as animal feed. However, using sorghum in food is gaining interest in the United States due to its gluten-free and bioactive compound-rich health properties. Sorghum can be used as a substitute for conventional cereals due to its high phenolics, fiber, and carotenoids, etc. [1]. These bioactive compounds have been linked with multiple health benefits, including cholesterol-lowering, antioxidants, slow digestibility, antiinflammatory, and anticarcinogenic properties [2]. As a gluten-free product, it can be a useful grain alternative for individuals with gluten sensitivity or celiac disease. Sorghum also contains high fiber content with a lower glycemic index compared with other grains, which could help reduce obesity and type-2 diabetes. On the other hand, compared with wheat, corn, and rice, sorghum has a lower sensory quality and reduced protein digestion [3]. One of the big possibilities for sorghum is cereal products, as expanded and extruded sorghum products are very popular due to its crispness and ease of use. These products are typically made with corn, rice, or wheat in countries such as the United States and Brazil. Even though sorghum has a lower cost and is easier to produce, only recently has it been used for this purpose. But sorghum-based products are showing good acceptability and have the potential to replace traditional cereals. The antioxidant activity of sorghum flours is also high because of high levels of phenolic compounds such as luteolinidin, apeiginidin, and 3-deoxyanthocyanins [3]. This is especially true of brown tannin-sorghum, which demonstrates excellent flavor, appearance, and texture [1].

Sorghum contains the highest phytochemical contents among cereals, including anthocyanins, carotenoids, phenolic acids, and condensed tannins, etc. Sorghum kernel is divided into three components: seed coat, endosperm, and embryo or germ. The seed coat is divided even further into the pericarp and testa [4]. These phytochemicals are usually contained in the pericarp and testa. Sorghum also typically contains provitamin A carotenoids, but the association of sorghum intake and vitamin A deficiency is shown to be contradictory. Studies conducted in Africa indicated various types of malnutrition including vitamin A deficiency when sorghum was the principle grain in the diet [5]. However, other studies have suggested that the sorghum consumed in Africa and Asia are critical sources of dietary carotenoids that might provide the needed

provitamin A [6]. These inconsistencies might be due to the different varieties of sorghum consumed [3]. Various sorghum varieties differ in the color of the pericarp and/or endosperm at a wide range from red to white. The pigmented phytochemicals such as anthocyanins and carotenoids, etc. contribute to these colors, which are considered to be a dependable indicator of sorghum varieties [3,7].

2 PERICARP PIGMENTS

Sorghum grains come in a wide variety of colors such as red, yellow, white, and black. These colors typically come from the pericarp, endosperm, and testa. The pericarp is the outer layers of the seed coat of the sorghum kernel. The phenotypic pigments of sorghum pericarp and endosperm are determined by genotypic factors [3,4]. Genetic factors control pericarp color, pericarp thickness, and secondary plant color [8,9]. The pericarp colors are controlled by the R and Y genes. The white color occurs when Y is homozygous recessive; yellow occurs when R and Y are homozygous recessive and homozygous dominant; and red occurs when both R and Y are dominant. Black sorghum is a special red sorghum that turns black in the presence of sunlight [8]. Normally, the endosperm is yellow or white, and the pericarp is usually red or black. The Z gene controls pericarp thickness, which doesn't affect the flavonoids levels [9].

The wide array of phenolic compounds is concentrated in the pericarp. Phenolic compounds are present in all sorghums; their composition, levels, and types are affected by the genotype. The phenolic compounds include phenolic acids, flavonoids, and condensed tannins. Sorghum flavonoids are 3-deoxyanthocyanins, orange luteolinidin, and yellow apigeninidin. The 3-deoxyanthocyanins are more stable than common anthocyanins, making them a potential natural food colorant [8]. Fig. 1 shows the chemical structures of the common anthocyanins presented in sorghum. Sorghums with red/purple secondary plant colors and sorghum with black pericarp have the highest levels of 3-deoxyanthocyanins, whereas red sorghum with tan secondary plant color have high apigenin and luteolin levels with red/purple secondary plant color [8].

Another important genotypic factor affecting pericarp color is other colored phytochemicals such as carotenoids. Carotenoids are one of the colored phytochemicals that may account for phenotypic color of sorghum pericarp [7]. These carotenoids include lutein, zeaxanthin, α -carotene, and β -carotene as shown in Fig. 1. Yellow pericarp/yellow endosperm sorghum accessions show the highest contents of carotenoids, followed by brown pericarp/yellow endosperm and white pericarp/yellow endosperm. The lowest carotenoids and provitamin A were found in the accessions with white pericarp/white endosperm [7]. It appears that the phenotypic diversity of sorghum pericarp colors is associated with the content of carotenoids and provitamin A, indicating a different impact of sorghum varieties on vitamin A deficiency and suggesting a possible prevention of vitamin A deficiency by breeding sorghum varieties with specific pericarp pigments [7]. This potential health benefits, as well as many others, are due in part to the various phytochemicals found in sorghum.




FIG. 1 Chemical structures of the common anthocyanins (left) and carotenoids (right) found in sorghum.

3 PHYTOCHEMICALS

Phytochemicals are biologically active compounds naturally found in plants. Sorghum contains various phytochemicals including phenolics, carotenoids, sterols, and policosanols, etc. Plant sterols and policosanols are mostly components of wax and oils, but phenolics are used as a natural defense of plants against pests and diseases [2]. Phenols, also known as polyphenols, in sorghum are classified into two categories: phenolic acids and flavonoids. Anthocyanins and condensed tannins fall under the flavonoid category. Only varieties with a pigmented testa have condensed tannins [10]. Carotenoids are also categorized as phytochemicals. Studies have indicated the potential health benefit of using sorghum for food consumption due to its bioactive compounds. Sorghum has the highest content of phenolic compounds among cereals [1]. Evidence suggests that sorghum consumption, for example, may reduce the risk of certain types of cancer in humans compared with other cereals, which could be due the high concentration of phytochemicals in the sorghum [11].

Phenolic acids are located in the pericarp, testa, and endosperm. They are generally classified into two types: free and bound phenolic acids. Free phenolic acids are found in the outer layers of the kernel, whereas the bound phenolic acids are associated within the cell walls. Free phenolic acids are soluble in nature and bound acids are insoluble, covalently bound to structural components of cells such as cellulose, hemicellulose, lignin, and pectin [12]. Phenolic compounds are created from the secondary metabolism of plants when they are subjected to stressful conditions such as infections and mechanical injuries. These compounds act as a natural defense mechanism in the plants by protecting them from pests and disease [13].

Tannins, also known as proanthocyanidins, are large molecular weight polyphenols present in sorghum testa. The testa is located between the pericarp and endosperm. Sorghum accessions with pigmented testa are classified as type II and III sorghums. These sorghums have dominant B1 and B2 genes. The B1 and B2 genes control the presence or absence of the pigmented testa layer, and both genes must be dominant for a pigmented testa to develop. When the spreader S gene is dominant concurrently with the dominant B1 and B2 genes, pericarp color becomes phenotypically brown [11]. Tannins bind to and reduce digestibility of various food/feed nutrients, such as proteins, carbohydrates, and other nutrients, thus limiting nutritional value and decreasing digestibility and negatively affecting productivity of livestock. This aspect of tannins has resulted in decades of breeding efforts to eliminate tannins from sorghum in the United States, whereas in other parts of the world where pest and diseases are common, tannin sorghums are still grown due to them being more tolerant of these conditions than nontannin varieties [11]. Also, due to tannin's ability to bind to free radicals, sorghums containing tannins have higher antioxidant capacity than sorghums that do not contain tannins [13].

Then there are the most-known phytochemicals found in sorghum, anthocyanins and carotenoids. These phytochemicals provide many of the colors found in sorghum's several varieties, but the most important aspects are the potential health benefits they can offer. To understand how these two phytochemicals contribute to numerous health benefits, it is imperative to know their makeup and overall function in sorghum. Therefore, the following sections go further into the specifics of these important phytochemicals and the roles they play in prevention of certain diseases and cancers.

4 ANTHOCYANINS

Anthocyanins are members of the flavonoid group of phytochemicals and are the major class of flavonoids studied in sorghum. There are more than 5000 flavonoids that include the groups anthocyanins, flavonols, flavones, flavanones, flavan-3-ols, and isoflavones [2]. These phytochemicals are also referred to as bioflavonoids due to their many roles in human health maintenance. Anthocyanins are one of the most important water-soluble plant pigments and are predominantly in grapes, wines, fruits, vegetables, nuts, and purple-fleshed sweet potatoes [14]. In general, this class of compounds contributes the blues, purples, and reds in plants. Anthocyanins are typically ingested as part of a complex mixture of flavonoids [15], and their intake has been projected to be nine times higher than any other dietary flavonoids [16]. They are synthesized by the flavonoid branch of the phenylpropanoid pathway through secondary metabolism in the plants [3]. Plants typically produce anthocyanins as a protective mechanism against environmental stress such as cold temperatures, drought, and ultraviolet (UV) light [16]. Among the more than 600 types of anthocyanins, the majority of anthocyanins found in food items usually consists of six anthocyanidins, including cyanidin, delphinidin, malvidin, pelargonidin, petunidin, and peonidin [17]. These exist in nature as mostly glycoside anthocyanins. Most studies related to anthocyanins are based on the compounds from fruit and vegetables. But due to a rising demand for cost-effective sources of natural, stable pigments, there is an increased interest in alternative sources of anthocyanins, that is, sources like cereal, corn, rice, and wheat. Pigmented sorghums are also potentially rich sources of unique anthocyanins [18].

Sorghum anthocyanins are called 3-deoxyanthocyanins, which are comprised of luteolinidin and apigeninidin, including their methoxylated derivatives of 5-methoxyluteolinidin and 7-methoxyapigeninidin [14]. The anthocyanins in sorghum are unique because they do not contain the hydroxyl group in the 3-position of the C-ring (Fig. 1). This feature




FIG. 2 The contents of total anthocyanins in four sorghum accessions with various pigments of pericarp: white (PI656079), yellow (PI229838), brown (PI221723), and red (PI297139). Data represent mean values \pm SD. Bars marked with different letters are statistically significant at $P < .05$, $n = 2$.

increases their stability at high pH compared with other anthocyanins, indicating these compounds can be used as potential natural food colorants. These anthocyanins are also called phytoalexins because they are produced as a response to mold or bacterial infection [10]. These compounds are more prevalent in purple-pigmented plants than in tan-pigmented plant sorghums. The two most common sorghum 3-deoxyanthocyanins are the yellow apigeninidin and the orange luteolinidin. These 3-deoxyanthocyanidins represent 36% to 50% of the total anthocyanin content in black and brown sorghum brans [3]. Sorghum with a black pericarp, which is concentrated in the bran, have the highest levels of 3-deoxyanthocyanins. The black sorghum bran has twice the level of anthocyanins as red and brown sorghum brans [10].

The profile of sorghum anthocyanins was recently identified and quantified in the selected 25 sorghum accessions with various phenotypic pericarp pigments [3]. As shown in Fig. 2, the high levels of total anthocyanins were found in red pericarp and then followed by brown pericarp and yellow pericarp. Sorghum accessions with white pericarp generally contained the least to undetectable amount. Although anthocyanins appeared to be associated with the pericarp color, however, a distinguishable diversity of anthocyanin contents was present among and between the phenotypic pericarp colors, suggesting other colorful phytochemicals such as carotenoids might have also contributed [3].

5 CAROTENOIDS

Carotenoids are a family of compounds of more than 600 fat-soluble pigments. They function in plants and photosynthetic bacteria as accessory pigments in photosynthesis. Carotenoids are responsible for many different colors in nature such as the red, yellow, and orange color. They play an important role in photosynthesis as light-harvesting pigments and protecting molecules from the formation of singlet oxygen. The most prevalent carotenoids are α -carotene, β -carotene, lycopene, lutein, zeaxanthin, and β -cryptoxanthin. The known carotenoid function in humans is vitamin A activity, but only provitamin A carotenoids have this function [7]. Carotenoids can be classified into two groups based on the functional groups. The first are xanthophylls, which has oxygen as the functional group, which includes lutein, zeaxanthin, and β -cryptoxanthin. The other group is carotene, which contains only a hydrocarbon chain without any function group. This group includes α -carotene, β -carotene, and lycopene [19]. Carotenoids typically contain 40 carbon atoms with an extensive system of conjugated double bonds. Carotenoids have one or two cyclic structures at the end of the conjugated chains. One of the exceptions to this is lycopene, an acyclic carotenoid with 11 conjugated double bonds with no retinol structure or no vitamin A activity [20]. Naturally occurring carotenoids are in the all-trans-configuration. The configuration of double bonds is either E or Z, which corresponds to trans or cis. Cis isomers are usually less thermodynamically stable than the trans isomers. Thus, trans forms of carotenoids are more prevalent in nature. Yet cis isomers are more soluble than trans and therefore are more readily absorbed and transported within cellular compartments [21]. Out of the six

common carotenoids, three of them, α -carotene, β -carotene, and β -cryptoxanthin, are precursors of vitamin A, meaning they can be converted into retinol, and thus are referred to as provitamin A carotenoids. The other three, lycopene, lutein, and zeaxanthin, have no vitamin A activity and are thus referred to as nonprovitamin A carotenoids [7]. Vitamin A is a family of molecules that contain a 20-carbon structure with a β -ionone ring and tetraene side chain with either a hydroxyl group (retinol), aldehyde group (retinal), carboxylic acid group (retinoic acid), or ester group (retinyl ester) at carbon-15 [15]. To convert provitamin A carotenoids to vitamin A involves cleavage of the central double bond, for instance, cleaving of the central double bond of β -carotene molecule yields two 20-carbon molecules of vitamin A [7].

Intestinal absorption of dietary carotenoids is enabled by the formation of bile acid micelles. Like other nonpolar lipids, the hydrocarbon backbone of carotenoids makes them insoluble in water and thus must be solubilized within micelles in the gastrointestinal tract to allow for absorption. Absorption requires that the carotenoids first be freed from the food matrix and solubilized in oil droplets. This transfer of carotenoids and other fat-soluble compounds from the food matrix into micelles during digestion is referred to as bioaccessibility. It is a necessary preliminary step so that the compounds become accessible for uptake by the intestinal mucosa. Carotenoids are then transferred to bile salt micelles, which are generated during the digestion of fat-soluble compounds including triglyceride, phospholipids, and cholesterol ester. Once inside mucosal cells, carotenoids must also be incorporated into chylomicrons and released into the lymphatics. The carotenoids and other fat-soluble compounds interact with brush border proteins for transfer to the cell interior. The bile salts in the micelles dissociate and are reabsorbed in the ileum [22,23]. Absorption of carotenoids from plant foods is relatively inefficient. This is due to numerous factors such as the physicochemical characteristics of carotenoids, physical and chemical barriers within the plant tissue, and processing and preparation of foods. These impede the solubilization of carotenoids emulsified in lipid droplets [22].

Foods rich in carotenoids are considered beneficial in the prevention of a variety of chronic diseases, including certain types of cancer and eye diseases such as macular degeneration. The most widely studied carotenoid is β -carotene, which is also the major carotenoid in our diet and in human blood and tissues. The major sources of dietary β -carotene include green leafy vegetables as well as orange and yellow fruits and vegetables. But there are factors besides just the type of food that are important in the bioavailability of β -carotene. These factors include cooking, chopping, and the presence of dietary fat, all of which improve bioavailability. Thus, the bioavailability of β -carotene from green leafy vegetables such as spinach is thought to be low [23]. Another important carotene is lycopene, which is derived from tomatoes and tomato products predominately. Lycopene bioavailability is greater from tomato paste compared with fresh tomatoes. Other dietary sources include dried apricots, guava, watermelon, papaya, and pink grapefruit. Dietary sources of lutein and zeaxanthin include spinach, kale, broccoli, peas, and brussels sprouts. Another good source is egg yolk due to the high bioavailability. There are no adverse effects of consuming large amounts of β -carotene or other carotenoids from foods except for carotenodermia. Carotenodermia is a harmless, biological effect of high carotenoid intake characterized by a yellowish discoloration of the skin due to an elevation of carotene concentrations [23]. When it comes to sorghum, the level of carotenoids is related to the pericarp pigment. The highest content of total carotenoids is found to be in sorghum accessions with yellow pericarp, although the highest β -carotene content is found in accessions with brown or yellow pericarp. Five carotenoids (α -carotene, β -carotene, lutein, zeaxanthin, and β -cryptoxanthin) were recently identified and quantitated, in the selected sorghum accessions with various pericarp pigments [7]. As shown in Fig. 3, the highest content of total carotenoids was found in the sorghum accessions with yellow pericarp/yellow endosperm, and followed by brown pericarp/yellow endosperm. The lowest carotenoids were found in the accessions with white pericarp/white endosperm and/or white pericarp/yellow endosperm. The results suggested a possible prevention of vitamin A deficiency by consuming selected sorghum varieties with yellow pericarp pigments [7].

6 HEALTH BENEFITS OF COLORED PHYTOCHEMICALS

One of the most studied aspects of phytochemicals in general is the antioxidant properties. Antioxidants come in many forms and are found in numerous food products. They are the primary phytochemical contributors to the prevention of certain diseases and cancers. The definition of antioxidants is based on several criteria. First, the substance needs to be found in the human diet; two, the content of the substances must have been measured in foods; and three, the substances in humans has to decrease the adverse effects of reactive species, such as reactive oxygen and nitrogen species in vivo [24]. Free radicals can be defined as molecules containing one or more unpaired electrons. They are mostly derived from oxygen, known as reactive oxygen species (ROS), or from nitrogen, known as reactive nitrogen species (RNS). Both ROS and RNS play a role as beneficial and harmful species. Overproduction of these species either by external or internal factors can result in a state called oxidative stress. This can cause damage to cell structures, including DNA, proteins, and membrane lipids. Antioxidants act on the biological systems to balance the toxic action of free radicals [25]. ROS include a group of oxidants




FIG. 3 The contents of total carotenoids in selected sorghum accessions with various pigments of pericarp and endosperm: WW (*white pericarp/white endosperm*), WY (*white pericarp/yellow endosperm*), BY (*brown pericarp/yellow endosperm*), and YY (*yellow pericarp/yellow endosperm*). Data represent mean values \pm SD. Bars marked with different letters are statistically significant at $P < .05$, $n=3$.

that consist of free radical and non-free radical oxygen intermediates. These include hydrogen peroxide (H_2O_2), superoxide (O_2^\bullet), singlet oxygen (1O_2), and the hydroxyl radical ($\bullet OH$) [26]. Another type of reactive radical derived from oxygen and can be formed in living systems is peroxy radicals (ROO^\bullet), the simplest form of the radical is perhydroxyl radical (HOO^\bullet). Hydroxyl radicals have a very high reactivity and a very short lifetime. This makes these radicals responsible for the damage to DNA, protein, and membranes and, due to their short lifetime, these radicals react close to the site of formation. Nitric oxide (NO) is a radical derived from nitrogen that also plays a role in living systems. It is a small molecule that contains an unpaired electron and is generated by nitric oxide synthase. NO acts as an important oxidative biological signaling molecule in a variety of physiological processes, such as neurotransmission, blood pressure regulation, smooth muscle relaxation, defense mechanisms, and immune regulation [21].

Anthocyanins have been extensively studied in fruits and vegetables due to their antioxidant properties. However, there is limited data on the types and levels of anthocyanins in cereals due to the belief that cereals are not a commercially significant source. In actuality, sorghum fractions have a higher anthocyanin activity *in vitro* than other cereals and fruits, and may have similar health benefits usually associated with fruits [11]. Various beneficial properties reported for anthocyanins are largely attributed to their antioxidant properties. Benefits include vasoprotective and antiinflammatory properties, anticancer, and chemoprotective properties. However, antioxidant activity alone is not a good predictor of potential biological benefits because the chemical structure of anthocyanins, such as the position, number, and types of substitutions, play an important role in the biological activity exerted. For instance, it has been reported that substitution on the B-ring of the anthocyanidins affected their ability to suppress carcinogenesis [16,18]. In both *in vitro* and *in vivo* research trails, anthocyanins have shown the ability to reduce cancer cell proliferation and inhibit tumor formation [14,15]. Anthocyanins' ability to interfere with the process of carcinogenesis is possibly linked to multiple mechanisms of action, which include inhibition of cyclooxygenase enzymes and antioxidant potential. Research has shown that significant anthocyanin concentration is effective against various stages of carcinogenesis, but the individual role of anthocyanins is not yet determined. This is due in part to the fact that anthocyanins were easily degraded during bioassays if separated from the stabilizing cofactors such as other phenolic compounds [15]. Typically, anthocyanins interact with other flavonoid components or anthocyanins to provide full potency. Anthocyanins almost never act independently when it comes to biological activity in the human body. Within a plant, phytochemical interaction is key, and thus any plant with anthocyanins includes a complex phytochemical mixture. Anthocyanins and other flavonoids are secondary products produced as defensive and protective agents that are used together to combat diseases and pests. This multiplicity of bioactive phytochemicals is also useful in humans and animals who can benefit from this interaction of flavonoids [15]. One of the larger roles of

anthocyanins, as well as other dietary flavonoids, is protection against cardiovascular disease (CVD), which is the most common cause of mortality among men and women. Studies have shown the beneficial effect of flavonoids, including anthocyanins, on the biomarkers of CVD risk which are NO, inflammation, and endothelial dysfunction. Inflammation is defined by heat, redness, and swelling, which plays a large role in the development of CVD. For anthocyanins, this protection is strongly linked to the oxidative stress protection that anthocyanins provide to endothelial cells. Also, there are several mechanisms that have been proposed to explain the *in vivo* antiinflammatory action of flavonoids such as anthocyanins. Anthocyanins isolates and anthocyanin-rich mixtures of flavonoids may possibly provide protection from DNA cleavage, estrogenic activity, enzyme inhibition, antiinflammatory activity, lipid peroxidation, and membrane strengthening [16].

Carotenoids are known to have vitamin A activity, but they are also thought to have possible antioxidant activity. They have been associated with various health effects such as decreased risk of macular degeneration and cataracts, and decreased risk of some cancers and cardiovascular events [23]. These beneficial effects are thought to be due to their role as antioxidants, which are a result of the conjugated double bond structures ability to delocalize unpaired electrons. This is the feature responsible for carotenoids' exceptional ability to quench singlet oxygen and terminate free radicals that have formed in tissues [21]. The presence of these numerous conjugated double bonds and functional end groups are crucial for the function of carotenoids, which is essential for light absorption in photosynthetic organisms. Carotenoids absorb radiant energy to chlorophyll molecules in a light-harvesting function during photosynthesis. This dissipates excess energy through the xanthophyll cycle and quenches excited-state chlorophylls directly. One of the factors responsible for oxidative damage to plant leaves is singlet oxygen, and carotenoids are the first line of defense by quenching them. This can be done physically by energy transfer or chemically by direct reaction with the radical. In plants, organelles such as chloroplast and peroxisome have an intense rate of electron flow that generates ROS as byproducts during respiration and photosynthesis. But they also host antioxidants to harmlessly dissipate ROS such as carotenoids [26]. Additionally, lutein and zeaxanthin may protect from eye disease due to their ability to absorb damaging blue light that enters the eye. The function of carotenoids is determined by their molecular properties such as size, structure, presence of functional group, and potential interaction with other carotenoids. Most importantly is the double bond conjugation linked to the antioxidant properties [21,23,26]. Because carotenoids are a part of a hydrophobic group of antioxidants, the major mechanism of this action is located within the lipid membranes. They are also associated with proteins or lipoprotein structures. This means that the local environment of carotenoids affects its properties and vice versa. The structural differences of carotenoids mean different types of interaction with the membrane lipids thus affecting membrane fluidity and thermostability in different ways [21]. One of the main possible health benefits of the antioxidant function of carotenoids is cancer prevention. One way is that carotenoids are shown to inhibit the growth of tumor cells by interfering at different phases of the cell cycle. Deregulated cell cycles are one of the major hallmarks of cancer cells, which means the cells lose their ability to regulate the cell cycle and control the rate of proliferation. Carotenoids modulate the cell cycle arrest by multiple mechanisms in cancer cells. One such carotenoid is lycopene, which has been reported to significantly suppress the proliferation of prostate cancer cells. Lycopene has also shown to produce an antiproliferative effect in human colon cancer and breast cancer cells [26]. Another hallmark of cancer cells is a defect in the apoptosis mechanism. Apoptosis is a form of programmed cell death. This defect is characterized by morphological changes including membrane blebbing and formation of apoptotic bodies. Cancer cells resist apoptosis by the expression of antiapoptotic proteins. Carotenoids possess chemopreventive activity by reducing cancer incidence through apoptosis. Another way cancer occurs, as well as atherosclerosis and diabetic retinopathy, is through the formation of tumors. Angiogenesis is fundamental for the growth and metastasis of tumors. Carotenoids, such as β -carotene and lycopene, have been shown to have excellent antiangiogenic properties [26]. Specifically, reports have shown the higher the intake of carotenoids, the lower the risk of lung cancer, the first leading cause of deaths worldwide. Also, high levels of carotenoids, such as α -carotene, trans- β -carotene, lutein, and zeaxanthin, have been reported to reduce the risk of prostate cancer, the second-most common cancer in men [26].

7 DISCUSSION

Anthocyanins and carotenoids were the two major phytochemicals addressed in this chapter because they are major components in the color and health benefits of sorghum. However, there are a few other phytochemicals mentioned as well that can be linked to the color and possible health benefits. For instance, condensed tannins, as previously stated, are polyphenols present in sorghum with pigment testas. Sorghum varieties with pigmented testas are classified as type II and type III sorghum. The difference between the two types is the S gene. The S gene, or spreader gene, spreads the pigment of the

testa to the pericarp. This means that if the gene is regressive, like in the type II sorghum, the pigment won't spread to the pericarp and thus the sorghum has a white appearance yet still containing polyphenols like tannins. If the gene is dominant, like in type III, the pigment will spread, which gives a white or red pericarp color, but an overall brownish red/white sorghum [4]. Like other polyphenols, tannins also show antioxidant properties as well as antiinflammatory, antihypertensive, anticancer, and anticarcinogenic properties. There has been a wide range of research conducted regarding the antioxidant properties of tannins but little has been done with condensed tannins in sorghum accessions with various testa pigments.

8 CONCLUSION

Sorghum is the fifth leading whole grain cereal crop in the world and is found primarily in tropical, subtropical, and arid regions such as Asia and Africa. In countries such as the United States and Brazil, sorghum is mainly used for animal feed. But sorghum use in food is on rise due to its functional and health properties. Most research on sorghum has been performed on its high phytochemical contents, including phenolic anthocyanins, polyphenolic tannins, and carotenoids that are mostly presented in the pericarp, endosperm, and testa. The major colored phytochemicals in sorghum are anthocyanins and carotenoids. The contents of anthocyanins appeared to be associated with the sorghum pericarp color, but a diversity of anthocyanin contents was present among and between the phenotypic pericarp colors. The content of total carotenoids was correlated with sorghum pericarp and endosperm pigments. Both anthocyanins and carotenoids are believed to be bioactive compounds and thus have potential health benefits including antioxidant, anticancer, vasoprotective, and antiinflammatory properties. There is also potential for more research to be performed on the other phytochemicals, such as condensed tannins, that contribute to sorghum color and provide potential health benefits. Along with many other functional and healthful properties, sorghum shows promise to be potentially functional in health benefit-promoted food products.

ACKNOWLEDGMENTS

This study was supported in part by a USDA Cooperative Project KS511-1001903 from the Agricultural Experiment Station, Kansas State University (Contribution #18-186-B from the Kansas Agricultural Experiment Station).

REFERENCES

- [1] Anunciacao P, Cardoso L, Gomes J, Lucia C, Carvalho C, Galdeano M, et al. Comparing sorghum and wheat whole grain breakfast cereals: sensorial acceptance and bioactive compound content. *Food Chem* 2017;221:984–9.
- [2] Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacol Ther* 2001;90:157–77.
- [3] Su X, Rhodes D, Xu J, Chen X, Davis H, Wang D, et al. Phenotypic diversity of anthocyanins in sorghum accessions with various pericarp pigments. *J Nutr Food Sci* 2017;7(4):1000610.
- [4] Salunkhe DK, Chavan JK, Kadam SS. Sorghum tannins. In: Salunkhe DK, editor. *Dietary tannins: consequences and remedies*. Boca Raton, FL: CRC Press; 1990. p. 31–44.
- [5] Adegboye OR, Smith C, Anang D, Musa H. Comparing and contrasting three cultural food customs from Nigeria and analyzing the nutrient content of diets from these cultures with the aim of proffering nutritional intervention. *Crit Rev Food Sci Nutr* 2016;56(15):2483–94.
- [6] Kean EG, Bordenave N, Ejeta G, Hamaker BR, Ferruzzi MG. Carotenoid bioaccessibility from whole grain and decorticated yellow endosperm sorghum porridge. *J Cereal Sci* 2011;54(3):450–9.
- [7] Shen Y, Su X, Rhodes D, Herald T, Xu J, Chen X, et al. The pigments of sorghum pericarp are associated with the contents of carotenoids and pro-vitamin A. *Int J Food Nutr Sci* 2017;6(3):48–56.
- [8] Dykes L, Seitz ML, Rooney LW, Rooney WL. Flavonoid composition of red sorghum genotypes. *Food Chem* 2009;116:313–7.
- [9] Dykes L, Peterson CG, Rooney LW, Rooney WL. Flavonoid composition of lemon-yellow sorghum genotypes. *Food Chem* 2010;128:173–9.
- [10] Dykes L, Rooney L. Sorghum and millet phenols and antioxidants. *J Cereal Sci* 2006;44:236–51.
- [11] Awika J, Rooney L. Sorghum phytochemicals and their potential impact on human health. *Phytochemistry* 2004;65:1199–221.
- [12] Duyi S, Das BA, Deka CS. Pigmented rice a potential source of bioactive compounds: a review. *Int J Food Sci Technol* 2017;52:1073–81.
- [13] Oliveira K, Queiroz V, Carlos L, Caradoso L, Pinheiro-Sant'Ana H, Anunciacao P, et al. Effect of the storage time and temperature on phenolic compounds of sorghum grain and flour. *Food Chem* 2017;216:390–8.
- [14] Lim S, Xu J, Kim J, Chen TY, Su X, Standard J, et al. Role of anthocyanin-enriched purple-fleshed sweetpotato P40 in colorectal cancer prevention. *Mol Nutr Food Res* 2013;57:1908–17.
- [15] Lilia M. Anthocyanins and human health: an in vitro investigative approach. *J Biomed Biotechnol* 2004;5:306–13.
- [16] Awika JM, Rooney LW, Waniska RD. Properties of 3-deoxyanthocyanins from sorghum. *J Agric Food Chem* 2004;52(14):4388–94.
- [17] Xu J, Su X, Lim S, Griffin J, Carey E, Katz B, et al. Characterization and stability of anthocyanins in purple-fleshed sweet potato P40. *Food Chem* 2015;186(11):90–6.

- [18] Wallace T. Anthocyanins in cardiovascular disease. *Adv Nutr* 2011;2:1–7.
- [19] Saini KR, Nile HS, Park WS. Carotenoids from fruits and vegetables: chemistry, analysis, occurrence, bioavailability and biological activities. *Food Res Int* 2015;76:735–50.
- [20] Institute of Medicine (US) Panel on Micronutrients. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. USA: National Academies Press; 2001. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK222318/>.
- [21] Jorova K, Valko M. Health protective effects of carotenoids and their interactions with other biological antioxidants. *Eur J Med Chem* 2013;70:102–10.
- [22] Kopec RE, Failla ML. Recent advances in the bioaccessibility and bioavailability for carotenoids and effects of other dietary lipophiles. *J Food Compos Anal* 2017; <https://doi.org/10.1016/j.jfca.2017.06.008>.
- [23] Krinsky IN, Johnson JE. Carotenoid actions and their relation to health and disease. *Mol Asp Med* 2005;26:459–516.
- [24] Institute of Medicine (US) Panel on Dietary Antioxidants and Related Compounds. Dietary reference intake for vitamin C, vitamin E, selenium, and carotenoids. USA: National Academy Press; 2000. Available from: <http://www.nap.edu/catalog/9810.html>.
- [25] Perea OC, Yen MG. Functional properties of carotenoids and human health. *Int J Food Prop* 2007;10:201–30.
- [26] Niranjana R, Gayathri R, Mol NS, Sugawara T, Hirata T, Miyashita K, et al. Carotenoids modulate the hallmarks of cancer cells. *J Funct Foods* 2015;18:968–85.