# Lectures on Quantum Computing and Completely Positive Maps

A formalisation in Isabelle/HOL of Vern Paulsen's Houston 2011 lecture notes

Dustin Bryant

formal statements and proofs: the theories listed in the table of contents

October 20, 2026

**Abstract**

This document is generated by Isabelle from the project theories. Every definition and theorem shown here has been checked by Isabelle/HOL (proofs are folded). The commentary restates Paulsen's lecture notes lecture by lecture (Day 1 = 22 August 2011, Day 34 = 18 November 2011): definitions and results in the notes' own language, each followed by the formal statement. The informal proofs of the notes are omitted; the formal proofs replace them. Where the formalisation deviates from the notes, the commentary says so. The companion file `NOTES_AUDIT.md` lists, for every lecture, what is formalised and what is not. Part VI goes beyond the notes: the functional calculus, phase estimation and the HHL algorithm, built on the same library.

## Contents

**1 Reading guide** **2**

**2 Conventions** **4**

**I Lectures 1–10: basic mathematics and the postulates of quantum mechanics** **4**

**3 Preliminaries: linear-space isomorphisms** **5**

**4 Day 1 (22 August 2011): the inner product** **6**
4.1 The Hilbert–Schmidt inner product on matrices . . . 6
4.2 Properties of the inner product . . . 7

**5 Day 2 (24 August 2011): bases, matrix units, trace, projections** **7**
5.1 The trace . . . 9
5.2 Matrix-unit algebra and cyclicity of the trace . . . 10
5.3 Orthogonal projections and Parseval's identity . . . 11

**6 Unitary algebra** **14**

**7 Tensor product infrastructure** **15**
7.1 Tensor of Identities . . . 15
7.2 Tensor Adjoint . . . 15
7.3 Mixed-Product Property . . . 15
7.4 Corollaries of the Mixed-Product Property . . . 15
7.5 Tensor Unitarity . . . 15

**8 Day 3 (31 August 2011): unitary matrices, Householder unitaries, Schur's theorem** **16**
8.1 Unitary matrices . . . 16
8.2 Householder unitaries . . . 17

**9 Day 4 (2 September 2011): Hermitian matrices; positive semidefinite and positive definite matrices** **23**
9.1 Hermitian matrices . . . 23
9.2 The spectral theorem for Hermitian matrices . . . 26
9.3 Positive semidefinite and positive definite matrices . . . 27
9.4 The positive square root . . . 30

**10 Day 5 (7 September 2011): tests for positivity; direct sums and partitioned matrices** **30**
10.1 Practical tests for positivity . . . 30
10.2 Direct sums of vector spaces and partitioned matrices . . . 31

**11 Day 6 (9 September 2011): tensor products** **32**

**12 Day 7 (12 September 2011): tensor products of Hilbert spaces; the postulates** **33**
12.1 Postulates of quantum mechanics . . . 34

**13 Day 8 (14 September 2011): the quantum game; projective measurements and expected values** **35**
13.1 Distinguishing states . . . 35
13.2 Projective measurements, expected values and self-adjoint operators . . . 37

**14 Day 9 (16 September 2011): positive operator-valued measures; composite systems; entanglement** **38**
14.1 Positive operator-valued measures . . . 38
14.2 Composite systems . . . 39
14.3 Entanglement . . . 40

**15 Day 10 (19 September 2011): superdense coding; classical and quantum gates** **40**
15.1 One-qubit gates . . . 40
15.2 Two- and three-qubit gates . . . 41
15.3 Superdense coding . . . 42
15.4 Day 2: the projection does not depend on the orthonormal basis . . . 43
15.5 Day 6: coordinates in a tensor product . . . 43
15.6 Day 8: non-orthogonal states cannot be distinguished . . . 45
15.7 Day 9: measuring one half of a Bell pair . . . 45
15.8 Day 11: SWAP from three CNOTs . . . 46
15.9 Day 5: a basis of $V \oplus W$ (D5-5) . . . 47

**16 Day 5: Cholesky's algorithm on the notes' two examples (D5-3)** **47**
16.1 One step of Cholesky's algorithm . . . 47
16.2 Small matrices entry by entry . . . 48
16.3 Example 1: a matrix that is not positive . . . 48
16.4 Example 2: a positive matrix, in two steps . . . 49

**17 Day 6: the Schmidt rank (D6-4)** **50**

**18 Days 10, 11 and 15: superdense coding, the CCNOT table, the partial trace over $A$** **51**
18.1 Superdense coding (D10-1) . . . 51
18.2 The CCNOT truth table (D11-2) . . . 52
18.3 The partial trace over $A$ (D15-6) . . . 53

**19 Days 4–29: the converse Hermitian tests, composite ensembles, Choi's four conditions, codes** **54**
19.1 Day 4: the converse directions for Hermitian matrices (D4-1) . . . 54
19.2 Choi's theorem with (2) isolated (D19-1) . . . 55
19.3 Composite ensembles (D14-5) . . . 55
19.4 Hamming distance and subspace codes (D29-4) . . . 56

**II Lectures 11–22 and 29–34: quantum computing, density matrices, completely positive maps, error correction** **57**

**20 Day 11 (21 September 2011): gates revisited, circuit diagrams, no-cloning** **57**
20.1 Cloning and no-cloning . . . 58

**21 Day 12 (23 September 2011): quantum parallelism** **59**

**22 Day 13 (26 September 2011): ensembles and density matrices** **59**
22.1 Ensembles . . . 59
22.2 Von Neumann's density matrix . . . 60
22.3 Probabilities via the trace . . . 61

**23 Day 14 (28 September 2011): density matrices, pure states, composite ensembles** **62**
23.1 Pure states . . . 63
23.2 Composite ensembles . . . 64
23.3 Partial traces: motivation . . . 64

**24 Day 15 (30 September 2011): traces, tensor products of operators, partial traces** **64**
24.1 The isomorphism $L(\mathcal{H}_A) \otimes L(\mathcal{H}_B) \cong L(\mathcal{H}_A \otimes \mathcal{H}_B)$ . . . 65
24.2 Partial traces . . . 67
24.3 Partial traces and measurements . . . 69

**25 Day 16 (3 October 2011): partial traces in block form; the postulates for density matrices** **69**

**26 Day 17 (5 October 2011): measurement maps, quantum noise, completely positive maps** **70**
26.1 Measurement maps . . . 70
26.2 Kraus maps, positive maps, complete positivity . . . 71

**27 Day 18 (7 October 2011): the theory of completely positive maps** **75**

**28 Day 19 (10 October 2011): Choi's theorem** **76**
28.1 From a positive Choi matrix to a Kraus representation . . . 77

**29 Day 20 (12 October 2011): positive functionals, adjoints, Choi rank** **80**
29.1 The adjoint of a Kraus map . . . 80
29.2 Complete positivity and the UCP/CPTP duality . . . 82
29.3 Positive functionals are completely positive . . . 82
29.4 Choi rank . . . 83

**30 Day 21 (14 October 2011): the Kraus rank bound; unitary freedom of representations** **83**

**31 Day 22 (17 October 2011): minimal Kraus representations, convex sets, UCP and CPTP** **84**
31.1 Convex sets and extreme points . . . 85
31.2 The convex sets UCP and CPTP . . . 85

**32 Day 29 (7 November 2011): classical error detection and correction** **86**
32.1 Binary strings, Hamming weight and distance . . . 86
32.2 Code distance . . . 87
32.3 The parity-check and repetition codes . . . 87

**33 Day 30 (9 November 2011): error probabilities; the three-qubit bit-flip code** **88**
33.1 The Pauli matrices . . . 89
33.2 Pauli magic (Day 32) . . . 90
33.3 The bit-flip channel (Day 31) . . . 90

**34 Day 31 (11 November 2011): the bit-flip code as a channel; fidelity** **91**
34.1 Fidelity . . . 91

**35 Day 32 (14 November 2011): the phase-flip code, the Shor code, Pauli magic** **92**
35.1 The phase-flip code . . . 92
35.2 Projections . . . 94
35.3 More on the repetition code . . . 94
35.4 Algebra of projections; towards Days 33 and 34 . . . 95

**36 Extensions beyond the notes** **96**
36.1 Rank-one projections . . . 96
36.2 The bit-flip channel at $p = 0$ and $p = 1$ . . . 97
36.3 The phase-flip channel . . . 97
36.4 Unitary-conjugation channels . . . 97
36.5 The bit-phase-flip channel and the identity channel . . . 98
36.6 Composition of channels . . . 99
36.7 Conjugation channels are completely positive . . . 100
36.8 Unitary-conjugation channels are UCP and CPTP . . . 101
36.9 The Pauli channels are UCP and CPTP . . . 101
36.10 Pauli matrices as Kraus operators . . . 102
36.11 Kraus maps of Kraus-operator lists are CPTP . . . 102
36.12 Concatenation of Kraus lists . . . 103
36.13 Fixed points of Pauli channels . . . 103
36.14 The maximally mixed state . . . 104
36.15 Fixed points of unitary-conjugation channels . . . 105
36.16 Channels preserve density matrices . . . 106
36.17 Pauli product identities . . . 106

**37 Small matrix and vector lemmas shared by the gap-closure theories** **106**

**38 Day 12: $H^{\otimes N}$ on $|0\cdots0\rangle$ and the counting probability (D12-1, D12-4) 109**

**39 Day 16: $H_A \otimes H_B \cong H_A^{(p)}$ and $L(H_A \otimes H_B) \cong M_p(L(H_A))$ (D16-2) 110**

**40 Days 17–18: densities to densities iff positive and trace-preserving (D17-4, D18-2) 113**

**41 Day 18: the transpose is positive but not 2-positive (D18-4) 113**

**42 Day 20: Choi's rank theorem $\mathrm{cr}(\Phi) = \mathrm{rank}\, P_\Phi$ (D20-4) 115**
42.1 $\mathrm{rank}(\sum_{l<L} w_l w_l^*) \le L$ . . . 115
42.2 Independent vectors inside the column space bound the rank from below . . . 116
42.3 The theorem . . . 116

**43 Day 21: unitary freedom of rank-one decompositions and Kraus representations (D21-3, D21-5, D21-6) 116**
43.1 The Gram matrix of an independent list is invertible; dual vectors . . . 116
43.2 D21-3: independent vectors give independent rank-one matrices . . . 117
43.3 Isometric mixing of columns preserves $VV^*$ . . . 117
43.4 D21-5: two rank-one decompositions are related by an isometry . . . 118
43.5 D21-6 (Choi 2): unitary freedom of Kraus representations . . . 118

**44 Day 21: ranges of positive matrices (D21-4) 119**
44.1 Day 4 revisited: positive definiteness and spanning (D4-8) . . . 119

**45 Day 22: linear independence bounds the number of Kraus operators (D22-6) 119**

**46 Day 22: a minimal Kraus representation is linearly independent (D22-1) 121**
46.1 Mixing a Kraus list by an isometry does not change the map . . . 121
46.2 A unitary with a prescribed first row . . . 121
46.3 D22-1 . . . 122

**47 Day 22: the adjoint map, an affine bijection from UCP onto CPTP (D22-5) 122**

**48 Day 22: $\mathrm{UCP}(M_n, M_d)$ is compact (D22-4) 124**
48.1 The notes' "Hamming [7,1,3]" code does not correct one error . . . 126
48.2 The Hamming [7,4,3] code corrects one error . . . 127
48.3 Probabilities of undetected and uncorrected errors (Day 30) . . . 127

**49 Day 30: the three-qubit bit-flip code and its recovery map (D30-3, D30-4) 128**
49.1 Errors followed by recovery . . . 132
49.2 Minimum fidelities (D31-4) . . . 133

**50 Day 32: the Pauli expansion identity for unitaries (D32-7) 133**

**51 Days 32–33: composition of Kraus maps and the general recovery proposition (D32-4, D33-2) 134**
51.1 Algebra of *foldr*-sums of matrices . . . 134
51.2 Composition of Kraus maps . . . 135
51.3 The general recovery proposition (D32-4) and its linear-combination form (D33-2) 136
51.4 Matrix sums acting on vectors . . . 136
51.5 Codes with orthogonal error images . . . 139
51.6 The $Y$-flip code . . . 140
51.7 Tensor products of matrices . . . 141

51.8 Single-qubit operators on nine qubits, and on a block of three . . . 141
51.9 The block vectors $|m\rangle \pm |7-m\rangle$ . . . 142
51.10 Code words and the images of the errors . . . 143
51.11 The 22 correctable errors . . . 145
51.12 The code space and its projection . . . 146
51.13 Proposition (Day 33): orthogonal images, and $Z_1 v = Z_2 v = Z_3 v$ . . . 146
51.14 Shor's theorem . . . 147

**52 Day 34: the Knill–Laflamme conditions (D34-1, D34-2, D34-3) 148**
52.1 Vectors in the range of a projection; rank-one decompositions inside $V$ . . . 148
52.2 Linearity facts for the composite map $\mathcal{R} \circ \mathbb{E}$ . . . 149
52.3 Knill–Laflamme, Theorem 1, necessity: $PE_i^* E_j P = \alpha_{ij} P$ . . . 149
52.4 Kraus lists consisting of multiples of one operator . . . 150
52.5 Theorem 2 . . . 150

**III Lectures 23–26: extreme points, operator systems and Arveson's theorem 152**

**53 Day 23 (19 October 2011): Choi's Theorem 3, extreme points of UCP and CPTP 152**
53.1 Extreme UCP maps . . . 153
53.2 UCP-Kraus representation . . . 154
53.3 Unital condition from the Kraus list . . . 155
53.4 UCP-Kraus existence . . . 155
53.5 Sanity facts about *kraus-products* . . . 155
53.6 Quadratic Kraus map (perturbation primitive) . . . 155
53.7 Lemma A: PSD coefficient matrix gives a CP perturbation . . . 159
53.8 Small-$\varepsilon$ PSD perturbation of the identity . . . 160
53.9 Symmetric / anti-hermitian split of the coefficient matrix . . . 160
53.10 Bridge: ¬ *mat-lin-indep-real* gives a real *A-orig* witness . . . 162
53.11 Auxiliary lemmas for the assembly . . . 164
53.12 Minimality and the distinctness lemma (Lemma B) . . . 164
53.13 CPTP-Kraus representation and extreme CPTP maps . . . 165
53.14 Vectorisation of Kraus operators . . . 166
53.15 Choi matrix of a Kraus map . . . 166
53.16 Choi matrix is linear in the map . . . 167
53.17 The Choi matrix of a *kraus-quad-map* . . . 167
53.18 CP Radon–Nikodym — the dominated-PSD kernel . . . 168
53.18.1 Diagonal pseudo-inverse list and its identity . . . 168
53.19 Linearity of *kraus-quad-map* in its matrix argument . . . 170
53.20 Choi matrix determines a linear CP map on carrier . . . 170
53.21 CP Radon–Nikodym . . . 171
53.22 Choi's Theorem 3 — reverse direction . . . 172
53.23 Trace pairing duality and TP-converse . . . 175
53.24 Choi's Theorem 3 — forward direction . . . 175
53.25 Choi's Theorem 3 (statement) . . . 175
53.26 Sanity / easy facts about extreme UCP . . . 177

**54 Day 23: Choi's Theorem 3 for unital maps (D23-1) 178**
54.1 Linear dependence tests . . . 179
54.2 Example 1: unitary conjugation is extreme . . . 180
54.3 Example 2: an average of two unitary conjugations . . . 180
54.4 Example 4: the diagonal map . . . 181
54.5 Example 3: $X \mapsto \mathrm{Tr}(X)\, I_d$ . . . 182
54.6 $M_p(OS)$ is an operator system . . . 183
54.7 Every element is a combination of four positives . . . 184
54.8 Finite sums of matrices . . . 184
54.9 $E_{ij} \otimes M$ and the block decomposition . . . 185
54.10The correspondence . . . 186
54.11Theorem (Arveson): completely positive iff $d$-positive iff $S_\Phi$ positive . . . 187
54.12Vectors . . . 189
54.13The operator norm . . . 189
54.14Unital functionals . . . 190
54.15Auxiliary facts . . . 192
54.16Matrices as points of the Euclidean space *lmat* . . . 193
54.17Extension of positive functionals . . . 193
54.18Arveson's extension theorem . . . 193

**55 Day 26: factoring a CP map through a unital or a trace-preserving one (D26-1) 194**

**IV Functional-analytic infrastructure (Lecture 25: Hahn–Banach) 195**

**56 Typed-dimension matrix wrapper 196**

**57 Typeclass instances 196**

**58 Real scalar multiplication and Hahn-Banach overloading 198**

**59 Real-vectorspace locale on the full type 198**

**60 The Hilbert-Schmidt norm on** (*complex*, *'m*, *'n*) *lmat* **198**

**61 Triangle inequality for the HS norm 199**

**62 Subspace, seminorm, normed-vectorspace instances 199**

**63 Complex scalar multiplication on lmat 200**

**64 Day 25 (26 October 2011): the Hahn–Banach theorem 201**

**65 Bounded complex-linear functionals on lmat 201**

**66 Real-part functional and linearform/continuous instances 201**

**67 Application of HOL/Hahn_Banach (norm-preserving extension) 202**

**68 Bohnenblust-Sobczyk: complex extension from real extension 202**

**69 The matrix-space Hahn-Banach theorem 203**

**70 Overview 204**

71 HS inner-product algebra 204
72 Trace lemmas 205
73 HS norm definition and basic facts 205
74 HS-norm convergence vs entry-wise convergence 207
75 Carathéodory's theorem for conic combinations 209
76 Overview 212
77 Real scalar multiplication and *real-vector* instance 212
78 Auxiliary: pointwise unwrapping (used by later instances) 212
79 Distance, uniformity, open: *metric-space* instance 213
80 Norm and *real-normed-vector* instance 214
81 Real inner product 214
82 Basis: matrix units carrying a basis vector 214
83 Basis set 215
84 Euclidean space instance 216
85 Heine-Borel 216
86 Dimension 216
87 Component-wise bounds via the L2 norm 216
88 Convergence: lmat sequence convergence implies entry convergence 217

V Lectures 26–28: separability and entanglement witnesses 217
89 Overview (Days 27–28 of the notes) 217
90 Typed PSD predicate 218
91 Typed tensor product 218
92 The separable cone 218
93 Step 1: PSD set is closed 219
94 Trace on lmat 219
95 Compactness of the unit-trace PSD slice 220
96 Tensor continuity 220
97 The unit-trace product slice 221

**98 Closedness of the separable cone** **221**

**99 The lmat inner product as a real-trace inner product** **224**

**100 Separating hyperplane for non-separable states** **224**

**101 Entanglement witness in real-trace form** **224**

**102 Hermitian entanglement witness** **225**

**103 Hermitian closure of the separable cone** **225**

**104 Main separability characterisation** **226**

**105 Density-matrix specialisation** **226**

**106 Days 26–28 (28 October – 2 November 2011): separability and entanglement witnesses** **229**
106.1 The separable cone (Day 27) . . . 229
106.2 Separable density matrices (Day 26) . . . 230
106.3 The separable cone is closed (Day 28) . . . 230
106.4 Separation (Days 27–28) . . . 230
106.5 Entanglement witnesses (Day 28) . . . 231
106.6 Tensor products with a matrix in the first factor . . . 232
106.7 The quadratic form of $\Phi_f$ . . . 233
106.8 Proposition (Day 27): $f(\mathrm{Sep}) \geq 0$ iff $\Phi_f$ is positive . . . 233
106.9 Proposition (Day 27): positive maps keep separable matrices positive . . . 233
106.10 Theorem (Day 28): the positive-map criterion . . . 234

**107 Day 27: entangled matrices detected by the transpose (D27-2)** **234**

**108 Day 28: the Horodecki–Peres criterion for two qubits (D28-5)** **236**

**VI Quantum Algorithms (beyond the notes)** **237**

**109 Functional calculus for Hermitian matrices** **237**
109.1 Spectral decompositions . . . 238
109.2 The definition . . . 238
109.3 Algebra . . . 239
109.4 Time evolution . . . 240
109.5 The inverse . . . 240

**110 Phase estimation** **241**
110.1 Sums of roots of unity . . . 241
110.2 The Fourier matrix . . . 241
110.3 Controlled powers and phase kickback . . . 242
110.4 The phase estimation circuit . . . 242
110.5 Measuring the register . . . 243
110.6 The success probability . . . 243

**111 The AFP Fourier transform circuit is the Fourier matrix** **244**

**112 The HHL algorithm** **245**
112.1 Block-diagonal controlled operations . . . 245
112.2 The circuit . . . 246
112.3 The circuit on an eigenvector . . . 247
112.4 All right-hand sides . . . 248
112.5 The success probability . . . 249
112.6 Non-Hermitian systems . . . 249

**113 HHL with inexact phase estimation** **250**
113.1 Register sums . . . 250
113.2 Phase estimation on a branch . . . 251
113.3 Error vectors of distinct eigenvectors are orthogonal . . . 253
113.4 Bounds on the branch distances and amplitudes . . . 254
113.5 The error bound . . . 254
113.6 From eigenvectors to every right-hand side . . . 256

**114 HHL with a sine-window phase register** **257**
114.1 The window amplitude . . . 257
114.2 Phase estimation with an arbitrary register state . . . 258
114.3 The sine window . . . 259
114.4 Counting lemmas . . . 260
114.5 Branch bounds for the window register . . . 260
114.6 The error bound on an eigenvector . . . 261
114.7 Every right-hand side . . . 262

**115 HHL with an approximate Hamiltonian simulation** **263**
115.1 Vector norms . . . 263
115.2 Closeness of block operators . . . 264
115.3 Tensoring with the ancilla . . . 265
115.4 Circuits with oracle calls . . . 266
115.5 Robustness: each call adds at most $\eta$ . . . 267
115.6 Controlled powers with $N-1$ calls . . . 267
115.7 HHL as a circuit with $2(N-1)$ calls . . . 267
115.8 The window is a unit vector . . . 268
115.9 End to end . . . 268

**116 Readout, amplitude amplification and the complete algorithm** **269**
116.1 Reading out the answer . . . 269
116.2 Normalisation is stable . . . 269
116.3 Post-selecting the ancilla . . . 269
116.4 The answer register . . . 270
116.5 Amplitude amplification . . . 270
116.6 The number of rounds . . . 271
116.7 Good and bad components of a state . . . 271
116.8 Reflections as diagonal sign matrices . . . 272
116.9 Amplitude amplification as a circuit . . . 272
116.10 The amplification theorem . . . 273
116.11 The complete algorithm . . . 274

**117 The filter-function algorithm of the HHL paper** **275**
117.1 Filter functions . . . 275
117.2 A $d$-level flag register . . . 277
117.3 Window weights with wrapped distances . . . 279
117.4 Post-selection on the flag register . . . 280

**118 Theorem 1 of the HHL paper for every right-hand side** **282**
118.1 Every right-hand side . . . 282
118.2 The flag gates exist . . . 286

**theory** *QC-Notes-Overview*
**imports**
*Vern-Paulsen-QC.Day23-ChoiExtremeUCP*
*Vern-Paulsen-QC.Day24-Separability*
*Vern-Paulsen-QC.Day12-Counting-Probability*
*Vern-Paulsen-QC.Day18-Positive-Maps*
*Vern-Paulsen-QC.Day22-Kraus-Independence*
*Vern-Paulsen-QC.Day28-Horodecki-Peres*
*Vern-Paulsen-QC.Day30-BitFlip-Code*
*Vern-Paulsen-QC.Day32-Pauli-Expansion*
*Vern-Paulsen-QC.Day21-Unitary-Freedom*
*Vern-Paulsen-QC.Day32-Recovery-Proposition*
*Vern-Paulsen-QC.Day22-Minimal-Kraus*
*Vern-Paulsen-QC.Day34-Knill-Laflamme*
*Vern-Paulsen-QC.Day20-Choi-Rank*
*Vern-Paulsen-QC.Day34-KL-Sufficiency*
*Vern-Paulsen-QC.Day33-Shor-Code*
*Vern-Paulsen-QC.Day17-Environment-Model*
*Vern-Paulsen-QC.Day27-Positive-Map-Criterion*
*Vern-Paulsen-QC.Day02-11-Supplement*
*Vern-Paulsen-QC.Day29-Classical-Codes*
*Vern-Paulsen-QC.Day31-Code-Fidelity*
*Vern-Paulsen-QC.Day32-Y-Code*
*Vern-Paulsen-QC.Day19-Choi-TP-Unital*
*Vern-Paulsen-QC.Day23-Extremality-Examples*
*Vern-Paulsen-QC.Day25-Unital-Functional-Norm*
*Vern-Paulsen-QC.Day26-CP-Factorization*
*Vern-Paulsen-QC.Day27-Entangled-Examples*
*Vern-Paulsen-QC.Day05-Cholesky-Examples*
*Vern-Paulsen-QC.Day10-15-Supplement*
*Vern-Paulsen-QC.Day04-29-Supplement*
*Vern-Paulsen-QC.Day21-Ranges*
*Vern-Paulsen-QC.Day06-Schmidt-Rank*
*Vern-Paulsen-QC.Day22-Adjoint-Maps*
*Vern-Paulsen-QC.Day23-Choi3-UCP*
*Vern-Paulsen-QC.Day16-Block-Form*
*Vern-Paulsen-QC.Day22-UCP-Compact*
*Vern-Paulsen-QC-Algorithms.HHL-Approximate*
*Vern-Paulsen-QC-Algorithms.HHL-Window*
*Vern-Paulsen-QC-Algorithms.HHL-Simulation*
*Vern-Paulsen-QC-Algorithms.HHL-Complete*
*Vern-Paulsen-QC-Algorithms.HHL-Filter*
*Vern-Paulsen-QC-Algorithms.HHL-Filter-Bounds*
*Vern-Paulsen-QC-Algorithms.QFT-Fourier*
**begin**

# 1 Reading guide

Paulsen's course had five parts: (I) basic mathematics, (II) an introduction to quantum computing, (III) the theory of completely positive maps, (IV) entanglement and entanglement witnesses, (V) quantum error correction. The formal development keeps the mathematics of the notes but is organised by theory, since later lectures reuse earlier infrastructure. The table gives the correspondence.

| Lectures | Topic |
|---|---|
| 1–2 | inner products, matrix units, trace, projections, Parseval |
| 3–10 | unitaries, Schur, Hermitian and positive matrices, tensor products, postulates, POVMs, gates |
| 11–22 | no-cloning, oracles, density matrices, partial traces, CP maps, Choi's theorem, adjoints, convexit |
| 23 | extreme points of CPTP/UCP maps (Choi's Theorem 3), CP Radon–Nikodym |
| 25 | Hahn–Banach on matrix spaces |
| – | Hilbert–Schmidt analysis, Carathéodory, typed matrices as a Euclidean space |
| 26–28 | separable cone, separating hyperplanes, entanglement witnesses |
| 29–32 | classical codes, bit/phase-flip channels, fidelity, Pauli algebra |
| 12 | $H^{\otimes N}\|0\cdots0\rangle$, the counting probability |
| 17 | the environment model of noise, $\mathrm{tr}_E(U(\rho\otimes\|\varphi\rangle\langle\varphi\|)U^*)=\sum W_i\rho W_i^*$ |
| 17–18 | densities to densities iff positive and TP; the transpose |
| 20 | $\mathrm{cr}(\Phi)=\operatorname{rank}P_\Phi$ |
| 21 | rank-one decompositions and Kraus representations up to an isometry (Choi 2) |
| 22 | $\mathrm{cr}(\Phi)\le n$ for extreme maps; minimal Kraus lists are independent |
| 24 | operator systems, $M_p(S)$, completely positive maps on $S$ |
| 24–25 | Arveson: CP $\iff$ $d$-positive $\iff$ $S_\Phi\ge0$; the correspondence $\Phi\leftrightarrow S_\Phi$ |
| 26 | Arveson's extension theorem; Kraus form on operator systems |
| 27–28 | positive maps and separability: $f(\mathrm{Sep})\ge0\iff\Phi_f$ positive, the positive-map criterion |
| 28 | the two-qubit Horodecki–Peres criterion |
| 30 | the three-qubit bit-flip code and its recovery theorem |
| 32–33 | Pauli expansion of unitaries; the general recovery proposition |
| 32–33 | the nine-qubit Shor code and Shor's theorem |
| 34 | Knill–Laflamme: states vs. $PXP$, Theorem 1 (both directions), closure |
| 2–11 | basis-independent projections, a basis of $V\oplus W$, tensor coordinates, non-orthogonal states, a Be |
| 29–30 | the notes' Hamming code, the $[7,4,3]$ code, undetected-error probabilities |
| 31 | the bit-flip code against independent flips: $R\circ E$, minimum fidelities |
| 32 | the code for $Y$ flips; codes with orthogonal error images |
| 19 | trace preservation and unitality read off the Choi matrix |
| 23 | extreme and non-extreme CPTP maps: $U^*XU$, averages, $\mathrm{Tr}(X)I/d$, diagonals |
| 25 | a unital functional is positive iff it has norm one |
| 5 | Sylvester's criterion, imported from the AFP (separate session and PDF, see Day 5) |
| 5 | Cholesky's algorithm on the notes' two $3\times3$ examples; the rank-one decomposition |
| 22 | the adjoint $\Phi^*$ of a linear map; $\Gamma(\Phi)=\Phi^*$ is an affine bijection UCP $\to$ CPTP |
| 22 | $\mathrm{UCP}(M_n,M_d)$ is compact (in Choi coordinates) |
| 23 | Choi's Theorem 3 for unital maps, via the adjoint |
| 16 | $H_A\otimes H_B\cong H_A^{(p)}$ isometrically; $X=\sum X_{ij}\otimes E_{ij}$; block action, partial traces |
| 6, 10 | the Schmidt rank: rank 1 iff a product state, entangled iff rank $>1$ |
| 10–15 | superdense coding table, the CCNOT truth table, the partial trace over $A$ |
| 4–29 | converse Hermitian tests, composite ensembles, Choi's four conditions, subspace codes |
| 4, 21 | $0\le ww^*\le P\Rightarrow w\in R(P)$; $R(\sum w_lw_l^*)=\mathrm{span}\{w_l\}$; $\sum v_iv_i^*>0$ iff the $v_i$ span |
| 26 | CP maps as $V^*\psi(X)V$ with $\psi$ UCP and as $\psi(VXV^*)$ with $\psi$ CPTP |
| 27 | the transpose detects $\left(\begin{smallmatrix}E_{ii}&E_{ij}\\E_{ji}&E_{jj}\end{smallmatrix}\right)$ and $(E_{ij})$ as entangled |
| – | functional calculus $g(A)$ for Hermitian $A$; $e^{itA}$, $A^{-1}$ |
| – | Fourier matrix, phase kickback, phase estimation (exact; $\ge4/\pi^2$) |
| – | the AFP circuit *ordered-QFT* is the Fourier matrix $F_{2^n}$ |
| – | the HHL algorithm: output state, $Ax=b$, success $\ge1/\kappa^2$, non-Hermitian $A$ |
| – | inexact phase estimation: the error identity; for the clamped circuit and every $b$, $\|\text{output}-\text{idea}$ |
| – | sine-window register: amplitudes decay like $1/(N\delta)^2$; for every $b$, $\|\text{output}-\text{ideal}\|^2\le(\frac{8\pi^2(2+\pi^2)}{N^2t_0^2C^2(1-}$ |
| – | circuits with oracle calls: HHL makes $2(N-1)$ calls of $e^{iAt_0}$; an oracle $V$ with $\|Vv-e^{iAt_0}v\|\le$ |
| – | readout by post-selection, stable to $2\delta/(C\|A^{-1}b\|)$; amplitude amplification ($k$ rounds give $\sin^2($ |
| – | the paper's filter functions and flag register; Lemma 2; Lemma 3 with the corrected constant $\pi^2$ |
| – | Theorem 1 of the HHL paper for every $b$: errors $O(\kappa/t)$, $t$ the total evolution time |

The theories of the second and third blocks (shared lemmas in `QC_Prelims`) close the gaps recorded by the first and second audits; the remaining unformalised items are listed in `NOTES_AUDIT.md`. The last block, Part VI, goes beyond the notes (session `Vern_Paulsen_QC_Algorithms`).

## 2 Conventions

- Vectors in $\mathbb{C}^n$ are *complex Matrix.vec* with *dim-vec* $v = n$; matrices in $M_{m,n}$ are *complex Matrix.mat* with $A \in$ *carrier-mat* $m$ $n$ (the Jordan-Normal-Form library). Dimensions are hypotheses, not types, except in the typed wrapper *lmat* used for the topological arguments of Part IV.
- The physicists' adjoint $X^\dagger = X^*$ is $A^\dagger$; $\langle y|x\rangle$ is $\langle y|x\rangle$ (conjugate-linear in the first slot, as in the notes); $|v\rangle\langle w|$ is $v \oslash w$; the Hilbert–Schmidt inner product $\langle Y|X\rangle = \mathrm{Tr}(Y^*X)$ is $\langle\langle$ $Y$ | $X$ $\rangle\rangle$; the matrix unit $E_{ij} \in M_{m,n}$ is $E[\![i,j$ ; $m,n]\!]$.
- Indices start at 0, as the physicists in the notes prefer: the canonical basis of $\mathbb{C}^n$ is $|0\rangle, \ldots, |n-1\rangle$, formally *unit-vec n i*.
- $\sum_i |v_i\rangle\langle v_i|$ for a list of vectors is *outer-sum n vs* $= VV^\dagger$ with $V$ the matrix whose columns are the $v_i$.
- Tensor products are Kronecker products: $v \otimes w$ and $A \bigotimes B$, with the first factor as the major index.
- Kraus sums $\sum_i W_i X W_i^\dagger$ over a list are *kraus-map-dim d Ws X*.

Verification status. All theories build with *quick-and-dirty* = *false*; there are no *sorry*s and no axioms beyond those of Isabelle/HOL and the imported libraries.

**end**

# Part I
# Lectures 1–10: basic mathematics and the postulates of quantum mechanics

**theory** *Basic-Math*
  **imports**
    *Quantum-Fourier-Transform.QFT*
    *Isabelle-Marries-Dirac.No-Cloning*
**begin**

Isabelle_Marries_Dirac names its gates *X*, *Y*, *Z*, *H*, *S*, *T*; hidden here so that these letters are variables. The gates remain *Quantum.X* etc.; the project names are *Pauli-X*, *Pauli-Y*, *Pauli-Z*, *Hadamard*, *Phase-S*. Theories importing this one must not import Isabelle_Marries_Dirac directly, as that undoes the hiding.

**hide-const** (**open**) *Quantum.X Quantum.Y Quantum.Z Quantum.H Quantum.S Quantum.T*

*No-Cloning* is imported for *cauchy-schwarz-ineq*; its simp rules (e.g. $v = w$ from inner products) slow simp on every vector equation.

**declare** *cauchy-schwarz-eq* [*simp del*] *cauchy-schwarz-col* [*simp del*] *eq-from-inner-prod* [*simp del*]

# 3 Preliminaries: linear-space isomorphisms

Matrix spaces are represented by carrier predicates on the single type *complex Matrix.mat*. The record below packages a carrier with its zero, addition and scalar multiplication, so that statements such as $L(\mathcal{H}_A) \otimes L(\mathcal{H}_B) \cong L(\mathcal{H}_A \otimes \mathcal{H}_B)$ (Day 15) can be made as explicit linear isomorphisms with explicit inverses.

**record** $('a, 'v)$ *linear-structure* =
  *ls-carrier* :: $'v$ *set*
  *ls-zero* :: $'v$
  *ls-add* :: $'v \Rightarrow 'v \Rightarrow 'v$
  *ls-smul* :: $'a \Rightarrow 'v \Rightarrow 'v$

**definition** *linear-structure-map* ::
  $('a, 'v)$ *linear-structure* $\Rightarrow$ $('a, 'w)$ *linear-structure* $\Rightarrow$ $('v \Rightarrow 'w) \Rightarrow$ *bool* **where**
  *linear-structure-map V W f* $\equiv$
    $f\ (ls\text{-}zero\ V) = ls\text{-}zero\ W \wedge$
    $(\forall x \in ls\text{-}carrier\ V.\ f\ x \in ls\text{-}carrier\ W) \wedge$
    $(\forall x \in ls\text{-}carrier\ V.\ \forall y \in ls\text{-}carrier\ V.$
      $f\ (ls\text{-}add\ V\ x\ y) = ls\text{-}add\ W\ (f\ x)\ (f\ y)) \wedge$
    $(\forall c\ x.\ x \in ls\text{-}carrier\ V \longrightarrow$
      $f\ (ls\text{-}smul\ V\ c\ x) = ls\text{-}smul\ W\ c\ (f\ x))$

**definition** *is-isomorphism-between* ::
  $('a, 'v)$ *linear-structure* $\Rightarrow$ $('a, 'w)$ *linear-structure* $\Rightarrow$ $('v \Rightarrow 'w) \Rightarrow$ *bool* **where**
  *is-isomorphism-between V W f* $\equiv$
    *linear-structure-map V W f* $\wedge$
    $(\exists g.$ *linear-structure-map W V g* $\wedge$
        $(\forall x \in ls\text{-}carrier\ V.\ g\ (f\ x) = x) \wedge$
        $(\forall y \in ls\text{-}carrier\ W.\ f\ (g\ y) = y))$

**definition** *is-isomorphic-to* ::
  $('a, 'v)$ *linear-structure* $\Rightarrow$ $('a, 'w)$ *linear-structure* $\Rightarrow$ *bool* **where**
  *is-isomorphic-to V W* $\equiv \exists f.$ *is-isomorphism-between V W f*

**lemma** *is-isomorphism-betweenD-map*:
  **assumes** *is-isomorphism-between V W f*
  **shows** *linear-structure-map V W f*
  ⟨*proof*⟩

**lemma** *is-isomorphism-betweenD-inverse*:
  **assumes** *is-isomorphism-between V W f*
  **obtains** *g* **where** *linear-structure-map W V g*
    **and** $\bigwedge x.\ x \in ls\text{-}carrier\ V \Longrightarrow g\ (f\ x) = x$
    **and** $\bigwedge y.\ y \in ls\text{-}carrier\ W \Longrightarrow f\ (g\ y) = y$
  ⟨*proof*⟩

**definition** *mat-linear-structure* :: *nat* $\Rightarrow$ *nat* $\Rightarrow$ (*complex*, *complex Matrix.mat*) *linear-structure* **where**
  *mat-linear-structure m n* $\equiv$
    $(\!|$ *ls-carrier* $=$ *carrier-mat m n*,
      *ls-zero* $= 0_m\ m\ n$,
      *ls-add* $= (\lambda x\ y.\ x + y)$,
      *ls-smul* $= (\lambda c\ x.\ c \cdot_m x)$ $|\!)$

# 4 Day 1 (22 August 2011): the inner product

Paulsen opens with the dictionary between classical and quantum computing: a bit is an element of $\mathbb{Z}_2$, a qubit a unit vector in $\mathbb{C}^2$; $N$ bits form $\mathbb{Z}_2^N$, $N$ qubits a unit vector in $\mathbb{C}^{2^N} \cong \mathbb{C}^2 \otimes \cdots \otimes \mathbb{C}^2$; operations are Boolean gates versus completely positive maps on $M_{2^N}$; information is a probability vector versus a positive semidefinite matrix of trace one. He then fixes the physicists' notation.

**Definition (notes).** $\mathbb{C}^n$ is the space of complex $n$-tuples with the inner product

$$\langle y|x\rangle = \sum_{i=1}^{n} \overline{y_i}\, x_i,$$

which is linear in the right-hand slot, conjugate linear in the left-hand slot ("sesquilinear"), and positive definite: $\langle x|x\rangle \geq 0$ with equality only for $x = 0$. The Euclidean length is $\|x\| = \sqrt{\langle x|x\rangle}$. A *Hilbert space* is a complex vector space with a sesquilinear positive definite form, complete for this norm.

The library's inner product $\langle y|x\rangle$ is exactly this form, with indices running from 0 to $n-1$.

**lemma** *inner-prod-expand*:
**assumes** *dim-vec* $u = n$ **and** *dim-vec* $v = n$
**shows** $\langle u|v\rangle = (\sum\ i \in \{0..< n\}.\ cnj(u\ \$\ i) * (v\ \$\ i))$
⟨*proof*⟩

## 4.1 The Hilbert–Schmidt inner product on matrices

**Definition (notes)** (Example)**.** The space $M_{n,k}$ of complex $n \times k$ matrices is a Hilbert space for

$$\langle (y_{ij}) \,|\, (x_{ij})\rangle = \sum_i \sum_j \overline{y_{ij}}\, x_{ij},$$

and $M_{n,k} \cong \mathbb{C}^{nk}$.

We write $\langle\langle y|x\rangle\rangle$ for this Hilbert–Schmidt (Frobenius) inner product; the double brackets distinguish it from the inner product of vectors.

**definition** *hs-inner-prod* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *complex* **where**
*hs-inner-prod* = (λ *y x*.
$\sum$ *i* ∈ {*0* ..< *dim-row x*}.
$\sum$ *j* ∈ {*0* ..< *dim-col x*}.
*cnj* (*y* $$ (*i*,*j*)) ∗ (*x* $$ (*i*,*j*)))

**notation** *hs-inner-prod* (⟨⟨ - | - ⟩⟩)

**lemma** *hs-inner-prod-expand*:
**assumes** *dim-row x* = *m* **and** *dim-col x* = *n*
**shows** ⟨⟨ *y* | *x* ⟩⟩ =
(∑ *i* ∈ {*0* ..< *m*}. ∑ *j* ∈ {*0* ..< *n*}. *cnj* (*y* $$ (*i*,*j*)) ∗ (*x* $$ (*i*,*j*)))
⟨*proof*⟩

**lemma** *hs-inner-prod-square*:
**assumes** *dim-row x* = *n* **and** *dim-col x* = *n*
**shows** ⟨⟨ *y* | *x* ⟩⟩ =
(∑ *i* ∈ {*0* ..< *n*}. ∑ *j* ∈ {*0* ..< *n*}. *cnj* (*y* $$ (*i*,*j*)) ∗ (*x* $$ (*i*,*j*)))
⟨*proof*⟩

### 4.2 Properties of the inner product

The properties listed in the notes, verified for the matrix inner product: conjugate symmetry, linearity in the right-hand slot, conjugate linearity in the left-hand slot, and positive definiteness.

**lemma** *hs-inner-prod-cnj*:
**assumes** *dim-row x = dim-row y* **and** *dim-col x = dim-col y*
**shows** ⟨⟨ *x* | *y* ⟩⟩ = *cnj* (⟨⟨ *y* | *x* ⟩⟩)
⟨*proof*⟩

**lemma** *hs-inner-prod-right-add*:
**assumes** *dim-row X1 = dim-row y* **and** *dim-col X1 = dim-col y*
**and** *dim-row X2 = dim-row y* **and** *dim-col X2 = dim-col y*
**shows** ⟨⟨ *y* | *X1 + X2* ⟩⟩ = ⟨⟨ *y* | *X1* ⟩⟩ + ⟨⟨ *y* | *X2* ⟩⟩
⟨*proof*⟩

**lemma** *hs-inner-prod-right-smult*:
**assumes** *dim-row x = dim-row y* **and** *dim-col x = dim-col y*
**shows** ⟨⟨ *y* | *c* ·$_m$ *x* ⟩⟩ = *c* ∗ ⟨⟨ *y* | *x* ⟩⟩
⟨*proof*⟩

**lemma** *hs-inner-prod-left-smult*:
**assumes** *dim-row x = dim-row y* **and** *dim-col x = dim-col y*
**shows** ⟨⟨ *c* ·$_m$ *y* | *x* ⟩⟩ = *cnj c* ∗ ⟨⟨ *y* | *x* ⟩⟩
⟨*proof*⟩

**lemma** *hs-inner-prod-self-real*:
**shows** ⟨⟨ *x* | *x* ⟩⟩ ∈ ℝ
⟨*proof*⟩

**lemma** *hs-inner-prod-self-nonneg*:
**shows** *Re* (⟨⟨ *x* | *x* ⟩⟩) ≥ *0*
⟨*proof*⟩

**lemma** *hs-inner-prod-eq-zero-imp*:
**assumes** ⟨⟨ *x* | *x* ⟩⟩ = *0*
**and** *i* < *dim-row x* **and** *j* < *dim-col x*
**shows** *x* $$ (*i*,*j*) = *0*
⟨*proof*⟩

# 5 Day 2 (24 August 2011): bases, matrix units, trace, projections

**Definition (notes)** (Examples)**.** $\mathbb{C}^n$ has the orthonormal basis $e_1, \ldots, e_n$ and every $x$ satisfies $x = \sum_i x_i e_i = \sum_i \langle e_i|x\rangle e_i$. The *matrix units* $E_{ij}$ (entry 1 in position $(i, j)$, 0 elsewhere) form an orthonormal basis of $M_{n,m}$, and

$$X = \sum_{i,j} X_{ij} E_{ij} = \sum_{i,j} \langle E_{ij}|X\rangle E_{ij}.$$

Formally $E[\![i,j\ ;\ m,n]\!]$ is the $m \times n$ matrix unit (indices first, dimensions second). The notes also recall the conjugate $\overline{X}$, the transpose $X^t$, the adjoint $X^* = \overline{X}^t$ (written $X^\dagger$ by physicists, formally $X^\dagger$) and the row-times-column rule for matrix products; these are library notions.

**definition** *mat-unit* :: *nat* ⇒ *nat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*mat-unit i j m n* = *Matrix.mat m n* (λ(*r*,*c*). *if r* = *i* ∧ *c* = *j then 1 else 0*)

**notation** *mat-unit* (*E*[![-,- ; -,-]!] [*1000*, *1000*, *1000*, *1000*] *999*)

**lemma** *mat-unit-dim-row* [*simp*]:
**fixes** *i j m n* :: *nat*
**shows** *dim-row* (*E*[![*i*,*j* ; *m*,*n*]!]) = *m*
⟨*proof*⟩

**lemma** *mat-unit-dim-col* [*simp*]:
**fixes** *i j m n* :: *nat*
**shows** *dim-col* (*E*[![*i*,*j* ; *m*,*n*]!]) = *n*
⟨*proof*⟩

**lemma** *mat-unit-index* [*simp*]:
**assumes** *r* < *m* **and** *c* < *n*
**shows** (*E*[![*i*,*j* ; *m*,*n*]!]) $$ (*r*,*c*) = (*if r* = *i* ∧ *c* = *j then 1 else 0*)
⟨*proof*⟩

Pairing with $E_{ij}$ extracts the $(i, j)$ entry: the matrix analogue of $\langle e_i|v\rangle = v_i$.

**lemma** *hs-inner-prod-mat-unit-left*:
**assumes** *i* < *dim-row x* **and** *j* < *dim-col x*
**shows** ⟨⟨ *E*[![*i*,*j* ; (*dim-row x*), (*dim-col x*)]!] | *x* ⟩⟩ = *x* $$ (*i*,*j*)
⟨*proof*⟩

**lemma** *mat-unit-orthonormal*:
**assumes** *i-lt*: *i* < *m*
**and** *j-lt*: *j* < *n*
**and** *k-lt*: *k* < *m*
**and** *l-lt*: *l* < *n*
**shows** ⟨⟨ *E*[![*i*,*j* ; *m*,*n*]!] | *E*[![*k*,*l* ; *m*,*n*]!] ⟩⟩ =
(*if i* = *k* ∧ *j* = *l then 1 else 0*)
⟨*proof*⟩

Expansion in the basis $\{E_{ij}\}$, stated entrywise because matrices of different dimensions cannot be added in the library.

**lemma** *mat-unit-expansion*:
**assumes** *r* < *dim-row x* **and** *c* < *dim-col x*
**shows** *x* $$ (*r*,*c*) =
(∑ *i* ∈ {*0*..<*dim-row x*}. ∑ *j* ∈ {*0*..<*dim-col x*}.
*x* $$ (*i*,*j*) ∗ (*mat-unit i j* (*dim-row x*) (*dim-col x*)) $$ (*r*,*c*))
⟨*proof*⟩

**lemma** *mat-unit-hs-expansion*:
**assumes** *r* < *dim-row x* **and** *c* < *dim-col x*
**shows** *x* $$ (*r*,*c*) =
(∑ *i* ∈ {*0*..<*dim-row x*}. ∑ *j* ∈ {*0*..<*dim-col x*}.
⟨⟨ *mat-unit i j* (*dim-row x*) (*dim-col x*) | *x* ⟩⟩ ∗
(*mat-unit i j* (*dim-row x*) (*dim-col x*)) $$ (*r*,*c*))

⟨*proof*⟩

## 5.1 The trace

**Definition (notes).** $\mathrm{Tr} : M_{n,n} \to \mathbb{C}$, $\mathrm{Tr}(X) = \sum_{i=1}^{n} x_{ii}$; we sometimes write $\mathrm{Tr}_n$.

The formal *Tr* sums the diagonal of any matrix. On matrix units, $\mathrm{Tr}(E_{ii}) = 1$ and $\mathrm{Tr}(E_{ij}) = 0$ for $i \neq j$, so the trace is the sum of the diagonal Hilbert–Schmidt coefficients.

**definition** *Tr* :: *complex Matrix.mat* ⇒ *complex* **where**
*Tr x* = (∑ *r* < *dim-row x*. *if r* < *dim-col x then x* $$ (*r*,*r*) *else 0*)

**lemma** *Tr-mat-unit*:
**assumes** *i-lt*: *i* < *m*
**and** *j-lt*: *j* < *n*
**shows** *Tr* (*E*⟦*i*,*j* ; *m*,*n*⟧) = (*if i* = *j then 1 else 0*)
⟨*proof*⟩

**corollary** *Tr-mat-unit-diag*:
**assumes** *i* < *n*
**shows** *Tr* (*E*⟦*i*,*i* ; *n*,*n*⟧) = *1*
⟨*proof*⟩

**corollary** *Tr-mat-unit-offdiag*:
**assumes** *i* < *m* **and** *j* < *n* **and** *i* ≠ *j*
**shows** *Tr* (*E*⟦*i*,*j* ; *m*,*n*⟧) = *0*
⟨*proof*⟩

**lemma** *Tr-hs-diagonal-sum*:
**shows** *Tr x* =
(∑ *r* < *dim-row x*.
*if r* < *dim-col x*
*then* ⟨⟨ *E*⟦*r*,*r* ; (*dim-row x*), (*dim-col x*)⟧ | *x* ⟩⟩
*else 0*)
⟨*proof*⟩

**corollary** *Tr-hs-diagonal-sum-square*:
**assumes** *rows*: *dim-row x* = *n*
**and** *cols*: *dim-col x* = *n*
**shows** *Tr x* = (∑ *r*<*n*. ⟨⟨ *E*⟦*r*,*r* ; *n*,*n*⟧ | *x* ⟩⟩)
⟨*proof*⟩

The trace is linear.

**lemma** *Tr-add*:
**assumes** *rows*: *dim-row x* = *dim-row y*
**and** *cols*: *dim-col x* = *dim-col y*
**shows** *Tr* (*x* + *y*) = *Tr x* + *Tr y*
⟨*proof*⟩

**lemma** *Tr-smult*: *Tr* (*a* ·$_m$ *x*) = *a* ∗ *Tr x*
⟨*proof*⟩

**corollary** *Tr-uminus*: *Tr* (− *x*) = − *Tr x*
⟨*proof*⟩

**corollary** *Tr-diff*:
**assumes** *rows*: *dim-row x* = *dim-row y*
**and** *cols*: *dim-col x* = *dim-col y*
**shows** *Tr* (*x* − *y*) = *Tr x* − *Tr y*
⟨*proof*⟩

**lemma** *Tr-zero-mat* [*simp*]:
**shows** *Tr* (*Matrix.mat m n* (λ(*r*,*c*). *0*)) = *0*
⟨*proof*⟩

## 5.2 Matrix-unit algebra and cyclicity of the trace

$E_{ij}^* = E_{ji}$, $E_{ij}E_{kl} = \delta_{jk}E_{il}$, $\mathrm{Tr}(AB) = \mathrm{Tr}(BA)$, $\mathrm{Tr}(U^*AU) = \mathrm{Tr}(A)$ for unitary $U$, and $\langle A|B\rangle = \mathrm{Tr}(A^*B)$: the facts about the trace that the notes use from Day 3 onwards.

**lemma** *mat-unit-dagger*:
**shows** (*E*⟦*i*,*j* ; *m*,*n*⟧)† = *E*⟦*j*,*i* ; *n*,*m*⟧
⟨*proof*⟩

**lemma** *mat-unit-mult-entry*:
**assumes** *r-lt*: *r* < *m*
**and** *c-lt*: *c* < *p*
**and** *j-lt*: *j* < *n*
**shows** (*E*⟦*i*,*j* ; *m*,*n*⟧ ∗ *E*⟦*k*,*l* ; *n*,*p*⟧) $$ (*r*,*c*) =
(*if j* = *k* ∧ *r* = *i* ∧ *c* = *l then 1 else 0*)
⟨*proof*⟩

**lemma** *Tr-cyclic*:
**assumes** *AB*: *dim-col A* = *dim-row B*
**and** *BA*: *dim-row A* = *dim-col B*
**shows** *Tr* (*A* ∗ *B*) = *Tr* (*B* ∗ *A*)
⟨*proof*⟩

**lemma** *Tr-unitary-conjugation*:
**assumes** *Uu*: *unitary U*
**and** *Arow*: *dim-row A* = *dim-row U*
**and** *Acol*: *dim-col A* = *dim-row U*
**shows** *Tr* ((*U*†) ∗ *A* ∗ *U*) = *Tr*(*A*)
⟨*proof*⟩

**lemma** *hs-inner-prod-as-trace*:
**assumes** *rows*: *dim-row A* = *dim-row B*
**and** *cols*: *dim-col A* = *dim-col B*
**shows** ⟨⟨ *A* | *B* ⟩⟩ = *Tr* ((*A*†) ∗ *B*)
⟨*proof*⟩

**corollary** *hs-inner-prod-self-as-trace*:
**shows** ⟨⟨ *A* | *A* ⟩⟩ = *Tr* ((*A*†) ∗ *A*)
⟨*proof*⟩

## 5.3 Orthogonal projections and Parseval's identity

**Definition (notes)** (Orthogonal projections)**.** Let $V \subseteq \mathbb{C}^n$ be a subspace with orthonormal basis $\{v_1, \ldots, v_k\}$ and put $P = \sum_{i=1}^{k} |v_i\rangle\langle v_i|$. Then $Px = \sum_i \langle v_i|x\rangle v_i \in V$ and $x - Px \perp V$, so every $x \in \mathbb{C}^n$ is uniquely $x = v + w$ with $v \in V$ and $w \in V^{\perp}$. Consequently $P$ does not depend on the orthonormal basis chosen for $V$, $Pv = v$ for $v \in V$, $P^2 = P$ and $P = P^*$: $P$ is *the* orthogonal projection onto $V$.

**Theorem (notes)** (Parseval)**.** If $\{v_1, \ldots, v_k\}$ is an orthonormal basis of $V$ and $v \in V$, then $\|v\|^2 = \sum_{i=1}^{k} |\langle v_i|v\rangle|^2$. For $V = \mathbb{C}^n$ this holds for every $v$.

**Theorem (notes).** If $P \in M_n$ satisfies $P^2 = P$ and $P = P^*$, and $V = \mathrm{range}(P)$, then $P$ is the orthogonal projection onto $V$.

Formally, an orthonormal list *vs* of vectors in $\mathbb{C}^n$ is *orthonormal-list n vs*, the matrix with columns $v_1, \ldots, v_k$ is *mat-of-cols n vs*, and $P = \sum_i |v_i\rangle\langle v_i| = VV^\dagger$ is *outer-sum n vs*. The results below are the projection formula, $V^\dagger V = I_k$, the completeness relation $VV^\dagger = I_n$ for a full basis, $P = P^\dagger$, $P^2 = P$, Parseval, the orthogonality of the residual, and the uniqueness of the decomposition. Independence of $P$ from the basis is *outer-sum-basis-independent* in theory Day02_11_Supplement; the characterisation of Hermitian idempotents as orthogonal projections is *projection-annihilates-orthogonal* in theory Quantum_Computing11.

**definition** *orthonormal-list* :: *nat* ⇒ *complex Matrix.vec list* ⇒ *bool* **where**
*orthonormal-list n vs* ≡
(∀ *v* ∈ *set vs*. *dim-vec v* = *n*) ∧
(∀ *i* < *length vs*. ∀ *j* < *length vs*.
⟨*vs* ! *i* | *vs* ! *j*⟩ = (*if i* = *j then 1 else 0*))

**lemma** *orthonormal-listD-dim*:
**assumes** *orthonormal-list n vs* **and** *i* < *length vs*
**shows** *dim-vec* (*vs* ! *i*) = *n*
⟨*proof*⟩

**lemma** *orthonormal-listD-inner*:
**assumes** *orthonormal-list n vs* **and** *i* < *length vs* **and** *j* < *length vs*
**shows** ⟨*vs* ! *i* | *vs* ! *j*⟩ = (*if i* = *j then 1 else 0*)
⟨*proof*⟩

**lemma** *orthonormal-listD-carrier*:
**assumes** *orthonormal-list n vs*
**shows** *set vs* ⊆ *carrier-vec n*
⟨*proof*⟩

**definition** *outer-sum* :: *nat* ⇒ *complex Matrix.vec list* ⇒ *complex Matrix.mat* **where**
*outer-sum n vs* = *mat-of-cols n vs* ∗ ((*mat-of-cols n vs*)†)

**lemma** *outer-sum-dim-row* [*simp*]: *dim-row* (*outer-sum n vs*) = *n*
⟨*proof*⟩

**lemma** *outer-sum-dim-col* [*simp*]: *dim-col* (*outer-sum n vs*) = *n*
⟨*proof*⟩

**lemma** *outer-sum-hermitian*:
**shows** (*outer-sum n vs*)† = *outer-sum n vs*
⟨*proof*⟩

**lemma** *cols-mat-dagger-left*:
**assumes** *orth*: *orthonormal-list n vs*
**shows** $(mat\text{-}of\text{-}cols\ n\ vs)^\dagger * (mat\text{-}of\text{-}cols\ n\ vs) = 1_m\ (length\ vs)$
⟨*proof*⟩

**lemma** *cols-mat-dagger-right*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *full*: *length vs* = *n*
**shows** $(mat\text{-}of\text{-}cols\ n\ vs) * ((mat\text{-}of\text{-}cols\ n\ vs)^\dagger) = 1_m\ n$
⟨*proof*⟩

**lemma** *completeness-relation*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *full*: *length vs* = *n*
**and** *i-lt*: $i < n$ **and** *j-lt*: $j < n$
**shows** $(\sum l{<}n.\ (vs\ !\ l)\ \$\ i * cnj\ ((vs\ !\ l)\ \$\ j)) = (if\ i = j\ then\ 1\ else\ 0)$
⟨*proof*⟩

**lemma** *outer-sum-index*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *i-lt*: $i < n$ **and** *j-lt*: $j < n$
**shows** $(outer\text{-}sum\ n\ vs)\ \$\$\ (i,j) =$
$(\sum l < length\ vs.\ (vs\ !\ l)\ \$\ i * cnj\ ((vs\ !\ l)\ \$\ j))$
⟨*proof*⟩

**lemma** *outer-sum-apply*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *xdim*: *dim-vec x* = *n*
**and** *i-lt*: $i < n$
**shows** $(outer\text{-}sum\ n\ vs *_v\ x)\ \$\ i = (\sum l < length\ vs.\ \langle vs\ !\ l \mid x\rangle * (vs\ !\ l)\ \$\ i)$
⟨*proof*⟩

**lemma** *outer-sum-idempotent*:
**assumes** *orth*: *orthonormal-list n vs*
**shows** *outer-sum n vs* ∗ *outer-sum n vs* = *outer-sum n vs*
⟨*proof*⟩

**lemma** *parseval*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *full*: *length vs* = *n*
**and** *vdim*: *dim-vec v* = *n*
**and** *i-lt*: $i < n$
**shows** $v\ \$\ i = (\sum l{<}n.\ \langle vs\ !\ l \mid v\rangle * (vs\ !\ l)\ \$\ i)$
⟨*proof*⟩

**corollary** *parseval-norm*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *full*: *length vs* = *n*
**and** *vdim*: *dim-vec v* = *n*
**shows** $\|v\|^2 = (\sum l{<}n.\ (cmod\ (\langle vs\ !\ l \mid v\rangle))^2)$
⟨*proof*⟩

The residual $x - Px$ is orthogonal to every $v_l$, because $\langle v_l|Px\rangle = \langle v_l|x\rangle$; hence the orthogonal decomposition $x = Px + (x - Px)$.

**lemma** *inner-prod-outer-sum*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *xdim*: *dim-vec x* = *n*
**and** *l-lt*: *l* < *length vs*
**shows** ⟨*vs* ! *l* | *outer-sum n vs* $*_v$ *x*⟩ = ⟨*vs* ! *l* | *x*⟩
⟨*proof*⟩

**lemma** *residual-orthogonal*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *xdim*: *dim-vec x* = *n*
**and** *l-lt*: *l* < *length vs*
**shows** ⟨*vs* ! *l* | *x* − *outer-sum n vs* $*_v$ *x*⟩ = *0*
⟨*proof*⟩

**corollary** *orthogonal-decomposition*:
**assumes** *xdim*: *dim-vec x* = *n*
**shows** *x* = *outer-sum n vs* $*_v$ *x* + (*x* − *outer-sum n vs* $*_v$ *x*)
⟨*proof*⟩

Uniqueness: if $x = v + w$ with $Pv = v$ and $w \perp v_l$ for all $l$, then $v = Px$ and $w = x - Px$.

**lemma** *outer-sum-orthogonal-zero*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *wdim*: *dim-vec w* = *n*
**and** *ortho*: ∀ *l* < *length vs*. ⟨*vs* ! *l* | *w*⟩ = *0*
**shows** *outer-sum n vs* $*_v$ *w* = *0*$_v$ *n*
⟨*proof*⟩

**lemma** *outer-sum-add*:
**assumes** *vdim*: *dim-vec v* = *n* **and** *wdim*: *dim-vec w* = *n*
**shows** *outer-sum n vs* $*_v$ (*v* + *w*) =
*outer-sum n vs* $*_v$ *v* + *outer-sum n vs* $*_v$ *w*
⟨*proof*⟩

**lemma** *orthogonal-decomposition-unique*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *xdim*: *dim-vec x* = *n*
**and** *vdim*: *dim-vec v* = *n*
**and** *wdim*: *dim-vec w* = *n*
**and** *decomp*: *x* = *v* + *w*

**and** *v-fixed*: *outer-sum n vs* $*_v$ *v* = *v*
**and** *w-ortho*: ∀ *l* < *length vs*. ⟨*vs* ! *l* | *w*⟩ = *0*
**shows** *v* = *outer-sum n vs* $*_v$ *x* **and** *w* = *x* − *outer-sum n vs* $*_v$ *x*
⟨*proof*⟩

# 6 Unitary algebra

Library-level consequences of unitarity ($U^\dagger U = I$ and $UU^\dagger = I$ in the library's definition) used throughout; Paulsen's Day 3 characterisations of unitaries are in theory Quantum_Computing_3_10.

**lemma** *unitary-square-dim*:
**assumes** *uA*: *unitary A*
**shows** *dim-row A* = *dim-col A*
⟨*proof*⟩

**lemma** *unitary-inv-l-dimcol*:
**assumes** *uA*: *unitary A*
**shows** ($A^\dagger$) * *A* = *1*$_m$ (*dim-col A*)
⟨*proof*⟩

**lemma** *unitary-inv-r-dimrow*:
**assumes** *uA*: *unitary A*
**shows** *A* * ($A^\dagger$) = *1*$_m$ (*dim-row A*)
⟨*proof*⟩

**lemma** *unitary-inv-l-dimrow*:
**assumes** *uA*: *unitary A*
**shows** ($A^\dagger$) * *A* = *1*$_m$ (*dim-row A*)
⟨*proof*⟩

**lemma** *unitary-inv-r-dimcol*:
**assumes** *uA*: *unitary A*
**shows** *A* * ($A^\dagger$) = *1*$_m$ (*dim-col A*)
⟨*proof*⟩

**lemma** *unitary-carrier-square-col*:
**assumes** *uA*: *unitary A*
**shows** *A* ∈ *carrier-mat* (*dim-col A*) (*dim-col A*)
⟨*proof*⟩

**lemma** *unitary-carrier-square-row*:
**assumes** *uA*: *unitary A*
**shows** *A* ∈ *carrier-mat* (*dim-row A*) (*dim-row A*)
⟨*proof*⟩

**lemma** *unitary-dagger*:
**assumes** *uA*: *unitary A*
**shows** *unitary* ($A^\dagger$)
⟨*proof*⟩

**lemma** *unitary-mult*:
**assumes** *UA*: *unitary A*
**and** *UB*: *unitary B*
**and** *dimA*: *A* ∈ *carrier-mat n n*
**and** *dimB*: *B* ∈ *carrier-mat n n*
**shows** *unitary* (*A* * *B*)
⟨*proof*⟩

**lemma** *unitary-mult2*:
**assumes** *UA*: *unitary A*
**and** *UB*: *unitary B*
**and** *dims*: *dim-row A* = *dim-row B*
**shows** *unitary* (*A* ∗ *B*)
⟨*proof*⟩

# 7 Tensor product infrastructure

Kronecker products of matrices, $A \bigotimes B$: the tensor of identities, the adjoint $(R \otimes T)^* = R^* \otimes T^*$, the mixed-product property $(R_1 \otimes T_1)(R_2 \otimes T_2) = R_1 R_2 \otimes T_1 T_2$ (both from Day 9), and unitarity of tensors of unitaries.

## 7.1 Tensor of Identities

**lemma** *tensor-mat-id*:
*Tensor.tensor-mat* (*1*$_m$ *n*) (*1*$_m$ *m*) = *1*$_m$ (*n* ∗ *m*)
⟨*proof*⟩

## 7.2 Tensor Adjoint

**lemma** *tensor-mat-adjoint*:
(*Tensor.tensor-mat A B*)$^\dagger$ = *Tensor.tensor-mat* (*A*$^\dagger$) (*B*$^\dagger$)
⟨*proof*⟩

## 7.3 Mixed-Product Property

**lemma** *tensor-mat-mult*:
**assumes** *dimAC*: *dim-col A* = *dim-row C*
**and** *dimBD*: *dim-col B* = *dim-row D*
**shows** *Tensor.tensor-mat A B* ∗ *Tensor.tensor-mat C D* =
*Tensor.tensor-mat* (*A* ∗ *C*) (*B* ∗ *D*)
⟨*proof*⟩

## 7.4 Corollaries of the Mixed-Product Property

**lemma** *tensor-adjoint-mult*:
(*Tensor.tensor-mat* (*A*$^\dagger$) (*B*$^\dagger$)) ∗ (*Tensor.tensor-mat A B*) =
*Tensor.tensor-mat* ((*A*$^\dagger$) ∗ *A*) ((*B*$^\dagger$) ∗ *B*)
⟨*proof*⟩

**lemma** *tensor-mult-adjoint*:
(*Tensor.tensor-mat A B*) ∗ (*Tensor.tensor-mat* (*A*$^\dagger$) (*B*$^\dagger$)) =
*Tensor.tensor-mat* (*A* ∗ (*A*$^\dagger$)) (*B* ∗ (*B*$^\dagger$))
⟨*proof*⟩

## 7.5 Tensor Unitarity

**lemma** *tensor-mat-unitary-carrier*:
**assumes** *dimA*: *A* ∈ *carrier-mat n n*
**and** *dimB*: *B* ∈ *carrier-mat m m*
**and** *uA*: *unitary A*
**and** *uB*: *unitary B*
**shows** *unitary* (*Tensor.tensor-mat A B*)
⟨*proof*⟩

**lemma** *tensor-mat-unitary*:
**assumes** *uA*: *unitary A*
**and** *uB*: *unitary B*
**shows** *unitary* (*Tensor.tensor-mat A B*)
⟨*proof*⟩

**lemma** *tensor-mat-unitary-one-left* [*simp*]:
**assumes** *uA*: *unitary A*
**shows** *unitary* (*Tensor.tensor-mat* (*1*$_m$ *n*) *A*)
⟨*proof*⟩

**lemma** *tensor-mat-unitary-one-right* [*simp*]:
**assumes** *uA*: *unitary A*
**shows** *unitary* (*Tensor.tensor-mat A* (*1*$_m$ *n*))
⟨*proof*⟩

**end**

**theory** *Quantum-Computing-3-10*
**imports** *Basic-Math Jordan-Normal-Form.Schur-Decomposition*
**begin**

# 8 Day 3 (31 August 2011): unitary matrices, Householder unitaries, Schur's theorem

## 8.1 Unitary matrices

**Definition (notes).** $U \in M_n$ is *unitary* if $U^*U = I$.

**Theorem (notes).** For $U \in M_n$ the following are equivalent: (a) $U$ is unitary; (b) $U$ is invertible and $U^{-1} = U^*$; (c) $UU^* = I$; (d) $U^*$ is unitary; (e) the columns of $U$ are orthonormal; (f) the rows of $U$ are orthonormal; (g) $\|Ux\| = \|x\|$ for all $x$ (isometry); (h) $\langle Ux|Uy\rangle = \langle x|y\rangle$ for all $x, y$.

The library predicate *unitary U* bundles (a) and (c). The lemmas below give (a)⇒(b), (d), (e), (f), (g), (h); (e)⇒(a) is *orthonormal-mat-unitary*, (g)⇒(a) and (h)⇒(a) are *unitary-of-isometry* and *unitary-of-inner-prod-preserving* (proved on Day 4, after the polarisation lemma); (d) is self-dual and (f) follows from (e) for $U^*$.

**lemma** *unitary-left-right*:
**assumes** *unitary U*
**shows** *U*$^\dagger$ ∗ *U* = *1*$_m$ (*dim-col U*) **and** *U* ∗ (*U*$^\dagger$) = *1*$_m$ (*dim-row U*)
⟨*proof*⟩

(b) $U$ is invertible with inverse $U^*$:

**lemma** *unitary-invertible*:
**assumes** *unitary U*
**shows** *invertible-mat U*
⟨*proof*⟩

(d) $U^*$ is unitary:

**lemma** *unitary-dagger-is-unitary*:
**assumes** *unitary U*
**shows** *unitary* (*U*$^\dagger$)
⟨*proof*⟩

(e), (f) the columns and the rows of $U$ are orthonormal:

**lemma** *unitary-cols-orthonormal*:
  **assumes** *U*: *unitary U*
    **and** *i*: *i* < *dim-col U*
    **and** *j*: *j* < *dim-col U*
  **shows** ⟨*Matrix.col U i* | *Matrix.col U j*⟩ = (*if i* = *j then 1 else 0* :: *complex*)
  ⟨*proof*⟩

**lemma** *unitary-rows-orthonormal*:
  **assumes** *U*: *unitary U*
    **and** *i*: *i* < *dim-row U*
    **and** *j*: *j* < *dim-row U*
  **shows** ⟨*Matrix.row U i* | *Matrix.row U j*⟩ = (*if i* = *j then 1 else 0* :: *complex*)
  ⟨*proof*⟩

(g), (h) $U$ is an isometry and preserves inner products:

**lemma** *unitary-isometry*:
  **assumes** *unitary U* **and** *dim-vec v* = *dim-col U*
  **shows** ‖*U* ∗*v v*‖ = ‖*v*‖
⟨*proof*⟩

**lemma** *unitary-preserves-inner-prod*:
  **assumes** *U*: *unitary U*
    **and** *u*: *dim-vec u* = *dim-col U*
    **and** *v*: *dim-vec v* = *dim-col U*
  **shows** ⟨*U* ∗*v u* | *U* ∗*v v*⟩ = ⟨*u* | *v*⟩
  ⟨*proof*⟩

## 8.2 Householder unitaries

**Definition (notes).** For $w \in \mathbb{C}^n$ with $\|w\| = 1$ let $P_w = |w\rangle\langle w|$, the projection onto span$\{w\}$, and $U_w = I - 2P_w$. Then $U_w^* U_w = (I - 2P_w)^2 = I - 4P_w + 4P_w^2 = I$, so $U_w$ is unitary: the *Householder unitary* of $w$.

**Remark.** Geometrically $U_w$ is the reflection through the hyperplane $\{w\}^\perp$.

Formally *householder w* is defined entrywise as $(\delta_{ij} - 2w_i\overline{w_j})$ and shown to equal $I - 2|w\rangle\langle w|$.

**abbreviation** *outer-prod* :: *complex Matrix.vec* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.mat*
    (**infixr** ⊘ *75*)
  **where** *v* ⊘ *w* ≡ |*v*⟩ ∗ *bra-vec w*

**lemma** *bra-vec-index-0*:
  **assumes** *j*: *j* < *dim-vec w*
  **shows** *bra-vec w* $$ (*0*,*j*) = *cnj* (*w* $ *j*)
  ⟨*proof*⟩

**lemma** *outer-prod-index*:
  **fixes** *v w* :: *complex Matrix.vec*
  **assumes** *i*: *i* < *dim-vec v*
    **and** *j*: *j* < *dim-vec w*
  **shows** (*v* ⊘ *w*) $$ (*i*,*j*) = (*v* $ *i*) ∗ *cnj* (*w* $ *j*)
⟨*proof*⟩

**lemma** *outer-prod-mult-vec-index*:
  **fixes** *v w x* :: *complex Matrix.vec*

**assumes** *xdim*: *dim-vec x* = *dim-vec w*
**and** *i*: *i* < *dim-vec v*
**shows** ((*v* ⊘ *w*) ∗$_v$ *x*) $ *i* = (*v* $ *i*) ∗ ⟨*w* | *x*⟩
⟨*proof*⟩

**lemma** *outer-prod-mult-vec*:
**fixes** *v w x* :: *complex Matrix.vec*
**assumes** *xdim*: *dim-vec x* = *dim-vec w*
**shows** (*v* ⊘ *w*) ∗$_v$ *x* = ⟨*w* | *x*⟩ ·$_v$ *v*
⟨*proof*⟩

**lemma** *smult-mat-mult-vec*:
**fixes** *A* :: *complex Matrix.mat* **and** *x* :: *complex Matrix.vec*
**assumes** *A*: *A* ∈ *carrier-mat nr nc*
**and** *x*: *x* ∈ *carrier-vec nc*
**shows** (*a* ·$_m$ *A*) ∗$_v$ *x* = *a* ·$_v$ (*A* ∗$_v$ *x*)
⟨*proof*⟩

**lemma** *cpx-vec-length-zero-iff*:
**fixes** *v* :: *complex Matrix.vec*
**shows** ‖*v*‖ = *0* ⟷ *v* = *0*$_v$ (*dim-vec v*)
⟨*proof*⟩

⟨*v*|*v*⟩ = 0 iff *v* = 0.

**lemma** *inner-prod-self-eq-zero-iff*:
**fixes** *v* :: *complex Matrix.vec*
**shows** ⟨*v* | *v*⟩ = *0* ⟷ *v* = *0*$_v$ (*dim-vec v*)
⟨*proof*⟩

**lemma** *cpx-vec-length-smult*:
**fixes** *a* :: *complex* **and** *v* :: *complex Matrix.vec*
**shows** ‖*a* ·$_v$ *v*‖ = *cmod a* ∗ ‖*v*‖
⟨*proof*⟩

**lemma** *inner-prod-right-smult′*:
**fixes** *u v* :: *complex Matrix.vec*
**shows** ⟨*u* | *a* ·$_v$ *v*⟩ = *a* ∗ ⟨*u* | *v*⟩
⟨*proof*⟩

**lemma** *inner-prod-left-smult′*:
**fixes** *u v* :: *complex Matrix.vec*
**assumes** *uv*: *dim-vec u* = *dim-vec v*
**shows** ⟨*a* ·$_v$ *u* | *v*⟩ = *cnj a* ∗ ⟨*u* | *v*⟩
⟨*proof*⟩

**lemma** *inner-prod-right-add′*:
**fixes** *u v w* :: *complex Matrix.vec*
**assumes** *vw*: *dim-vec v* = *dim-vec w*
**shows** ⟨*u* | *v* + *w*⟩ = ⟨*u* | *v*⟩ + ⟨*u* | *w*⟩
⟨*proof*⟩

**lemma** *inner-prod-right-minus′*:
**fixes** *u v w* :: *complex Matrix.vec*
**assumes** *vw*: *dim-vec v* = *dim-vec w*
**shows** ⟨*u* | *v* − *w*⟩ = ⟨*u* | *v*⟩ − ⟨*u* | *w*⟩

⟨*proof*⟩

**lemma** *inner-prod-left-minus'*:
**fixes** *u v w* :: *complex Matrix.vec*
**assumes** *uv*: *dim-vec u* = *dim-vec v*
**and** *uw*: *dim-vec u* = *dim-vec w*
**shows** ⟨*u* − *v* | *w*⟩ = ⟨*u* | *w*⟩ − ⟨*v* | *w*⟩
⟨*proof*⟩

**lemma** *dagger-of-smult*:
**fixes** *γ* :: *complex*
**and** *U* :: *complex Matrix.mat*
**shows** (*γ* ·$_m$ *U*)$^\dagger$ = (*cnj γ*) ·$_m$ (*U*$^\dagger$)
⟨*proof*⟩

The adjoint of an $m \times n$ matrix is $n \times m$.

**lemma** *dagger-carrier*:
**assumes** *M* ∈ *carrier-mat m n*
**shows** (*M*$^\dagger$) ∈ *carrier-mat n m*
⟨*proof*⟩

**definition** *householder* :: *complex Matrix.vec* ⇒ *complex Matrix.mat* **where**
*householder w* ≡
*Matrix.mat* (*dim-vec w*) (*dim-vec w*)
(λ(*i*,*j*). (*if i* = *j then 1 else 0*) − *2* ∗ (*w* $ *i*) ∗ *cnj* (*w* $ *j*))

**notation** *householder* (*U*$_H$)

**lemma** *householder-index*:
**assumes** *i* < *dim-vec w* **and** *j* < *dim-vec w*
**shows** (*householder w*) $$ (*i*,*j*) =
(*if i* = *j then 1 else 0*) − *2* ∗ (*w* $ *i*) ∗ *cnj* (*w* $ *j*)
⟨*proof*⟩

**lemma** *householder-outer-prod*:
*householder w* = *1*$_m$ (*dim-vec w*) − *2* ·$_m$ (*w* ⊘ *w*)
⟨*proof*⟩

**lemma** *householder-hermitian*:
**shows** (*householder w*)$^\dagger$ = *householder w*
⟨*proof*⟩

**lemma** *householder-unitary*:
**assumes** *norm-w*: ‖*w*‖ = (*1* :: *real*)
**shows** *unitary* (*householder w*)
⟨*proof*⟩

**lemma** *householder-mult-vec-index*:
**fixes** *w x* :: *complex Matrix.vec*
**assumes** *xdim*: *dim-vec x* = *dim-vec w*
**and** *i*: *i* < *dim-vec w*
**shows** (*householder w* ∗$_v$ *x*) $ *i* =
*x* $ *i* − *2* ∗ (*w* $ *i*) ∗ ⟨*w* | *x*⟩
⟨*proof*⟩

**lemma** *householder-mult-vec*:
**fixes** $w\ x$ :: *complex Matrix.vec*
**assumes** *xdim*: *dim-vec* $x$ = *dim-vec* $w$
**shows** *householder* $w *_v x$ =
$x - (2 * \langle w \mid x \rangle) \cdot_v w$
⟨*proof*⟩

**Lemma (notes)** (Schur)**.** Given $x, y \in \mathbb{C}^n$ with $\|x\| = \|y\|$ there are $w \in \mathbb{C}^n$ with $\|w\| = 1$ and $\theta$ such that $U_w x = e^{i\theta} y$.

The formal statement writes $e^{i\theta}$ as *Complex* (*cos* $\vartheta$) (*sin* $\vartheta$); the case $x = 0$ is treated separately.

**lemma** *schur-lemma-zero-case*:
**fixes** $x\ y$ :: *complex Matrix.vec*
**assumes** *n-pos*: $n > 0$
**and** *xdim*: *dim-vec* $x = n$
**and** *ydim*: *dim-vec* $y = n$
**and** *norm-eq*: $\|x\| = \|y\|$
**and** *x-zero-norm*: $\|x\| = 0$
**shows** $\exists\, w$ :: *complex Matrix.vec*. $\exists\, \vartheta$ :: *real*.
*dim-vec* $w = n \wedge \|w\| = 1 \wedge$
*householder* $w *_v x$ = *Complex* (*cos* $\vartheta$) (*sin* $\vartheta$) $\cdot_v y$
⟨*proof*⟩

**thm** *of-real-divide*[**where** $x$ = *cmod* ($\langle y \mid x \rangle$) ^ 2 **and** $y$ = *cmod* ($\langle y \mid x \rangle$), *symmetric*]

**lemma** *schur-lemma*:
**fixes** $x\ y$ :: *complex Matrix.vec*
**assumes** *n-pos*: $n > 0$
**and** *xdim*: *dim-vec* $x = n$
**and** *ydim*: *dim-vec* $y = n$
**and** *norm-eq*: $\|x\| = \|y\|$
**shows** $\exists\, w$ :: *complex Matrix.vec*. $\exists\, \vartheta$ :: *real*.
*dim-vec* $w = n \wedge \|w\| = 1 \wedge$
*householder* $w *_v x$ = *Complex* (*cos* $\vartheta$) (*sin* $\vartheta$) $\cdot_v y$
⟨*proof*⟩

**Theorem (notes)** (Schur)**.** For every $A \in M_n$ there is a unitary $U$ such that $U^* A U$ is upper triangular.

The notes prove this by induction, peeling off one eigenvector with the Householder unitary of Schur's lemma and recursing on the $(n-1) \times (n-1)$ block. The formal proof follows the same route: *schur-triangular-similar* is the similarity version available from the Jordan-Normal-Form library, and the block-matrix lemmas that follow assemble the unitary step by step.

**lemma** *schur-triangular-similar*:
**fixes** $A$ :: *complex Matrix.mat*
**assumes** *sq*: $A \in$ *carrier-mat n n*
**shows** $\exists\, T$ :: *complex Matrix.mat*.
$T \in$ *carrier-mat n n* $\wedge$
*upper-triangular* $T \wedge$
*similar-mat* $A\ T$
⟨*proof*⟩

**lemma** *cpx-mat-cnj-four-block*:
**assumes** *A*: $A \in$ *carrier-mat nr1 nc1*
**and** *B*: $B \in$ *carrier-mat nr1 nc2*

**and** $C$: $C \in$ *carrier-mat nr2 nc1*
**and** $D$: $D \in$ *carrier-mat nr2 nc2*
**shows** $(\textit{four-block-mat}\ A\ B\ C\ D)^\star =$
$\textit{four-block-mat}\ (A^\star)\ (B^\star)\ (C^\star)\ (D^\star)$
⟨*proof*⟩

**lemma** *dagger-four-block-mat-complex*:
**assumes** $A$: $A \in$ *carrier-mat nr1 nc1*
**and** $B$: $B \in$ *carrier-mat nr1 nc2*
**and** $C$: $C \in$ *carrier-mat nr2 nc1*
**and** $D$: $D \in$ *carrier-mat nr2 nc2*
**shows** $(\textit{four-block-mat}\ A\ B\ C\ D)^\dagger =$
$\textit{four-block-mat}\ (A^\dagger)\ (C^\dagger)\ (B^\dagger)\ (D^\dagger)$
⟨*proof*⟩

**lemma** *dagger-zero-mat*[*simp*]:
$(0_m\ m\ n :: \textit{complex Matrix.mat})^\dagger = 0_m\ n\ m$
⟨*proof*⟩

**lemma** *block-diag-mult*:
**fixes** $A\ B\ D\ E ::$ *complex Matrix.mat*
**assumes** $A$: $A \in$ *carrier-mat n n*
**and** $B$: $B \in$ *carrier-mat n n*
**and** $D$: $D \in$ *carrier-mat m m*
**and** $E$: $E \in$ *carrier-mat m m*
**shows** $\textit{four-block-mat}\ A\ (0_m\ n\ m)\ (0_m\ m\ n)\ D *$
$\textit{four-block-mat}\ B\ (0_m\ n\ m)\ (0_m\ m\ n)\ E =$
$\textit{four-block-mat}\ (A * B)\ (0_m\ n\ m)\ (0_m\ m\ n)\ (D * E)$
⟨*proof*⟩

**lemma** *block-diag-unitary*:
**fixes** $U\ V ::$ *complex Matrix.mat*
**assumes** $U$: $U \in$ *carrier-mat n n*
**and** $V$: $V \in$ *carrier-mat m m*
**and** $uU$: *unitary U*
**and** $uV$: *unitary V*
**shows** $\textit{unitary}\ (\textit{four-block-mat}\ U\ (0_m\ n\ m)\ (0_m\ m\ n)\ V)$
⟨*proof*⟩

**lemma** *complex-matrix-has-eigenvector*:
**fixes** $A ::$ *complex Matrix.mat*
**assumes** *sq*: $A \in$ *carrier-mat n n*
**and** *n-pos*: $n > 0$
**shows** $\exists\, e\ v.$ *eigenvector A v e*
⟨*proof*⟩

**lemma** *orthonormal-mat-unitary*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *full*: *length vs* $= n$
**shows** $\textit{unitary}\ (\textit{mat-of-cols}\ n\ vs)$
⟨*proof*⟩

**lemma** *eigenvec-normalize*:
**fixes** $A ::$ *complex Matrix.mat* **and** $v ::$ *complex Matrix.vec*
**assumes** *sq*: $A \in$ *carrier-mat n n*
**and** *vdim*: $v \in$ *carrier-vec n*
**and** *vne*: $v \neq 0_v\ n$
**and** *ev*: $A *_v\ v = e \cdot_v v$

**defines** $v_1 \equiv (1 \;/\; \|v\|) \cdot_v v$
**shows** $A *_v v_1 = e \cdot_v v_1$ **and** $\|v_1\| = 1$
⟨*proof*⟩

**definition** *vec-normalize* :: *complex Matrix.vec* ⇒ *complex Matrix.vec* **where**
*vec-normalize v* = (*if v* = $0_v$ (*dim-vec v*) *then v else* (1 / ‖*v*‖) ·$_v$ *v*)

**lemma** *normalized-vec-norm*:
**assumes** *vdim*: $v \in$ *carrier-vec n*
**and** *vne*: $v \neq 0_v$ *n*
**shows** ‖*vec-normalize v*‖ = 1
⟨*proof*⟩

**lemma** *four-block-mat-mult-left-block-diag*:
**fixes** *U* :: *complex Matrix.mat*
**and** *A11* :: *complex Matrix.mat*
**and** *A12* :: *complex Matrix.mat*
**and** *A21* :: *complex Matrix.mat*
**and** *A22* :: *complex Matrix.mat*
**assumes** *U*: *U* ∈ *carrier-mat n n*
**and** *A11*: *A11* ∈ *carrier-mat 1 1*
**and** *A12*: *A12* ∈ *carrier-mat 1 n*
**and** *A21*: *A21* ∈ *carrier-mat n 1*
**and** *A22*: *A22* ∈ *carrier-mat n n*
**shows**
(*four-block-mat* ($1_m$ 1 :: *complex Matrix.mat*)
($0_m$ 1 *n* :: *complex Matrix.mat*)
($0_m$ *n* 1 :: *complex Matrix.mat*)
*U*)
*
(*four-block-mat A11 A12 A21 A22*)
=
*four-block-mat A11 A12* (*U* ∗ *A21*) (*U* ∗ *A22*)
⟨*proof*⟩

**lemma** *cscalar-prod-is-inner-prod-swap*:
**fixes** *u v* :: *complex Matrix.vec*
**assumes** *u* ∈ *carrier-vec n* **and** *v* ∈ *carrier-vec n*
**shows** *u* ·*c v* = ⟨*v*|*u*⟩
⟨*proof*⟩

**lemma** *vec-normalize-dim*[*simp*]:
*dim-vec* (*vec-normalize v*) = *dim-vec v*
⟨*proof*⟩

**lemma** *vec-normalize-carrier*[*simp*]:
**assumes** *v* ∈ *carrier-vec n*
**shows** *vec-normalize v* ∈ *carrier-vec n*
⟨*proof*⟩

**lemma** *vec-normalize-unit*:
**assumes** *vdim*: *v* ∈ *carrier-vec n* **and** *vnorm*: ‖*v*‖ = 1
**shows** *vec-normalize v* = *v*
⟨*proof*⟩

Any unit vector extends to an orthonormal basis (needed for the induction).

**lemma** *unit-vec-extends-to-onb*:
**fixes** *v* :: *complex Matrix.vec*

**assumes** *vdim*: $v \in$ *carrier-vec* $n$
**and** *vnorm*: $\|v\| = 1$
**shows** $\exists$ *vs*. *orthonormal-list* $n$ *vs* $\wedge$ *length vs* $= n \wedge$ *vs* ! $0 = v$
⟨*proof*⟩

**lemma** *schur-induct*:
$\forall A ::$ *complex Matrix.mat*.
$A \in$ *carrier-mat* $n\ n \longrightarrow$
$(\exists U ::$ *complex Matrix.mat*. $\exists T ::$ *complex Matrix.mat*.
$U \in$ *carrier-mat* $n\ n \wedge$
$T \in$ *carrier-mat* $n\ n \wedge$
*unitary* $U \wedge$
*upper-triangular* $T \wedge$
$(((U^\dagger) * A) * U) = T)$
⟨*proof*⟩

**theorem** *schur-theorem*:
**fixes** $A ::$ *complex Matrix.mat*
**assumes** *sq*: $A \in$ *carrier-mat* $n\ n$
**shows** $\exists U ::$ *complex Matrix.mat*. $\exists T ::$ *complex Matrix.mat*.
$U \in$ *carrier-mat* $n\ n \wedge$
$T \in$ *carrier-mat* $n\ n \wedge$
*unitary* $U \wedge$
*upper-triangular* $T \wedge$
$(((U^\dagger) * A) * U) = T$
⟨*proof*⟩

**Remark.** The characteristic polynomial of $A$ is $p_A(t) = \det(tI - A)$, a polynomial of degree $n$.

**Corollary (notes).** $\mathrm{Tr}(A) = \lambda_1 + \cdots + \lambda_n$, where the $\lambda_j$ are the eigenvalues of $A$.

Formally the eigenvalues are listed as the roots of the characteristic polynomial, $p_A(t) = \prod_j (t - \lambda_j)$, read off the diagonal of the Schur form.

**corollary** *trace-eq-sum-eigenvalues*:
**fixes** $A ::$ *complex Matrix.mat*
**assumes** *sq*: $A \in$ *carrier-mat* $n\ n$
**obtains** *eigs* :: *complex list*
**where** *length eigs* $= n$
**and** *Tr A* $=$ *sum-list eigs*
**and** *char-poly A* $= (\prod e \leftarrow$ *eigs*. [:$-e$, *1*:])
⟨*proof*⟩

# 9 Day 4 (2 September 2011): Hermitian matrices; positive semidefinite and positive definite matrices

## 9.1 Hermitian matrices

**Definition (notes).** $H \in M_n$ is *Hermitian* (self-adjoint) if $H = H^*$.

**Theorem (notes).** The following are equivalent: (a) $H$ is Hermitian; (b) there is a unitary $U$ with $U^*HU = D$ diagonal with real entries; (c) $\mathbb{C}^n$ has an orthonormal basis of eigenvectors of $H$ with real eigenvalues; (d) $\langle x|Hx\rangle \in \mathbb{R}$ for all $x \in \mathbb{C}^n$.

Formally *hermitian H* is $H^\dagger = H$; (a)⇒(b) is the spectral theorem *hermitian-spectral-decomp* below, (a)⇔(d) is *hermitian-iff-real-quad-form*, and (c) is read off from (b): the columns of $U$

are an orthonormal eigenbasis. The proof of (d)⇒(a) needs the following lemma, which the notes note fails over $\mathbb{R}$ (take $T = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$).

**definition** *hermitian* :: *complex Matrix.mat* ⇒ *bool* **where**
*hermitian A* ≡ $A^\dagger = A$

**lemma** *hermitian-square*:
**assumes** *hermitian A*
**shows** *dim-row A* = *dim-col A*
⟨*proof*⟩

**lemma** *hermitian-index*:
**assumes** *hermitian A*
**and** *i* < *dim-row A* **and** *j* < *dim-col A*
**shows** *A* $$ (*i*,*j*) = *cnj* (*A* $$ (*j*,*i*))
⟨*proof*⟩

**lemma** *inner-T-double-sum*:
**fixes** *T* :: *complex Matrix.mat* **and** *u v* :: *complex Matrix.vec*
**assumes** *T-carr*: *T* ∈ *carrier-mat n n*
**and** *udim*: *dim-vec u* = *n* **and** *vdim*: *dim-vec v* = *n*
**shows** ⟨*u* | *T* ∗$_v$ *v*⟩ =
(∑ *k* = *0*..<*n*. ∑ *l* = *0*..<*n*. *cnj* (*u* $ *k*) ∗ *T* $$ (*k*, *l*) ∗ *v* $ *l*)
⟨*proof*⟩

**lemma** *inner-unit-T-unit*:
**fixes** *T* :: *complex Matrix.mat*
**assumes** *T-carr*: *T* ∈ *carrier-mat n n*
**and** *p*: *p* < *n* **and** *q*: *q* < *n*
**shows** ⟨*unit-vec n p* | *T* ∗$_v$ *unit-vec n q*⟩ = *T* $$ (*p*, *q*)
⟨*proof*⟩

**lemma** *sum-indicator-T-indicator*:
**fixes** *T* :: *complex Matrix.mat*
**assumes** *p*: *p* < *n* **and** *q*: *q* < *n*
**shows** (∑ *k* = *0*..<*n*. ∑ *l* = *0*..<*n*.
(*if k* = *p then* (*1* :: *complex*) *else 0*) ∗ *T* $$ (*k*, *l*) ∗
(*if l* = *q then* (*1* :: *complex*) *else 0*))
= *T* $$ (*p*, *q*)
⟨*proof*⟩

**lemma** *polarization-helper*:
**fixes** *T* :: *complex Matrix.mat* **and** *a b c d* :: *complex*
**assumes** *sq*: *T* ∈ *carrier-mat n n*
**and** *i*: *i* < *n* **and** *j*: *j* < *n*
**shows** ⟨*a* ·$_v$ *unit-vec n i* + *b* ·$_v$ *unit-vec n j*
| *T* ∗$_v$ (*c* ·$_v$ *unit-vec n i* + *d* ·$_v$ *unit-vec n j*)⟩
= *cnj a* ∗ *c* ∗ *T* $$ (*i*, *i*) + *cnj a* ∗ *d* ∗ *T* $$ (*i*, *j*)
+ *cnj b* ∗ *c* ∗ *T* $$ (*j*, *i*) + *cnj b* ∗ *d* ∗ *T* $$ (*j*, *j*)
⟨*proof*⟩

**Lemma (notes).** For $T \in M_n$, if $\langle x|Tx\rangle = 0$ for all $x \in \mathbb{C}^n$, then $T = 0$.

**lemma** *quadratic-form-zero*:

**fixes** $T$ :: *complex Matrix.mat*
**assumes** *sq*: $T \in$ *carrier-mat n n*
**and** *zero*: $\forall x$ :: *complex Matrix.vec*. *dim-vec* $x = n \longrightarrow \langle x \mid T *_v x\rangle = 0$
**shows** $T = 0_m\ n\ n$
⟨*proof*⟩

**lemma** *mult-mat-vec-index*:
**fixes** $A$ :: $'a$::{*comm-ring-1*} *Matrix.mat*
**and** $x$ :: $'a$ *Matrix.vec*
**assumes** $A$: $A \in$ *carrier-mat m n*
**and** $x$: $x \in$ *carrier-vec n*
**and** $j$: $j < m$
**shows** $(A *_v x)\ \$\ j = (\sum i = 0..<n.\ A\ \$\$\ (j,\ i) * x\ \$\ i)$
⟨*proof*⟩

The adjoint moves across the inner product, $\langle x|My\rangle = \langle M^\dagger x|y\rangle$: the defining property of $M^\dagger$.

**lemma** *inner-prod-adjoint-swap*:
**fixes** $M$ :: *complex Matrix.mat*
**and** $x\ y$ :: *complex Matrix.vec*
**assumes** $M$: $M \in$ *carrier-mat n m*
**and** $x$: *dim-vec* $x = n$
**and** $y$: *dim-vec* $y = m$
**shows** $\langle x \mid M *_v y\rangle = \langle M^\dagger *_v x \mid y\rangle$
⟨*proof*⟩

Sums and real multiples of Hermitian matrices are Hermitian; the identity is Hermitian.

**lemma** *hermitian-plus*:
**fixes** $A\ B$ :: *complex Matrix.mat*
**assumes** *Aherm*: *hermitian A* **and** *Bherm*: *hermitian B*
**and** *Ac*: $A \in$ *carrier-mat k k* **and** *Bc*: $B \in$ *carrier-mat k k*
**shows** *hermitian* $(A + B)$
⟨*proof*⟩

**lemma** *hermitian-identity*:
*hermitian* $(1_m\ k$ :: *complex Matrix.mat*$)$
⟨*proof*⟩

**lemma** *hermitian-real-scale*:
**fixes** $A$ :: *complex Matrix.mat*
**assumes** *hermitian A* $A \in$ *carrier-mat k k*
**shows** *hermitian* (*complex-of-real d* $\cdot_m$ $A$)
⟨*proof*⟩

The converses (g)⇒(a) and (h)⇒(a) of the Day 3 theorem on unitaries, postponed until the lemma $\langle x|Tx\rangle = 0\ \forall x \Rightarrow T = 0$ was available: a square matrix that preserves lengths, or inner products, is unitary (the library proves the length-preserving case in its own matrix notation as *length-preserving-is-unitary*).

**lemma** *unitary-of-isometry*:
**fixes** $U$ :: *complex Matrix.mat*
**assumes** $U$: $U \in$ *carrier-mat n n*
**and** *iso*: $\bigwedge v$ :: *complex Matrix.vec*. *dim-vec* $v = n \Longrightarrow \|U *_v v\| = \|v\|$
**shows** *unitary U*
⟨*proof*⟩

**lemma** *unitary-of-inner-prod-preserving*:
**fixes** $U$ :: *complex Matrix.mat*
**assumes** $U$: $U \in$ *carrier-mat n n*

**and** *pres*: $\bigwedge u\ v ::$ *complex Matrix.vec*. *dim-vec* $u = n \Longrightarrow$ *dim-vec* $v = n \Longrightarrow$
$\langle U *_v u \mid U *_v v\rangle = \langle u \mid v\rangle$
**shows** *unitary U*
⟨*proof*⟩

**lemma** *hermitian-iff-real-quad-form*:
**assumes** *sq*: $A \in$ *carrier-mat n n*
**shows** *hermitian* $A \longleftrightarrow (\forall x ::$ *complex Matrix.vec*. *dim-vec* $x = n \longrightarrow \langle x \mid A *_v x\rangle \in \mathbb{R})$
⟨*proof*⟩

## 9.2 The spectral theorem for Hermitian matrices

(a)⇒(b) of the theorem above: by Schur, $U^*HU = T$ is upper triangular, and $T^* = (U^*HU)^* = T$ forces $T$ to be diagonal with real entries. *diag-mat-of-real D* is the complex diagonal matrix with the real list $D$ on its diagonal.

**definition** *diag-mat-of-real* :: *real list* ⇒ *complex Matrix.mat* **where**
*diag-mat-of-real D* =
*Matrix.mat* (*length D*) (*length D*)
(λ(*i*,*j*). *if i* = *j then complex-of-real* (*D* ! *i*) *else 0*)

**lemma** *dim-row-diag-mat-of-real*[*simp*]: *dim-row* (*diag-mat-of-real D*) = *length D*
⟨*proof*⟩

**lemma** *dim-col-diag-mat-of-real*[*simp*]: *dim-col* (*diag-mat-of-real D*) = *length D*
⟨*proof*⟩

**lemma** *diag-mat-of-real-carrier*[*simp*]:
*diag-mat-of-real D* ∈ *carrier-mat* (*length D*) (*length D*)
⟨*proof*⟩

**lemma** *diag-mat-of-real-index*[*simp*]:
**assumes** *i* < *length D* **and** *j* < *length D*
**shows** *diag-mat-of-real D* $$ (*i*, *j*) =
(*if i* = *j then complex-of-real* (*D* ! *i*) *else 0*)
⟨*proof*⟩

**lemma** *diag-mat-of-real-mult-vec-index*:
**fixes** *y* :: *complex Matrix.vec*
**assumes** *ydim*: *dim-vec y* = *length D*
**and** *i*: *i* < *length D*
**shows** (*diag-mat-of-real D* $*_v$ *y*) $ *i* = *complex-of-real* (*D* ! *i*) ∗ *y* $ *i*
⟨*proof*⟩

**lemma** *diag-mat-of-real-quad-Re-nonneg*:
**fixes** *y* :: *complex Matrix.vec*
**assumes** *ydim*: *dim-vec y* = *length D*
**and** *D-nonneg*: $\bigwedge i.$ *i* < *length D* ⟹ 0 ≤ *D* ! *i*
**shows** 0 ≤ *Re* ⟨*y* | *diag-mat-of-real D* $*_v$ *y*⟩
⟨*proof*⟩

**theorem** *hermitian-spectral-decomp*:
**fixes** *H* :: *complex Matrix.mat*
**assumes** *sq*: *H* ∈ *carrier-mat n n*
**and** *herm*: *hermitian H*
**shows** ∃(*U* :: *complex Matrix.mat*) (*D* :: *real list*).
*U* ∈ *carrier-mat n n* ∧
*unitary U* ∧

*length D = n ∧*
*(U†) ∗ H ∗ U = diag-mat-of-real D*
⟨*proof*⟩

**corollary** *hermitian-real-eigenvalues*:
**fixes** *H* :: *complex Matrix.mat*
**assumes** *sq*: *H* ∈ *carrier-mat n n*
**and** *herm*: *hermitian H*
**and** *eig*: *H ∗v v = lam ·v v*
**and** *v-nonzero*: *v ≠ 0v n*
**and** *vdim*: *dim-vec v = n*
**shows** *lam* ∈ ℝ
⟨*proof*⟩

## 9.3 Positive semidefinite and positive definite matrices

**Definition (notes).** $P \in M_n$ is *positive semidefinite*, $P \geq 0$, if $\langle x|Px\rangle \geq 0$ for all $x \in \mathbb{C}^n$, and *positive definite*, $P > 0$, if $\langle x|Px\rangle > 0$ for all $x \neq 0$.

By part (d) of the previous theorem, $P \geq 0$ implies $P = P^*$. The formal predicates *psd* and *pd* carry Hermiticity as an explicit conjunct together with the real-part inequality; by *hermitian-iff-real-quad-form* this is equivalent to the notes' definition.

**Theorem (notes).** $P \geq 0$ if and only if $P = P^*$ and all eigenvalues of $P$ are non-negative.

**Corollary (notes).** $P > 0$ if and only if $P = P^*$ and all eigenvalues of $P$ are strictly positive.

**Theorem (notes).** $P \geq 0$ if and only if $P = \sum_{i=1}^{m} |v_i\rangle\langle v_i|$ for some vectors $v_1, \ldots, v_m$.

Key: if $P = \sum_i |v_i\rangle\langle v_i|$ then $\langle x|Px\rangle = \sum_i |\langle v_i|x\rangle|^2$.

**Theorem (notes).** Let $P = \sum_{i=1}^{m} |v_i\rangle\langle v_i| \in M_n$. Then $P > 0$ if and only if $\mathrm{span}\{v_1, \ldots, v_m\} = \mathbb{C}^n$.

The notes' form of the definition is *psd-iff-quad-form-nonneg*; the corollary on positive definite matrices is *pd-iff-pos-eigenvalues* (proved after the square root); the last theorem is formalised with "the $v_i$ span $\mathbb{C}^n$" phrased as "no nonzero $x$ is orthogonal to every $v_i$".

**definition** *psd* :: *complex Matrix.mat ⇒ bool* **where**
*psd P ≡ hermitian P ∧*
*(∀ x :: complex Matrix.vec. dim-vec x = dim-row P ⟶*
*Re (⟨x | P ∗v x⟩) ≥ 0)*

**definition** *pd* :: *complex Matrix.mat ⇒ bool* **where**
*pd P ≡ hermitian P ∧*
*(∀ x :: complex Matrix.vec. dim-vec x = dim-row P ⟶*
*x ≠ 0v (dim-row P) ⟶*
*Re (⟨x | P ∗v x⟩) > 0)*

**notation** *psd* (- ≥M 0)
**notation** *pd* (- >M 0)

**lemma** *psd-hermitian*: *psd P ⟹ hermitian P*
⟨*proof*⟩

The notes' definition, without the explicit Hermiticity conjunct: $P \geq 0$ iff every quadratic form $\langle x|Px\rangle$ is a non-negative real number.

**lemma** *psd-iff-quad-form-nonneg*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** *sq*: $P \in$ *carrier-mat n n*
**shows** *psd* $P \longleftrightarrow$
$(\forall\, x ::$ *complex Matrix.vec. dim-vec* $x = n \longrightarrow$
$\langle x \mid P *_v x\rangle \in \mathbb{R} \wedge 0 \leq$ *Re* $(\langle x \mid P *_v x\rangle))$
⟨*proof*⟩

**lemma** *psd-MdagM*:
**assumes** *M*: $M \in$ *carrier-mat n m*
**shows** *psd* $(M^\dagger * M)$
⟨*proof*⟩

$P \geq 0$ iff $P$ is Hermitian with non-negative eigenvalues:

**theorem** *psd-iff-nonneg-eigenvalues*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** *sq*: $P \in$ *carrier-mat n n*
**shows** *psd* $P \longleftrightarrow$
$($*hermitian* $P \wedge$
$(\forall\, mu ::$ *complex*. $\forall\, v ::$ *complex Matrix.vec.*
$v \neq 0_v\ n \wedge$ *dim-vec* $v = n \wedge P *_v v = mu \cdot_v v \longrightarrow$ *Re* $mu \geq 0))$
⟨*proof*⟩

$P \geq 0$ iff $P = \sum_i |v_i\rangle\langle v_i|$. First the direction $\Leftarrow$:

**lemma** *outer-sum-psd*:
**assumes** *len*: *length vs* $= k$
**and** *vdim*: $\forall\, v \in$ *set vs*. *dim-vec* $v = n$
**shows** *psd* (*outer-sum n vs*)
⟨*proof*⟩

**lemma** *hermitian-spectral-decomp-eq*:
**assumes** *Uc*: $U \in$ *carrier-mat n n*
**and** *Pcar*: $P \in$ *carrier-mat n n*
**and** *unit*: *unitary* $U$
**and** *spec*: $U^\dagger * P * U =$ *diag-mat-of-real* $D$
**shows** $P = U *$ *diag-mat-of-real* $D * (U^\dagger)$
⟨*proof*⟩

**lemma** *hermitian-spectral-decomp-right*:
**assumes** *Uc*: $U \in$ *carrier-mat n n*
**and** *Pcar*: $P \in$ *carrier-mat n n*
**and** *unit*: *unitary* $U$
**and** *spec*: $U^\dagger * P * U =$ *diag-mat-of-real* $D$
**shows** $P * U = U *$ *diag-mat-of-real* $D$
⟨*proof*⟩

The direction $\Rightarrow$, from the spectral theorem: $P = \sum_j \lambda_j |u_j\rangle\langle u_j| = \sum_j |\lambda_j^{1/2} u_j\rangle\langle\lambda_j^{1/2} u_j|$.

The spectral theorem for positive matrices: $P = U\,\mathrm{diag}(D)\,U^\dagger$ with $D \geq 0$.

**lemma** *psd-spectral-decomp-nonneg*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** *sq*: $P \in$ *carrier-mat n n*
**and** *pos*: *psd* $P$
**shows** $\exists\,(U ::$ *complex Matrix.mat*$)$ $(D ::$ *real list*$)$.
$U \in$ *carrier-mat n n* $\wedge$ *unitary* $U \wedge$ *length* $D = n \wedge$
$(\forall\, i < n.\ 0 \leq D\ !\ i) \wedge$

$P = U * \mathit{diag\text{-}mat\text{-}of\text{-}real}\ D * (U^\dagger)$
⟨*proof*⟩

**theorem** *psd-kraus-decomp*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** *sq*: $P \in \mathit{carrier\text{-}mat}\ n\ n$
**and** *pos*: *psd* $P$
**shows** $\exists\, vs :: \mathit{complex\ Matrix.vec\ list}.$
$\mathit{length}\ vs = n \wedge$
$(\forall\, v \in \mathit{set}\ vs.\ \mathit{dim\text{-}vec}\ v = n) \wedge$
$\mathit{outer\text{-}sum}\ n\ vs = P$
⟨*proof*⟩

$P = \sum_i |v_i\rangle\langle v_i|$ is positive definite iff the $v_i$ span $\mathbb{C}^n$:

Entries of $\sum_i |v_i\rangle\langle v_i|$ for an arbitrary list of vectors.

**lemma** *outer-sum-entry-sum-list*:
**assumes** *vs-dim*: $\forall\, v \in \mathit{set}\ vs.\ \mathit{dim\text{-}vec}\ v = n$
**and** *i-lt*: $i < n$ **and** *j-lt*: $j < n$
**shows** $\mathit{outer\text{-}sum}\ n\ vs\ \$\$\ (i, j) =$
$\mathit{sum\text{-}list}\ (\mathit{map}\ (\lambda v.\ v\ \$\ i * \mathit{cnj}\ (v\ \$\ j))\ vs)$
⟨*proof*⟩

**lemma** *mat-of-cols-dagger-mult-vec-index*:
**assumes** *xdim*: $\mathit{dim\text{-}vec}\ x = n$
**and** *vs-dim*: $\forall\, v \in \mathit{set}\ vs.\ \mathit{dim\text{-}vec}\ v = n$
**and** *i*: $i < \mathit{length}\ vs$
**shows** $(((\mathit{mat\text{-}of\text{-}cols}\ n\ vs)^\dagger) *_v x)\ \$\ i = \langle vs\ !\ i \mid x\rangle$
⟨*proof*⟩

**lemma** *inner-prod-outer-sum-general*:
**assumes** *xdim*: $\mathit{dim\text{-}vec}\ x = n$
**and** *vs-dim*: $\forall\, v \in \mathit{set}\ vs.\ \mathit{dim\text{-}vec}\ v = n$
**shows** $\langle x \mid \mathit{outer\text{-}sum}\ n\ vs *_v x\rangle =$
$\langle((\mathit{mat\text{-}of\text{-}cols}\ n\ vs)^\dagger) *_v x \mid$
$((\mathit{mat\text{-}of\text{-}cols}\ n\ vs)^\dagger) *_v x\rangle$
⟨*proof*⟩

**lemma** *diag-mat-of-real-hermitian*:
**shows** *hermitian* (*diag-mat-of-real D*)
⟨*proof*⟩

$U\,\mathrm{diag}(D)\,U^\dagger$ is positive whenever the real entries of $D$ are non-negative.

**lemma** *psd-U-diag-Udag*:
**fixes** $U$ :: *complex Matrix.mat* **and** $D$ :: *real list*
**assumes** *Uc*: $U \in \mathit{carrier\text{-}mat}\ n\ n$
**and** *Dlen*: $\mathit{length}\ D = n$
**and** *Dnn*: $\bigwedge i.\ i < n \Longrightarrow 0 \le D\ !\ i$
**shows** $\mathit{psd}\ (U * \mathit{diag\text{-}mat\text{-}of\text{-}real}\ D * (U^\dagger))$
⟨*proof*⟩

**lemma** *diag-mat-of-real-sqrt-square*:
**assumes** *D-nonneg*: $\bigwedge i.\ i < \mathit{length}\ D \Longrightarrow 0 \le D\ !\ i$
**shows** $\mathit{diag\text{-}mat\text{-}of\text{-}real}\ (\mathit{map}\ \mathit{sqrt}\ D) * \mathit{diag\text{-}mat\text{-}of\text{-}real}\ (\mathit{map}\ \mathit{sqrt}\ D) = \mathit{diag\text{-}mat\text{-}of\text{-}real}\ D$
⟨*proof*⟩

**theorem** *pd-iff-spanning*:
**fixes** $P$ :: *complex Matrix.mat*

**assumes** *sq*: $P \in$ *carrier-mat n n*
**and** *decomp*: $P =$ *outer-sum n vs*
**and** *vs-dim*: $\forall\, v \in$ *set vs*. *dim-vec* $v = n$
**shows** *pd* $P \longleftrightarrow$
$(\forall\, x ::$ *complex Matrix.vec*. *dim-vec* $x = n \longrightarrow x \neq 0_v\ n \longrightarrow$
$(\exists\, v \in$ *set vs*. $\langle v \mid x\rangle \neq 0))$
⟨*proof*⟩

### 9.4 The positive square root

**Remark** (Day 8)**.** Given $P \geq 0$ with $U^*PU = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$, $\lambda_i \geq 0$, put $P^{1/2} = U\,\mathrm{diag}(\lambda_1^{1/2}, \ldots, \lambda_n^{1/2})U^*$. Then $P^{1/2} \geq 0$ and $(P^{1/2})^2 = P$.

*psd-sqrt P* is defined as a choice of such a square root; *psd-sqrt-exists* is the existence statement and *psd-sqrt-props* the defining properties.

**definition** *psd-sqrt* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*psd-sqrt P* ≡ (*SOME Q*. *Q* ∈ *carrier-mat* (*dim-row P*) (*dim-row P*) ∧ *psd Q* ∧ *Q* ∗ *Q* = *P*)

**theorem** *psd-sqrt-exists*:
**fixes** *P* :: *complex Matrix.mat*
**assumes** *sq*: *P* ∈ *carrier-mat n n*
**and** *pos*: *psd P*
**shows** ∃ *Q* :: *complex Matrix.mat*.
*Q* ∈ *carrier-mat n n* ∧ *psd Q* ∧ *Q* ∗ *Q* = *P*
⟨*proof*⟩
**lemma** *psd-sqrt-props*:
**fixes** *P* :: *complex Matrix.mat*
**assumes** *sq*: *P* ∈ *carrier-mat n n*
**and** *pos*: *psd P*
**shows** *psd-sqrt P* ∈ *carrier-mat n n*
**and** *psd* (*psd-sqrt P*)
**and** *psd-sqrt P* ∗ *psd-sqrt P* = *P*
⟨*proof*⟩

The corollary of Day 4: $P > 0$ iff $P = P^*$ and all eigenvalues are strictly positive.

**lemma** *pd-iff-pos-eigenvalues*:
**fixes** *P* :: *complex Matrix.mat*
**assumes** *sq*: *P* ∈ *carrier-mat n n*
**shows** *pd P* ⟷
(*hermitian P* ∧
(∀ *mu* :: *complex*. ∀ *v* :: *complex Matrix.vec*.
*v* ≠ $0_v$ *n* ∧ *dim-vec v* = *n* ∧ *P* ∗$_v$ *v* = *mu* ·$_v$ *v* ⟶ 0 < *Re mu*))
⟨*proof*⟩

## 10 Day 5 (7 September 2011): tests for positivity; direct sums and partitioned matrices

### 10.1 Practical tests for positivity

Given $P = P^*$, how does one decide whether $P \geq 0$ or $P > 0$? The notes give two tests. (1) Sylvester's criterion, in terms of the leading principal submatrices $P_k = (p_{ij})_{1 \leq i,j \leq k}$ (the upper left $k \times k$ corners):

**Theorem (notes)** (Sylvester's criterion)**.** Let $P = P^* \in M_n$. Then $P > 0$ if and only if $\det P_k > 0$ for $k = 1, \ldots, n$.

Each $\det P_k$ is real, as $P_k$ is Hermitian. The same test with $\geq 0$ does *not* characterise $P \geq 0$: $\begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}$ has leading minors 0 and 0 but is not positive. For $n = 2$ the criterion says $\begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} > 0$ iff $a > 0$ and $ac - |b|^2 > 0$. The notes state it without proof. It is not reproved in this development: it is imported from the AFP entry *Two Hermitian Results* and restated for *pd* and *hermitian* as `pd_iff_leading_minors_pos` (for $P \in M_n$ Hermitian, $P >_M 0$ iff $\det P_k > 0$ for all $k \leq n$) in the theory `Day05_Sylvester_Criterion`. That theory forms the separate session `Vern_Paulsen_QC_Sylvester` with its own document (`Sylvester_Criterion.pdf`), because the AFP entry builds on HOL-Analysis, HOL-Probability and QHLProver, which this document's sessions do not load. (2) Cholesky's algorithm:

**Theorem (notes).** If $P = (p_{ij})$ is $n \times n$ with $P = P^*$, then $P \geq 0$ if and only if $P - \left(\frac{p_{i1}p_{1j}}{p_{11}}\right)_{ij} \geq 0$.

The subtracted matrix agrees with $P$ in the first row and column, so the difference is really an $(n-1) \times (n-1)$ matrix and the test can be repeated; when $P \geq 0$ the algorithm also writes $P$ as a sum of rank-one matrices. The formal theorem adds the hypothesis $p_{11} > 0$, which the notes leave implicit (for $p_{11} = 0$ the quotient is undefined).

**theorem** *cholesky-reduction*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** *sq*: $P \in$ *carrier-mat n n*
**and** *herm*: *hermitian P*
**and** *n-pos*: $0 < n$
**and** *p11-pos*: $P \$\$ (0,0) > 0$
**shows** *psd P* $\longleftrightarrow$
*psd* (*Matrix.mat* $(n-1)$ $(n-1)$
$(\lambda(i,j).\ P \$\$ (i+1, j+1) -$
$P \$\$ (i+1, 0) * cnj\ (P \$\$ (j+1, 0)) / P \$\$ (0,0)))$
⟨*proof*⟩

## 10.2 Direct sums of vector spaces and partitioned matrices

**Definition (notes).** The *direct sum* of vector spaces $V, W$ is $V \oplus W = \{(v, w) : v \in V, w \in W\}$ with componentwise operations and inner product $\langle (v_1, w_1) | (v_2, w_2) \rangle = \langle v_1 | v_2 \rangle_V + \langle w_1 | w_2 \rangle_W$.

**Proposition (notes).** If $v_1, \ldots, v_n$ is a basis of $V$ and $w_1, \ldots, w_k$ a basis of $W$, then $(v_1, 0), \ldots, (v_n, 0), (0, w_1), \ldots$ is a basis of $V \oplus W$; hence $\dim(V \oplus W) = \dim V + \dim W$.

Key examples: $\mathbb{C}^{n+k} = \mathbb{C}^n \oplus \mathbb{C}^k$, and a linear map $T : \mathbb{C}^{n_1+k_1} \to \mathbb{C}^{n_2+k_2}$ has the block form $T = \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix}$ with $T\binom{v}{w} = \binom{T_{11}v+T_{12}w}{T_{21}v+T_{22}w}$. Formally the direct sum of vectors is the concatenation $v \oplus_v w$, and block matrices are the library's *four-block-mat*.

**definition** *direct-sum-vec* :: *complex Matrix.vec* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.vec* (**infixr** $\oplus_v$ *65*) **where**
$v \oplus_v w \equiv$ *Matrix.vec* (*dim-vec v + dim-vec w*) ($\lambda i$. *if* $i <$ *dim-vec v then v* \$ *i else w* \$ ($i -$ *dim-vec v*))

**lemma** *direct-sum-vec-dim* [*simp*]:
*dim-vec* $(v \oplus_v w) =$ *dim-vec v + dim-vec w*
⟨*proof*⟩

**lemma** *direct-sum-vec-left* [*simp*]:
$i <$ *dim-vec v* $\Longrightarrow$ $(v \oplus_v w)$ \$ $i = v$ \$ $i$
⟨*proof*⟩

**lemma** *direct-sum-vec-right* [*simp*]:

*i* < *dim-vec w* ⟹ (*v* ⊕*v w*) $ (*dim-vec v* + *i*) = *w* $ *i*
⟨*proof*⟩

**lemma** *direct-sum-inner-prod*:
**assumes** *vd1*: *dim-vec v1* = *m* **and** *vd2*: *dim-vec v2* = *m*
**and** *wd1*: *dim-vec w1* = *p* **and** *wd2*: *dim-vec w2* = *p*
**shows** ⟨*v1* ⊕*v w1* | *v2* ⊕*v w2*⟩ = ⟨*v1* | *v2*⟩ + ⟨*w1* | *w2*⟩
⟨*proof*⟩

**corollary** *direct-sum-dim*:
*dim-vec* (*v* ⊕*v w*) = *dim-vec v* + *dim-vec w*
⟨*proof*⟩

**lemma** *block-mat-apply*:
**assumes** *dim-vec v* = *k1* **and** *dim-vec w* = *k2*
**and** *T11* ∈ *carrier-mat n1 k1* **and** *T12* ∈ *carrier-mat n1 k2*
**and** *T21* ∈ *carrier-mat n2 k1* **and** *T22* ∈ *carrier-mat n2 k2*
**shows** *four-block-mat T11 T12 T21 T22* ∗*v* (*v* ⊕*v w*) =
(*T11* ∗*v v* + *T12* ∗*v w*) ⊕*v* (*T21* ∗*v v* + *T22* ∗*v w*)
⟨*proof*⟩

# 11 Day 6 (9 September 2011): tensor products

**Definition (notes).** $B : X \times Y \to Z$ is *bilinear* if $B(x_1 + x_2, y) = B(x_1, y) + B(x_2, y)$, $B(x, y_1 + y_2) = B(x, y_1) + B(x, y_2)$ and $B(\lambda x, y) = B(x, \lambda y) = \lambda B(x, y)$.

The tensor product $X \otimes Y$ is the span of elementary tensors $x \otimes y$ with the universal property: every bilinear $B$ factors as $B(x, y) = L_B(x \otimes y)$ for a unique linear $L_B$. Key relations: $(x_1 + x_2) \otimes y = x_1 \otimes y + x_2 \otimes y$, $x \otimes (y_1 + y_2) = x \otimes y_1 + x \otimes y_2$, $\lambda(x \otimes y) = (\lambda x) \otimes y = x \otimes (\lambda y)$.

**Theorem (notes).** If $\{e_1, \ldots, e_k\}$ and $\{f_1, \ldots, f_m\}$ are bases of $X$ and $Y$, then $\{e_i \otimes f_j\}$ is a basis of $X \otimes Y$; hence $\dim(X \otimes Y) = \dim X \cdot \dim Y$.

**Proposition (notes).** With bases as above, every $u \in X \otimes Y$ is uniquely $u = \sum_j x_j \otimes f_j$ with $x_j \in X$, and uniquely $u = \sum_i e_i \otimes y_i$ with $y_i \in Y$; consequently $X \otimes Y \cong X \oplus \cdots \oplus X$ ($m$ copies).

**Definition (notes).** The *Schmidt rank* of $u \in X \otimes Y$ is the least number of elementary tensors in an expression for $u$.

The formalisation uses the concrete model: $\mathbb{C}^n \otimes \mathbb{C}^m = \mathbb{C}^{nm}$ with $(v \otimes w)_{im+j} = v_i w_j$ (*tensor-vec v w* from the library). The three key relations are the bilinearity lemmas below; the product basis theorem is *tensor-onb* (Day 7); the unique decomposition along a basis of $Y$ is not stated for vectors (its operator analogue is in theory Quantum_Computing11).

**lemma** *tensor-vec-dim*:
**assumes** *vd*: *dim-vec v* = *n dim-vec w* = *m*
**shows** *dim-vec* (*tensor-vec v w*) = *n* ∗ *m*
⟨*proof*⟩

**lemma** *tensor-vec-index*:
**assumes** *dim-vec v* = *n dim-vec w* = *m i* < *n* ∗ *m*
**shows** *tensor-vec v w* $ *i* = *v* $ (*i div m*) ∗ *w* $ (*i mod m*)
⟨*proof*⟩

**lemma** *tensor-vec-bilinear-left*:
**assumes** *vd*: *dim-vec v1 = n dim-vec v2 = n dim-vec w = m*
**shows** *tensor-vec (v1 + v2) w = tensor-vec v1 w + tensor-vec v2 w*
⟨*proof*⟩

**lemma** *tensor-vec-bilinear-right*:
**assumes** *vd*: *dim-vec v = n dim-vec w1 = m dim-vec w2 = m*
**shows** *tensor-vec v (w1 + w2) = tensor-vec v w1 + tensor-vec v w2*
⟨*proof*⟩

**lemma** *tensor-vec-smult-left*:
**assumes** *vd*: *dim-vec v = n dim-vec w = m*
**shows** *tensor-vec (α ·$_v$ v) w = α ·$_v$ tensor-vec v w*
⟨*proof*⟩

**lemma** *tensor-vec-smult-right*:
**assumes** *vd*: *dim-vec v = n dim-vec w = m*
**shows** *tensor-vec v (α ·$_v$ w) = α ·$_v$ tensor-vec v w*
⟨*proof*⟩

The Schmidt rank, the least length of a representation by elementary tensors. It is *LEAST*, not *Min* of the set of lengths: that set is infinite (zero terms can be added), and *Min* is unspecified on infinite sets. Its properties (rank 1 iff a product state) are in theory Day06_Schmidt_Rank.

**definition** *schmidt-rank* :: *complex Matrix.vec* ⇒ *nat* ⇒ *nat* **where**
*schmidt-rank u m* ≡
*LEAST k*. ∃ *vs* :: *complex Matrix.vec list*. ∃ *ws* :: *complex Matrix.vec list*.
*length vs = k* ∧ *length ws = k* ∧
(∀ *i* < *k*. *dim-vec (vs ! i) = dim-vec u div m*) ∧
(∀ *i* < *k*. *dim-vec (ws ! i) = m*) ∧
*u = foldr* (λ(*v*,*w*) *acc*. *tensor-vec v w + acc*)
(*zip vs ws*) (*0*$_v$ (*dim-vec u*))

# 12 Day 7 (12 September 2011): tensor products of Hilbert spaces; the postulates

**Remark.** For Hilbert spaces $\mathcal{H}, \mathcal{K}$ set $\langle h_1 \otimes k_1 | h_2 \otimes k_2 \rangle = \langle h_1 | h_2 \rangle_{\mathcal{H}} \langle k_1 | k_2 \rangle_{\mathcal{K}}$.

**Theorem (notes).** This extends to a sesquilinear form on $\mathcal{H} \otimes \mathcal{K}$ with $\langle u | u \rangle = 0$ only for $u = 0$, hence to an inner product; in finite dimensions $\mathcal{H} \otimes \mathcal{K}$ with this inner product is the Hilbert space tensor product (in infinite dimensions one completes).

**Proposition (notes).** If $\{h_i\}$ and $\{k_j\}$ are orthonormal bases of $\mathcal{H}$ and $\mathcal{K}$, then $\{h_i \otimes k_j\}$ is an orthonormal basis of $\mathcal{H} \otimes \mathcal{K}$.

In physics notation $e_i \otimes e_j = |ij\rangle$, and for $N$ qubits $e_{i_1} \otimes \cdots \otimes e_{i_N} = |i_1 \cdots i_N\rangle$ with the multi-index $I = (i_1, \ldots, i_N) \in \mathbb{Z}_2^N$. In the concrete model positivity of the form is automatic; the multiplicativity of the inner product and the orthonormal product basis are proved below.

**lemma** *tensor-inner-prod*:
**assumes** *dim-vec h1 = n dim-vec h2 = n*

*dim-vec k1 = m dim-vec k2 = m*
**shows** ⟨*tensor-vec h1 k1* | *tensor-vec h2 k2*⟩ =
⟨*h1* | *h2*⟩ ∗ ⟨*k1* | *k2*⟩
⟨*proof*⟩

**lemma** *length-concat-map-upt*:
*length* (*concat* (*map* (λ*i*. *map* (λ*j*. *f i j*) [*0*..<*m*]) [*0*..<*n*])) = *n* ∗ *m*
⟨*proof*⟩

**lemma** *nth-concat-map-upt*:
**assumes** *p-lt*: *p* < *n* ∗ *m*
**shows** *concat* (*map* (λ*i*. *map* (λ*j*. *f i j*) [*0*..<*m*]) [*0*..<*n*]) ! *p* =
*f* (*p div m*) (*p mod m*)
⟨*proof*⟩

**lemma** *tensor-onb*:
**assumes** *hs-onb*: *orthonormal-list n hs*
**and** *hs-len*: *length hs* = *n*
**and** *ks-onb*: *orthonormal-list m ks*
**and** *ks-len*: *length ks* = *m*
**shows** *orthonormal-list* (*n*∗*m*)
[*tensor-vec* (*hs* ! *i*) (*ks* ! *j*). *i* ← [*0*..<*n*], *j* ← [*0*..<*m*]]
⟨*proof*⟩

## 12.1 Postulates of quantum mechanics

**Postulate (notes)** (1)**.** To each isolated physical system corresponds a Hilbert space, the *state space*; the state of the system is completely described by a unit vector in it, the *state vector*.

**Postulate (notes)** (2)**.** The time evolution of a closed system from $t_1$ to $t_2$ is described by a unitary $U : \mathcal{H} \to \mathcal{H}$: a system in state $\psi$ at $t_1$ is in state $U\psi$ at $t_2$.

**Postulate (notes)** (3)**.** Quantum measurements are described by a collection $\{M_m\}$ of *measurement operators* on $\mathcal{H}$. If the system is in state $\psi$, the probability of observing $m$ is $p_m(\psi) = \langle\psi|M_m^* M_m\psi\rangle = \|M_m\psi\|^2$; since the probabilities sum to one, $\sum_m M_m^* M_m = I$. After observing $m$ the state is $M_m\psi/p_m(\psi)^{1/2}$.

Formally a state is *quantum-state n ψ* (a unit vector in $\mathbb{C}^n$), evolution is *evolve U ψ*, a measurement system is a list *Ms* with *measurement-system n Ms* (all $n \times n$ with $\sum_m M_m^\dagger M_m = I$), and $p_m(\psi)$ is *meas-prob M ψ*.

**definition** *quantum-state* :: *nat* ⇒ *complex Matrix.vec* ⇒ *bool* **where**
*quantum-state n ψ* ≡ *dim-vec ψ* = *n* ∧ ‖*ψ*‖ = *1*

**lemma** *quantum-state-norm1*:
**assumes** *quantum-state n ψ*
**shows** ‖*ψ*‖ = *1*
⟨*proof*⟩

**lemma** *quantum-state-dim*:
**assumes** *quantum-state n ψ*
**shows** *dim-vec ψ* = *n*
⟨*proof*⟩

**definition** *evolve* :: *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.vec* **where**
*evolve U ψ* ≡ *U* ∗*v ψ*

**lemma** *unitary-preserves-state*:
**assumes** *unitary U* **and** *quantum-state n ψ* **and** *dim-col U = n*
**shows** *quantum-state* (*dim-row U*) (*evolve U ψ*)
⟨*proof*⟩

**definition** *measurement-system* :: *nat* ⇒ *complex Matrix.mat list* ⇒ *bool* **where**
*measurement-system n Ms* ≡
(∀ *M* ∈ *set Ms*. *M* ∈ *carrier-mat n n*) ∧
*foldr* (λ*M acc*. (*M*†) ∗ *M* + *acc*) *Ms* (*0*$_m$ *n n*) = *1*$_m$ *n*

**definition** *meas-prob* :: *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *real* **where**
*meas-prob M ψ* ≡ ‖*M* ∗$_v$ *ψ*‖$^2$

**lemma** *measurement-probs-sum-one*:
**assumes** *ms*: *measurement-system n Ms*
**and** *st*: *quantum-state n ψ*
**shows** (∑ *M* ← *Ms*. *meas-prob M ψ*) = *1*
⟨*proof*⟩

$p_m(\psi) = \langle\psi|M_m^* M_m\psi\rangle$, the form in which the notes state Postulate 3.

**lemma** *meas-prob-quad-form*:
**fixes** *M* :: *complex Matrix.mat* **and** *ψ* :: *complex Matrix.vec*
**assumes** *M*: *M* ∈ *carrier-mat n n* **and** *ψ*: *dim-vec ψ* = *n*
**shows** *meas-prob M ψ* = *Re* ⟨*ψ* | (*M*† ∗ *M*) ∗$_v$ *ψ*⟩
⟨*proof*⟩

After outcome $m$ the state becomes $M_m\psi/\|M_m\psi\|$:

**definition** *post-meas-state* :: *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.vec* **where**
*post-meas-state M ψ* ≡ (*1* / *complex-of-real* (‖*M* ∗$_v$ *ψ*‖)) ·$_v$ (*M* ∗$_v$ *ψ*)

**Postulate (notes)** (4, Day 9)**.** The state space of a composite system is the tensor product of the state spaces of its components; if the $i$-th component is in state $\psi_i$, the system is in state $\psi_1 \otimes \cdots \otimes \psi_n$.

**lemma** *tensor-of-states-is-state*:
**assumes** *quantum-state n ψ1* **and** *quantum-state m ψ2*
**shows** *quantum-state* (*n*∗*m*) (*tensor-vec ψ1 ψ2*)
⟨*proof*⟩

# 13 Day 8 (14 September 2011): the quantum game; projective measurements and expected values

## 13.1 Distinguishing states

Alice has two states $\psi_1, \psi_2$ known to Bob; she sends one. Can Bob build a measurement system that decides with certainty which one he received? Formally one wants $M_1, M_2$ with $\|M_1\psi_2\| = 0$, $\|M_1\psi_1\|^2 = 1$, $\|M_2\psi_1\| = 0$, $\|M_2\psi_2\|^2 = 1$.

Case I, $\psi_1 \perp \psi_2$: take $M_1 = |\psi_1\rangle\langle\psi_1|$, $M_2 = |\psi_2\rangle\langle\psi_2|$ and $M_3 = I - M_1 - M_2$; this distinguishes with certainty (formalised below).

Case II, $\psi_1 \not\perp \psi_2$: write $\psi_2 = \alpha\psi_1 + \beta\gamma$ with $\gamma \perp \psi_1$ a unit vector, so $\alpha \neq 0$; any measurement system with $\|M_1\psi_1\| = 1$ satisfies $|\alpha|^2 \le \|M_1\psi_2\|^2$, so outcome 1 also occurs with probability at least $|\alpha|^2$ when $\psi_2$ is sent (*nonorthogonal-outcome-prob* in theory Day02_11_Supplement).

**Theorem (notes).** Given linearly independent states $\psi_1, \ldots, \psi_n$ there is a measurement system $M_0, \ldots, M_n$ such that if outcome $i \geq 1$ occurs then $\psi_i$ was received.

The notes' proof takes $E_i$ the projection onto $\mathrm{span}\{\psi_j : j \neq i\}^\perp$, $M_i = E_i/\sqrt{n}$, and $M_0 = (I - \sum_i M_i^* M_i)^{1/2}$; the remark that follows observes that the inconclusive outcome 0 then has probability at least $(n-1)/n$. Formally this is *lin-indep-states-unambiguous* below.

**lemma** *outer-proj-self*:
  **assumes** *q*: *quantum-state n ψ*
  **shows** *outer-prod ψ ψ ∗v ψ = ψ*
⟨*proof*⟩

**lemma** *outer-proj-orth-zero*:
  **assumes** *q1*: *quantum-state n ψ1*
    **and** *q2*: *quantum-state n ψ2*
    **and** *orth*: ⟨*ψ1 | ψ2*⟩ = *0*
  **shows** *outer-prod ψ1 ψ1 ∗v ψ2 = 0v n*
⟨*proof*⟩

**lemma** *outer-proj-idem*:
  **assumes** *q*: *quantum-state n ψ*
  **shows** (*outer-prod ψ ψ*) ∗ (*outer-prod ψ ψ*) = *outer-prod ψ ψ*
⟨*proof*⟩

**lemma** *outer-proj-hermitian*:
  **assumes** *dim-vec ψ = n*
  **shows** (*outer-prod ψ ψ*)† = *outer-prod ψ ψ*
⟨*proof*⟩

**lemma** *outer-proj-cross-zero*:
  **assumes** *q1*: *quantum-state n ψ1*
    **and** *q2*: *quantum-state n ψ2*
    **and** *orth*: ⟨*ψ1 | ψ2*⟩ = *0*
  **shows** (*outer-prod ψ1 ψ1*) ∗ (*outer-prod ψ2 ψ2*) = *0m n n*
⟨*proof*⟩

Case I: orthogonal states can be distinguished perfectly.

**lemma** *orthogonal-states-distinguishable*:
  **fixes** *ψ1 ψ2* :: *complex Matrix.vec*
  **assumes** *q1*: *quantum-state n ψ1*
    **and** *q2*: *quantum-state n ψ2*
    **and** *orth*: ⟨*ψ1 | ψ2*⟩ = *0*
  **shows** ∃ *Ms* :: *complex Matrix.mat list*.
      *measurement-system n Ms* ∧
      (∃ *M1 M2*. *M1* ∈ *set Ms* ∧ *M2* ∈ *set Ms* ∧
        *meas-prob M1 ψ1* = *1* ∧ *meas-prob M2 ψ1* = *0* ∧
        *meas-prob M1 ψ2* = *0* ∧ *meas-prob M2 ψ2* = *1*)
⟨*proof*⟩

The theorem on linearly independent states. Linear independence of a list of vectors is injectivity of the matrix with those columns; the formal statement produces the measurement system $M_0, \ldots, M_k$ in the order $M_1, \ldots, M_k, M_0$ (outcome $i < k$ identifies $\psi_i$, outcome $k$ is inconclusive). The proof follows the notes with one change: instead of the projections $E_i$ onto $\{\psi_j : j \neq i\}^\perp$ it uses the dual vectors $\phi_i$ with $\langle \phi_i | \psi_j \rangle = \delta_{ij}$ (columns of $V(V^*V)^{-1}$) and the operators $M_i = \sqrt{c}\,|e_0\rangle\langle\phi_i|$, where $c > 0$ makes $I - c\sum_i |\phi_i\rangle\langle\phi_i|$ positive; $M_0$ is its square root.

**definition** *lin-indep-vecs* :: *nat* ⇒ *complex Matrix.vec list* ⇒ *bool* **where**
  *lin-indep-vecs n vs* ≡ (∀ *v* ∈ *set vs*. *dim-vec v = n*) ∧

$(\forall\, x ::$ *complex Matrix.vec*. $x \in$ *carrier-vec* (*length vs*) $\longrightarrow$
*mat-of-cols n vs* $*_v$ $x = 0_v$ *n* $\longrightarrow$ $x = 0_v$ (*length vs*))

**lemma** *outer-prod-carrier*:
**fixes** *v w* :: *complex Matrix.vec*
**assumes** *dim-vec v* = *n* **and** *dim-vec w* = *m*
**shows** *v* $\oslash$ *w* $\in$ *carrier-mat n m*
⟨*proof*⟩

**lemma** *outer-prod-dagger*:
**fixes** *v w* :: *complex Matrix.vec*
**shows** $(v \oslash w)^\dagger = w \oslash v$
⟨*proof*⟩

**lemma** *foldr-add-mat-carrier*:
**fixes** *f* :: *'a* $\Rightarrow$ *complex Matrix.mat*
**assumes** $\forall\, x \in$ *set xs*. *f x* $\in$ *carrier-mat n n*
**shows** *foldr* ($\lambda x$ *acc*. *f x* + *acc*) *xs* ($0_m$ *n n*) $\in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *foldr-add-mat-index*:
**fixes** *f* :: *'a* $\Rightarrow$ *complex Matrix.mat*
**assumes** $\forall\, x \in$ *set xs*. *f x* $\in$ *carrier-mat n n* **and** *i* < *n* **and** *j* < *n*
**shows** *foldr* ($\lambda x$ *acc*. *f x* + *acc*) *xs* ($0_m$ *n n*) $$ (*i*, *j*) = *sum-list* (*map* ($\lambda x$. *f x* $$ (*i*, *j*)) *xs*)
⟨*proof*⟩

**lemma** *foldr-MdagM-shift*:
**assumes** *Ms-c*: $\forall\, M \in$ *set Ms*. *M* $\in$ *carrier-mat n n* **and** *z*: *z* $\in$ *carrier-mat n n*
**shows** *foldr* ($\lambda M$ *acc*. $(M^\dagger) * M$ + *acc*) *Ms z*
= *foldr* ($\lambda M$ *acc*. $(M^\dagger) * M$ + *acc*) *Ms* ($0_m$ *n n*) + *z*
⟨*proof*⟩

**theorem** *lin-indep-states-unambiguous*:
**fixes** $\psi s$ :: *complex Matrix.vec list*
**assumes** *qs*: $\forall\, \psi \in$ *set* $\psi s$. *quantum-state n* $\psi$
**and** *li*: *lin-indep-vecs n* $\psi s$
**and** *k*: *length* $\psi s$ = *k* **and** *kpos*: *0* < *k*
**shows** $\exists$ *Ms* :: *complex Matrix.mat list*.
*measurement-system n Ms* $\wedge$ *length Ms* = *k* + *1* $\wedge$
$(\forall\, i{<}k.\ \forall\, j{<}k.\ i \neq j \longrightarrow$ *meas-prob* (*Ms* ! *i*) ($\psi s$ ! *j*) = *0*) $\wedge$
$(\forall\, i{<}k.\ 0 <$ *meas-prob* (*Ms* ! *i*) ($\psi s$ ! *i*))
⟨*proof*⟩

## 13.2 Projective measurements, expected values and self-adjoint operators

If a measurement $\{M_m\}$ has real outcomes $\lambda_m$, the expected value of the outcome in state $\psi$ is $E = \sum_m p_m(\psi)\lambda_m = \langle\psi|(\sum_m \lambda_m M_m^* M_m)\psi\rangle = \langle\psi|H\psi\rangle$ with $H = \sum_m \lambda_m M_m^* M_m$ self-adjoint: expected values of observables are quadratic forms of Hermitian matrices (*expected-value-eq-quad-form* below).

Formally: a measurement whose operator is the rank-one projection $|\gamma\rangle\langle\gamma|$ has $p(\psi) = |\langle\gamma|\psi\rangle|^2$ (the proposition of Day 12); an *observable* with outcomes $\lambda_m$ attached to the measurement operators $M_m$ is $H = \sum_m \lambda_m M_m^* M_m$, which is Hermitian, and the expected outcome $\sum_m \lambda_m p_m(\psi)$ equals $\langle\psi|H\psi\rangle$.

**lemma** *rank-one-meas-prob*:
**fixes** $\gamma$ $\psi$ :: *complex Matrix.vec*
**assumes** *g*: *quantum-state n* $\gamma$ **and** $\psi$: *dim-vec* $\psi$ = *n*

**shows** *meas-prob* $(\gamma \oslash \gamma)\ \psi = (cmod\ \langle\gamma \mid \psi\rangle)^2$
⟨*proof*⟩

**definition** *observable* :: *nat* ⇒ (*real* × *complex Matrix.mat*) *list* ⇒ *complex Matrix.mat* **where**
*observable n lMs* ≡
*foldr* (λ(*l*, *M*) *acc*. *complex-of-real l* ·$_m$ ((*M*†) ∗ *M*) + *acc*) *lMs* (*0*$_m$ *n n*)

**definition** *expected-value* :: (*real* × *complex Matrix.mat*) *list* ⇒ *complex Matrix.vec* ⇒ *real* **where**
*expected-value lMs ψ* ≡ (∑ (*l*, *M*) ← *lMs*. *l* ∗ *meas-prob M ψ*)

**lemma** *observable-Nil* [*simp*]: *observable n* [] = *0*$_m$ *n n*
⟨*proof*⟩

**lemma** *observable-Cons* [*simp*]:
*observable n* ((*l*, *M*) # *lMs*) = *complex-of-real l* ·$_m$ ((*M*†) ∗ *M*) + *observable n lMs*
⟨*proof*⟩

**lemma** *observable-carrier*:
**assumes** ∀ (*l*, *M*) ∈ *set lMs*. *M* ∈ *carrier-mat n n*
**shows** *observable n lMs* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *observable-hermitian*:
**assumes** ∀ (*l*, *M*) ∈ *set lMs*. *M* ∈ *carrier-mat n n*
**shows** *hermitian* (*observable n lMs*)
⟨*proof*⟩

**lemma** *expected-value-eq-quad-form*:
**fixes** *ψ* :: *complex Matrix.vec*
**assumes** *ψ*: *dim-vec ψ* = *n*
**and** *lMs*: ∀ (*l*, *M*) ∈ *set lMs*. *M* ∈ *carrier-mat n n*
**shows** *expected-value lMs ψ* = *Re* ⟨*ψ* | *observable n lMs* ∗$_v$ *ψ*⟩
⟨*proof*⟩

# 14 Day 9 (16 September 2011): positive operator-valued measures; composite systems; entanglement

## 14.1 Positive operator-valued measures

If two measurement systems satisfy $M_m^* M_m = \tilde{M}_m^* \tilde{M}_m$ for all $m$ then $p_m = \tilde{p}_m$: only the operators $M_m^* M_m$ matter, and $\sum_m M_m^* M_m = I$. Conversely if $P_m \geq 0$ with $\sum_m P_m = I$ then $M_m = P_m^{1/2}$ is a measurement system.

**Definition (notes).** A *positive operator-valued measure* (POVM) is a set $\{P_m\}$ of positive operators with $\sum_m P_m = I$.

**definition** *povm* :: *nat* ⇒ *complex Matrix.mat list* ⇒ *bool* **where**
*povm n Ps* ≡
(∀ *P* ∈ *set Ps*. *P* ∈ *carrier-mat n n* ∧ *psd P*) ∧
*foldr* (λ*P acc*. *P* + *acc*) *Ps* (*0*$_m$ *n n*) = *1*$_m$ *n*

**lemma** *measurement-induces-povm*:
**assumes** *ms*: *measurement-system n Ms*
**shows** *povm n* (*map* (λ*M*. (*M*†) ∗ *M*) *Ms*)
⟨*proof*⟩

**lemma** *psd-sqrt-measurement-factor*:
**assumes** *Pcar*: $P \in$ *carrier-mat n n*
**and** *Ppsd*: *psd P*
**shows** $\exists M.\ M \in$ *carrier-mat n n* $\wedge\ M^\dagger * M = P$
⟨*proof*⟩

**lemma** *psd-sqrt-measurement-factors*:
**assumes** *Ps*: $\forall P \in$ *set Ps*. $P \in$ *carrier-mat n n* $\wedge$ *psd P*
**shows** $\exists$ *Ms*. $(\forall M \in$ *set Ms*. $M \in$ *carrier-mat n n*$) \wedge$
*map* $(\lambda M.\ M^\dagger * M)$ *Ms* = *Ps*
⟨*proof*⟩

**lemma** *povm-from-sqrt*:
**assumes** *pv*: *povm n Ps*
**shows** $\exists$ *Ms*. *measurement-system n Ms* $\wedge$
*map* $(\lambda M.\ M^\dagger * M)$ *Ms* = *Ps*
⟨*proof*⟩

**lemma** *povm-prob*:
**assumes** *pv*: *povm n Ps* **and** *st*: *quantum-state n* $\psi$
**and** *Pin*: $P \in$ *set Ps*
**shows** *Re* $(\langle \psi \mid P *_v \psi \rangle) \geq 0$
⟨*proof*⟩

## 14.2 Composite systems

Postulate 4 was stated with Day 7. The notes' example: photons in states $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ and $\frac{1}{\sqrt{2}}(|0\rangle + i|1\rangle)$ form the pair $\frac{1}{2}(|00\rangle + i|01\rangle + |10\rangle + i|11\rangle) \in \mathbb{C}^2 \otimes \mathbb{C}^2$, a vector of Schmidt rank 1. Two photons described by a unit vector $\psi \in \mathbb{C}^2 \otimes \mathbb{C}^2$ can be regarded as two separate subsystems iff $\psi = \psi_1 \otimes \psi_2$ iff the Schmidt rank of $\psi$ is 1; otherwise $\psi$ is *entangled*, which does happen in nature.

Tensor products of operators: for $R : \mathcal{H} \to \mathcal{H}$ and $T : \mathcal{K} \to \mathcal{K}$ there is a unique $R \otimes T$ with $(R \otimes T)(\sum_l h_l \otimes k_l) = \sum_l Rh_l \otimes Tk_l$, and (1) $(R_1 \otimes T_1)(R_2 \otimes T_2) = R_1R_2 \otimes T_1T_2$, (2) $(R \otimes T)^* = R^* \otimes T^*$ (both proved in Basic_Math for the Kronecker product).

**Remark.** If lab A has the measurement system $\{M_m\}$ on $\mathcal{H}$ and lab B has state space $\mathcal{K}$, the measurement system of the composite is $\{M_m \otimes I_\mathcal{K}\}$ and $p^{AB}_m(\psi \otimes \phi) = \|(M_m \otimes I)(\psi \otimes \phi)\|^2 = p^A_m(\psi)$.

**lemma** *tensor-mat-vec-mult*:
**assumes** *A*: $A \in$ *carrier-mat nr nc*
**and** *B*: $B \in$ *carrier-mat mr mc*
**and** *v*: *dim-vec v* = *nc*
**and** *w*: *dim-vec w* = *mc*
**shows** $(A \bigotimes B) *_v$ *tensor-vec v w* = *tensor-vec* $(A *_v v)$ $(B *_v w)$
⟨*proof*⟩

**lemma** *tensor-mat-vec-mult-right-id*:
**assumes** *A*: $A \in$ *carrier-mat n n*
**and** *v*: *dim-vec v* = *n*
**and** *w*: *dim-vec w* = *m*
**shows** $(A \bigotimes 1_m\ m) *_v$ *tensor-vec v w* = *tensor-vec* $(A *_v v)$ *w*
⟨*proof*⟩

**lemma** *composite-measurement-prob*:
**fixes** $M$ :: *complex Matrix.mat* **and** $\psi$ :: *complex Matrix.vec* **and** $\varphi$ :: *complex Matrix.vec*
**assumes** *psi-st*: *quantum-state n* $\psi$ **and** *phi-st*: *quantum-state m* $\varphi$
**and** *M-sq*: $M \in$ *carrier-mat n n*
**shows** *meas-prob* $(M \bigotimes 1_m\ m)$ *(tensor-vec* $\psi$ $\varphi$*)* = *meas-prob M* $\psi$
⟨*proof*⟩

### 14.3 Entanglement

A state $\psi \in \mathcal{H}_A \otimes \mathcal{H}_B$ is *entangled* if it is not a product state $u \otimes v$. The Bell state $\gamma = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ is entangled. The notes' first application of entanglement (A and B share $\gamma$; A measures $|0\rangle\langle 0|, |1\rangle\langle 1|$ and gets each outcome with probability $\frac{1}{2}$, after which B's measurement is determined) is *bell-measurement* in theory Day02_11_Supplement.

**definition** *entangled* :: *nat* ⇒ *nat* ⇒ *complex Matrix.vec* ⇒ *bool* **where**
*entangled n m* $\psi \equiv$
*dim-vec* $\psi = n * m \wedge$
$\neg$ $(\exists\, u\ v.$ *dim-vec u* $= n \wedge$ *dim-vec v* $= m \wedge \psi =$ *tensor-vec u v*$)$

**definition** *bell-state* :: *complex Matrix.vec* **where**
*bell-state* ≡ *(1 / csqrt 2)* $\cdot_v$
*(tensor-vec (unit-vec 2 0) (unit-vec 2 0) +*
*tensor-vec (unit-vec 2 1) (unit-vec 2 1))*

**lemma** *bell-state-norm*:
‖*bell-state*‖ = *1*
⟨*proof*⟩

**lemma** *bell-state-is-quantum-state*:
*quantum-state 4 bell-state*
⟨*proof*⟩

**lemma** *bell-state-entangled*:
*entangled 2 2 bell-state*
⟨*proof*⟩

## 15 Day 10 (19 September 2011): superdense coding; classical and quantum gates

### 15.1 One-qubit gates

The one-qubit gates of the notes:

$$X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ (NOT)}, \quad Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad H = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad S = \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}.$$

$H$ is the Hadamard gate; the notes write "$H^2 = X$" here and correct it to $H^2 = I$ on Day 11, which is what is proved below. $H$ and $S$ together generate all $2 \times 2$ unitaries (not formalised). Formally $H$ is *Hadamard* (the library's *Quantum.H*) and $S$ is *Phase-S*.

**definition** *Hadamard* :: *complex Matrix.mat* **where**
*Hadamard* ≡ *Quantum.H*

**lemma** *csqrt-two*: *csqrt (2*::*complex)* = *(sqrt 2* :: *complex)*
⟨*proof*⟩

**lemma** *hadamard-squared*:
*Hadamard* $*$ *Hadamard* $=$ $1_m$ *2*
⟨*proof*⟩

**lemma** *hadamard-e0*:
*Hadamard* $*_v$ *unit-vec 2 0* $=$
(*1 / csqrt 2*) $\cdot_v$ (*unit-vec 2 0* $+$ *unit-vec 2 1*)
⟨*proof*⟩

**lemma** *hadamard-e1*:
*Hadamard* $*_v$ *unit-vec 2 1* $=$
(*1 / csqrt 2*) $\cdot_v$ (*unit-vec 2 0* $-$ *unit-vec 2 1*)
⟨*proof*⟩

The phase gate $S = \mathrm{diag}(1, i)$ is unitary.

**definition** *Phase-S* :: *complex Matrix.mat* **where**
*Phase-S* $\equiv$ *Matrix.mat 2 2* ($\lambda$(*i*,*j*). *if i* $=$ *0* $\wedge$ *j* $=$ *0 then 1 else if i* $=$ *1* $\wedge$ *j* $=$ *1 then* i *else 0*)

**lemma** *Phase-S-carrier* [*simp*]: *Phase-S* $\in$ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *Phase-S-dagger-S*: *Phase-S*$^\dagger$ $*$ *Phase-S* $=$ $1_m$ *2*
⟨*proof*⟩

**lemma** *Phase-S-S-dagger*: *Phase-S* $*$ (*Phase-S*$^\dagger$) $=$ $1_m$ *2*
⟨*proof*⟩

**lemma** *Phase-S-unitary*: *unitary Phase-S*
⟨*proof*⟩

## 15.2 Two- and three-qubit gates

The notes tabulate the classical gates AND, OR, XOR, NAND, NOR (Peirce: NAND generates all Boolean operations; Sheffer: so does NOR). No two-bit classical gate is a quantum gate, since it maps two dimensions to one and is not reversible. The controlled NOT acts by $|a, b\rangle \mapsto |a, b \oplus a\rangle$:

$$\mathrm{CNOT} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

which is unitary, and Toffoli's CCNOT (identity on the first six basis vectors, swapping $|110\rangle$ and $|111\rangle$) is a reversible gate generating all Boolean operations; it is defined in theory Quantum_Computing11.

**lemma** *CNOT-unitary*: *unitary CNOT*
⟨*proof*⟩

**lemma** *CNOT-00*:
*CNOT* $*_v$ (*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 0*)) $=$
*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 0*)
⟨*proof*⟩

**lemma** *CNOT-10*:
*CNOT* $*_v$ (*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 0*)) $=$

*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 1*)
⟨*proof*⟩

**lemma** *CNOT-01*:
*CNOT* $*_v$ (*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 1*)) =
*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 1*)
⟨*proof*⟩

**lemma** *CNOT-11*:
*CNOT* $*_v$ (*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 1*)) =
*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 0*)
⟨*proof*⟩

**lemma** *tensor-unit-vec-2-00*:
*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 0*) = *unit-vec 4 0*
⟨*proof*⟩

**lemma** *tensor-unit-vec-2-01*:
*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 1*) = *unit-vec 4 1*
⟨*proof*⟩

**lemma** *tensor-unit-vec-2-10*:
*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 0*) = *unit-vec 4 2*
⟨*proof*⟩

**lemma** *tensor-unit-vec-2-11*:
*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 1*) = *unit-vec 4 3*
⟨*proof*⟩

## 15.3 Superdense coding

A and B share $\gamma = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. Applying nothing, $\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}$, $\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}$ or $\begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}$ to its qubit, A turns $\gamma$ into one of the four orthonormal Bell states $\frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle)$, $\frac{1}{\sqrt{2}}(|10\rangle \pm |01\rangle)$, so by sending one qubit A communicates two bits; with $2m$ shared qubits, $4^m$ messages. Below: the four Bell states and their orthonormality (the action table itself is not formalised).

**definition** *bell-basis* :: *complex Matrix.vec list* **where**
*bell-basis* ≡
[ (*1/csqrt 2*) $\cdot_v$ (*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 0*) +
*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 1*)),
(*1/csqrt 2*) $\cdot_v$ (*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 0*) −
*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 1*)),
(*1/csqrt 2*) $\cdot_v$ (*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 0*) +
*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 1*)),
(*1/csqrt 2*) $\cdot_v$ (*tensor-vec* (*unit-vec 2 0*) (*unit-vec 2 1*) −
*tensor-vec* (*unit-vec 2 1*) (*unit-vec 2 0*)) ]

**lemma** *bell-basis-orthonormal*: *orthonormal-list 4 bell-basis*
⟨*proof*⟩

**end**
**theory** *Day02-11-Supplement*
**imports** *Quantum-Computing11*
**begin**

Supplementary results for Days 2–11 recorded as gaps by the first audit.

## 15.4 Day 2: the projection does not depend on the orthonormal basis

**definition** *col-span* :: *nat* ⇒ *complex Matrix.vec list* ⇒ *complex Matrix.vec set* **where**
*col-span n vs* = {*mat-of-cols n vs* ∗$_v$ *c* | *c*. *dim-vec c* = *length vs*}

**lemma** *outer-sum-fixes-col-span*:
**assumes** *orth*: *orthonormal-list n vs* **and** *w*: *w* ∈ *col-span n vs*
**shows** *outer-sum n vs* ∗$_v$ *w* = *w*
⟨*proof*⟩

**lemma** *mem-col-span-self*:
**assumes** ∀ *v* ∈ *set vs*. *dim-vec v* = *n* **and** *j*: *j* < *length vs*
**shows** *vs* ! *j* ∈ *col-span n vs*
⟨*proof*⟩

**lemma** *outer-sum-absorbs*:
**assumes** *ov*: *orthonormal-list n vs* **and** *ow*: *orthonormal-list n ws*
**and** *sub*: *set ws* ⊆ *col-span n vs*
**shows** *outer-sum n vs* ∗ *outer-sum n ws* = *outer-sum n ws*
⟨*proof*⟩

D2-8: two orthonormal bases of the same subspace give the same projection.

**theorem** *outer-sum-basis-independent*:
**assumes** *ov*: *orthonormal-list n vs* **and** *ow*: *orthonormal-list n ws*
**and** *span*: *col-span n vs* = *col-span n ws*
**shows** *outer-sum n vs* = *outer-sum n ws*
⟨*proof*⟩

## 15.5 Day 6: coordinates in a tensor product

$\sum_j x_j \otimes y_j$ in $\mathbb{C}^N$, written as in *schmidt-rank*.

**definition** *tensor-combination* :: *nat* ⇒ *complex Matrix.vec list* ⇒ *complex Matrix.vec list* ⇒ *complex Matrix.vec* **where**
*tensor-combination N xs ys* = *foldr* (λ(*x*,*y*) *acc*. *tensor-vec x y* + *acc*) (*zip xs ys*) (*0*$_v$ *N*)

**lemma** *tensor-combination-dim* [*simp*]: *dim-vec* (*tensor-combination N xs ys*) = *N*
⟨*proof*⟩

**lemma** *tensor-combination-index*:
**assumes** ∀ *x* ∈ *set xs*. *dim-vec x* = *k* ∀ *y* ∈ *set ys*. *dim-vec y* = *m length xs* = *length ys p* < *k* ∗ *m*
**shows** *tensor-combination* (*k* ∗ *m*) *xs ys* $ *p* = (∑*j*<*length xs*. *xs* ! *j* $ (*p div m*) ∗ *ys* ! *j* $ (*p mod m*))
⟨*proof*⟩

**lemma** *idx-bound*: *i* < *k* ⟹ *l* < (*m*::*nat*) ⟹ *i* ∗ *m* + *l* < *k* ∗ *m*
⟨*proof*⟩

**lemma** *idx-div-mod*: *l* < (*m*::*nat*) ⟹ (*i* ∗ *m* + *l*) *div m* = *i* ∧ (*i* ∗ *m* + *l*) *mod m* = *l*
⟨*proof*⟩

A basis $f_1, \ldots, f_m$ of $\mathbb{C}^m$ is a list whose column matrix $F$ is invertible; the coordinates use $G = F^{-1}$.

**definition** *right-coord* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *nat* ⇒ *complex Matrix.vec* **where**
*right-coord k m G u j* = *Matrix.vec k* (λ*i*. ∑ *l*<*m*. *u* $ (*i* ∗ *m* + *l*) ∗ *G* $$ (*j*,*l*))

**definition** *left-coord* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *nat* ⇒ *complex Matrix.vec* **where**

*left-coord k m Hm u i = Matrix.vec m (λl. ∑ r<k. Hm $$ (i,r) * u $ (r * m + l))*

**lemma** *basis-inverse*:
**fixes** *fs* :: *complex Matrix.vec list*
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *inv*: *invertible-mat* (*mat-of-cols m fs*)
**obtains** *G* **where** *G* ∈ *carrier-mat m m G * mat-of-cols m fs = 1*$_m$ *m mat-of-cols m fs * G = 1*$_m$ *m*
⟨*proof*⟩

**lemma** *inv-entry-left*:
**fixes** *fs* :: *complex Matrix.vec list*
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *G*: *G* ∈ *carrier-mat m m*
**and** *GF*: *G * mat-of-cols m fs = 1*$_m$ *m* **and** *j*: *j < m* **and** *j'*: *j' < m*
**shows** (∑ *l<m*. *fs ! j' $ l * G $$ (j,l)*) = (*if j = j' then 1 else 0*)
⟨*proof*⟩

**lemma** *inv-entry-right*:
**fixes** *fs* :: *complex Matrix.vec list*
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *G*: *G* ∈ *carrier-mat m m*
**and** *FG*: *mat-of-cols m fs * G = 1*$_m$ *m* **and** *r*: *r < m* **and** *l*: *l < m*
**shows** (∑ *j<m*. *fs ! j $ r * G $$ (j,l)*) = (*if r = l then 1 else 0*)
⟨*proof*⟩

**lemma** *right-coord-unique*:
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *G*: *G* ∈ *carrier-mat m m*
**and** *GF*: *G * mat-of-cols m fs = 1*$_m$ *m*
**and** *xs*: *length xs = m* ∀ *x* ∈ *set xs*. *dim-vec x = k* **and** *j*: *j < m*
**shows** *right-coord k m G* (*tensor-combination* (*k * m*) *xs fs*) *j = xs ! j*
⟨*proof*⟩

**lemma** *right-coord-exists*:
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *G*: *G* ∈ *carrier-mat m m*
**and** *FG*: *mat-of-cols m fs * G = 1*$_m$ *m* **and** *u*: *dim-vec u = k * m*
**shows** *tensor-combination* (*k * m*) (*map* (*right-coord k m G u*) [*0..<m*]) *fs = u*
⟨*proof*⟩

D6-3 (i): for a basis $f_1, \ldots, f_m$ of $\mathbb{C}^m$, every $u \in \mathbb{C}^k \otimes \mathbb{C}^m$ is $\sum_j x_j \otimes f_j$ for unique $x_1, \ldots, x_m \in \mathbb{C}^k$.

**theorem** *tensor-right-coordinates*:
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *inv*: *invertible-mat* (*mat-of-cols m fs*)
**and** *u*: *dim-vec u = k * m*
**shows** ∃!*xs*. *length xs = m* ∧ (∀ *x* ∈ *set xs*. *dim-vec x = k*) ∧ *u = tensor-combination* (*k * m*) *xs fs*
⟨*proof*⟩

**lemma** *left-coord-unique*:
**assumes** *ed*: ∀ *e* ∈ *set es*. *dim-vec e = k length es = k* **and** *Hm*: *Hm* ∈ *carrier-mat k k*
**and** *HE*: *Hm * mat-of-cols k es = 1*$_m$ *k*
**and** *ys*: *length ys = k* ∀ *y* ∈ *set ys*. *dim-vec y = m* **and** *i*: *i < k*
**shows** *left-coord k m Hm* (*tensor-combination* (*k * m*) *es ys*) *i = ys ! i*
⟨*proof*⟩

**lemma** *left-coord-exists*:
**assumes** *ed*: ∀ *e* ∈ *set es*. *dim-vec e = k length es = k* **and** *Hm*: *Hm* ∈ *carrier-mat k k*
**and** *EH*: *mat-of-cols k es * Hm = 1*$_m$ *k* **and** *u*: *dim-vec u = k * m*
**shows** *tensor-combination* (*k * m*) *es* (*map* (*left-coord k m Hm u*) [*0..<k*]) *= u*
⟨*proof*⟩

D6-3 (ii): likewise $u = \sum_i e_i \otimes y_i$ for unique $y_1, \ldots, y_k \in \mathbb{C}^m$.

**theorem** *tensor-left-coordinates*:
**assumes** *ed*: ∀ *e* ∈ *set es*. *dim-vec e = k length es = k* **and** *inv*: *invertible-mat* (*mat-of-cols k es*)

**and** *u*: *dim-vec u = k * m*
**shows** ∃!*ys*. *length ys = k* ∧ (∀ *y* ∈ *set ys*. *dim-vec y = m*) ∧ *u = tensor-combination* (*k * m*) *es ys*
⟨*proof*⟩

Corollary: $\mathbb{C}^k \otimes \mathbb{C}^m \cong \mathbb{C}^k \oplus \cdots \oplus \mathbb{C}^k$ ($m$ times) by the linear bijection $(x_1, \ldots, x_m) \mapsto \sum_j x_j \otimes f_j$, and $\cong \mathbb{C}^m \oplus \cdots \oplus \mathbb{C}^m$ ($k$ times) likewise; the identification depends on the chosen basis.

**lemma** *tensor-combination-add-left*:
**assumes** *l*: *length xs = length fs length xs' = length fs*
**and** *d*: ∀ *x* ∈ *set xs*. *dim-vec x = k* ∀ *x* ∈ *set xs'*. *dim-vec x = k* **and** *fd*: ∀ *y* ∈ *set fs*. *dim-vec y = m*
**shows** *tensor-combination* (*k * m*) (*map2* (+) *xs xs'*) *fs = tensor-combination* (*k * m*) *xs fs* + *tensor-combination* (*k * m*) *xs' fs*
⟨*proof*⟩

**lemma** *tensor-combination-smult-left*:
**assumes** *l*: *length xs = length fs* **and** *d*: ∀ *x* ∈ *set xs*. *dim-vec x = k* **and** *fd*: ∀ *y* ∈ *set fs*. *dim-vec y = m*
**shows** *tensor-combination* (*k * m*) (*map* (λ*x*. *a* ·$_v$ *x*) *xs*) *fs = a* ·$_v$ *tensor-combination* (*k * m*) *xs fs*
⟨*proof*⟩

**corollary** *tensor-iso-direct-sum*:
**assumes** *fd*: ∀ *f* ∈ *set fs*. *dim-vec f = m length fs = m* **and** *inv*: *invertible-mat* (*mat-of-cols m fs*)
**shows** *bij-betw* (λ*xs*. *tensor-combination* (*k * m*) *xs fs*)
{*xs*. *length xs = m* ∧ (∀ *x* ∈ *set xs*. *dim-vec x = k*)} (*carrier-vec* (*k * m*))
⟨*proof*⟩

## 15.6 Day 8: non-orthogonal states cannot be distinguished

**lemma** *meas-sum-norm*:
**assumes** *ms*: *measurement-system n Ms* **and** *v*: *dim-vec v = n*
**shows** (∑ *l*<*length Ms*. ‖*Ms* ! *l* *$_v$ *v*‖²) = ‖*v*‖²
⟨*proof*⟩

**lemma** *inner-minus-smult*:
**assumes** *d1*: *dim-vec u = n* **and** *d2*: *dim-vec w = n*
**shows** ⟨*w* − *a* ·$_v$ *u* | *w* − *a* ·$_v$ *u*⟩ = ⟨*w*|*w*⟩ − *a* * ⟨*w*|*u*⟩ − *cnj a* * ⟨*u*|*w*⟩ + *a* * *cnj a* * ⟨*u*|*u*⟩
⟨*proof*⟩

**lemma** *norm-sq-minus-component*:
**assumes** *d1*: *dim-vec u = n* **and** *d2*: *dim-vec w = n* **and** *u*: ‖*u*‖ = *1*
**shows** ‖*w* − ⟨*u*|*w*⟩ ·$_v$ *u*‖² = ‖*w*‖² − (*cmod* ⟨*u*|*w*⟩)²
⟨*proof*⟩

D8-2: if outcome $i$ certainly occurs on $\psi_1$, it occurs on $\psi_2$ with probability at least $|\langle\psi_1|\psi_2\rangle|^2$.

**theorem** *nonorthogonal-outcome-prob*:
**assumes** *ms*: *measurement-system n Ms* **and** *q1*: *quantum-state n ψ1* **and** *q2*: *quantum-state n ψ2*
**and** *i*: *i < length Ms* **and** *p1*: *meas-prob* (*Ms* ! *i*) *ψ1 = 1*
**shows** (*cmod* ⟨*ψ1*|*ψ2*⟩)² ≤ *meas-prob* (*Ms* ! *i*) *ψ2*
⟨*proof*⟩

**corollary** *nonorthogonal-not-distinguishable*:
**assumes** *ms*: *measurement-system n Ms* **and** *q1*: *quantum-state n ψ1* **and** *q2*: *quantum-state n ψ2*
**and** *no*: ⟨*ψ1*|*ψ2*⟩ ≠ *0* **and** *i*: *i < length Ms* **and** *p1*: *meas-prob* (*Ms* ! *i*) *ψ1 = 1*
**shows** *meas-prob* (*Ms* ! *i*) *ψ2 > 0*
⟨*proof*⟩

## 15.7 Day 9: measuring one half of a Bell pair

**definition** *e0* :: *complex Matrix.vec* **where** *e0 = unit-vec 2 0*

**definition** *e1* :: *complex Matrix.vec* **where** *e1 = unit-vec 2 1*
**definition** *proj0* :: *complex Matrix.mat* **where** *proj0 = e0 ⊘ e0*
**definition** *proj1* :: *complex Matrix.mat* **where** *proj1 = e1 ⊘ e1*

**lemma** *e-dims* [*simp*]: *dim-vec e0 = 2 dim-vec e1 = 2*
⟨*proof*⟩

**lemma** *e-inner* [*simp*]: ⟨*e0*|*e0*⟩ = *1* ⟨*e0*|*e1*⟩ = *0* ⟨*e1*|*e0*⟩ = *0* ⟨*e1*|*e1*⟩ = *1*
⟨*proof*⟩

**lemma** *proj-carrier* [*simp*]: *proj0 ∈ carrier-mat 2 2 proj1 ∈ carrier-mat 2 2*
⟨*proof*⟩

**lemma** *proj-apply* [*simp*]:
*proj0 *$_v$ *e0 = e0 proj0 *$_v$ *e1 = 0*$_v$ *2 proj1 *$_v$ *e0 = 0*$_v$ *2 proj1 *$_v$ *e1 = e1*
⟨*proof*⟩

**lemma** *bell-state-e*: *bell-state = (1 / csqrt 2) ·*$_v$ *(tensor-vec e0 e0 + tensor-vec e1 e1)*
⟨*proof*⟩

**lemma** *tensor-zero-left*: *dim-vec w = 2 ⟹ tensor-vec (0*$_v$ *2) w = 0*$_v$ *4*
⟨*proof*⟩

**lemma** *tensor-zero-right*: *dim-vec v = 2 ⟹ tensor-vec v (0*$_v$ *2) = 0*$_v$ *4*
⟨*proof*⟩

**lemma** *A-meas-bell*:
*(proj0 ⨂ 1*$_m$ *2) *$*_v$* *bell-state = (1 / csqrt 2) ·*$_v$ *tensor-vec e0 e0*
*(proj1 ⨂ 1*$_m$ *2) *$*_v$* *bell-state = (1 / csqrt 2) ·*$_v$ *tensor-vec e1 e1*
⟨*proof*⟩

**lemma** *tensor-e-norm*: ‖*tensor-vec e0 e0*‖ = *1* ‖*tensor-vec e1 e1*‖ = *1*
⟨*proof*⟩

D9-6: A measures $|0\rangle\langle 0|, |1\rangle\langle 1|$ on the shared state $\gamma$: each outcome has probability $\frac{1}{2}$, and $\gamma$ collapses to $e_0 \otimes e_0$ resp. $e_1 \otimes e_1$, after which B's measurement $I \otimes M_m$ gives A's outcome with probability 1.

**theorem** *bell-measurement*:
*meas-prob (proj0 ⨂ 1*$_m$ *2) bell-state = 1/2*
*meas-prob (proj1 ⨂ 1*$_m$ *2) bell-state = 1/2*
*post-meas-state (proj0 ⨂ 1*$_m$ *2) bell-state = tensor-vec e0 e0*
*post-meas-state (proj1 ⨂ 1*$_m$ *2) bell-state = tensor-vec e1 e1*
*meas-prob (1*$_m$ *2 ⨂ proj0) (tensor-vec e0 e0) = 1*
*meas-prob (1*$_m$ *2 ⨂ proj1) (tensor-vec e1 e1) = 1*
⟨*proof*⟩

**lemma** *proj-explicit*:
*proj0 = Matrix.mat 2 2 (λ(i,j). if i = 0 ∧ j = 0 then 1 else 0)*
*proj1 = Matrix.mat 2 2 (λ(i,j). if i = 1 ∧ j = 1 then 1 else 0)*
⟨*proof*⟩

**lemma** *proj-measurement-system*: *measurement-system 2* [*proj0*, *proj1*]
⟨*proof*⟩

## 15.8 Day 11: SWAP from three CNOTs

**lemma** *less4-cases*: (*i*::*nat*) < *4* ⟷ *i = 0 ∨ i = 1 ∨ i = 2 ∨ i = 3*
⟨*proof*⟩

CNOT′ is CNOT controlled by the second qubit.

**definition** *CNOT′* :: *complex Matrix.mat* **where**
*CNOT′* = *Matrix.mat 4 4* (λ(*i*,*j*). *if* (*i*,*j*) ∈ {(*0*,*0*),(*1*,*3*),(*2*,*2*),(*3*,*1*)} *then 1 else 0*)

D11-3: SWAP = CNOT · CNOT′ · CNOT (*SWAP* is the AFP's).

**theorem** *SWAP-three-CNOTs*: *SWAP* = *CNOT* ∗ *CNOT′* ∗ *CNOT*
⟨*proof*⟩

**lemma** *SWAP-basis*:
**assumes** *a* < *2 b* < *2*
**shows** *SWAP* ∗$_v$ *tensor-vec* (*unit-vec 2 a*) (*unit-vec 2 b*) = *tensor-vec* (*unit-vec 2 b*) (*unit-vec 2 a*)
⟨*proof*⟩

**lemma** *CNOT′-basis*:
**assumes** *a* < *2 b* < *2*
**shows** *CNOT′* ∗$_v$ *tensor-vec* (*unit-vec 2 a*) (*unit-vec 2 b*) = *tensor-vec* (*unit-vec 2* ((*a* + *b*) *mod 2*)) (*unit-vec 2 b*)
⟨*proof*⟩

## 15.9 Day 5: a basis of $V \oplus W$ (D5-5)

A basis of $\mathbb{C}^n$ is a list of $n$ vectors whose column matrix is invertible, as in *tensor-right-coordinates*. The notes: if $v_1, \ldots, v_n$ and $w_1, \ldots, w_k$ are bases of $V$ and $W$ then $(v_i, 0)$, $(0, w_j)$ is a basis of $V \oplus W$, so $\dim(V \oplus W) = \dim V + \dim W$. The column matrix is block diagonal.

**definition** *direct-sum-basis* :: *nat* ⇒ *nat* ⇒ *complex Matrix.vec list* ⇒ *complex Matrix.vec list* ⇒ *complex Matrix.vec list* **where**
*direct-sum-basis n k vs ws* = *map* (λ*v*. *v* ⊕$_v$ *0*$_v$ *k*) *vs* @ *map* (λ*w*. *0*$_v$ *n* ⊕$_v$ *w*) *ws*

**lemma** *direct-sum-basis-block*:
**assumes** *vs*: ∀ *v* ∈ *set vs*. *dim-vec v* = *n length vs* = *n* **and** *ws*: ∀ *w* ∈ *set ws*. *dim-vec w* = *k length ws* = *k*
**shows** *mat-of-cols* (*n* + *k*) (*direct-sum-basis n k vs ws*)
= *four-block-mat* (*mat-of-cols n vs*) (*0*$_m$ *n k*) (*0*$_m$ *k n*) (*mat-of-cols k ws*)
⟨*proof*⟩

**theorem** *direct-sum-basis-is-basis*:
**assumes** *vs*: ∀ *v* ∈ *set vs*. *dim-vec v* = *n length vs* = *n* **and** *iv*: *invertible-mat* (*mat-of-cols n vs*)
**and** *ws*: ∀ *w* ∈ *set ws*. *dim-vec w* = *k length ws* = *k* **and** *iw*: *invertible-mat* (*mat-of-cols k ws*)
**shows** ∀ *u* ∈ *set* (*direct-sum-basis n k vs ws*). *dim-vec u* = *n* + *k*
**and** *length* (*direct-sum-basis n k vs ws*) = *n* + *k*
**and** *invertible-mat* (*mat-of-cols* (*n* + *k*) (*direct-sum-basis n k vs ws*))
⟨*proof*⟩

**end**
**theory** *Day05-Cholesky-Examples*
**imports** *QC-Prelims*
**begin**

# 16 Day 5: Cholesky's algorithm on the notes' two examples (D5-3)

## 16.1 One step of Cholesky's algorithm

For $P = (p_{ij})$ with $p_{11} \neq 0$ the notes subtract $R = (\frac{p_{i1}p_{1j}}{p_{11}})_{ij}$; $P - R$ vanishes in the first row and column, and what remains is the $(n-1) \times (n-1)$ matrix *chol-next P*.

**definition** *chol-R* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*chol-R P* = *Matrix.mat* (*dim-row P*) (*dim-row P*) (λ(*i*, *j*). *P* $$ (*i*, *0*) ∗ *P* $$ (*0*, *j*) / *P* $$ (*0*, *0*))

**definition** *chol-next* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*chol-next P* = *Matrix.mat* (*dim-row P* − *1*) (*dim-row P* − *1*) (λ(*i*, *j*). (*P* − *chol-R P*) $$ (*i* + *1*, *j* + *1*))

**lemma** *chol-R-carrier* [*simp*]: *chol-R P* ∈ *carrier-mat* (*dim-row P*) (*dim-row P*)
⟨*proof*⟩

**lemma** *hermitian-entry*:
**assumes** *hermitian P P* ∈ *carrier-mat n n i* < *n j* < *n*
**shows** *P* $$ (*i*, *j*) = *cnj* (*P* $$ (*j*, *i*))
⟨*proof*⟩

The test: for $P = P^*$ with $p_{11} > 0$, $P \geq 0$ iff the next matrix is $\geq 0$ (from *cholesky-reduction*).

**lemma** *chol-test*:
**assumes** *P*: *P* ∈ *carrier-mat n n* **and** *h*: *hermitian P* **and** *n*: *0* < *n* **and** *p*: *P* $$ (*0*, *0*) > *0*
**shows** *psd P* ⟷ *psd* (*chol-next P*)
⟨*proof*⟩

## 16.2 Small matrices entry by entry

**lemma** *mat3-eqI*:
**assumes** *A* ∈ *carrier-mat 3 3 B* ∈ *carrier-mat 3 3*
**and** ⋀*i j*. *i* ∈ {*0*, *1*, *2*} ⟹ *j* ∈ {*0*, *1*, *2*} ⟹ *A* $$ (*i*, *j*) = *B* $$ (*i*, *j*)
**shows** *A* = *B*
⟨*proof*⟩

**lemma** *mat2-eqI*:
**assumes** *A* ∈ *carrier-mat 2 2 B* ∈ *carrier-mat 2 2*
**and** ⋀*i j*. *i* ∈ {*0*, *1*} ⟹ *j* ∈ {*0*, *1*} ⟹ *A* $$ (*i*, *j*) = *B* $$ (*i*, *j*)
**shows** *A* = *B*
⟨*proof*⟩

## 16.3 Example 1: a matrix that is not positive

The notes: for $P = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ 3 & 5 & 7 \end{pmatrix}$, $R = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{pmatrix}$ and $P - R = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & -2 \end{pmatrix}$, so $P \not\geq 0$. Each matrix is checked below; the last step is the test vector $(1, 1)$ for the remaining $2 \times 2$ block.

**definition** *notes-P1* :: *complex Matrix.mat* **where**
*notes-P1* = *mat-of-rows-list 3* [[*1*, *2*, *3*], [*2*, *4*, *5*], [*3*, *5*, *7*]]

**lemma** *notes-P1-carrier*: *notes-P1* ∈ *carrier-mat 3 3*
⟨*proof*⟩

**lemma** *notes-P1-hermitian*: *hermitian notes-P1*
⟨*proof*⟩

**lemma** *notes-P1-R*: *chol-R notes-P1* = *mat-of-rows-list 3* [[*1*, *2*, *3*], [*2*, *4*, *6*], [*3*, *6*, *9*]]
⟨*proof*⟩

**lemma** *notes-P1-diff*: *notes-P1* − *chol-R notes-P1* = *mat-of-rows-list 3* [[*0*, *0*, *0*], [*0*, *0*, −*1*], [*0*, −*1*, −*2*]]
⟨*proof*⟩

**lemma** *notes-P1-next*: *chol-next notes-P1 = mat-of-rows-list 2 [[0, −1], [−1, −2]]*
⟨*proof*⟩

**lemma** *not-psd-witness*:
**assumes** *A*: *A ∈ carrier-mat n n* **and** *x*: *dim-vec x = n* **and** *neg*: *Re (Quantum.inner-prod x (A ∗v x)) < 0*
**shows** *¬ psd A*
⟨*proof*⟩

$\begin{pmatrix} 0 & -1 \\ -1 & -2 \end{pmatrix}$ has the value $-4$ at $(1,1)$, so it is not positive.

**lemma** *notes-P1-next-not-psd*: *¬ psd (mat-of-rows-list 2 [[0, −1], [−1, −2]])*
⟨*proof*⟩

**theorem** *notes-example-1*: *hermitian notes-P1 ∧ ¬ psd notes-P1*
⟨*proof*⟩

## 16.4 Example 2: a positive matrix, in two steps

The notes: for $P = \begin{pmatrix} 4 & 2 & 3 \\ 2 & 5 & 6 \\ 3 & 6 & 8 \end{pmatrix}$, $R = \frac{1}{4}\begin{pmatrix} 16 & 8 & 12 \\ 8 & 4 & 6 \\ 12 & 6 & 9 \end{pmatrix}$ and $P - R = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 4 & \frac{9}{2} \\ 0 & \frac{9}{2} & \frac{23}{4} \end{pmatrix}$. Repeating on $Q = \begin{pmatrix} 4 & \frac{9}{2} \\ \frac{9}{2} & \frac{23}{4} \end{pmatrix}$: $Q - \frac{1}{4}\begin{pmatrix} 4 \\ \frac{9}{2} \end{pmatrix}\begin{pmatrix} 4 & \frac{9}{2} \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & \frac{11}{16} \end{pmatrix}$, so $P \geq 0$.

**definition** *notes-P2* :: *complex Matrix.mat* **where**
*notes-P2 = mat-of-rows-list 3 [[4, 2, 3], [2, 5, 6], [3, 6, 8]]*

**lemma** *notes-P2-carrier*: *notes-P2 ∈ carrier-mat 3 3*
⟨*proof*⟩

**lemma** *notes-P2-hermitian*: *hermitian notes-P2*
⟨*proof*⟩

**lemma** *notes-P2-R*: *chol-R notes-P2 = (1/4) ·m mat-of-rows-list 3 [[16, 8, 12], [8, 4, 6], [12, 6, 9]]*
⟨*proof*⟩

**lemma** *notes-P2-diff*: *notes-P2 − chol-R notes-P2 = mat-of-rows-list 3 [[0, 0, 0], [0, 4, 9/2], [0, 9/2, 23/4]]*
⟨*proof*⟩

**definition** *notes-Q2* :: *complex Matrix.mat* **where**
*notes-Q2 = mat-of-rows-list 2 [[4, 9/2], [9/2, 23/4]]*

**lemma** *notes-P2-next*: *chol-next notes-P2 = notes-Q2*
⟨*proof*⟩

**lemma** *notes-Q2-carrier*: *notes-Q2 ∈ carrier-mat 2 2*
⟨*proof*⟩

**lemma** *notes-Q2-hermitian*: *hermitian notes-Q2*
⟨*proof*⟩

**lemma** *notes-Q2-R*: *chol-R notes-Q2 = (1/4) ·m mat-of-rows-list 2 [[16, 18], [18, 81/4]]*
⟨*proof*⟩

**lemma** *notes-Q2-diff*: *notes-Q2 − chol-R notes-Q2 = mat-of-rows-list 2 [[0, 0], [0, 11/16]]*

⟨*proof*⟩

The last matrix is $(\frac{11}{16}) = \frac{11}{16}\,|e\rangle\langle e|$, which is positive.

**lemma** *notes-Q2-next*: *chol-next notes-Q2 = complex-of-real (11/16)* $\cdot_m$ *(Matrix.vec 1 (λ-. 1)* $\oslash$ *Matrix.vec 1 (λ-. 1))*
⟨*proof*⟩

**lemma** *notes-Q2-next-psd*: *psd (chol-next notes-Q2)*
⟨*proof*⟩

**theorem** *notes-example-2*: *hermitian notes-P2* $\wedge$ *psd notes-P2*
⟨*proof*⟩

The notes' remark: the algorithm writes $P$ as a sum of rank-one matrices, $P = \frac{1}{4}|v_1\rangle\langle v_1| + \frac{1}{4}|v_2\rangle\langle v_2| + \frac{11}{16}|e_3\rangle\langle e_3|$ with $v_1 = (4, 2, 3)$, $v_2 = (0, 4, \frac{9}{2})$, $e_3 = (0, 0, 1)$ (the first columns of $P$, of $P - R$ and of the last step).

**lemma** *outer-prod-of-list*:
*vec-of-list l* $\oslash$ *vec-of-list l = Matrix.mat (length l) (length l) (λ(i, j). l ! i * cnj (l ! j))*
⟨*proof*⟩

**lemma** *notes-P2-rank-one-sum*:
*notes-P2 = complex-of-real (1/4)* $\cdot_m$ *(vec-of-list [4, 2, 3]* $\oslash$ *vec-of-list [4, 2, 3])*
*+ complex-of-real (1/4)* $\cdot_m$ *(vec-of-list [0, 4, 9/2]* $\oslash$ *vec-of-list [0, 4, 9/2])*
*+ complex-of-real (11/16)* $\cdot_m$ *(vec-of-list [0, 0, 1]* $\oslash$ *vec-of-list [0, 0, 1])*
⟨*proof*⟩

**end**
**theory** *Day06-Schmidt-Rank*
**imports** *Day02-11-Supplement*
**begin**

# 17 Day 6: the Schmidt rank (D6-4)

The notes define the Schmidt rank $\mathrm{rank}_S(u)$ of $u \in X \otimes Y$ as the least number of elementary tensors in an expression for $u$, and use on Day 10 that a two-photon state $\psi$ is a product $\psi_1 \otimes \psi_2$ iff $\mathrm{rank}_S(\psi) = 1$; otherwise it is entangled. Here: every $u \in \mathbb{C}^n \otimes \mathbb{C}^m$ has an expression of length $n$, so the least length exists; $\mathrm{rank}_S(u) = 0$ iff $u = 0$; for $u \neq 0$, $\mathrm{rank}_S(u) = 1$ iff $u$ is a product and $u$ is entangled iff $\mathrm{rank}_S(u) > 1$; the Bell state has Schmidt rank 2.

A representation of $u \in \mathbb{C}^n \otimes \mathbb{C}^m$ by $k$ elementary tensors.

**definition** *schmidt-repr* :: *nat* ⇒ *nat* ⇒ *complex Matrix.vec* ⇒ *nat* ⇒ *bool* **where**
*schmidt-repr n m u k* ⟷ (∃ *vs ws. length vs = k* ∧ *length ws = k* ∧
(∀ *i < k. dim-vec (vs ! i) = n)* ∧ (∀ *i < k. dim-vec (ws ! i) = m)* ∧
*u = tensor-combination (n * m) vs ws*)

**lemma** *schmidt-rank-eq*:
**assumes** *u*: *dim-vec u = n * m* **and** *m*: *0 < m*
**shows** *schmidt-rank u m = (LEAST k. schmidt-repr n m u k)*
⟨*proof*⟩

Every $u$ is a sum of $n$ elementary tensors, $u = \sum_i e_i \otimes u_i$ with $u_i = (u_{im+j})_j$, so the least length exists.

**lemma** *schmidt-repr-exists*:
**assumes** *u*: *dim-vec u = n * m*
**shows** *schmidt-repr n m u n*

⟨*proof*⟩

**lemma** *schmidt-rank-repr*:
**assumes** *u*: *dim-vec u* = *n* ∗ *m* **and** *m*: *0* < *m*
**shows** *schmidt-repr n m u* (*schmidt-rank u m*) **and** ⋀*k*. *schmidt-repr n m u k* ⟹ *schmidt-rank u m* ≤ *k*
⟨*proof*⟩

**lemma** *schmidt-repr-0*: *schmidt-repr n m u 0* ⟷ *u* = $0_v$ (*n* ∗ *m*)
⟨*proof*⟩

**lemma** *schmidt-repr-1*:
*schmidt-repr n m u 1* ⟷ (∃ *v w*. *dim-vec v* = *n* ∧ *dim-vec w* = *m* ∧ *u* = *tensor-vec v w*)
⟨*proof*⟩

**theorem** *schmidt-rank-zero-iff*:
**assumes** *u*: *dim-vec u* = *n* ∗ *m* **and** *m*: *0* < *m*
**shows** *schmidt-rank u m* = *0* ⟷ *u* = $0_v$ (*n* ∗ *m*)
⟨*proof*⟩

The notes: $\psi = \psi_1 \otimes \psi_2$ iff $\text{rank}_S(\psi) = 1$ (for $\psi \neq 0$).

**theorem** *schmidt-rank-one-iff-product*:
**assumes** *u*: *dim-vec u* = *n* ∗ *m* **and** *m*: *0* < *m* **and** *u0*: *u* ≠ $0_v$ (*n* ∗ *m*)
**shows** *schmidt-rank u m* = *1* ⟷ (∃ *v w*. *dim-vec v* = *n* ∧ *dim-vec w* = *m* ∧ *u* = *tensor-vec v w*)
⟨*proof*⟩

**corollary** *entangled-iff-schmidt-rank*:
**assumes** *u*: *dim-vec u* = *n* ∗ *m* **and** *m*: *0* < *m* **and** *u0*: *u* ≠ $0_v$ (*n* ∗ *m*)
**shows** *entangled n m u* ⟷ *1* < *schmidt-rank u m*
⟨*proof*⟩

**corollary** *schmidt-rank-bell-state*: *schmidt-rank bell-state 2* = *2*
⟨*proof*⟩

**end**
**theory** *Day10-15-Supplement*
**imports** *QC-Prelims*
**begin**

# 18 Days 10, 11 and 15: superdense coding, the CCNOT table, the partial trace over $A$

## 18.1 Superdense coding (D10-1)

Lab A and Lab B share $\gamma = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$, Lab A holding the first qubit. Lab A multiplies its qubit by $I$, $\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}$, $\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}$ or $\begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}$ and sends it to Lab B, which then holds $\frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle)$ or $\frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle)$: four orthonormal states, so one qubit carries two bits. The four matrices are $I$, $Z$, $X$ and $ZX$ (*sd-op-paulis*).

**lemma** *less4*: *i* < (*4*::*nat*) ⟷ *i* = *0* ∨ *i* = *1* ∨ *i* = *2* ∨ *i* = *3* ⟨*proof*⟩

**lemma** *tensor-I2-apply*:
**fixes** *A* :: *complex Matrix.mat* **and** *v* :: *complex Matrix.vec*
**assumes** *A*: *A* ∈ *carrier-mat 2 2* **and** *v*: *dim-vec v* = *4* **and** *i*: *i* < *4*
**shows** (*Tensor.tensor-mat A* ($1_m$ *2*) ∗$_v$ *v*) $ *i* =

$(\sum j{<}4.\ A\ \$\$\ (i\ div\ 2,\ j\ div\ 2) * (if\ i\ mod\ 2 = j\ mod\ 2\ then\ 1\ else\ 0) * v\ \$\ j)$
⟨*proof*⟩

**definition** *superdense-gamma* :: *complex Matrix.vec* **where**
*superdense-gamma* = *bell-basis* ! 0

Lab A applies one of the notes' four matrices to its qubit.

**definition** *sd-op* :: *nat* ⇒ *complex Matrix.mat* **where**
*sd-op k* = (*if k* = 0 *then mat-of-rows-list* 2 [[1, 0], [0, 1]]
*else if k* = 1 *then mat-of-rows-list* 2 [[1, 0], [0, −1]]
*else if k* = 2 *then mat-of-rows-list* 2 [[0, 1], [1, 0]]
*else mat-of-rows-list* 2 [[0, 1], [−1, 0]])

**lemma** *sd-op-carrier*: *sd-op k* ∈ *carrier-mat* 2 2
⟨*proof*⟩

**lemma** *sd-op-paulis*:
*sd-op* 0 = $1_m$ 2 *sd-op* 1 = *Pauli-Z sd-op* 2 = *Pauli-X sd-op* 3 = *Pauli-Z* ∗ *Pauli-X*
⟨*proof*⟩

The notes' table: the four results are the four Bell states.

**lemma** *superdense-coding-0*: *Tensor.tensor-mat* (*sd-op* 0) ($1_m$ 2) $*_v$ *superdense-gamma* = *bell-basis* ! 0
⟨*proof*⟩

**lemma** *superdense-coding-1*: *Tensor.tensor-mat* (*sd-op* 1) ($1_m$ 2) $*_v$ *superdense-gamma* = *bell-basis* ! 1
⟨*proof*⟩

**lemma** *superdense-coding-2*: *Tensor.tensor-mat* (*sd-op* 2) ($1_m$ 2) $*_v$ *superdense-gamma* = *bell-basis* ! 2
⟨*proof*⟩

**lemma** *superdense-coding-3*: *Tensor.tensor-mat* (*sd-op* 3) ($1_m$ 2) $*_v$ *superdense-gamma* = *bell-basis* ! 3
⟨*proof*⟩

**theorem** *superdense-coding*:
**assumes** *k*: *k* < 4
**shows** *Tensor.tensor-mat* (*sd-op k*) ($1_m$ 2) $*_v$ *superdense-gamma* = *bell-basis* ! *k*
⟨*proof*⟩

**corollary** *superdense-coding-orthonormal*:
*orthonormal-list* 4 (*map* (λ*k*. *Tensor.tensor-mat* (*sd-op k*) ($1_m$ 2) $*_v$ *superdense-gamma*) [0..<4])
⟨*proof*⟩

## 18.2 The CCNOT truth table (D11-2)

**lemma** *tensor-unit-vec*:
**assumes** *i*: *i* < *n* **and** *j*: *j* < *m*
**shows** *tensor-vec* (*unit-vec n i*) (*unit-vec m j*) = (*unit-vec* (*n* ∗ *m*) (*i* ∗ *m* + *j*) :: *complex Matrix.vec*)
⟨*proof*⟩

**lemma** *CCNOT-unit-vec*:
**assumes** *k*: *k* < 8
**shows** *CCNOT* $*_v$ *unit-vec* 8 *k* = *unit-vec* 8 (*if k* = 6 *then* 7 *else if k* = 7 *then* 6 *else k*)
⟨*proof*⟩

On basis states $|a, b, c\rangle = e_a \otimes e_b \otimes e_c$: when $a = 0$ nothing happens; when $a = 1$, CNOT acts on $b, c$.

**theorem** *CCNOT-basis*:
**assumes** *a*: $a < 2$ **and** *b*: $b < 2$ **and** *c*: $c < 2$
**shows** $CCNOT *_v$ *tensor-vec* (*unit-vec 2 a*) (*tensor-vec* (*unit-vec 2 b*) (*unit-vec 2 c*)) =
*tensor-vec* (*unit-vec 2 a*) (*tensor-vec* (*unit-vec 2 b*) (*unit-vec 2* (*if* $a = 1 \wedge b = 1$ *then* $1 - c$ *else* $c$)))
⟨*proof*⟩

**corollary** *CCNOT-truth-table*:
**fixes** *e* :: *nat* ⇒ *complex Matrix.vec*
**defines** $e \equiv \lambda a.$ *unit-vec 2 a*
**shows** $CCNOT *_v$ *tensor-vec* (*e 0*) (*tensor-vec* (*e 0*) (*e 0*)) = *tensor-vec* (*e 0*) (*tensor-vec* (*e 0*) (*e 0*))
**and** $CCNOT *_v$ *tensor-vec* (*e 0*) (*tensor-vec* (*e 0*) (*e 1*)) = *tensor-vec* (*e 0*) (*tensor-vec* (*e 0*) (*e 1*))
**and** $CCNOT *_v$ *tensor-vec* (*e 0*) (*tensor-vec* (*e 1*) (*e 0*)) = *tensor-vec* (*e 0*) (*tensor-vec* (*e 1*) (*e 0*))
**and** $CCNOT *_v$ *tensor-vec* (*e 0*) (*tensor-vec* (*e 1*) (*e 1*)) = *tensor-vec* (*e 0*) (*tensor-vec* (*e 1*) (*e 1*))
**and** $CCNOT *_v$ *tensor-vec* (*e 1*) (*tensor-vec* (*e 0*) (*e 0*)) = *tensor-vec* (*e 1*) (*tensor-vec* (*e 0*) (*e 0*))
**and** $CCNOT *_v$ *tensor-vec* (*e 1*) (*tensor-vec* (*e 0*) (*e 1*)) = *tensor-vec* (*e 1*) (*tensor-vec* (*e 0*) (*e 1*))
**and** $CCNOT *_v$ *tensor-vec* (*e 1*) (*tensor-vec* (*e 1*) (*e 0*)) = *tensor-vec* (*e 1*) (*tensor-vec* (*e 1*) (*e 1*))
**and** $CCNOT *_v$ *tensor-vec* (*e 1*) (*tensor-vec* (*e 1*) (*e 1*)) = *tensor-vec* (*e 1*) (*tensor-vec* (*e 1*) (*e 0*))
⟨*proof*⟩

## 18.3 The partial trace over $A$ (D15-6)

Reordering the basis: for a bijection $\sigma$ of $\{0, \ldots, N-1\}$ the matrix $(x_{\sigma(r)\sigma(s)})$ has the trace of $X$ and is Hermitian, resp. positive, with $X$.

**definition** *perm-conj* :: *nat* ⇒ (*nat* ⇒ *nat*) ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*perm-conj N σ X* = *Matrix.mat N N* (λ(*r*, *s*). *X* $$ (*σ r*, *σ s*))

**lemma** *perm-conj-carrier* [*simp*]: *perm-conj N σ X* ∈ *carrier-mat N N*
⟨*proof*⟩

**lemma** *perm-conj-dims* [*simp*]: *dim-row* (*perm-conj N σ X*) = *N dim-col* (*perm-conj N σ X*) = *N*
⟨*proof*⟩

**lemma** *perm-conj-trace*:
**assumes** *σ*: *bij-betw σ* {..<*N*} {..<*N*} **and** *X*: *X* ∈ *carrier-mat N N*
**shows** *Tr* (*perm-conj N σ X*) = *Tr X*
⟨*proof*⟩

**lemma** *perm-conj-hermitian*:
**assumes** *σ*: ⋀*r*. *r* < *N* ⟹ *σ r* < *N* **and** *X*: *X* ∈ *carrier-mat N N* **and** *h*: *hermitian X*
**shows** *hermitian* (*perm-conj N σ X*)
⟨*proof*⟩

**lemma** *perm-conj-psd*:
**assumes** *σ*: *bij-betw σ* {..<*N*} {..<*N*} **and** *X*: *X* ∈ *carrier-mat N N* **and** *ps*: *psd X*
**shows** *psd* (*perm-conj N σ X*)
⟨*proof*⟩

Exchanging the factors: with $\sigma(im + k) = kp + i$ $(i < p,\ k < m)$, tracing out $A$ is tracing out $B$ after the exchange.

**definition** *swap-index* :: *nat* ⇒ *nat* ⇒ *nat* ⇒ *nat* **where**
*swap-index m p r* = (*r mod m*) ∗ *p* + *r div m*

**lemma** *swap-index-less*:
**assumes** $r < m * p$ **shows** *swap-index m p r* $< m * p$
⟨*proof*⟩

**lemma** *swap-index-bij*: *bij-betw* (*swap-index m p*) {..<$m * p$} {..<$m * p$}
⟨*proof*⟩

**lemma** *partial-trace-A-as-B*:
**assumes** *X*: $X \in$ *carrier-mat* $(m * p)$ $(m * p)$
**shows** *partial-trace-A m p X* = *partial-trace-B p m* (*perm-conj* $(m * p)$ (*swap-index m p*) *X*)
⟨*proof*⟩

**theorem** *partial-trace-A-trace*:
**assumes** *X*: $X \in$ *carrier-mat* $(m * p)$ $(m * p)$ **and** *m*: $0 < m$
**shows** *Tr* (*partial-trace-A m p X*) = *Tr X*
⟨*proof*⟩

**theorem** *partial-trace-A-hermitian*:
**assumes** *X*: $X \in$ *carrier-mat* $(m * p)$ $(m * p)$ **and** *h*: *hermitian X*
**shows** *hermitian* (*partial-trace-A m p X*)
⟨*proof*⟩

**theorem** *partial-trace-A-is-density*:
**assumes** *dm*: *density-matrix* $(m * p)$ *X* **and** *m*: $0 < m$
**shows** *density-matrix p* (*partial-trace-A m p X*)
⟨*proof*⟩

**end**
**theory** *Day04-29-Supplement*
**imports** *Day27-Entangled-Examples*
**begin**

# 19 Days 4–29: the converse Hermitian tests, composite ensembles, Choi's four conditions, codes

Completions of variant rows of the audit: on Day 4 the directions (b)⇒(a) and (c)⇒(a) of the characterisation of Hermitian matrices (D4-1); on Day 14 the density of a composite ensemble (D14-5); on Day 19 Choi's theorem with the $n$-positivity condition (2) stated in its own right (D19-1); on Day 29 the triangle inequality, the distance of a subspace code and unique decoding (D29-4).

## 19.1 Day 4: the converse directions for Hermitian matrices (D4-1)

**lemma** *unitary-real-diag-imp-hermitian*:
**assumes** *U*: $U \in$ *carrier-mat n n* **and** *u*: *unitary U* **and** *H*: $H \in$ *carrier-mat n n*
**and** *D*: *length D* = *n* **and** *spec*: $U^\dagger * H * U =$ *diag-mat-of-real D*
**shows** *hermitian H*
⟨*proof*⟩

**lemma** *eigenbasis-real-imp-hermitian*:
**assumes** *H*: $H \in$ *carrier-mat n n* **and** *orth*: *orthonormal-list n vs* **and** *len*: *length vs* = *n*
**and** *eig*: $\bigwedge i.\ i < n \Longrightarrow H *_v vs\ !\ i =$ *complex-of-real* (*lam* ! *i*) $\cdot_v$ *vs* ! *i* **and** *lam*: *length lam* = *n*
**shows** *hermitian H*
⟨*proof*⟩

## 19.2 Choi's theorem with (2) isolated (D19-1)

**lemma** *choi-seed-psd*: *psd* (*choi-seed n*)
⟨*proof*⟩

**lemma** *choi-psd-if-n-positive*:
**assumes** *n*: $0 < n$ **and** *d*: $0 < d$ **and** *pp*: *p-positive n d n* Φ
**shows** *psd* (*choi-matrix n d* Φ)
⟨*proof*⟩

**theorem** *choi-four-conditions*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: $\bigwedge A\ B.\ A \in$ *carrier-mat n n* $\Longrightarrow B \in$ *carrier-mat n n* $\Longrightarrow$ Φ $(A + B) =$ Φ $A +$ Φ $B$
**and** *smult*: $\bigwedge c\ A.\ A \in$ *carrier-mat n n* $\Longrightarrow$ Φ $(c \cdot_m A) = c \cdot_m$ Φ $A$
**and** *range*: $\forall A.\ A \in$ *carrier-mat n n* $\longrightarrow$ Φ $A \in$ *carrier-mat d d*
**and** *n*: $0 < n$ **and** *d*: $0 < d$
**shows** (*completely-positive n d* Φ $\longleftrightarrow$ *p-positive n d n* Φ)
$\wedge$ (*p-positive n d n* Φ $\longleftrightarrow$ *psd* (*choi-matrix n d* Φ))
$\wedge$ (*psd* (*choi-matrix n d* Φ) $\longleftrightarrow$
(∃ *Ws*. (∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*) ∧ (∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* = *kraus-map-dim d Ws A*)))
⟨*proof*⟩

## 19.3 Composite ensembles (D14-5)

$|\psi \otimes \varphi\rangle\langle\psi \otimes \varphi| = |\psi\rangle\langle\psi| \otimes |\varphi\rangle\langle\varphi|$.

**lemma** *outer-sum-tensor*:
**assumes** ψ: *dim-vec* ψ = *n* **and** φ: *dim-vec* φ = *m*
**shows** *outer-sum* (*n* ∗ *m*) [*tensor-vec* ψ φ] = *outer-sum n* [ψ] ⨂ *outer-sum m* [φ]
⟨*proof*⟩

For ensembles $\{p_i, \psi_i\}$ on $\mathcal{H}_A$ and $\{q_j, \varphi_j\}$ on $\mathcal{H}_B$, the composite ensemble is $\{p_i q_j, \psi_i \otimes \varphi_j\}$.

**definition** *product-ensemble* :: (*real* × *complex Matrix.vec*) *list* ⇒ (*real* × *complex Matrix.vec*) *list* ⇒ (*real* × *complex Matrix.vec*) *list* **where**
*product-ensemble EA EB* = *map* (λ((*p*, ψ), (*q*, φ)). (*p* ∗ *q*, *tensor-vec* ψ φ)) (*List.product EA EB*)

**lemma** *sum-list-product*:
**fixes** *f* :: *'a* ⇒ *real* **and** *g* :: *'b* ⇒ *real*
**shows** (∑ (*x*, *y*) ← *List.product xs ys*. *f x* ∗ *g y*) = (∑ *x* ← *xs*. *f x*) ∗ (∑ *y* ← *ys*. *g y*)
⟨*proof*⟩

**lemma** *product-ensemble-ensemble*:
**assumes** *A*: *ensemble n EA* **and** *B*: *ensemble m EB*
**shows** *ensemble* (*n* ∗ *m*) (*product-ensemble EA EB*)
⟨*proof*⟩

The notes' composite density $\sum_{i,j} p_i q_j\, |\psi_i\rangle\langle\psi_i| \otimes |\varphi_j\rangle\langle\varphi_j|$ is the density of the composite ensemble, hence a density matrix.

**definition** *composite-density* :: *nat* ⇒ *nat* ⇒ (*real* × *complex Matrix.vec*) *list* ⇒ (*real* × *complex Matrix.vec*) *list* ⇒ *complex Matrix.mat* **where**
*composite-density n m EA EB* = *foldr* (λ((*p*, ψ), (*q*, φ)) *acc*.
*complex-of-real* (*p* ∗ *q*) ·$_m$ (*outer-sum n* [ψ] ⨂ *outer-sum m* [φ]) + *acc*) (*List.product EA EB*) (0$_m$ (*n* ∗ *m*) (*n* ∗ *m*))

**lemma** *composite-density-eq*:
**assumes** *A*: *ensemble n EA* **and** *B*: *ensemble m EB*
**shows** *composite-density n m EA EB* = *ensemble-density* (*n* ∗ *m*) (*product-ensemble EA EB*)

⟨*proof*⟩

**theorem** *composite-density-is-dm*:
**assumes** *A*: *ensemble n EA* **and** *B*: *ensemble m EB*
**shows** *density-matrix* (*n* ∗ *m*) (*composite-density n m EA EB*)
⟨*proof*⟩

## 19.4 Hamming distance and subspace codes (D29-4)

**definition** *bit-xor* :: *nat list* ⇒ *nat list* ⇒ *nat list* **where**
*bit-xor v w* = *map2* (λ*a b*. (*a* + *b*) *mod 2*) *v w*

**lemma** *hamming-dist-xor*: *hamming-dist v w* = *hamming-weight* (*bit-xor v w*)
⟨*proof*⟩

**lemma** *hamming-weight-Cons'*: *hamming-weight* (*a* # *v*) = (*if a* ≠ *0 then Suc* (*hamming-weight v*) *else hamming-weight v*)
⟨*proof*⟩

The triangle inequality for words of bits of equal length.

**lemma** *hamming-dist-triangle*:
**assumes** *length u* = *length v length v* = *length w*
**and** ∀ *x* ∈ *set u*. *x* < *2* ∀ *x* ∈ *set v*. *x* < *2* ∀ *x* ∈ *set w*. *x* < *2*
**shows** *hamming-dist u w* ≤ *hamming-dist u v* + *hamming-dist v w*
⟨*proof*⟩

For a subspace code (closed under addition mod 2) the distance is the least weight of a nonzero code word.

**definition** *zero-word* :: *nat* ⇒ *nat list* **where**
*zero-word m* = *replicate m 0*

**lemma** *bit-xor-self*: *bit-xor v v* = *zero-word* (*length v*)
⟨*proof*⟩

**lemma** *bit-xor-zero*: ∀ *x* ∈ *set v*. *x* < *2* ⟹ *bit-xor v* (*zero-word* (*length v*)) = *v*
⟨*proof*⟩

**lemma** *bit-xor-eq-zero-iff*:
**assumes** *length v* = *length w* ∀ *x* ∈ *set v*. *x* < *2* ∀ *x* ∈ *set w*. *x* < *2*
**shows** *bit-xor v w* = *zero-word* (*length v*) ⟷ *v* = *w*
⟨*proof*⟩

**lemma** *hamming-weight-zero-word* [*simp*]: *hamming-weight* (*zero-word m*) = *0*
⟨*proof*⟩

**lemma** *hamming-weight-eq-zero-iff*: ∀ *x* ∈ *set v*. *x* < *2* ⟹ *hamming-weight v* = *0* ⟷ *v* = *zero-word* (*length v*)
⟨*proof*⟩

**definition** *subspace-code* :: *nat* ⇒ *nat list set* ⇒ *bool* **where**
*subspace-code m S* ⟷ (∀ *v* ∈ *S*. *length v* = *m* ∧ (∀ *x* ∈ *set v*. *x* < *2*))
∧ (∀ *v* ∈ *S*. ∀ *w* ∈ *S*. *bit-xor v w* ∈ *S*)

**lemma** *bit-xor-bits*: ∀ *x* ∈ *set* (*bit-xor v w*). *x* < *2*
⟨*proof*⟩

**lemma** *length-bit-xor* [*simp*]: *length* (*bit-xor v w*) = *min* (*length v*) (*length w*)
⟨*proof*⟩

**theorem** *subspace-code-distance*:
**assumes** *S*: *subspace-code m S* **and** *fin*: *finite S* **and** *two*: $\exists\, v \in S.\ \exists\, w \in S.\ v \neq w$
**shows** *code-distance S* = *Min* {*hamming-weight v* | *v*. $v \in S \wedge v \neq$ *zero-word m*}
⟨*proof*⟩

A code of distance at least $2t + 1$ corrects $t$ errors: a received word is within distance $t$ of at most one code word.

**theorem** *corrects-errors-unique-decoding*:
**assumes** *ce*: *corrects-errors S t* **and** *fin*: *finite S*
**and** *S*: $\forall\, v \in S.$ *length* $v = m \wedge (\forall\, x \in$ *set v*. $x < 2)$
**and** *r*: *length* $r = m\ \forall\, x \in$ *set r*. $x < 2$
**and** *c*: $c \in S$ *hamming-dist r c* $\leq t$ **and** *c'*: $c' \in S$ *hamming-dist r c'* $\leq t$
**shows** $c = c'$
⟨*proof*⟩

**end**

# Part II

# Lectures 11–22 and 29–34: quantum computing, density matrices, completely positive maps, error correction

**theory** *Quantum-Computing11*
**imports**
*Vern-Paulsen-QC-3-10.Basic-Math*
*Vern-Paulsen-QC-3-10.Quantum-Computing-3-10*
**begin**

## 20 Day 11 (21 September 2011): gates revisited, circuit diagrams, no-cloning

The lecture opens with two corrections: $H^2 = I$ (not $X$), with $He_0 = (e_0 + e_1)/\sqrt{2}$, $Xe_0 = e_1$, $Xe_1 = e_0$, $He_1 = (e_0 - e_1)/\sqrt{2}$, so that $X$ and $H$ are reflections; and the truth tables of CNOT ($|a, b\rangle \mapsto |a, b + a\rangle$, arithmetic in $\mathbb{Z}_2$) and of CCNOT ($|a, b, c\rangle \mapsto |a, b, c + ab\rangle$). Circuit diagrams are introduced; the CNOT with control on the second wire has matrix $\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$, and three alternating CNOTs compose to the swap of the two wires (*SWAP-three-CNOTs* in theory Day02_11_Supplement).

**Definition (notes).** The Toffoli gate CCNOT is the $8 \times 8$ permutation matrix that fixes the first six basis vectors and exchanges $|110\rangle$ and $|111\rangle$.

**definition** *CCNOT* :: *complex Matrix.mat* **where**
*CCNOT* ≡ *Matrix.mat 8 8*
(λ(*i*,*j*). *if* ($i = 6 \wedge j = 7$) ∨ ($i = 7 \wedge j = 6$)

*then 1*
*else if* $i = j \wedge i < 6$ *then 1 else 0*)

**lemma** *CCNOT-dagger* [*simp*]:
$CCNOT^\dagger = CCNOT$
⟨*proof*⟩

**lemma** *CCNOT-dim-row* [*simp*]:
*dim-row CCNOT* = *8*
⟨*proof*⟩

**lemma** *CCNOT-dim-col* [*simp*]:
*dim-col CCNOT* = *8*
⟨*proof*⟩

**lemma** *CCNOT-inv* [*simp*]:
$CCNOT * CCNOT = 1_m\ 8$
⟨*proof*⟩

**lemma** *CCNOT-unitary*: *unitary CCNOT*
⟨*proof*⟩

## 20.1 Cloning and no-cloning

CNOT clones the computational basis: $U(e_a \otimes e_0) = e_a \otimes e_a$, and more generally $|i_1 \cdots i_n\rangle \otimes |0 \cdots 0\rangle \mapsto |i_1 \cdots i_n\rangle \otimes |i_1 \cdots i_n\rangle$ for some unitary. But for $\psi = \alpha e_0 + \beta e_1$ with $\alpha\beta \neq 0$, $U(\psi \otimes e_0) \neq \psi \otimes \psi$.

**Theorem (notes)** (No-cloning)**.** There is no state $\phi$ and unitary $U$ on $\mathcal{H} \otimes \mathcal{H}$ with $U(\psi \otimes \phi) = \psi \otimes \psi$ for all $\psi \in \mathcal{H}$.

The notes argue with $-\psi$; the formal proof (for $\dim \mathcal{H} = n \geq 2$ and unit vectors $\psi$) uses two states with inner product neither 0 nor 1.

**definition** *is-cloner* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *bool* **where**
*is-cloner n U* $\varphi \equiv$
$U \in$ *carrier-mat* ($n\hat{}2$) ($n\hat{}2$) $\wedge$ *unitary U* $\wedge$ *dim-vec* $\varphi = n \wedge$
($\forall \psi$. *quantum-state n* $\psi \longrightarrow$
$U *_v$ *tensor-vec* $\psi$ $\varphi$ = *tensor-vec* $\psi$ $\psi$)

Two states whose inner product is neither 0 nor 1 exist once $n \geq 2$.

**lemma** *two-states-nontrivial-inner-prod*:
**assumes** *n*: *2* ≤ *n*
**obtains** $\psi_1$ $\psi_2$ :: *complex Matrix.vec*
**where** *quantum-state n* $\psi_1$ **and** *quantum-state n* $\psi_2$
**and** $\langle \psi_1 \mid \psi_2 \rangle \neq$ (*0* :: *real*) **and** $\langle \psi_1 \mid \psi_2 \rangle \neq$ (*1* :: *real*)
⟨*proof*⟩

**theorem** *no-cloning*:
**assumes** *n2*: *2* ≤ *n*
**shows** ¬ ($\exists$ *U* $\varphi$. *is-cloner n U* $\varphi$)
⟨*proof*⟩

# 21 Day 12 (23 September 2011): quantum parallelism

Since $He_0 = (e_0 + e_1)/\sqrt{2}$,

$$H^{\otimes N}(e_0 \otimes \cdots \otimes e_0) = \left(\tfrac{1}{\sqrt{2}}\right)^N \sum_{J \in \mathbb{Z}_2^N} e_J = \left(\tfrac{1}{\sqrt{2}}\right)^N \sum_J |J\rangle.$$

Only the case $N = 1$ is stated here; the general identity is *hadamard-uniform-N* in *Day12-Counting-Probability* (with *tensor-pow-mat* for $H^{\otimes N}$).

**Definition (notes)** (Oracle)**.** For $f : \mathbb{Z}_2^N \to \mathbb{Z}_2$, the vectors $e_J \otimes e_{i+f(J)}$ form an orthonormal basis of $(\mathbb{C}^2)^{\otimes N} \otimes \mathbb{C}^2$, so there is a unitary $U_f$ with $U_f(e_J \otimes e_i) = e_J \otimes e_{i+f(J)}$, the *oracle* for $f$.

**Proposition (notes).** For a unit vector $\gamma$, the measurement $P_0 = |\gamma\rangle\langle\gamma|$, $P_1 = I - P_0$ has $p_0(\psi) = |\langle\gamma|\psi\rangle|^2$ and $p_1(\psi) = 1 - |\langle\gamma|\psi\rangle|^2$.

Application: with $x = (He_0)^{\otimes N}$, $U_f(x \otimes e_0) = 2^{-N/2} \sum_J e_J \otimes e_{f(J)}$, and measuring against $\gamma = x \otimes e_0$ gives $\langle\gamma|U_f(x \otimes e_0)\rangle = M/2^N$ with $M = \#\{J : f(J) = 0\}$, hence $p_0 = (M/2^N)^2$: repeating the experiment estimates $M$ (a Bernoulli trial). The existence of the oracle is formalised below; the proposition is *rank-one-meas-prob* (Day 8); the counting identity is *counting-probability* in *Day12-Counting-Probability*.

**lemma** *hadamard-uniform-1*:
*Hadamard* $*_v$ *unit-vec 2 0* =
*(1/csqrt 2)* $\cdot_v$ *(unit-vec 2 0 + unit-vec 2 1)*
⟨*proof*⟩

**definition** *oracle-unitary* :: *(nat ⇒ nat) ⇒ nat ⇒ bool* **where**
*oracle-unitary f N* ≡
∃ *U* :: *complex Matrix.mat*.
*U* ∈ *carrier-mat (2^(N+1)) (2^(N+1))* ∧
*unitary U* ∧
(∀ *J* < *2^N*. ∀ *i* < *2*.
*U* $*_v$ *(tensor-vec (unit-vec (2^N) J) (unit-vec 2 i)) =*
*tensor-vec (unit-vec (2^N) J) (unit-vec 2 ((i + f J) mod 2))*)

**theorem** *oracle-exists*:
**fixes** *f* :: *nat ⇒ nat*
**assumes** *f-range*: ∀ *J* < *2^(N* :: *nat)*. *f J* < *2*
**shows** *oracle-unitary f N*
⟨*proof*⟩

$M = \#\{J < 2^N : f(J) = 0\}$, the quantity estimated by the counting experiment.

**definition** *count-zeros* :: *(nat ⇒ nat) ⇒ nat ⇒ nat* **where**
*count-zeros f N* ≡ *card* {*J*. *J* < *2^N* ∧ *f J* = *0*}

# 22 Day 13 (26 September 2011): ensembles and density matrices

## 22.1 Ensembles

Motivation: measuring $\psi = a|0\rangle + b|1\rangle$ with $M_0 = |0\rangle\langle 0|$, $M_1 = |1\rangle\langle 1|$ leaves the state $\frac{a}{|a|}|0\rangle$ with probability $|a|^2$ and $\frac{b}{|b|}|1\rangle$ with probability $|b|^2$; a later measurement $\{\tilde{M}_\alpha\}$ then has outcome probabilities $p_\alpha = |a|^2\|\tilde{M}_\alpha \frac{a}{|a|}|0\rangle\|^2 + |b|^2\|\tilde{M}_\alpha \frac{b}{|b|}|1\rangle\|^2$.

**Definition (notes).** An *ensemble* $\{p_i, \psi_i\}_{i=1}^{L}$ is a set of states with probabilities $p_i \geq 0$, $\sum_i p_i = 1$. For a measurement system $\{M_\alpha\}$ the probability of outcome $\alpha$ is $p_\alpha(\{p_i, \psi_i\}) = \sum_i p_i \|M_\alpha \psi_i\|^2$.

A second source of ensembles: a unit vector $\psi = \sum_l \phi_l \otimes f_l \in \mathcal{H}_A \otimes \mathcal{H}_E$ (environment $\mathcal{H}_E$ with orthonormal basis $f_l$) behaves, for measurements $M_\alpha \otimes I$, exactly like the ensemble $\{\|\phi_l\|^2, \phi_l/\|\phi_l\|\}$.

**definition** *ensemble* :: *nat* ⇒ (*real* × *complex Matrix.vec*) *list* ⇒ *bool* **where**
*ensemble n E* ≡
(∀ (*p*, *ψ*) ∈ *set E*. *p* ≥ *0* ∧ *quantum-state n ψ*) ∧
(∑ (*p*, -) ← *E*. *p*) = *1*

## 22.2 Von Neumann's density matrix

Measurements cannot distinguish $\psi$ from $e^{i\theta}\psi$, and $|e^{i\theta}\psi\rangle\langle e^{i\theta}\psi| = |\psi\rangle\langle\psi| = (\alpha_i \overline{\alpha_j})$: the object attached to a state is the projection, not the vector. For any $\gamma$, $\|\gamma\|^2 = \mathrm{Tr}\,|\gamma\rangle\langle\gamma|$, and

$$p_\alpha(\psi) = \|M_\alpha\psi\|^2 = \mathrm{Tr}\big((M_\alpha^* M_\alpha)(\psi\psi^*)\big), \qquad p_\alpha(\{p_l, \psi_l\}) = \mathrm{Tr}\Big((M_\alpha^* M_\alpha)\sum_l p_l \psi_l \psi_l^*\Big).$$

**Definition (notes).** $P = \sum_l p_l |\psi_l\rangle\langle\psi_l|$ is the *density matrix* of the ensemble.

Formally *pure-density* $\psi$ is $|\psi\rangle\langle\psi|$, *ensemble-density n E* is $P$, and *density-matrix n ϱ* is the Day 14 characterisation ($\rho \geq 0$, $\mathrm{Tr}\,\rho = 1$).

**definition** *density-matrix* :: *nat* ⇒ *complex Matrix.mat* ⇒ *bool* **where**
*density-matrix n ϱ* ≡ *ϱ* ∈ *carrier-mat n n* ∧ *psd ϱ* ∧ *Tr ϱ* = *1*

**definition** *pure-density* :: *complex Matrix.vec* ⇒ *complex Matrix.mat* **where**
*pure-density ψ* ≡
*let n* = *dim-vec ψ in*
*outer-sum n* [*ψ*]

**lemma** *Tr-outer-sum-single*:
**assumes** *dim*: *dim-vec ψ* = *n*
**shows** *Tr* (*outer-sum n* [*ψ*]) = ⟨*ψ* | *ψ*⟩
⟨*proof*⟩

**lemma** *pure-density-is-density-matrix*:
**assumes** *quantum-state n ψ*
**shows** *density-matrix n* (*pure-density ψ*)
⟨*proof*⟩

**definition** *ensemble-density* :: *nat* ⇒ (*real* × *complex Matrix.vec*) *list* ⇒ *complex Matrix.mat* **where**
*ensemble-density n E* ≡
*foldr* (λ(*p*,*ψ*) *acc*. *complex-of-real p* ·$_m$ *outer-sum n* [*ψ*] + *acc*)
*E* (*0*$_m$ *n n*)

**lemma** *psd-zero-mat* [*simp*]:
*psd* (*0*$_m$ *n n*)
⟨*proof*⟩

**lemma** *psd-smult-nonneg*:
**assumes** *c*: *0* ≤ *c*
**and** *A*: *psd A* **and** *Ac*: *A* ∈ *carrier-mat n n*

**shows** *psd* (*complex-of-real c* $\cdot_m$ *A*)
⟨*proof*⟩

**lemma** *psd-add-psd*:
**assumes** *A*: *psd A* **and** *B*: *psd B*
**and** *Ac*: *A* ∈ *carrier-mat n n* **and** *Bc*: *B* ∈ *carrier-mat n n*
**shows** *psd* (*A* + *B*)
⟨*proof*⟩

**lemma** *ensemble-density-Nil* [*simp*]:
*ensemble-density n* [] = $0_m$ *n n*
⟨*proof*⟩

**lemma** *ensemble-density-Cons*:
*ensemble-density n* ((*p*, *ψ*) # *E*) =
*complex-of-real p* $\cdot_m$ *outer-sum n* [*ψ*] + *ensemble-density n E*
⟨*proof*⟩

**lemma** *ensemble-density-carrier*:
**assumes** ∀(*p*, *ψ*) ∈ *set E*. *quantum-state n ψ*
**shows** *ensemble-density n E* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *ensemble-density-psd-aux*:
**assumes** ∀(*p*, *ψ*) ∈ *set E*. *p* ≥ *0* ∧ *quantum-state n ψ*
**shows** *psd* (*ensemble-density n E*)
⟨*proof*⟩

**lemma** *Tr-zero-mat′* [*simp*]:
*Tr* ($0_m$ *n n*) = *0*
⟨*proof*⟩

**lemma** *ensemble-density-Tr*:
**assumes** ∀(*p*, *ψ*) ∈ *set E*. *quantum-state n ψ*
**shows** *Tr* (*ensemble-density n E*) = *complex-of-real* (∑ (*p*, -) ← *E*. *p*)
⟨*proof*⟩

**theorem** *ensemble-density-is-dm*:
**assumes** *E*: *ensemble n E*
**shows** *density-matrix n* (*ensemble-density n E*)
⟨*proof*⟩

## 22.3 Probabilities via the trace

$p_\alpha(\psi) = \|M_\alpha\psi\|^2 = \mathrm{Tr}(M_\alpha^* M_\alpha |\psi\rangle\langle\psi|)$. For an ensemble with density matrix $\rho$, $p_\alpha = \mathrm{Tr}(M_\alpha^* M_\alpha \rho) = \langle M_\alpha^* M_\alpha | \rho \rangle$ (*ensemble-prob-trace-eq* below).

**lemma** *pure-state-prob-trace*:
**fixes** *M* :: *complex Matrix.mat* **and** *ψ* :: *complex Matrix.vec*
**assumes** *st*: *quantum-state n ψ*
**and** *Msq*: *M* ∈ *carrier-mat n n*
**shows** *meas-prob M ψ* = *Re* (*Tr* ((*M*†) ∗ *M* ∗ *outer-sum n* [*ψ*]))
⟨*proof*⟩

For an ensemble: $p_\alpha(\{p_i, \psi_i\}) = \sum_i p_i \|M_\alpha \psi_i\|^2 = \mathrm{Tr}(M_\alpha^* M_\alpha \rho)$.

**lemma** *ensemble-prob-trace-eq*:

**fixes** $M$ :: *complex Matrix.mat*
**assumes** $M$: $M \in$ *carrier-mat n n*
**and** $E$: $\forall (p, \psi) \in$ *set E. quantum-state n* $\psi$
**shows** $(\sum (p, \psi) \leftarrow E.\ p * \textit{meas-prob}\ M\ \psi)$
$= \textit{Re}\ (\textit{Tr}\ (((M^\dagger) * M) * \textit{ensemble-density}\ n\ E))$
⟨*proof*⟩

**lemma** *Tr-psd-Re-nonneg*:
**assumes** $P$: *psd P* **and** $Pc$: $P \in$ *carrier-mat n n*
**shows** $0 \leq \textit{Re}\ (\textit{Tr}\ P)$
⟨*proof*⟩

**lemma** *ensemble-prob-trace*:
**fixes** *Ms* :: *complex Matrix.mat list*
**assumes** *ms*: *measurement-system n Ms*
**and** *dm*: *density-matrix n* $\varrho$
**and** *Min*: $M \in$ *set Ms*
**shows** $\textit{Re}\ (\textit{Tr}\ ((M^\dagger) * M * \varrho)) \geq 0$
⟨*proof*⟩

# 23 Day 14 (28 September 2011): density matrices, pure states, composite ensembles

**Proposition (notes).** $\rho \in L(\mathcal{H}_A)$ is the density matrix of some ensemble if and only if $\rho \geq 0$ and $\operatorname{Tr}\rho = 1$.

**Definition (notes).** $\rho$ is a *density matrix* when $\rho \geq 0$ and $\operatorname{Tr}\rho = 1$. The ensemble $\{\lambda_i, \psi_i\}$ of eigenvalues and an orthonormal eigenbasis is the *spectral ensemble*.

Many ensembles have the same density matrix: writing $\rho = XX^* = \sum_i c_i c_i^*$ with columns $c_i$ gives the ensemble $\{\|c_i\|^2, c_i/\|c_i\|\}$.

**theorem** *density-matrix-characterisation*:
**fixes** $\varrho$ :: *complex Matrix.mat*
**assumes** *sq*: $\varrho \in$ *carrier-mat n n*
**shows** *density-matrix n* $\varrho \longleftrightarrow$ *psd* $\varrho \wedge$ *Tr* $\varrho = 1$
⟨*proof*⟩

**lemma** *ensemble-density-carrier-dim*:
**shows** *ensemble-density n E* $\in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *outer-sum-singleton-index*:
**assumes** $r$: $r < n$ **and** $c$: $c < n$
**shows** *outer-sum n* $[\psi]$ $\$\$\ (r, c) = \psi\ \$\ r * \textit{cnj}\ (\psi\ \$\ c)$
⟨*proof*⟩

Day 13: $|e^{i\theta}\psi\rangle\langle e^{i\theta}\psi| = |\psi\rangle\langle\psi|$; the density matrix forgets the phase.

**lemma** *pure-density-smult-unit*:
**fixes** $\psi$ :: *complex Matrix.vec* **and** $c$ :: *complex*
**assumes** $c$: *cmod c = 1*
**shows** *pure-density* $(c \cdot_v \psi) =$ *pure-density* $\psi$
⟨*proof*⟩

**lemma** *ensemble-density-index*:
**assumes** $r$: $r < n$ **and** $c$: $c < n$

**shows** *ensemble-density n E $$ (r, c) =*
*(∑ (p, ψ) ← E. complex-of-real p * ψ $ r * cnj (ψ $ c))*
⟨*proof*⟩

The spectral ensemble: every density matrix comes from an ensemble.

**theorem** *spectral-ensemble-exists*:
**fixes** *ϱ :: complex Matrix.mat*
**assumes** *dm*: *density-matrix n ϱ*
**shows** *∃ E. ensemble n E ∧ ensemble-density n E = ϱ*
⟨*proof*⟩

**corollary** *density-matrix-iff-ensemble*:
**fixes** *ϱ :: complex Matrix.mat*
**shows** *density-matrix n ϱ ⟷*
*(∃ E. ensemble n E ∧ ensemble-density n E = ϱ)*
⟨*proof*⟩

## 23.1 Pure states

If $\rho = |\psi\rangle\langle\psi| = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ then for $\gamma \perp \psi$, $0 = \langle\gamma|\rho\gamma\rangle = \sum_i p_i |\langle\psi_i|\gamma\rangle|^2$, so every $\psi_i$ with $p_i > 0$ is parallel to $\psi$: $\psi_i = \beta_i \psi$ with $|\beta_i| = 1$ and $|\psi_i\rangle\langle\psi_i| = |\psi\rangle\langle\psi|$. These density matrices are the *pure states*.

**definition** *pure-state-density :: nat ⇒ complex Matrix.mat ⇒ bool* **where**
*pure-state-density n ϱ ≡*
*density-matrix n ϱ ∧*
*(∃ ψ. quantum-state n ψ ∧ ϱ = pure-density ψ)*

**lemma** *pure-state-density-is-density*:
**assumes** *pure-state-density n ϱ*
**shows** *density-matrix n ϱ*
⟨*proof*⟩

**lemma** *pure-density-is-pure-state-density*:
**assumes** *quantum-state n ψ*
**shows** *pure-state-density n (pure-density ψ)*
⟨*proof*⟩

A companion fact not in the notes: a density matrix is pure iff it is idempotent.

**theorem** *pure-density-iff-idempotent*:
**fixes** *ϱ :: complex Matrix.mat*
**assumes** *dm*: *density-matrix n ϱ*
**shows** *pure-state-density n ϱ ⟷ ϱ * ϱ = ϱ*
⟨*proof*⟩

**lemma** *outer-sum-singleton-mult-vec*:
**assumes** *ψ-dim*: *dim-vec ψ = n* **and** *γ-dim*: *dim-vec γ = n*
**shows** *outer-sum n [ψ] *v γ = ⟨ψ|γ⟩ ·v ψ*
⟨*proof*⟩

**lemma** *ensemble-density-quad-form*:
**assumes** *E-dim*: *∀ (p, ψ) ∈ set E. dim-vec ψ = n*
**and** *γ-dim*: *dim-vec γ = n*
**shows** *⟨γ | ensemble-density n E *v γ⟩ =*
*(∑ (p, ψ) ← E. complex-of-real p * ⟨ψ|γ⟩ * cnj ⟨ψ|γ⟩)*
⟨*proof*⟩

**theorem** *pure-ensemble-parallel*:
**fixes** $\psi$ :: *complex Matrix.vec*
**assumes** *st*: *quantum-state n* $\psi$
**and** *ens*: *ensemble n E*
**and** *dens*: *ensemble-density n E* = *pure-density* $\psi$
**and** *mem*: $(p, \psi_i) \in$ *set E*
**and** *ppos*: $0 < p$
**shows** $\exists \beta.\ \psi_i = \beta \cdot_v \psi$
⟨*proof*⟩

### 23.2 Composite ensembles

For ensembles $\{p_i, \psi_i\}$ in lab A and $\{q_l, \phi_l\}$ in lab B the composite is the ensemble $\{p_i q_l, \psi_i \otimes \phi_l\}$ with density matrix $\rho = \sum_{i,l} p_i q_l |\psi_i\rangle\langle\psi_i| \otimes |\phi_l\rangle\langle\phi_l|$, because $|\psi \otimes \phi\rangle\langle\psi \otimes \phi| = |\psi\rangle\langle\psi| \otimes |\phi\rangle\langle\phi|$ (checked on elementary tensors $h \otimes k$). The identity is formalised below; that $\rho_1 \otimes \rho_2$ is again a density matrix is *density-matrix-tensor* (Day 15).

**lemma** *tensor-pure-density*:
**fixes** $\psi$ $\varphi$ :: *complex Matrix.vec*
**assumes** $\psi$-*dim*: *dim-vec* $\psi$ = *m*
**and** $\varphi$-*dim*: *dim-vec* $\varphi$ = *p*
**and** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
**shows** *pure-density* (*tensor-vec* $\psi$ $\varphi$) =
(*pure-density* $\psi$) ⨂ (*pure-density* $\varphi$)
⟨*proof*⟩

### 23.3 Partial traces: motivation

For a density matrix $\rho \in L(\mathcal{H}_A \otimes \mathcal{H}_B)$ and measurements $\{M_\alpha\}$ in lab A, which act as $M_\alpha \otimes I$, $p_\alpha(\rho) = \mathrm{Tr}((M_\alpha^* M_\alpha \otimes I)\rho)$. The notes announce a density matrix $\rho_A \in L(\mathcal{H}_A)$ with $\mathrm{Tr}_{\mathcal{H}_A \otimes \mathcal{H}_B}((M^*M \otimes I)\rho) = \mathrm{Tr}_{\mathcal{H}_A}(M^* M \rho_A)$ for all $M$; it is constructed on Day 15.

## 24 Day 15 (30 September 2011): traces, tensor products of operators, partial traces

**Proposition (notes).** For any orthonormal basis $\{u_1, \ldots, u_n\}$ of $\mathbb{C}^n$, $\mathrm{tr}(A) = \sum_i \langle u_i | A u_i \rangle$.

**theorem** *trace-via-onb*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *sq*: *A* ∈ *carrier-mat n n*
**and** *orth*: *orthonormal-list n vs*
**and** *full*: *length vs* = *n*
**shows** *Tr A* = (∑ *i*<*n*. ⟨*vs* ! *i* | *A* ∗$_v$ *vs* ! *i*⟩)
⟨*proof*⟩

**Proposition (notes).** $\mathrm{tr}(R \otimes T) = \mathrm{tr}(R)\,\mathrm{tr}(T)$.

**theorem** *trace-tensor*:
**fixes** *R T* :: *complex Matrix.mat*
**assumes** *Rsq*: *R* ∈ *carrier-mat m m*
**and** *Tsq*: *T* ∈ *carrier-mat p p*
**and** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
**shows** *Tr* (*R* ⨂ *T*) = *Tr R* ∗ *Tr T*
⟨*proof*⟩

## 24.1 The isomorphism $L(\mathcal{H}_A) \otimes L(\mathcal{H}_B) \cong L(\mathcal{H}_A \otimes \mathcal{H}_B)$

**Proposition (notes).** In finite dimensions the map $\Gamma : L(\mathcal{H}_A) \otimes L(\mathcal{H}_B) \to L(\mathcal{H}_A \otimes \mathcal{H}_B)$, $R \otimes T \mapsto R \otimes T$, is a vector-space isomorphism (both sides have dimension $m^2p^2$ and $\Gamma$ is injective).

Formally, $L(\mathcal{H}_A) \otimes L(\mathcal{H}_B)$ is represented in coordinates as $M_{m^2,p^2}$ and $\Gamma$ as the reindexing $M_{m^2,p^2} \to M_{mp}$; it is an explicit linear isomorphism with explicit inverse.

Postulate $4'$ (Day 16) and the composite ensembles of Day 14: tensor products of positive matrices are positive, and tensor products of density matrices are density matrices.

**lemma** *psd-tensor*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A*: *A* ∈ *carrier-mat m m* **and** *B*: *B* ∈ *carrier-mat p p*
**and** *pA*: *psd A* **and** *pB*: *psd B*
**shows** *psd* (*A* ⊗ *B*)
⟨*proof*⟩

**lemma** *density-matrix-tensor*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *dA*: *density-matrix m A* **and** *dB*: *density-matrix p B*
**and** *m*: *0* < *m* **and** *p*: *0* < *p*
**shows** *density-matrix* (*m* ∗ *p*) (*A* ⊗ *B*)
⟨*proof*⟩

**proposition** *L-tensor-dim*:
**fixes** *m p* :: *nat*
**shows** *m*^2 ∗ *p*^2 = (*m* ∗ *p*)^2
⟨*proof*⟩

**definition** *operator-tensor-iso-map* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*operator-tensor-iso-map m p c* ≡
*Matrix.mat* (*m*∗*p*) (*m*∗*p*)
(λ(*r*,*s*). *c* $$ ((*r div p*) ∗ *m* + (*s div p*), (*r mod p*) ∗ *p* + (*s mod p*)))

**lemma** *operator-tensor-iso-map-carrier* [*simp*]:
*operator-tensor-iso-map m p c* ∈ *carrier-mat* (*m*∗*p*) (*m*∗*p*)
⟨*proof*⟩

**lemma** *operator-tensor-iso-map-entry*:
**assumes** *r* < *m*∗*p* **and** *s* < *m*∗*p*
**shows** *operator-tensor-iso-map m p c* $$ (*r*,*s*) =
*c* $$ ((*r div p*) ∗ *m* + (*s div p*), (*r mod p*) ∗ *p* + (*s mod p*))
⟨*proof*⟩

**lemma** *operator-tensor-iso-map-row-bound*:
**fixes** *m p r s* :: *nat*
**assumes** *p-pos*: *0* < *p* **and** *r* < *m*∗*p* **and** *s* < *m*∗*p*
**shows** (*r div p*) ∗ *m* + (*s div p*) < *m*∗*m*
⟨*proof*⟩

**lemma** *operator-tensor-iso-map-col-bound*:
**fixes** *p r s* :: *nat*
**assumes** *p-pos*: *0* < *p*
**shows** (*r mod p*) ∗ *p* + (*s mod p*) < *p*∗*p*
⟨*proof*⟩

**definition** *operator-tensor-iso-inv* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**

*operator-tensor-iso-inv m p x* ≡
*Matrix.mat* (*m*∗*m*) (*p*∗*p*)
(λ(*a*,*b*). *x* $$ ((*a div m*) ∗ *p* + (*b div p*), (*a mod m*) ∗ *p* + (*b mod p*)))

**definition** *operator-tensor-coordinate-space* ::
*nat* ⇒ *nat* ⇒ (*complex*, *complex Matrix.mat*) *linear-structure* **where**
*operator-tensor-coordinate-space m p* ≡ *mat-linear-structure* (*m*∗*m*) (*p*∗*p*)

**definition** *bipartite-operator-space* ::
*nat* ⇒ *nat* ⇒ (*complex*, *complex Matrix.mat*) *linear-structure* **where**
*bipartite-operator-space m p* ≡ *mat-linear-structure* (*m*∗*p*) (*m*∗*p*)

**lemma** *operator-tensor-iso-inv-carrier* [*simp*]:
*operator-tensor-iso-inv m p x* ∈ *carrier-mat* (*m*∗*m*) (*p*∗*p*)
⟨*proof*⟩

**lemma** *operator-tensor-iso-inv-entry*:
**assumes** *a* < *m*∗*m* **and** *b* < *p*∗*p*
**shows** *operator-tensor-iso-inv m p x* $$ (*a*,*b*) =
*x* $$ ((*a div m*) ∗ *p* + (*b div p*), (*a mod m*) ∗ *p* + (*b mod p*))
⟨*proof*⟩

**lemma** *operator-tensor-iso-inv-row-bound*:
**fixes** *m p a b* :: *nat*
**assumes** *m-pos*: *0* < *m* **and** *p-pos*: *0* < *p* **and** *a* < *m*∗*m* **and** *b* < *p*∗*p*
**shows** (*a div m*) ∗ *p* + (*b div p*) < *m*∗*p*
⟨*proof*⟩

**lemma** *operator-tensor-iso-inv-col-bound*:
**fixes** *m p a b* :: *nat*
**assumes** *m-pos*: *0* < *m* **and** *p-pos*: *0* < *p*
**shows** (*a mod m*) ∗ *p* + (*b mod p*) < *m*∗*p*
⟨*proof*⟩

**lemma** *operator-tensor-iso-map-add*:
**assumes** *p-pos*: *0* < *p*
**and** *x*: *x* ∈ *carrier-mat* (*m*∗*m*) (*p*∗*p*)
**and** *y*: *y* ∈ *carrier-mat* (*m*∗*m*) (*p*∗*p*)
**shows** *operator-tensor-iso-map m p* (*x* + *y*) =
*operator-tensor-iso-map m p x* + *operator-tensor-iso-map m p y*
⟨*proof*⟩

**lemma** *operator-tensor-iso-map-zero*:
**assumes** *p-pos*: *0* < *p*
**shows** *operator-tensor-iso-map m p* (*0*$_m$ (*m*∗*m*) (*p*∗*p*)) = *0*$_m$ (*m*∗*p*) (*m*∗*p*)
⟨*proof*⟩

**lemma** *operator-tensor-iso-map-smult*:
**assumes** *p-pos*: *0* < *p*
**and** *x*: *x* ∈ *carrier-mat* (*m*∗*m*) (*p*∗*p*)
**shows** *operator-tensor-iso-map m p* (*c* ·$_m$ *x*) =
*c* ·$_m$ *operator-tensor-iso-map m p x*
⟨*proof*⟩

**lemma** *operator-tensor-iso-inv-add*:
**assumes** *m-pos*: *0* < *m*
**and** *p-pos*: *0* < *p*
**and** *x*: *x* ∈ *carrier-mat* (*m*∗*p*) (*m*∗*p*)

**and** *y*: $y \in$ *carrier-mat* $(m{*}p)$ $(m{*}p)$
**shows** *operator-tensor-iso-inv m p* $(x + y) =$
*operator-tensor-iso-inv m p x + operator-tensor-iso-inv m p y*
⟨*proof*⟩

**lemma** *operator-tensor-iso-inv-zero*:
**assumes** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
**shows** *operator-tensor-iso-inv m p* $(0_m\ (m{*}p)\ (m{*}p)) = 0_m\ (m{*}m)\ (p{*}p)$
⟨*proof*⟩

**lemma** *operator-tensor-iso-inv-smult*:
**assumes** *m-pos*: $0 < m$
**and** *p-pos*: $0 < p$
**and** *x*: $x \in$ *carrier-mat* $(m{*}p)$ $(m{*}p)$
**shows** *operator-tensor-iso-inv m p* $(c \cdot_m x) =$
$c \cdot_m$ *operator-tensor-iso-inv m p x*
⟨*proof*⟩

**proposition** *operator-tensor-space-isomorphism*:
**assumes** *m-pos*: $0 < m$
**and** *p-pos*: $0 < p$
**shows** *is-isomorphism-between*
(*operator-tensor-coordinate-space m p*)
(*bipartite-operator-space m p*)
(*operator-tensor-iso-map m p*)
⟨*proof*⟩

**corollary** *operator-tensor-space-isomorphic*:
**assumes** $0 < m$ **and** $0 < p$
**shows** *is-isomorphic-to*
(*operator-tensor-coordinate-space m p*)
(*bipartite-operator-space m p*)
⟨*proof*⟩

## 24.2 Partial traces

**Definition (notes).** Identify $L(\mathcal{H}_A \otimes \mathcal{H}_B) = L(\mathcal{H}_A) \otimes L(\mathcal{H}_B)$ and define $\mathrm{tr}_B(\sum_l R_l \otimes T_l) = \sum_l \mathrm{tr}(T_l) R_l \in L(\mathcal{H}_A)$ and $\mathrm{tr}_A(\sum_l R_l \otimes T_l) = \sum_l \mathrm{tr}(R_l) T_l \in L(\mathcal{H}_B)$; also written $X^A = \mathrm{tr}_B(X)$, $X^B = \mathrm{tr}_A(X)$. Well defined by the universal property, since $(R, T) \mapsto \mathrm{tr}(T)R$ is bilinear.

The formal definitions use the block formulas of Day 16: an $mp \times mp$ matrix is indexed by pairs $(i, k)$, $i < m$, $k < p$, with flat index $ip + k$ ($\mathcal{H}_A$ is the major factor, matching $(A \otimes B)_{(r,c)} = A_{(r \div p,\, c \div p)} B_{(r \bmod p,\, c \bmod p)}$), and $(\mathrm{tr}_B\, X)_{ij} = \sum_{k<p} X_{ip+k,\, jp+k}$, $(\mathrm{tr}_A\, X)_{ij} = \sum_{k<m} X_{kp+i,\, kp+j}$. The defining property on elementary tensors is proved below.

**record** *'lab qspace* =
*qdim* :: *nat*
*qlabel* :: *'lab*

**record** *'lab bipartite-space* =
*left-space* :: *'lab qspace*
*right-space* :: *'lab qspace*

**definition** *partial-trace-B* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*partial-trace-B m p x* ≡
*Matrix.mat m m* (λ(*i*, *j*). ∑ *k*<*p*. *x* $$ (*i* ∗ *p* + *k*, *j* ∗ *p* + *k*))

**definition** *partial-trace-A* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*partial-trace-A m p x* ≡
*Matrix.mat p p* (λ(*i*, *j*). ∑ *k*<*m*. *x* $$ (*k* ∗ *p* + *i*, *k* ∗ *p* + *j*))

**definition** *partial-trace-on* ::
*'lab qspace* ⇒ *'lab bipartite-space* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*partial-trace-on s ab x* ≡
(*if s* = *right-space ab then*
*partial-trace-B* (*qdim* (*left-space ab*)) (*qdim* (*right-space ab*)) *x*
*else if s* = *left-space ab then*
*partial-trace-A* (*qdim* (*left-space ab*)) (*qdim* (*right-space ab*)) *x*
*else* $0_m$ *0 0*)

**notation** *partial-trace-on* (*tr*_ - [*1000*, *1000*, *1000*] *1000*)

**lemma** *partial-trace-on-right*:
**fixes** *S* :: *'lab qspace* **and** *AB* :: *'lab bipartite-space*
**assumes** *S* = *right-space AB*
**shows** $tr_S{}^{AB}\ x$ =
*partial-trace-B* (*qdim* (*left-space AB*)) (*qdim* (*right-space AB*)) *x*
⟨*proof*⟩

**lemma** *partial-trace-on-left*:
**fixes** *S* :: *'lab qspace* **and** *AB* :: *'lab bipartite-space*
**assumes** *S* = *left-space AB*
**and** *left-space AB* ≠ *right-space AB*
**shows** $tr_S{}^{AB}\ x$ =
*partial-trace-A* (*qdim* (*left-space AB*)) (*qdim* (*right-space AB*)) *x*
⟨*proof*⟩

**lemma** *dim-row-partial-trace-B* [*simp*]: *dim-row* (*partial-trace-B m p x*) = *m*
⟨*proof*⟩

**lemma** *dim-col-partial-trace-B* [*simp*]: *dim-col* (*partial-trace-B m p x*) = *m*
⟨*proof*⟩

**lemma** *partial-trace-B-carrier* [*simp*]:
*partial-trace-B m p x* ∈ *carrier-mat m m*
⟨*proof*⟩

**lemma** *dim-row-partial-trace-A* [*simp*]: *dim-row* (*partial-trace-A m p x*) = *p*
⟨*proof*⟩

**lemma** *dim-col-partial-trace-A* [*simp*]: *dim-col* (*partial-trace-A m p x*) = *p*
⟨*proof*⟩

**lemma** *partial-trace-A-carrier* [*simp*]:
*partial-trace-A m p x* ∈ *carrier-mat p p*
⟨*proof*⟩

**lemma** *partial-trace-B-tensor*:
**fixes** *R T* :: *complex Matrix.mat*
**assumes** *Rsq*: *R* ∈ *carrier-mat m m*
**and** *Tsq*: *T* ∈ *carrier-mat p p*
**and** *m-pos*: *0* < *m* **and** *p-pos*: *0* < *p*
**shows** *partial-trace-B m p* (*R* ⊗ *T*) = *Tr T* ·$_m$ *R*
⟨*proof*⟩

**lemma** *partial-trace-A-tensor*:
**fixes** $R$ $T$ :: *complex Matrix.mat*
**assumes** *Rsq*: $R \in$ *carrier-mat m m*
**and** *Tsq*: $T \in$ *carrier-mat p p*
**and** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
**shows** *partial-trace-A m p* $(R \bigotimes T) =$ *Tr R* $\cdot_m$ $T$
⟨*proof*⟩

### 24.3 Partial traces and measurements

Writing $\rho = \sum_l R_l \otimes T_l$,

$$p_\alpha(\rho) = \operatorname{tr}((M_\alpha^* M_\alpha \otimes I)\rho) = \sum_l \operatorname{tr}(M_\alpha^* M_\alpha R_l)\operatorname{tr}(T_l) = \operatorname{tr}\big(M_\alpha^* M_\alpha \operatorname{tr}_B(\rho)\big) = p_\alpha(\rho^A) :$$

lab A only sees $\rho^A$, and symmetrically lab B only sees $\rho^B$. Check: $\operatorname{tr}(\rho^A) = \operatorname{tr}\rho = 1$, and $\rho \geq 0$ implies $\rho^A \geq 0$.

## 25 Day 16 (3 October 2011): partial traces in block form; the postulates for density matrices

With matrix units $E_{ij}$ of $L(\mathcal{H}_B)$, every $X \in L(\mathcal{H}_A) \otimes L(\mathcal{H}_B)$ is uniquely $X = \sum_{i,j} X_{ij} \otimes E_{ij}$ with $X_{ij} \in L(\mathcal{H}_A)$, and $\operatorname{tr}_B(X) = \sum_i X_{ii}$, $\operatorname{tr}_A(X) = (\operatorname{tr} X_{ij})_{ij}$; symmetrically for $X = \sum E_{kl} \otimes Y_{kl}$. Choosing a basis $f_1, \ldots, f_p$ of $\mathcal{H}_B$ identifies $\mathcal{H}_A \otimes \mathcal{H}_B \cong \mathcal{H}_A^{(p)} = \mathcal{H}_A \oplus \cdots \oplus \mathcal{H}_A$ and $L(\mathcal{H}_A \otimes \mathcal{H}_B) \cong M_p(L(\mathcal{H}_A))$, block matrices $(X_{ij})$ acting by $(X_{ij})(h_j)_j = (\sum_j X_{ij} h_j)_i$; in block notation $\operatorname{tr}_B((X_{ij})) = X_{11} + \cdots + X_{pp}$ and $\operatorname{tr}_A((X_{ij})) = (\operatorname{tr} X_{ij})$.

Reformulated postulates. $1'$: the state of a system is a density matrix $\rho \in L(\mathcal{H}_A)$. $2'$: closed evolution is $\rho \mapsto U\rho U^*$. $3'$: a measurement system $\{M_\alpha\}$ with $\sum M_\alpha^* M_\alpha = I$ yields outcome $\alpha$ with probability $\operatorname{tr}(M_\alpha^* M_\alpha \rho)$. $4'$: systems with densities $\rho_1, \rho_2$ compose to $\rho_1 \otimes \rho_2$ on $\mathcal{H}_A \otimes \mathcal{H}_B$.

**lemma** *partial-trace-B-trace*:
**fixes** $X$ :: *complex Matrix.mat*
**assumes** *Xsq*: $X \in$ *carrier-mat* $(m * p)$ $(m * p)$
**and** *p-pos*: $0 < p$
**shows** *Tr* (*partial-trace-B m p X*) = *Tr X*
⟨*proof*⟩

**lemma** *partial-trace-B-hermitian*:
**fixes** $X$ :: *complex Matrix.mat*
**assumes** *Xsq*: $X \in$ *carrier-mat* $(m * p)$ $(m * p)$
**and** *herm*: *hermitian X*
**shows** *hermitian* (*partial-trace-B m p X*)
⟨*proof*⟩

**lemma** *partial-trace-B-is-density*:
**fixes** $X$ :: *complex Matrix.mat*
**assumes** *dm*: *density-matrix* $(m * p)$ $X$
**and** *p-pos*: $0 < p$
**shows** *density-matrix m* (*partial-trace-B m p X*)
⟨*proof*⟩

Key identity: measurements in lab A only see $\rho^A = \operatorname{tr}_B \rho$.

**theorem** *partial-trace-measurement*:

**fixes** *meas* :: *complex Matrix.mat* **and** $\varrho$ :: *complex Matrix.mat*
**assumes** *M-sq*: *meas* ∈ *carrier-mat m m*
**and** *dm*: *density-matrix* (*m* ∗ *p*) $\varrho$
**and** *p-pos*: *0* < *p*
**shows** *Tr* (((*meas*† ∗ *meas*) ⨂ *1*$_m$ *p*) ∗ $\varrho$) =
*Tr* (*meas*† ∗ *meas* ∗ *partial-trace-B m p* $\varrho$)
⟨*proof*⟩

# 26 Day 17 (5 October 2011): measurement maps, quantum noise, completely positive maps

## 26.1 Measurement maps

Measuring $\psi$ with $\{M_\alpha\}$ and keeping the outcome as an ensemble gives the density matrix $\sum_\alpha p_\alpha(\psi)|M_\alpha\psi/\|M_\alpha\psi\|\rangle\langle M_\alpha\psi/\|M_\alpha\psi\|| = \sum_\alpha M_\alpha(\psi\psi^*)M_\alpha^*$; for an ensemble with density $\rho_1$ one obtains $M(\rho_1) = \sum_\alpha M_\alpha\rho_1 M_\alpha^*$: a linear map on $L(\mathcal{H}_A)$ sending density matrices to density matrices.

Classical noise flips a bit with some probability; probability vectors evolve by stochastic matrices (non-negative entries, columns summing to one; doubly stochastic if rows do too). Quantum noise: with an environment $\mathcal{H}_E$ in state $|\phi\rangle$, the system $\rho\otimes|\phi\rangle\langle\phi|$ evolves by a unitary $U = (U_{ij})$ on $\mathcal{H}_A \otimes \mathcal{H}_E$ and lab A sees

$$\mathrm{tr}_E(U(\rho\otimes|\phi\rangle\langle\phi|)U^*) = \sum_i W_i\rho W_i^*, \qquad W_i = \sum_k U_{ik}\alpha_k, \qquad \sum_i W_i^*W_i = I.$$

Measurement by an intruder and environmental noise are maps of the same form $\rho \mapsto \sum_\alpha X_\alpha\rho X_\alpha^*$ with $\sum X_\alpha^*X_\alpha = I$. The environment computation is *environment-model* in *Day17-Environment-Model*.

Third way, axiomatic: a quantum event is a linear $\Phi : L(\mathcal{H}_A) \to L(\mathcal{H}_A)$.

**Proposition (notes).** $\Phi$ maps density matrices to density matrices iff $\Phi$ is positive ($\rho \geq 0 \Rightarrow \Phi(\rho) \geq 0$) and trace preserving ($\mathrm{tr}\,\Phi(X) = \mathrm{tr}\,X$); then $\Phi \otimes I_{L(\mathcal{H}_B)}$ also maps density matrices to density matrices for every $\mathcal{H}_B$.

**Definition (notes).** $\Phi$ is *completely positive* if $\Phi\otimes I_{L(\mathcal{H}_B)}$ is positive for every finite-dimensional $\mathcal{H}_B$.

**Theorem (notes)** (Choi–Kraus)**.** $\Phi : L(\mathcal{H}_A) \to L(\mathcal{H}_A)$ is completely positive and trace preserving iff $\Phi(X) = \sum_i E_i X E_i^*$ for matrices $E_i$ with $\sum_i E_i^*E_i = I$.

The proposition is *density-preserving-iff-positive-tp* in *Day18-Positive-Maps*; complete positivity is defined below and the Choi–Kraus theorem is *choi-kraus-cptp* on Day 19.

**definition** *meas-map* :: *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*meas-map Ms* $\varrho$ ≡
*foldr* (λ*M acc*. *M* ∗ $\varrho$ ∗ (*M*†) + *acc*) *Ms* (*0*$_m$ (*dim-row* $\varrho$) (*dim-col* $\varrho$))

**lemma** *meas-map-Nil* [*simp*]:
*meas-map* [] $\varrho$ = *0*$_m$ (*dim-row* $\varrho$) (*dim-col* $\varrho$)
⟨*proof*⟩

**lemma** *meas-map-Cons* [*simp*]:
*meas-map* (*M* # *Ms*) $\varrho$ = *M* ∗ $\varrho$ ∗ (*M*†) + *meas-map Ms* $\varrho$
⟨*proof*⟩

**lemma** *meas-map-carrier*:
**assumes** *Ms-c*: ∀ *M* ∈ *set Ms*. *M* ∈ *carrier-mat n n*

**and** $\varrho$-*c*: $\varrho \in$ *carrier-mat n n*
**shows** *meas-map Ms* $\varrho \in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *meas-map-psd*:
**assumes** *Ms-c*: $\forall M \in$ *set Ms*. $M \in$ *carrier-mat n n*
**and** $\varrho$-*c*: $\varrho \in$ *carrier-mat n n*
**and** $\varrho$-*psd*: *psd* $\varrho$
**shows** *psd* (*meas-map Ms* $\varrho$)
⟨*proof*⟩

**lemma** *foldr-MdagM-carrier*:
**assumes** *Ms-c*: $\forall M \in$ *set Ms*. $M \in$ *carrier-mat n n*
**shows** *foldr* $(\lambda M\ acc.\ (M^\dagger) * M + acc)$ *Ms* $(0_m\ n\ n) \in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *meas-map-trace-aux*:
**assumes** *Ms-c*: $\forall M \in$ *set Ms*. $M \in$ *carrier-mat n n*
**and** $\varrho$-*c*: $\varrho \in$ *carrier-mat n n*
**shows** *Tr* (*meas-map Ms* $\varrho$) =
*Tr* (*foldr* $(\lambda M\ acc.\ (M^\dagger) * M + acc)$ *Ms* $(0_m\ n\ n) * \varrho$)
⟨*proof*⟩

**lemma** *meas-map-density*:
**assumes** *ms*: *measurement-system n Ms*
**and** *dm*: *density-matrix n* $\varrho$
**shows** *density-matrix n* (*meas-map Ms* $\varrho$)
⟨*proof*⟩

## 26.2 Kraus maps, positive maps, complete positivity

A *Kraus map* $\rho \mapsto \sum_i W_i \rho W_i^\dagger$ is *kraus-map-dim d Ws*; the list *Ws* is a list of *Kraus operators* (*kraus-operators n d Ws*) when $\sum_i W_i^\dagger W_i = I_n$. Positive, trace-preserving and unital maps are defined on the carrier $M_n$. The amplification $\Phi^{(p)} = \Phi \otimes \mathrm{id}_p$ acts on block matrices by $(X_{ab}) \mapsto (\Phi(X_{ab}))$ (*amplification n d p* $\Phi$), and $\Phi$ is $p$-positive when $\Phi^{(p)}$ is positive, completely positive when it is $p$-positive for all $p$. The main result of the subsection: Kraus maps are completely positive.

**definition** *kraus-map-dim* :: *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*kraus-map-dim d Ws* $\varrho \equiv$
*foldr* $(\lambda W\ acc.\ W * \varrho * (W^\dagger) + acc)$ *Ws* $(0_m\ d\ d)$

**definition** *kraus-map* :: *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*kraus-map Ws* $\varrho \equiv$ *kraus-map-dim* (*dim-row* $\varrho$) *Ws* $\varrho$

**definition** *kraus-operators* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat list* ⇒ *bool* **where**
*kraus-operators n d Ws* ≡
$(\forall W \in$ *set Ws*. $W \in$ *carrier-mat d n*$) \wedge$
*foldr* $(\lambda W\ acc.\ (W^\dagger) * W + acc)$ *Ws* $(0_m\ n\ n) = 1_m\ n$

**lemma** *kraus-map-dim-Nil* [*simp*]:
*kraus-map-dim d* [] $\varrho = 0_m\ d\ d$
⟨*proof*⟩

**lemma** *kraus-map-dim-Cons* [*simp*]:
*kraus-map-dim d* (*W* # *Ws*) $\varrho$ =
$W * \varrho * (W^\dagger)$ + *kraus-map-dim d Ws* $\varrho$

⟨*proof*⟩

**lemma** *kraus-map-dim-carrier*:
**assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**and** *ϱ-c*: *ϱ* ∈ *carrier-mat n n*
**shows** *kraus-map-dim d Ws ϱ* ∈ *carrier-mat d d*
⟨*proof*⟩

**lemma** *kraus-map-dim-psd*:
**assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**and** *ϱ-c*: *ϱ* ∈ *carrier-mat n n*
**and** *ϱ-psd*: *psd ϱ*
**shows** *psd* (*kraus-map-dim d Ws ϱ*)
⟨*proof*⟩

**definition** *positive-map* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*positive-map n d* Φ ≡
(∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* ∈ *carrier-mat d d*) ∧
(∀ *P*. *P* ∈ *carrier-mat n n* ⟶ *psd P* ⟶ *psd* (Φ *P*))

**lemma** *kraus-map-dim-positive-map*:
**assumes** *Ws*: *kraus-operators n d Ws*
**shows** *positive-map n d* (*kraus-map-dim d Ws*)
⟨*proof*⟩

**definition** *trace-preserving* :: *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*trace-preserving n* Φ ≡
∀ *X*. *X* ∈ *carrier-mat n n* ⟶ *Tr* (Φ *X*) = *Tr X*

**definition** *unital-map* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*unital-map n d* Φ ≡ Φ (1$_m$ *n*) = 1$_m$ *d*

**definition** *block-of-mat* :: *nat* ⇒ *complex Matrix.mat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*block-of-mat n x a b* ≡
*Matrix.mat n n* (λ(*r*,*c*). *x* $$ (*a*∗*n* + *r*, *b*∗*n* + *c*))

**lemma** *block-of-mat-carrier* [*simp*]:
*block-of-mat n X a b* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *block-of-mat-index*:
**assumes** *r* < *n* **and** *c* < *n*
**shows** *block-of-mat n X a b* $$ (*r*,*c*) = *X* $$ (*a*∗*n* + *r*, *b*∗*n* + *c*)
⟨*proof*⟩

**lemma** *block-of-mat-single*:
**assumes** *X*: *X* ∈ *carrier-mat n n*
**shows** *block-of-mat n X* 0 0 = *X*
⟨*proof*⟩

**definition** *amplification* :: *nat* ⇒ *nat* ⇒ *nat* ⇒
(*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*amplification n d p* Φ *x* ≡
*Matrix.mat* (*p*∗*d*) (*p*∗*d*)
(λ(*i*,*j*). Φ (*block-of-mat n x* (*i div d*) (*j div d*)) $$
(*i mod d*, *j mod d*))

**lemma** *amplification-carrier* [*simp*]:

*amplification n d p Φ X ∈ carrier-mat (p∗d) (p∗d)*
⟨*proof*⟩

**lemma** *amplification-index*:
**assumes** *i* < *p∗d* **and** *j* < *p∗d*
**shows** *amplification n d p Φ X $$ (i,j)* =
Φ (*block-of-mat n X* (*i div d*) (*j div d*)) $$
(*i mod d*, *j mod d*)
⟨*proof*⟩

**lemma** *amplification-one*:
**assumes** *X*: *X* ∈ *carrier-mat n n*
**and** *range*: Φ *X* ∈ *carrier-mat d d*
**shows** *amplification n d 1* Φ *X* = Φ *X*
⟨*proof*⟩

**lemma** *tensor-id-kraus-carrier*:
**assumes** *W*: *W* ∈ *carrier-mat d n*
**shows** (*1*$_m$ *p* ⨂ *W*) ∈ *carrier-mat* (*p∗d*) (*p∗n*)
⟨*proof*⟩

**lemma** *tensor-id-kraus-list-carrier*:
**assumes** *Ws*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**shows** ∀ *L* ∈ *set* (*map* (λ*W*. *1*$_m$ *p* ⨂ *W*) *Ws*).
*L* ∈ *carrier-mat* (*p∗d*) (*p∗n*)
⟨*proof*⟩

**lemma** *tensor-id-kraus-index*:
**assumes** *W*: *W* ∈ *carrier-mat d n*
**and** *r*: *r* < *p∗d*
**and** *c*: *c* < *p∗n*
**and** *dpos*: *0* < *d*
**and** *npos*: *0* < *n*
**shows** (*1*$_m$ *p* ⨂ *W*) $$ (*r*,*c*) =
(*if r div d* = *c div n then W $$* (*r mod d*, *c mod n*) *else 0*)
⟨*proof*⟩

**lemma** *sum-block-delta*:
**fixes** *f* :: *nat* ⇒ *complex*
**assumes** *a*: *a* < *p*
**and** *npos*: *0* < *n*
**shows** (∑ *t* < *p∗n*. *if a* = *t div n then f* (*t mod n*) *else 0*) =
(∑ *r* < *n*. *f r*)
⟨*proof*⟩

**lemma** *sum-block-delta-index*:
**fixes** *h* :: *nat* ⇒ *complex*
**assumes** *a*: *a* < *p*
**and** *npos*: *0* < *n*
**shows** (∑ *t* < *p∗n*. *if a* = *t div n then h t else 0*) =
(∑ *r* < *n*. *h* (*a∗n* + *r*))
⟨*proof*⟩

**lemma** *tensor-id-kraus-left-mult-index*:
**assumes** *W*: *W* ∈ *carrier-mat d n*
**and** *x*: *x* ∈ *carrier-mat* (*p∗n*) *q*
**and** *i*: *i* < *p∗d*
**and** *s*: *s* < *q*

**and** *dpos*: $0 < d$
**and** *npos*: $0 < n$
**shows** $((1_m\ p \bigotimes W) * x)\ \$\$\ (i,s) =$
$(\sum t < n.\ W\ \$\$\ (i\ mod\ d,\ t) * x\ \$\$\ ((i\ div\ d) * n + t,\ s))$
⟨*proof*⟩

**lemma** *tensor-id-kraus-right-mult-index*:
**assumes** *W*: $W \in carrier\text{-}mat\ d\ n$
**and** *y*: $y \in carrier\text{-}mat\ q\ (p*n)$
**and** *i*: $i < q$
**and** *j*: $j < p*d$
**and** *dpos*: $0 < d$
**and** *npos*: $0 < n$
**shows** $(y * (1_m\ p \bigotimes (W^\dagger)))\ \$\$\ (i,j) =$
$(\sum s < n.\ y\ \$\$\ (i,\ (j\ div\ d) * n + s) * (W^\dagger)\ \$\$\ (s,\ j\ mod\ d))$
⟨*proof*⟩

**lemma** *kraus-term-index*:
**assumes** *W*: $W \in carrier\text{-}mat\ d\ n$
**and** *B*: $B \in carrier\text{-}mat\ n\ n$
**and** *i*: $i < d$
**and** *j*: $j < d$
**shows** $(W * B * (W^\dagger))\ \$\$\ (i,j) =$
$(\sum s < n.\ (\sum t < n.\ W\ \$\$\ (i,t) * B\ \$\$\ (t,s)) * (W^\dagger)\ \$\$\ (s,j))$
⟨*proof*⟩

**lemma** *kraus-amplification-single*:
**assumes** *W*: $W \in carrier\text{-}mat\ d\ n$
**and** *x*: $x \in carrier\text{-}mat\ (p*n)\ (p*n)$
**and** *dpos*: $0 < d$
**and** *npos*: $0 < n$
**shows** $amplification\ n\ d\ p\ (\lambda\varrho.\ W * \varrho * (W^\dagger))\ x =$
$((1_m\ p \bigotimes W) * x) * (1_m\ p \bigotimes (W^\dagger))$
⟨*proof*⟩

**lemma** *amplification-zero-map*:
**assumes** *dpos*: $0 < d$
**shows** $amplification\ n\ d\ p\ (\lambda\varrho.\ 0_m\ d\ d)\ x = 0_m\ (p*d)\ (p*d)$
⟨*proof*⟩

**lemma** *amplification-add*:
**assumes** *dpos*: $0 < d$
**and** *F*: $\bigwedge a\ b.\ F\ (block\text{-}of\text{-}mat\ n\ x\ a\ b) \in carrier\text{-}mat\ d\ d$
**and** *G*: $\bigwedge a\ b.\ G\ (block\text{-}of\text{-}mat\ n\ x\ a\ b) \in carrier\text{-}mat\ d\ d$
**shows** $amplification\ n\ d\ p\ (\lambda\varrho.\ F\ \varrho + G\ \varrho)\ x =$
$amplification\ n\ d\ p\ F\ x + amplification\ n\ d\ p\ G\ x$
⟨*proof*⟩

**lemma** *amplification-kraus-map-dim*:
**assumes** *Ws*: $\forall W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n$
**and** *x*: $x \in carrier\text{-}mat\ (p*n)\ (p*n)$
**and** *dpos*: $0 < d$
**and** *npos*: $0 < n$
**shows** $amplification\ n\ d\ p\ (kraus\text{-}map\text{-}dim\ d\ Ws)\ x =$
$kraus\text{-}map\text{-}dim\ (p*d)\ (map\ (\lambda W.\ 1_m\ p \bigotimes W)\ Ws)\ x$
⟨*proof*⟩

**definition** *p-positive* :: $nat \Rightarrow nat \Rightarrow nat \Rightarrow (complex\ Matrix.mat \Rightarrow complex\ Matrix.mat) \Rightarrow bool$

**where**
*p-positive n d p Φ ≡*
*∀ X :: complex Matrix.mat. X ∈ carrier-mat (p∗n) (p∗n) ⟶ psd X ⟶*
*psd (amplification n d p Φ X)*

**definition** *completely-positive :: nat ⇒ nat ⇒ (complex Matrix.mat ⇒ complex Matrix.mat) ⇒ bool*
**where**
*completely-positive n d Φ ≡ ∀ p > 0. p-positive n d p Φ*

**theorem** *kraus-is-cp*:
**assumes** *Ws*: *kraus-operators n d Ws*
**and** *npos*: *0 < n*
**and** *dpos*: *0 < d*
**shows** *completely-positive n d (kraus-map-dim d Ws)*
⟨*proof*⟩

# 27 Day 18 (7 October 2011): the theory of completely positive maps

**Proposition (notes).** Every $X \in M_n$ is $X = (P_1 - P_2) + i(P_3 - P_4)$ with $P_i \geq 0$: $\mathrm{span}(M_n^+) = M_n$.

(Write $X = H + iK$ with $H, K$ Hermitian and split each spectral decomposition into its non-negative and negative eigenvalues.)

**Definition (notes).** A linear $\Phi : M_n \to M_d$ is *positive* if $P \geq 0$ implies $\Phi(P) \geq 0$; it is *p-positive* if $\Phi^{(p)} : M_p(M_n) \to M_p(M_d)$, $(X_{ij}) \mapsto (\Phi(X_{ij}))$, is positive, and *completely positive* if it is $p$-positive for every $p$.

**Proposition (notes).** A linear $\Phi : M_n \to M_d$ maps density matrices to density matrices iff it is positive and trace preserving.

Example (notes): the transpose $\Phi(A) = A^t$ on $M_2$ is positive but not 2-positive, since $\Phi^{(2)}$ sends the positive matrix $(E_{ij})_{i,j}$ to a matrix of determinant $-1$. The decomposition into positives is formalised below; the proposition is *density-preserving-iff-positive-tp* and the example *transpose-not-2-positive*, both in *Day18-Positive-Maps*.

**lemma** *diag-mat-of-real-hermitian*:
*((diag-mat-of-real D)†) = diag-mat-of-real D*
⟨*proof*⟩

**lemma** *hermitian-decomp-psd*:
**fixes** *H :: complex Matrix.mat*
**assumes** *sq*: *H ∈ carrier-mat n n*
**and** *herm*: *hermitian H*
**shows** *∃ P1 P2 :: complex Matrix.mat.*
*P1 ∈ carrier-mat n n ∧ P2 ∈ carrier-mat n n ∧*
*psd P1 ∧ psd P2 ∧*
*H = P1 − P2*
⟨*proof*⟩

**lemma** *matrix-decomp-psd*:
**fixes** *X :: complex Matrix.mat*
**assumes** *sq*: *X ∈ carrier-mat n n*
**shows** *∃ P1 P2 P3 P4 :: complex Matrix.mat.*
*(∀ i ∈ {0..3}. [P1,P2,P3,P4] ! i ∈ carrier-mat n n ∧ psd ([P1,P2,P3,P4] ! i)) ∧*

$X = (P1 - P2) + \mathrm{i} \cdot_m (P3 - P4)$
⟨*proof*⟩

# 28 Day 19 (10 October 2011): Choi's theorem

**Theorem (notes)** (Choi, 1975)**.** For linear $\Phi : M_n \to M_d$ the following are equivalent: (1) $\Phi$ is completely positive; (2) $\Phi$ is $n$-positive; (3) $P_\Phi = (\Phi(E_{ij}))_{i,j} \geq 0$ in $M_n(M_d)$; (4) there are $d \times n$ matrices $B_1, \ldots, B_K$ with $\Phi(X) = \sum_k B_k X B_k^*$.

Notes: (1) $\sum_k B_k X B_k^*$ is a *Choi–Kraus representation*; (2) $P_\Phi$ is the *Choi–Jamiołkowski matrix*; (3) $CP(M_n, M_d) \leftrightarrow M_n(M_d)^+$ via $\Phi \leftrightarrow P_\Phi$; (4) $\Phi$ is trace preserving iff $\mathrm{tr}(\Phi(E_{ij})) = \delta_{ij}$; (5) $\Phi$ is unital iff $\sum_i \Phi(E_{ii}) = I_d$.
Formally $P_\Phi$ is *choi-matrix n d* $\Phi$ (an $nd \times nd$ matrix built from the blocks $\Phi(E_{ij})$); *choi-theorem* is (1)⇔(3), *kraus-rep-of-psd-choi* is (3)⇒(4), and *kraus-is-cp* is (4)⇒(1). The step (2)⇒(3) is the observation that $P_\Phi = \Phi^{(n)}((E_{ij}))$ with $(E_{ij}) \geq 0$ (*choi-matrix-as-amplification*). Linearity of $\Phi$ is assumed on the carrier $M_n$ (additivity and homogeneity hypotheses). The trace-preserving case is *choi-kraus-cptp*; notes (4) and (5) in Choi-matrix form are *choi-trace-preserving-iff* and *choi-unital-iff* in theory Day19_Choi_TP_Unital.

**definition** *choi-matrix* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* **where**
*choi-matrix n d* Φ ≡
*Matrix.mat* (*n*∗*d*) (*n*∗*d*)
(λ(*r*,*c*). Φ (*mat-unit* (*r div d*) (*c div d*) *n n*) $$ (*r mod d*, *c mod d*))

**notation** *choi-matrix* ($P_\Phi$)

**definition** *choi-seed* :: *nat* ⇒ *complex Matrix.mat* **where**
*choi-seed n* ≡
*Matrix.mat* (*n*∗*n*) (*n*∗*n*)
(λ(*r*,*c*). *mat-unit* (*r div n*) (*c div n*) *n n* $$ (*r mod n*, *c mod n*))

**lemma** *choi-seed-carrier* [*simp*]:
*choi-seed n* ∈ *carrier-mat* (*n*∗*n*) (*n*∗*n*)
⟨*proof*⟩

**lemma** *choi-seed-block*:
**assumes** *npos*: *n* > *0*
**and** *a*: *a* < *n*
**and** *b*: *b* < *n*
**shows** *block-of-mat n* (*choi-seed n*) *a b* = *mat-unit a b n n*
⟨*proof*⟩

**lemma** *choi-matrix-as-amplification*:
**assumes** *npos*: *n* > *0*
**and** *dpos*: *d* > *0*
**shows** *choi-matrix n d* Φ = *amplification n d n* Φ (*choi-seed n*)
⟨*proof*⟩

**lemma** *choi-psd-if-cp*:
**assumes** *npos*: *n* > *0*
**and** *dpos*: *d* > *0*
**and** *cp*: *completely-positive n d* Φ
**shows** *psd* (*choi-matrix n d* Φ)
⟨*proof*⟩

## 28.1 From a positive Choi matrix to a Kraus representation

The notes' proof of (3)⇒(4): write $P_\Phi = \sum_k v_k v_k^*$ with $v_k \in \mathbb{C}^{nd}$, cut each $v_k$ into $n$ blocks $w_1^k, \ldots, w_n^k \in \mathbb{C}^d$, and let $B_k = [w_1^k | \cdots | w_n^k]$; then $B_k E_{ij} B_k^* = w_i^k (w_j^k)^*$ and $\Phi(X) = \sum_{ij} x_{ij} \Phi(E_{ij}) = \sum_k B_k X B_k^*$ by linearity.

Complete positivity from a Kraus list without the trace-preservation constraint: the argument is the same as ⟦*kraus-operators ?n ?d ?Ws*; *0 < ?n*; *0 < ?d*⟧ ⟹ *completely-positive ?n ?d* (*kraus-map-dim ?d ?Ws*), which only uses the carrier hypothesis.

**lemma** *kraus-cp-no-tp*:
  **assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
    **and** *npos*: *0 < n*
    **and** *dpos*: *0 < d*
  **shows** *completely-positive n d* (*kraus-map-dim d Ws*)
⟨*proof*⟩

Linearity of the Kraus-sum map in its matrix argument.

**lemma** *kraus-map-dim-add*:
  **assumes** *A-c*: *A* ∈ *carrier-mat n n* **and** *B-c*: *B* ∈ *carrier-mat n n*
    **and** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
  **shows** *kraus-map-dim d Ws* (*A* + *B*) =
      *kraus-map-dim d Ws A* + *kraus-map-dim d Ws B*
  ⟨*proof*⟩

**lemma** *kraus-map-dim-smult*:
  **assumes** *A-c*: *A* ∈ *carrier-mat n n*
    **and** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
  **shows** *kraus-map-dim d Ws* (*c* ·$_m$ *A*) = *c* ·$_m$ *kraus-map-dim d Ws A*
  ⟨*proof*⟩

**lemma** *kraus-map-dim-zero* [*simp*]:
  **assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
  **shows** *kraus-map-dim d Ws* (*0*$_m$ *n n*) = *0*$_m$ *d d*
  ⟨*proof*⟩

Enumeration of [*0*..<*n*] × [*0*..<*n*] as a list of pairs.

**definition** *mat-unit-pairs* :: *nat* ⇒ (*nat* × *nat*) *list* **where**
  *mat-unit-pairs n* ≡
    *concat* (*map* (λ*a*. *map* (*Pair a*) [*0*..<*n*]) [*0*..<*n*])

**lemma** *mat-unit-pairs-set*:
  *set* (*mat-unit-pairs n*) = {*p*. *fst p* < *n* ∧ *snd p* < *n*}
  ⟨*proof*⟩

**lemma** *mat-unit-pairs-distinct*: *distinct* (*mat-unit-pairs n*)
  ⟨*proof*⟩

Generic pair-indexed linear combination of $f\ (E_{ab})$ contributions, weighted by the entries of $A$.

**primrec** *pair-comb* ::
  (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* ⇒
   *nat* ⇒ *nat* ⇒ (*nat* × *nat*) *list* ⇒ *complex Matrix.mat* **where**
  *pair-comb f A n d* [] = *0*$_m$ *d d* |
  *pair-comb f A n d* (*p* # *ps*) =
    *A* $$ *p* ·$_m$ *f* (*mat-unit* (*fst p*) (*snd p*) *n n*) + *pair-comb f A n d ps*

**lemma** *pair-comb-carrier*:
  **assumes** *f-car*: ⋀*a b*. *a* < *n* ⟹ *b* < *n* ⟹ *f* (*mat-unit a b n n*) ∈ *carrier-mat d d*

**and** *ps-valid*: $\forall\, p \in set\ ps.\ fst\ p < n \wedge snd\ p < n$
**shows** *pair-comb f A n d ps* $\in$ *carrier-mat d d*
⟨*proof*⟩

**lemma** *pair-comb-entry*:
**assumes** *f-car*: $\bigwedge a\ b.\ a < n \Longrightarrow b < n \Longrightarrow f\ (mat\text{-}unit\ a\ b\ n\ n) \in carrier\text{-}mat\ d\ d$
**and** *ps-valid*: $\forall\, p \in set\ ps.\ fst\ p < n \wedge snd\ p < n$
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < d$
**shows** *pair-comb f A n d ps* $\$\$\ (r,c) =$
$(\sum p \leftarrow ps.\ A\ \$\$\ p * f\ (mat\text{-}unit\ (fst\ p)\ (snd\ p)\ n\ n)\ \$\$\ (r,c))$
⟨*proof*⟩

For the identity-like choice $f = id$, the pair combination reconstructs $A$ when $ps$ enumerates all basis positions.

**lemma** *pair-comb-id-reconstructs*:
**assumes** *A-car*: $A \in carrier\text{-}mat\ n\ n$
**shows** *pair-comb* $(\lambda X.\ X)$ *A n n* (*mat-unit-pairs n*) = *A*
⟨*proof*⟩

For a linear map $f : M_n \to M_d$, applying $f$ to $A$ equals the pair combination of $f$ evaluated on the basis.

**lemma** *linear-eq-pair-comb*:
**fixes** *f* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *f-add*: $\bigwedge X\ Y.\ X \in carrier\text{-}mat\ n\ n \Longrightarrow Y \in carrier\text{-}mat\ n\ n \Longrightarrow$
$f\ (X + Y) = f\ X + f\ Y$
**and** *f-smult*: $\bigwedge c\ X.\ X \in carrier\text{-}mat\ n\ n \Longrightarrow f\ (c \cdot_m X) = c \cdot_m f\ X$
**and** *f-car*: $\bigwedge X.\ X \in carrier\text{-}mat\ n\ n \Longrightarrow f\ X \in carrier\text{-}mat\ d\ d$
**and** *A-car*: $A \in carrier\text{-}mat\ n\ n$
**and** *npos*: $n > 0$ **and** *dpos*: $d > 0$
**shows** *f A* = *pair-comb f A n d* (*mat-unit-pairs n*)
⟨*proof*⟩

Conjugation $W\ E_{ab}\ W^\dagger$ has entry $W_{ia} \cdot^\dagger W_{jb}$.

**lemma** *conjugation-mat-unit-entry*:
**assumes** *Wc*: $W \in carrier\text{-}mat\ d\ n$
**and** *a-lt*: $a < n$ **and** *b-lt*: $b < n$
**and** *i-lt*: $i < d$ **and** *j-lt*: $j < d$
**shows** $(W * mat\text{-}unit\ a\ b\ n\ n * (W^\dagger))\ \$\$\ (i,\ j) =$
$W\ \$\$\ (i,\ a) * cnj\ (W\ \$\$\ (j,\ b))$
⟨*proof*⟩

Entry-wise formula for *kraus-map-dim* as a sum over the list.

**lemma** *kraus-map-dim-entry-sum-list*:
**assumes** *Ws-c*: $\forall\, W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $\varrho \in carrier\text{-}mat\ n\ n$
**and** *i-lt*: $i < d$ **and** *j-lt*: $j < d$
**shows** *kraus-map-dim d Ws* $\varrho$ $\$\$\ (i,\ j) =$
*sum-list* (*map* $(\lambda W.\ (W * \varrho * (W^\dagger))\ \$\$\ (i,\ j))$ *Ws*)
⟨*proof*⟩

Kraus representation extracted from a psd Choi matrix: reshape each column of a psd factorization of $P_\Phi$ into a Kraus operator, and the resulting list agrees with $\Phi$ on all of $M_n$.

**lemma** *kraus-rep-of-psd-choi*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: $\bigwedge A\ B.\ A \in carrier\text{-}mat\ n\ n \Longrightarrow B \in carrier\text{-}mat\ n\ n \Longrightarrow$
$\Phi\ (A + B) = \Phi\ A + \Phi\ B$

**and** *smult*: $\bigwedge c\ A.\ A \in carrier\text{-}mat\ n\ n \Longrightarrow \Phi\ (c \cdot_m A) = c \cdot_m \Phi\ A$
**and** *range*: $\forall A.\ A \in carrier\text{-}mat\ n\ n \longrightarrow \Phi\ A \in carrier\text{-}mat\ d\ d$
**and** *npos*: $n > 0$
**and** *dpos*: $d > 0$
**and** *psdP*: $psd\ (choi\text{-}matrix\ n\ d\ \Phi)$
**shows** $\exists\ Ws :: complex\ Matrix.mat\ list.$
$length\ Ws = n * d \wedge$
$(\forall\ W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n) \wedge$
$(\forall A.\ A \in carrier\text{-}mat\ n\ n \longrightarrow \Phi\ A = kraus\text{-}map\text{-}dim\ d\ Ws\ A)$
⟨*proof*⟩

**theorem** *choi-theorem*:
**fixes** $\Phi :: complex\ Matrix.mat \Rightarrow complex\ Matrix.mat$
**assumes** *linear*: $\bigwedge A\ B.\ A \in carrier\text{-}mat\ n\ n \Longrightarrow B \in carrier\text{-}mat\ n\ n \Longrightarrow$
$\Phi\ (A + B) = \Phi\ A + \Phi\ B$
**and** *smult*: $\bigwedge c\ A.\ A \in carrier\text{-}mat\ n\ n \Longrightarrow \Phi\ (c \cdot_m A) = c \cdot_m \Phi\ A$
**and** *range*: $\forall A.\ A \in carrier\text{-}mat\ n\ n \longrightarrow \Phi\ A \in carrier\text{-}mat\ d\ d$
**and** *npos*: $n > 0$
**and** *dpos*: $d > 0$
**shows** *completely-positive* $n\ d\ \Phi \longleftrightarrow$
$psd\ (choi\text{-}matrix\ n\ d\ \Phi)$
⟨*proof*⟩

Accumulated $\sum_i\ {W_i}^\dagger\ W_i$ over a Kraus list.

**definition** *sum-wdag-w* :: *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* **where**
*sum-wdag-w n Ws* ≡
$foldr\ (\lambda W\ acc.\ (W^\dagger) * W + acc)\ Ws\ (0_m\ n\ n)$

**lemma** *sum-wdag-w-Nil* [*simp*]: *sum-wdag-w n* [] = $0_m\ n\ n$
⟨*proof*⟩

**lemma** *sum-wdag-w-Cons*:
*sum-wdag-w n* $(W\ \#\ Ws) = (W^\dagger) * W +$ *sum-wdag-w n Ws*
⟨*proof*⟩

**lemma** *sum-wdag-w-carrier*:
**assumes** $\forall\ W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n$
**shows** *sum-wdag-w n Ws* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *kraus-operators-altdef*:
*kraus-operators n d Ws* ⟷
$(\forall\ W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n) \wedge$ *sum-wdag-w n Ws* = $1_m\ n$
⟨*proof*⟩

**lemma** *Tr-kraus-map-dim-as-wdagw*:
**fixes** $\varrho :: complex\ Matrix.mat$
**assumes** *Ws-c*: $\forall\ W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n$
**and** *X-c*: $\varrho \in carrier\text{-}mat\ n\ n$
**shows** $Tr\ (kraus\text{-}map\text{-}dim\ d\ Ws\ \varrho) = Tr\ (sum\text{-}wdag\text{-}w\ n\ Ws * \varrho)$
⟨*proof*⟩

The $\sum_i\ {W_i}^\dagger\ W_i$ matrix entry equals a sum-list of $cnj(W_{ika}) \cdot W_{ikb}$ contributions (needed to recover $1_n$ from trace preservation).

**lemma** *sum-wdag-w-entry*:
**assumes** *Ws-c*: $\forall\ W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n$
**and** *a-lt*: $a < n$ **and** *b-lt*: $b < n$
**shows** *sum-wdag-w n Ws* $$ $(a,\ b) =$

*sum-list* (*map* (λ*W*. ((*W*†) ∗ *W*) $$ (*a*, *b*)) *Ws*)
⟨*proof*⟩

Pairing with $E_{ab}$ extracts the $(b,a)$ entry under the trace.

**lemma** *Tr-mult-mat-unit*:
**assumes** *M-c*: *M* ∈ *carrier-mat n n*
**and** *a-lt*: *a* < *n* **and** *b-lt*: *b* < *n*
**shows** *Tr* (*M* ∗ *mat-unit a b n n*) = *M* $$ (*b*, *a*)
⟨*proof*⟩

**theorem** *choi-kraus-cptp*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: ⋀*A B*. *A* ∈ *carrier-mat n n* ⟹ *B* ∈ *carrier-mat n n* ⟹
Φ (*A* + *B*) = Φ *A* + Φ *B*
**and** *smult*: ⋀*c A*. *A* ∈ *carrier-mat n n* ⟹ Φ (*c* ·*m A*) = *c* ·*m* Φ *A*
**and** *range*: ∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* ∈ *carrier-mat d d*
**and** *npos*: *n* > *0*
**and** *dpos*: *d* > *0*
**shows** (*completely-positive n d* Φ ∧ *trace-preserving n* Φ) ⟷
(∃ *Ws* :: *complex Matrix.mat list*.
*kraus-operators n d Ws* ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ *X* = *kraus-map-dim d Ws X*))
⟨*proof*⟩

# 29 Day 20 (12 October 2011): positive functionals, adjoints, Choi rank

**Proposition (notes).** For linear $\phi : M_n \to \mathbb{C}$: $\phi$ positive $\Leftrightarrow$ $\phi$ completely positive $\Leftrightarrow$ $(\phi(E_{ij})) \in M_n^+$.

**Proposition (notes).** Let $\Phi : M_n \to M_d$ with Hilbert–Schmidt adjoint $\Phi^* : M_d \to M_n$ ($\langle Y|X\rangle = \mathrm{tr}(Y^*X)$). (1) If $\Phi(X) = \sum_i A_i X A_i^*$ then $\Phi^*(Y) = \sum_i A_i^* Y A_i$. (2) $\Phi$ is completely positive iff $\Phi^*$ is. (3) $\Phi$ is unital and completely positive iff $\Phi^*$ is completely positive and trace preserving.

**Definition (notes).** The *Choi rank* of a completely positive $\Phi$ is $\mathrm{cr}(\Phi) = \min\{L : \Phi(X) = \sum_{l=1}^{L} B_l X B_l^*\}$.

**Theorem (notes)** (Choi)**.** $\mathrm{cr}(\Phi) = \mathrm{rank}(P_\Phi)$.

The functional proposition, the adjoint proposition (for Kraus maps) and the definition of the Choi rank are formalised below; the rank theorem is *choi-rank-eq-rank* in *Day20-Choi-Rank* (its lemma $\mathrm{rank}((E_{ij})) = 1$ is not stated separately), and the bound $\mathrm{cr}(\Phi) \le nd$ is below.

## 29.1 The adjoint of a Kraus map

For $\Phi(X) = \sum_i A_i X A_i^\dagger$ with $A_i \in M_{d,n}$, the adjoint is $\Phi^*(Y) = \sum_i A_i^\dagger Y A_i$ (*kraus-adjoint-dim n As*; the output dimension $n$ is kept explicit for the empty list).

**definition** *kraus-adjoint-dim* ::
*nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*kraus-adjoint-dim n As y* ≡
*foldr* (λ*A acc*. (*A*†) ∗ *y* ∗ *A* + *acc*) *As* (*0m n n*)

**lemma** *kraus-adjoint-dim-Nil* [*simp*]:
**fixes** *Y* :: *complex Matrix.mat*

**shows** *kraus-adjoint-dim n* [] *Y* = $0_m$ *n n*
⟨*proof*⟩

**lemma** *kraus-adjoint-dim-Cons* [*simp*]:
**fixes** *Y* :: *complex Matrix.mat*
**shows** *kraus-adjoint-dim n* (*A* # *As*) *Y* =
($A^\dagger$) ∗ *Y* ∗ *A* + *kraus-adjoint-dim n As Y*
⟨*proof*⟩

**lemma** *kraus-adjoint-dim-carrier*:
**fixes** *Y* :: *complex Matrix.mat*
**assumes** *As-c*: ∀ *A* ∈ *set As*. *A* ∈ *carrier-mat d n*
**and** *Y-c*: *Y* ∈ *carrier-mat d d*
**shows** *kraus-adjoint-dim n As Y* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *kraus-adjoint-dim-as-kraus-map-dim*:
**fixes** *Y* :: *complex Matrix.mat*
**assumes** *As-c*: ∀ *A* ∈ *set As*. *A* ∈ *carrier-mat d n*
**and** *Y-c*: *Y* ∈ *carrier-mat d d*
**shows** *kraus-adjoint-dim n As Y* =
*kraus-map-dim n* (*map* (λ*A*. ($A^\dagger$)) *As*) *Y*
⟨*proof*⟩

The adjoint identity $\langle Y|\Phi(X)\rangle = \langle \Phi^*(Y)|X\rangle$, by induction on the Kraus list and cyclicity of the trace.

**lemma** *hs-inner-prod-left-add*:
**assumes** *d1r*: *dim-row Y1* = *dim-row X* **and** *d1c*: *dim-col Y1* = *dim-col X*
**and** *d2r*: *dim-row Y2* = *dim-row X* **and** *d2c*: *dim-col Y2* = *dim-col X*
**shows** ⟨⟨ *Y1* + *Y2* | *X* ⟩⟩ = ⟨⟨ *Y1* | *X* ⟩⟩ + ⟨⟨ *Y2* | *X* ⟩⟩
⟨*proof*⟩

**lemma** *hs-inner-prod-right-zero*:
⟨⟨ *y* | $0_m$ *m n* ⟩⟩ = *0*
⟨*proof*⟩

**lemma** *hs-inner-prod-left-zero*:
**fixes** *X* :: *complex Matrix.mat*
**assumes** *X-c*: *X* ∈ *carrier-mat m n*
**shows** ⟨⟨ $0_m$ *m n* | *X* ⟩⟩ = *0*
⟨*proof*⟩

**lemma** *hs-inner-prod-kraus-term*:
**fixes** *A X Y* :: *complex Matrix.mat*
**assumes** *A-c*: *A* ∈ *carrier-mat d n*
**and** *X-c*: *X* ∈ *carrier-mat n n*
**and** *Y-c*: *Y* ∈ *carrier-mat d d*
**shows** ⟨⟨ *Y* | *A* ∗ *X* ∗ ($A^\dagger$) ⟩⟩ = ⟨⟨ ($A^\dagger$) ∗ *Y* ∗ *A* | *X* ⟩⟩
⟨*proof*⟩

**theorem** *kraus-adjoint-correct*:
**fixes** *As* :: *complex Matrix.mat list*
**and** *X* :: *complex Matrix.mat*
**and** *Y* :: *complex Matrix.mat*
**assumes** *As-c*: ∀ *A* ∈ *set As*. *A* ∈ *carrier-mat d n*
**and** *X-c*: *X* ∈ *carrier-mat n n*

**and** *Y-c*: $Y \in$ *carrier-mat d d*
**shows** $\langle\langle$ *Y* | *kraus-map-dim d As X* $\rangle\rangle$
$= \langle\langle$ *kraus-adjoint-dim n As Y* | *X* $\rangle\rangle$
⟨*proof*⟩

## 29.2 Complete positivity and the UCP/CPTP duality

$\Phi^*$ is the Kraus map of the adjoint operators, hence completely positive, and $\Phi$ is unital iff $\Phi^*$ is trace preserving: $\sum_i A_i A_i^* = I$ in both cases, using that $\mathrm{tr}(BY) = \mathrm{tr}(Y)$ for all $Y$ forces $B = I$.

**theorem** *adjoint-cp-from-kraus*:
**assumes** *As-c*: $\forall A \in$ *set As*. $A \in$ *carrier-mat d n*
**and** *npos*: $0 < n$
**and** *dpos*: $0 < d$
**shows** *completely-positive d n* (*kraus-adjoint-dim n As*)
⟨*proof*⟩

**lemma** *kraus-map-dim-one-eq-sum-wdag-w-dagger*:
**fixes** *As* :: *complex Matrix.mat list*
**assumes** *As-c*: $\forall A \in$ *set As*. $A \in$ *carrier-mat d n*
**shows** *kraus-map-dim d As* ($1_m$ *n*)
$=$ *sum-wdag-w d* (*map* ($\lambda A$. ($A^\dagger$)) *As*)
⟨*proof*⟩

**theorem** *cp-unital-iff-adjoint-tp*:
**assumes** *As-c*: $\forall A \in$ *set As*. $A \in$ *carrier-mat d n*
**and** *npos*: $0 < n$
**and** *dpos*: $0 < d$
**shows** *unital-map n d* (*kraus-map-dim d As*)
$\longleftrightarrow$ *trace-preserving d* (*kraus-adjoint-dim n As*)
⟨*proof*⟩

## 29.3 Positive functionals are completely positive

The case $d = 1$ of Choi's theorem: for $P = (P_{ij}) \in M_r(M_n)^+$ and $v = (\alpha_i) \in \mathbb{C}^r$, $\langle v|(\phi(P_{ij}))v\rangle = \phi(\sum_{ij} \overline{\alpha_i}\alpha_j P_{ij}) = \phi(Y^* P Y) \geq 0$ with $Y$ the block column $(\alpha_i I)_i$.

**lemma** *phi-entry-double-sum*:
**fixes** $\varphi$ :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *f-add*: $\bigwedge X\ Y$. $X \in$ *carrier-mat n n* $\Longrightarrow$ $Y \in$ *carrier-mat n n*
$\Longrightarrow \varphi\ (X + Y) = \varphi\ X + \varphi\ Y$
**and** *f-smult*: $\bigwedge c\ X$. $X \in$ *carrier-mat n n* $\Longrightarrow \varphi\ (c \cdot_m X) = c \cdot_m \varphi\ X$
**and** *f-car*: $\bigwedge X$. $X \in$ *carrier-mat n n* $\Longrightarrow \varphi\ X \in$ *carrier-mat d d*
**and** *M-c*: $M \in$ *carrier-mat n n*
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < d$
**and** *npos*: $0 < n$ **and** *dpos*: $0 < d$
**shows** $\varphi\ M$ \$\$ $(r, c) =$
$(\sum i{<}n.\ \sum j{<}n.\ M$ \$\$ $(i, j) * \varphi$ (*mat-unit i j n n*) \$\$ $(r, c))$
⟨*proof*⟩

**theorem** *positive-scalar-is-cp*:
**fixes** $\varphi$ :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *linear*: $\bigwedge A\ B$. $A \in$ *carrier-mat n n* $\Longrightarrow$ $B \in$ *carrier-mat n n*
$\Longrightarrow \varphi\ (A + B) = \varphi\ A + \varphi\ B$
**and** *smult*: $\bigwedge c\ A$. $A \in$ *carrier-mat n n* $\Longrightarrow \varphi\ (c \cdot_m A) = c \cdot_m \varphi\ A$
**and** *range*: $\forall A$. $A \in$ *carrier-mat n n* $\longrightarrow \varphi\ A \in$ *carrier-mat 1 1*
**and** *pos*: *positive-map n 1* $\varphi$
**and** *npos*: $0 < n$

**shows** *completely-positive n 1 φ*
⟨*proof*⟩

### 29.4 Choi rank

*choi-rank n d Φ* is the least length of a Kraus list representing $\Phi$ on the carrier. The theorem $\mathrm{cr}(\Phi) = \mathrm{rank}(P_\Phi)$ is *choi-rank-eq-rank* in *Day20-Choi-Rank*; the bound $\mathrm{cr}(\Phi) \le nd$ follows from the Kraus-rank bound of Day 21.

**definition** *choi-rank* ::
*nat ⇒ nat ⇒ (complex Matrix.mat ⇒ complex Matrix.mat) ⇒ nat* **where**
*choi-rank n d Φ ≡*
*Inf { length As | As :: complex Matrix.mat list.*
*(∀ A ∈ set As. A ∈ carrier-mat d n)*
*∧ (∀ X. X ∈ carrier-mat n n ⟶ Φ X = kraus-map-dim d As X) }*

## 30 Day 21 (14 October 2011): the Kraus rank bound; unitary freedom of representations

**Corollary (notes).** A completely positive $\Phi : M_n \to M_d$ has a Kraus representation $\Phi(X) = \sum_{l=1}^{K} A_l X A_l^*$ with $K \le nd$.

The notes then record a conjecture (every CP map is an average of $n$ UCP maps of Choi rank $\le d$) and develop the uniqueness theory:

**Lemma (notes).** If $T : \mathbb{C}^r \to \mathbb{C}^m$ is one-to-one then $T^*$ is onto.

**Proposition (notes).** If $v_1, \dots, v_r \in \mathbb{C}^k$ are linearly independent then so are $\{v_i v_j^*\}_{i,j} \subseteq M_k$.

**Lemma (notes).** If $0 \le ww^* \le P$ then $w \in \mathrm{range}(P)$; hence for $P = \sum_l w_l w_l^* \ge 0$, $\mathrm{range}(P) = \mathrm{span}\{w_l\}$.

**Theorem (notes).** If $P = \sum_{j=1}^{r} v_j v_j^* = \sum_{l=1}^{m} w_l w_l^*$ with $\mathrm{rank}\, P = r$, there is $U = (\alpha_{ij})_{m \times r}$ with $w_i = \sum_j \alpha_{ij} v_j$ and $U^*U = I_r$.

**Theorem (notes)** (Choi 2)**.** If $\Phi$ is completely positive with $\mathrm{cr}(\Phi) = r$ and $\Phi(X) = \sum_{j=1}^{r} V_j X V_j^* = \sum_{l=1}^{m} W_l X W_l^*$, then $W_i = \sum_j \alpha_{ij} V_j$ for a unique $U = (\alpha_{ij})$ with $U^*U = I_r$.

Only the Kraus rank bound is formalised here. The kernel form of the lemma ($\ker A \subseteq \ker B$ when $0 \le B \le A$) is proved in theory Day23 and used there to prove Choi's Theorem 3 without the unitary-freedom theorem; the remaining statements are audit items D21-3, D21-5, D21-6.

**corollary** *kraus-rank-bound*:
**fixes** *Φ :: complex Matrix.mat ⇒ complex Matrix.mat*
**assumes** *linear*: *⋀A B. A ∈ carrier-mat n n ⟹ B ∈ carrier-mat n n*
*⟹ Φ (A + B) = Φ A + Φ B*
**and** *smult*: *⋀c A. A ∈ carrier-mat n n ⟹ Φ (c ·m A) = c ·m Φ A*
**and** *range*: *∀ A. A ∈ carrier-mat n n ⟶ Φ A ∈ carrier-mat d d*
**and** *cp*: *completely-positive n d Φ*
**and** *npos*: *0 < n*
**and** *dpos*: *0 < d*
**shows** *∃ As :: complex Matrix.mat list.*
*length As ≤ n ∗ d ∧*
*(∀ A ∈ set As. A ∈ carrier-mat d n) ∧*
*(∀ X. X ∈ carrier-mat n n ⟶ Φ X = kraus-map-dim d As X)*
⟨*proof*⟩

Dimension bookkeeping for $T^\dagger$ (the onto statement is not formalised).

**lemma** *adjoint-image-in-dom*:
**fixes** $T$ :: *complex Matrix.mat*
**assumes** *sq*: $T \in$ *carrier-mat m r*
**shows** $\forall w$ :: *complex Matrix.vec*. *dim-vec* $w = m$
$\longrightarrow$ *dim-vec* $(T^\dagger *_v w) = r$
⟨*proof*⟩

$\mathrm{cr}(\Phi) \leq nd$: the defining set of the Choi rank is non-empty.

**corollary** *choi-rank-le*:
**fixes** $\Phi$ :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *linear*: $\bigwedge A\ B.\ A \in$ *carrier-mat n n* $\Longrightarrow B \in$ *carrier-mat n n*
$\Longrightarrow \Phi\ (A + B) = \Phi\ A + \Phi\ B$
**and** *smult*: $\bigwedge c\ A.\ A \in$ *carrier-mat n n* $\Longrightarrow \Phi\ (c \cdot_m A) = c \cdot_m \Phi\ A$
**and** *range*: $\forall A.\ A \in$ *carrier-mat n n* $\longrightarrow \Phi\ A \in$ *carrier-mat d d*
**and** *cp*: *completely-positive n d* $\Phi$
**and** *npos*: $0 < n$
**and** *dpos*: $0 < d$
**shows** *choi-rank n d* $\Phi \leq n * d$
⟨*proof*⟩

# 31 Day 22 (17 October 2011): minimal Kraus representations, convex sets, UCP and CPTP

**Proposition (notes).** If $\Phi$ is completely positive, $\mathrm{cr}(\Phi) = r$ and $\Phi(X) = \sum_{i=1}^{r} V_i^* X V_i$, then $V_1, \ldots, V_r$ are linearly independent.

(Not formalised: on Day 23 linear independence of the Kraus operators is a hypothesis.)

**Definition (notes).** $C \subseteq V$ is *convex* if $tv_1 + (1-t)v_2 \in C$ for $v_1, v_2 \in C$, $0 \leq t \leq 1$; $v \in C$ is *extreme* if $v = \frac{1}{2}(v_1 + v_2)$ with $v_1, v_2 \in C$ forces $v = v_1 = v_2$. The *convex hull* of $E$ is the set of convex combinations of points of $E$; a map between convex sets is *affine* if it preserves convex combinations.

**Theorem (notes)** (Krein–Milman)**.** A compact convex subset of a finite-dimensional real vector space is the convex hull of its extreme points.

Examples: $UCP(M_n, M_d)$, the unital completely positive maps, is convex ($P_{t\Phi_1+(1-t)\Phi_2} = tP_{\Phi_1} + (1-t)P_{\Phi_2} \geq 0$) and compact (closed; bounded because $0 \leq \Phi(E_{ii}) \leq I$ gives $|r_{ij}| \leq 1$ for the entries of $P_\Phi$); $CPTP(M_d, M_n)$ is convex and compact, being the image of $UCP(M_n, M_d)$ under the affine bijection $\Gamma(\Phi) = \Phi^*$.

**Theorem (notes)** (Choi 3)**.** If $\Phi : M_n \to M_d$ is unital and completely positive, $\Phi(X) = \sum_{i=1}^{r} V_i^* X V_i$ with $r = \mathrm{cr}(\Phi)$, then $\Phi$ is an extreme point of $UCP(M_n, M_d)$ iff $\{V_i^* V_j : 1 \leq i, j \leq r\}$ is linearly independent in $M_d$.

**Corollary (notes).** If $\Phi$ is extreme then $\mathrm{cr}(\Phi) \leq d$. If $\Phi : M_d \to M_n$ is CPTP with $\Phi(X) = \sum_{i=1}^{r} V_i X V_i^*$, $r = \mathrm{cr}(\Phi)$, then $\Phi$ is extreme in $CPTP(M_d, M_n)$ iff $\{V_i^* V_j\}$ is linearly independent.

Convexity of UCP and CPTP is proved below; compactness and the affine bijection are not formalised; the corollary $\mathrm{cr}(\Phi) \leq d$ is *extreme-CPTP-choi-rank-le* in *Day22-Kraus-Independence*, and Choi's Theorem 3 is proved in theory Day23 in its CPTP form. Krein–Milman is available in HOL-Analysis (*Krein-Milman-Minkowski*).

## 31.1 Convex sets and extreme points

**definition** *convex-set-mat* :: *complex Matrix.mat set* ⇒ *bool* **where**
*convex-set-mat C* ≡
∀ *A* ∈ *C*. ∀ *B* ∈ *C*. ∀ *t* :: *real*. 0 ≤ *t* ∧ *t* ≤ 1 ⟶
*complex-of-real t* ·$_m$ *A*
+ (1 − *complex-of-real t*) ·$_m$ *B* ∈ *C*

**definition** *extreme-point-mat* ::
*complex Matrix.mat set* ⇒ *complex Matrix.mat* ⇒ *bool* **where**
*extreme-point-mat C v* ≡
*v* ∈ *C* ∧
¬ (∃ *v1* ∈ *C*. ∃ *v2* ∈ *C*. *v1* ≠ *v* ∧ *v* ≠ *v2* ∧
*v* = (1/2 :: *complex*) ·$_m$ *v1* + (1/2 :: *complex*) ·$_m$ *v2*)

**definition** *affine-map-mat* ::
(*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*affine-map-mat tmap* ≡
∀ *A B*. ∀ *t* :: *real*. 0 ≤ *t* ∧ *t* ≤ 1 ⟶
*tmap* (*complex-of-real t* ·$_m$ *A*
+ (1 − *complex-of-real t*) ·$_m$ *B*) =
*complex-of-real t* ·$_m$ *tmap A*
+ (1 − *complex-of-real t*) ·$_m$ *tmap B*

## 31.2 The convex sets UCP and CPTP

**definition** *UCP* ::
*nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) *set* **where**
*UCP n d* ≡ {Φ. *completely-positive n d* Φ ∧ *unital-map n d* Φ}

**definition** *CPTP* ::
*nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) *set* **where**
*CPTP n d* ≡ {Φ. *completely-positive n d* Φ ∧ *trace-preserving n* Φ}

**lemma** *amplification-smult*:
**assumes** *dpos*: 0 < *d*
**and** *F-block*: ⋀*a b*. *F* (*block-of-mat n x a b*) ∈ *carrier-mat d d*
**shows** *amplification n d p* (λϱ. *c* ·$_m$ *F* ϱ) *x*
= *c* ·$_m$ *amplification n d p F x*
⟨*proof*⟩

UCP is convex. The range hypotheses are explicit because *completely-positive* does not constrain the values of $\Phi_i$ off the carrier.

**lemma** *UCP-convex*:
**fixes** *t* :: *real*
**assumes** *t-nn*: 0 ≤ *t*
**and** *t-le*: *t* ≤ 1
**and** Φ*1-in*: Φ*1* ∈ *UCP n d*
**and** Φ*2-in*: Φ*2* ∈ *UCP n d*
**and** *r1*: ∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ*1 X* ∈ *carrier-mat d d*
**and** *r2*: ∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ*2 X* ∈ *carrier-mat d d*
**and** *dpos*: 0 < *d*
**shows** (λ*X*. *complex-of-real t* ·$_m$ Φ*1 X*
+ (1 − *complex-of-real t*) ·$_m$ Φ*2 X*) ∈ *UCP n d*
⟨*proof*⟩

CPTP is convex.

**lemma** *CPTP-convex*:
**fixes** *t* :: *real*
**assumes** *t-nn*: $0 \leq t$
**and** *t-le*: $t \leq 1$
**and** *Φ1-in*: $\Phi 1 \in CPTP\ n\ d$
**and** *Φ2-in*: $\Phi 2 \in CPTP\ n\ d$
**and** *r1*: $\forall X.\ X \in carrier\text{-}mat\ n\ n \longrightarrow \Phi 1\ X \in carrier\text{-}mat\ d\ d$
**and** *r2*: $\forall X.\ X \in carrier\text{-}mat\ n\ n \longrightarrow \Phi 2\ X \in carrier\text{-}mat\ d\ d$
**and** *dpos*: $0 < d$
**shows** $(\lambda X.\ complex\text{-}of\text{-}real\ t \cdot_m \Phi 1\ X + (1 - complex\text{-}of\text{-}real\ t) \cdot_m \Phi 2\ X) \in CPTP\ n\ d$
⟨*proof*⟩

# 32 Day 29 (7 November 2011): classical error detection and correction

Days 23–28 are formalised in the theories Day23_ChoiExtremeUCP, Separable_Cone and Day24_Separability.

To transmit $v \in \mathbb{Z}_2^n$ through a channel that may flip bits (+1 in $\mathbb{Z}_2$), one encodes $v \mapsto \phi(v) \in \mathbb{Z}_2^{n+k}$, transmits, and decodes. Examples: repetition $v \mapsto (v, v)$ detects one error but corrects none and doubles the cost; the parity check $(a_1, \ldots, a_n) \mapsto (a_1, \ldots, a_n; a_1 + \cdots + a_n)$ detects one error cheaply; the Hamming $[7, 1, 3]$ code detects and corrects one error (theory Day29_Classical_Codes: the code as written detects one error but cannot correct it; the $[7, 4, 3]$ code corrects one).

**Definition (notes).** The *Hamming weight* $\|w\|$ of $w \in \mathbb{Z}_2^m$ is its number of non-zero entries; the weight of a code $S \subseteq \mathbb{Z}_2^m$ is $\min\{\|v - w\| : v \neq w \in S\}$, the least number of errors turning one code word into another; for a subspace code it is $\min\{\|v\| : 0 \neq v \in S\}$.

Decoding amounts to choosing a left inverse of $\phi$ (different left inverses correct differently); the best binary codes are cyclic codes, ideals in $\mathbb{Z}_2[x]/\langle x^m - 1\rangle$ (not formalised).

## 32.1 Binary strings, Hamming weight and distance

**definition** *hamming-weight* :: *nat list* ⇒ *nat* **where**
*hamming-weight v* ≡ *length* (*filter* (λ*b*. *b* ≠ *0*) *v*)

**definition** *hamming-dist* :: *nat list* ⇒ *nat list* ⇒ *nat* **where**
*hamming-dist v w* ≡ *hamming-weight* (*map2* (λ*a b*. (*a* + *b*) *mod 2*) *v w*)

**lemma** *hamming-weight-Nil* [*simp*]: *hamming-weight* [] = *0*
⟨*proof*⟩

**lemma** *hamming-weight-Cons*:
*hamming-weight* (*a* # *v*)
= (*if a* ≠ *0 then Suc* (*hamming-weight v*) *else hamming-weight v*)
⟨*proof*⟩

**lemma** *hamming-dist-Nil* [*simp*]:
*hamming-dist* [] [] = *0*
⟨*proof*⟩

**lemma** *hamming-dist-Cons*:
*hamming-dist* (*a* # *v*) (*b* # *w*)
= (*if* (*a* + *b*) *mod 2* ≠ *0*

*then Suc (hamming-dist v w)*
*else hamming-dist v w)*
⟨*proof*⟩

**lemma** *hamming-dist-symm*:
**assumes** *length v = length w*
**shows** *hamming-dist v w = hamming-dist w v*
⟨*proof*⟩

**lemma** *hamming-dist-zero*:
**assumes** *lenw*: *length v = length w*
**and** *bits-v*: ∀ *x* ∈ *set v*. *x < 2*
**and** *bits-w*: ∀ *x* ∈ *set w*. *x < 2*
**shows** *hamming-dist v w = 0* ⟷ *v = w*
⟨*proof*⟩

**lemma** *hamming-dist-self* [*simp*]:
*hamming-dist v v = 0*
⟨*proof*⟩

## 32.2 Code distance

A code of distance at least $2t+1$ can correct $t$ errors (recorded as the definition *corrects-errors*; the decoding theorem itself is not formalised).

**definition** *code-distance* :: *nat list set* ⇒ *nat* **where**
*code-distance code* ≡
*if* (∃ *v* ∈ *code*. ∃ *w* ∈ *code*. *v* ≠ *w*)
*then Min* {*hamming-dist v w* | *v w*. *v* ∈ *code* ∧ *w* ∈ *code* ∧ *v* ≠ *w*}
*else 0*

**definition** *corrects-errors* :: *nat list set* ⇒ *nat* ⇒ *bool* **where**
*corrects-errors code t* ≡ *code-distance code* ≥ *2*∗*t + 1*

## 32.3 The parity-check and repetition codes

Parity check: append the sum of the bits; any single error changes the code word. Repetition doubles Hamming distances.

**definition** *parity-bit* :: *nat list* ⇒ *nat* **where**
*parity-bit v* ≡ *foldr* (λ*a b*. (*a + b*) *mod 2*) *v 0*

**definition** *parity-check-encode* :: *nat list* ⇒ *nat list* **where**
*parity-check-encode v* ≡ *v* @ [*parity-bit v*]

**lemma** *length-parity-check-encode* [*simp*]:
*length* (*parity-check-encode v*) = *length v + 1*
⟨*proof*⟩

**lemma** *hamming-weight-pos-iff*:
*0 < hamming-weight v* ⟷ (∃ *x* ∈ *set v*. *x* ≠ *0*)
⟨*proof*⟩

**lemma** *hamming-dist-pos-iff*:
**assumes** *lenw*: *length v = length w*
**and** *bits-v*: ∀ *x* ∈ *set v*. *x < 2*
**and** *bits-w*: ∀ *x* ∈ *set w*. *x < 2*
**shows** *0 < hamming-dist v w* ⟷ *v* ≠ *w*
⟨*proof*⟩

**lemma** *parity-bit-lt-2*: *parity-bit v < 2*
⟨*proof*⟩

**lemma** *parity-check-encode-binary*:
**assumes** ∀ *x* ∈ *set v*. *x < 2*
**shows** ∀ *x* ∈ *set* (*parity-check-encode v*). *x < 2*
⟨*proof*⟩

**lemma** *parity-check-detects-one-error*:
**assumes** *lenE*: *length e = length v + 1*
**and** *bits-v*: ∀ *x* ∈ *set v*. *x < 2*
**and** *bits-e*: ∀ *x* ∈ *set e*. *x < 2*
**and** *weight*: *hamming-weight e = 1*
**shows** *parity-check-encode v*
≠ *map2* (λ*a b*. (*a + b*) *mod 2*) (*parity-check-encode v*) *e*
⟨*proof*⟩

**definition** *repetition-encode* :: *nat list* ⇒ *nat list* **where**
*repetition-encode v* ≡ *v* @ *v*

**lemma** *length-repetition-encode* [*simp*]:
*length* (*repetition-encode v*) = *2* ∗ *length v*
⟨*proof*⟩

**lemma** *hamming-dist-repetition-doubles*:
**assumes** *length v = length w*
**shows** *hamming-dist* (*repetition-encode v*) (*repetition-encode w*)
= *2* ∗ *hamming-dist v w*
⟨*proof*⟩

# 33 Day 30 (9 November 2011): error probabilities; the three-qubit bit-flip code

If a bit flips with probability $p$ (independently), the repetition code on three bits leaves an error undetected with probability $3p^2(1-p)^4 + 3p^4(1-p)^2 + p^6$, the parity code with probability $\binom{4}{2}p^2(1-p)^2 + p^4$ (the former is smaller for $p < \frac{1}{2}$), and the majority-rule code $0 \mapsto 000$, $1 \mapsto 111$ decodes wrongly with probability $3p^2(1-p)+p^3$ (theory Day29_Classical_Codes; the comparison holds for all $0 < p < 1$).

Quantum error correction: one cannot clone, but basis vectors can be copied and measurements can decode. With $X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ the allowed errors are qubit flips $X \otimes I \otimes I$, $I \otimes X \otimes I$, .... The *three-qubit bit-flip code* encodes $a|0\rangle + b|1\rangle \mapsto a|000\rangle + b|111\rangle$ (two CNOTs). Decoding: (1) syndrome measurement with $P_0 = |000\rangle\langle 000| + |111\rangle\langle 111|$, $P_1 = |100\rangle\langle 100| + |011\rangle\langle 011|$, $P_2 = |010\rangle\langle 010| + |101\rangle\langle 101|$, $P_3 = |001\rangle\langle 001| + |110\rangle\langle 110|$, $\sum P_i^2 = I$, which leaves a state with at most one flip unchanged and reveals where the flip occurred; (2) recovery $R(Y) = P_0 Y P_0 + (X \otimes I \otimes I) P_1 Y P_1 (X \otimes I \otimes I) + \cdots$, a completely positive trace-preserving map with $R(|\psi_i\rangle\langle\psi_i|) = |\psi\rangle\langle\psi|$ after zero or one error. The code's recovery theorem is *bit-flip-code-recovery* in *Day30-BitFlip-Code* (syndrome projections, $\sum P_i = I$, $R$ CPTP); the Pauli matrices and the bit-flip channel are below.

## 33.1 The Pauli matrices

The notes use two sign conventions for $Y$: $\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ on Day 10 (the library's $Y$) and $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$ on Day 32. *Pauli-Y* follows Day 32; both are Hermitian unitary involutions, and the Pauli identities below carry the corresponding signs. *Pauli-X* and *Pauli-Z* are the library's *Quantum.X* and *Quantum.Z*.

**definition** *Pauli-X* :: *complex Matrix.mat* **where**
*Pauli-X* ≡ *Matrix.mat 2 2*
(λ(*i*,*j*). *if* (*i* = *0* ∧ *j* = *1*) ∨ (*i* = *1* ∧ *j* = *0*) *then 1 else 0*)

**definition** *Pauli-Y* :: *complex Matrix.mat* **where**
*Pauli-Y* ≡ *Matrix.mat 2 2*
(λ(*i*,*j*). *if i* = *0* ∧ *j* = *1 then* i
*else if i* = *1* ∧ *j* = *0 then* −i *else 0*)

**definition** *Pauli-Z* :: *complex Matrix.mat* **where**
*Pauli-Z* ≡ *Matrix.mat 2 2*
(λ(*i*,*j*). *if i* = *0* ∧ *j* = *0 then 1*
*else if i* = *1* ∧ *j* = *1 then* −*1 else 0*)

**lemma** *Pauli-X-carrier* [*simp*]: *Pauli-X* ∈ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *Pauli-Y-carrier* [*simp*]: *Pauli-Y* ∈ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *Pauli-Z-carrier* [*simp*]: *Pauli-Z* ∈ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *Pauli-X-dagger*: *Pauli-X*$^\dagger$ = *Pauli-X*
⟨*proof*⟩

**lemma** *Pauli-X-squared*: *Pauli-X* ∗ *Pauli-X* = *1*$_m$ *2*
⟨*proof*⟩

**lemma** *Pauli-X-unitary*: *unitary Pauli-X*
⟨*proof*⟩

**lemma** *Pauli-Y-dagger*: *Pauli-Y*$^\dagger$ = *Pauli-Y*
⟨*proof*⟩

**lemma** *Pauli-Y-squared*: *Pauli-Y* ∗ *Pauli-Y* = *1*$_m$ *2*
⟨*proof*⟩

**lemma** *Pauli-Y-unitary*: *unitary Pauli-Y*
⟨*proof*⟩

**lemma** *Pauli-Z-dagger*: *Pauli-Z*$^\dagger$ = *Pauli-Z*
⟨*proof*⟩

**lemma** *Pauli-Z-squared*: *Pauli-Z* ∗ *Pauli-Z* = *1*$_m$ *2*
⟨*proof*⟩

**lemma** *Pauli-Z-unitary*: *unitary Pauli-Z*
⟨*proof*⟩

## 33.2 Pauli magic (Day 32)

**Proposition (notes).** $I, X, Y, Z$ are orthogonal in $M_2$ and all have 2-norm $\sqrt{2}$.

**proposition** *pauli-orthogonal*:
**shows** $\langle\langle\ 1_m\ 2 \mid \textit{Pauli-X}\ \rangle\rangle = 0$
**and** $\langle\langle\ 1_m\ 2 \mid \textit{Pauli-Y}\ \rangle\rangle = 0$
**and** $\langle\langle\ 1_m\ 2 \mid \textit{Pauli-Z}\ \rangle\rangle = 0$
**and** $\langle\langle\ \textit{Pauli-X} \mid \textit{Pauli-Y}\ \rangle\rangle = 0$
**and** $\langle\langle\ \textit{Pauli-X} \mid \textit{Pauli-Z}\ \rangle\rangle = 0$
**and** $\langle\langle\ \textit{Pauli-Y} \mid \textit{Pauli-Z}\ \rangle\rangle = 0$
⟨*proof*⟩

**proposition** *pauli-norm*:
**shows** $\langle\langle\ 1_m\ 2 \mid (1_m\ 2 :: \textit{complex Matrix.mat})\ \rangle\rangle = 2$
**and** $\langle\langle\ \textit{Pauli-X} \mid \textit{Pauli-X}\ \rangle\rangle = 2$
**and** $\langle\langle\ \textit{Pauli-Y} \mid \textit{Pauli-Y}\ \rangle\rangle = 2$
**and** $\langle\langle\ \textit{Pauli-Z} \mid \textit{Pauli-Z}\ \rangle\rangle = 2$
⟨*proof*⟩

## 33.3 The bit-flip channel (Day 31)

Doing nothing to $\psi$ while a flip occurs with probability $p$ produces the ensemble $\{1-p, \psi\}, \{p, X\psi\}$, i.e. the error map $E(Y) = (1-p)Y + pXYX^*$, which is completely positive and trace preserving.

**definition** *bit-flip-channel* ::
*real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*bit-flip-channel p ϱ* ≡
*complex-of-real* (1 − *p*) ·$_m$ *ϱ*
+ *complex-of-real p* ·$_m$ (*Pauli-X* ∗ *ϱ* ∗ (*Pauli-X*$^\dagger$))

**lemma** *bit-flip-channel-carrier*:
**assumes** *ϱ* ∈ *carrier-mat 2 2*
**shows** *bit-flip-channel p ϱ* ∈ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *bit-flip-channel-trace-preserving*:
**fixes** *p* :: *real*
**assumes** *ϱ-c*: *ϱ* ∈ *carrier-mat 2 2*
**shows** *Tr* (*bit-flip-channel p ϱ*) = *Tr ϱ*
⟨*proof*⟩

**lemma** *bit-flip-channel-psd*:
**fixes** *p* :: *real*
**assumes** *p-nn*: 0 ≤ *p*
**and** *p-le-1*: *p* ≤ 1
**and** *ϱ-c*: *ϱ* ∈ *carrier-mat 2 2*
**and** *ϱ-psd*: *psd ϱ*
**shows** *psd* (*bit-flip-channel p ϱ*)
⟨*proof*⟩

# 34 Day 31 (11 November 2011): the bit-flip code as a channel; fidelity

Encoding $\tilde{\psi} = a|000\rangle + b|111\rangle$ and flipping each qubit independently with probability $p$ gives the error map $E(Y) = (1-p)^3 Y + p(1-p)^2 \sum_j (X_j) Y (X_j)^* + \cdots + p^3 (X \otimes X \otimes X) Y (X \otimes X \otimes X)^*$, and with the recovery $R$ of Day 30,

$$R \circ E(Y) = \big((1-p)^3 + 3p(1-p)^2\big) Y + \big(3p^2(1-p) + p^3\big)(X \otimes X \otimes X) Y (X \otimes X \otimes X).$$

The code helps iff $(1-p)^3 + 3p(1-p)^2 \geq 1-p$ iff $p \leq \frac{1}{2}$ (theory Day31_Code_Fidelity, where the formula for $R \circ E$ holds for $Y$ supported on the code space). Below: the encoding isometry.

**definition** *encode-mat* :: *complex Matrix.mat* **where**
*encode-mat* ≡ *Matrix.mat 8 2* (λ(*i*,*j*).
*if i = 0 ∧ j = 0 then 1*
*else if i = 7 ∧ j = 1 then 1*
*else 0*)

**lemma** *encode-mat-carrier* [*simp*]: *encode-mat ∈ carrier-mat 8 2*
⟨*proof*⟩

## 34.1 Fidelity

A state $\psi$ perturbed to $\psi' = a\psi + b\psi^\perp$ is close to $\psi$ when $|a|$ is close to 1, so the *fidelity* of states is $F = |\langle\psi'|\psi\rangle|$; for the density matrices $\rho = |\psi\rangle\langle\psi|$, $\rho' = |\psi'\rangle\langle\psi'|$ one has $\langle\rho|\rho'\rangle = \mathrm{Tr}(\rho\rho') = |\langle\psi'|\psi\rangle|^2$, so $F = \sqrt{\mathrm{Tr}(\rho\rho')}$. Error correction is compared by $F = \sqrt{\langle\psi|R \circ E(|\psi\rangle\langle\psi|)\psi\rangle}$, minimised or averaged over $\psi$: doing nothing gives $\min F = \sqrt{1-p}$, the bit-flip code gives $\min F = \sqrt{(1-p)^3 + 3p(1-p)^2}$ (theory Day31_Code_Fidelity). Formally *fidelity* $\psi$ $\varrho$ is $\sqrt{\Re\langle\psi|\rho\psi\rangle}$.

**definition** *fidelity* ::
*complex Matrix.vec* ⇒ *complex Matrix.mat* ⇒ *real* **where**
*fidelity* ψ ϱ ≡ *sqrt* (*Re* (⟨ψ | ϱ *$_v$* ψ⟩))

**lemma** *fidelity-def-eq*:
*fidelity* ψ ϱ = *sqrt* (*Re* (⟨ψ | ϱ *$_v$* ψ⟩))
⟨*proof*⟩

**lemma** *fidelity-pure-states*:
**fixes** ψ ψ′ :: *complex Matrix.vec*
**assumes** ψ-*qs*: *quantum-state n* ψ
**and** ψ′-*qs*: *quantum-state n* ψ′
**shows** *fidelity* ψ ((*mat-of-cols n* [ψ′]) ∗ ((*mat-of-cols n* [ψ′])†))
= *cmod* (⟨ψ | ψ′⟩)
⟨*proof*⟩

**lemma** *fidelity-self*:
**fixes** ψ :: *complex Matrix.vec*
**assumes** ψ-*qs*: *quantum-state n* ψ
**shows** *fidelity* ψ ((*mat-of-cols n* [ψ]) ∗ ((*mat-of-cols n* [ψ])†)) = *1*
⟨*proof*⟩

**lemma** *fidelity-orthogonal*:
**fixes** ψ ψ′ :: *complex Matrix.vec*

**assumes** *ψ-qs*: *quantum-state n ψ*
**and** *ψ′-qs*: *quantum-state n ψ′*
**and** *ortho*: $\langle \psi \mid \psi' \rangle = 0$
**shows** *fidelity ψ* ((*mat-of-cols n* [*ψ′*]) ∗ ((*mat-of-cols n* [*ψ′*])†)) = *0*
⟨*proof*⟩

# 35 Day 32 (14 November 2011): the phase-flip code, the Shor code, Pauli magic

## 35.1 The phase-flip code

With $Z|0\rangle = |0\rangle$, $Z|1\rangle = -|1\rangle$ and $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$ one has $|+\rangle \perp |-\rangle$, $Z|+\rangle = |-\rangle$, $Z|-\rangle = |+\rangle$: $Z$ acts on $\{|+\rangle, |-\rangle\}$ as the bit flip acts on $\{|0\rangle, |1\rangle\}$. Encoding $|0\rangle \mapsto |++\rangle$, $|1\rangle \mapsto |--\rangle$ gives the three-qubit *phase-flip code*, which detects and corrects phase flips exactly as the bit-flip code corrects bit flips; with $\beta = Y|+\rangle$ (orthogonal to $|+\rangle$, $Y\beta = |+\rangle$) the same works for $Y$.

The *Shor code* encodes $|0\rangle \mapsto |0_L\rangle = 2^{-3/2}(|000\rangle + |111\rangle)^{\otimes 3}$ and $|1\rangle \mapsto |1_L\rangle = 2^{-3/2}(|000\rangle - |111\rangle)^{\otimes 3}$ in $\mathbb{C}^{2^9}$.

**Theorem (notes)** (Shor)**.** There is a completely positive trace-preserving $R : M_{2^9} \to M_{2^9}$ such that for every $2 \times 2$ unitary $U$, every position $j$, $\tilde{U} = I \otimes \cdots \otimes U \otimes \cdots \otimes I$, and every $\psi = a|0_L\rangle + b|1_L\rangle$, $R(\tilde{U}|\psi\rangle\langle\psi|\tilde{U}^*) = |\psi\rangle\langle\psi|$.

**Proposition (notes).** If $V \subseteq \mathbb{C}^n$ is a subspace and $U_1, \ldots, U_t$ are unitaries with $U_i V \perp U_j V$ for $i \neq j$, then there is a CPTP $R$ with $R(U_i|\psi\rangle\langle\psi|U_i^*) = |\psi\rangle\langle\psi|$ for all $\psi \in V$ and all $i$ (take $R(X) = \sum_i U_i^* P_i X P_i U_i$ with $P_i$ the projection onto $U_i V$, $P_0 = I - \sum P_i$, $U_0 = I$).

The $3 \cdot 9 + 1$ *1-Pauli* operators $I \otimes \cdots \otimes U \otimes \cdots \otimes I$ ($U$ a Pauli matrix) map $V = \mathrm{span}\{|0_L\rangle, |1_L\rangle\}$ to mutually orthogonal subspaces (up to the coincidences $Z_1 v = Z_2 v = Z_3 v$ etc.). The proposition is *recovery-single-error* in *Day32-Recovery-Proposition*, together with its linear-combination form *recovery-linear-combination* (Day 33); the Pauli expansion identity is *pauli-expansion-unitary* in *Day32-Pauli-Expansion*. The Shor code and Shor's theorem are *Day33-Shor-Code* (*shor-code-theorem*). As first stated, the orthogonality claim is false ($Z_1 \neq Z_2$ but $Z_1|0_L\rangle = Z_2|0_L\rangle$, *shor-notes-Z1-Z2*); Day 33 corrects it.

**definition** *ket-plus* :: *complex Matrix.vec* **where**
*ket-plus* ≡ (*1 / sqrt 2* :: *complex*) ·$_v$ (*unit-vec 2 0 + unit-vec 2 1*)

**definition** *ket-minus* :: *complex Matrix.vec* **where**
*ket-minus* ≡ (*1 / sqrt 2* :: *complex*) ·$_v$ (*unit-vec 2 0 − unit-vec 2 1*)

**lemma** *ket-plus-carrier* [*simp*]: *ket-plus* ∈ *carrier-vec 2*
⟨*proof*⟩

**lemma** *ket-minus-carrier* [*simp*]: *ket-minus* ∈ *carrier-vec 2*
⟨*proof*⟩

**lemma** *ket-plus-dim* [*simp*]: *dim-vec ket-plus = 2*
⟨*proof*⟩

**lemma** *ket-minus-dim* [*simp*]: *dim-vec ket-minus = 2*
⟨*proof*⟩

**lemma** *ket-plus-index-0*: *ket-plus* $ *0 = 1 / sqrt 2*
⟨*proof*⟩

**lemma** *ket-plus-index-1*: *ket-plus $ 1 = 1 / sqrt 2*
⟨*proof*⟩

**lemma** *ket-minus-index-0*: *ket-minus $ 0 = 1 / sqrt 2*
⟨*proof*⟩

**lemma** *ket-minus-index-1*: *ket-minus $ 1 = − 1 / sqrt 2*
⟨*proof*⟩

**lemma** *ket-plus-unit*: *‖ket-plus‖ = 1*
⟨*proof*⟩

**lemma** *ket-minus-unit*: *‖ket-minus‖ = 1*
⟨*proof*⟩

**lemma** *ket-plus-minus-orthog*: *⟨ket-plus | ket-minus⟩ = 0*
⟨*proof*⟩

**lemma** *Pauli-Z-ket-plus*: *Pauli-Z ∗v ket-plus = ket-minus*
⟨*proof*⟩

**lemma** *Pauli-Z-ket-minus*: *Pauli-Z ∗v ket-minus = ket-plus*
⟨*proof*⟩

**lemma** *Pauli-X-ket-plus*: *Pauli-X ∗v ket-plus = ket-plus*
⟨*proof*⟩

**lemma** *Pauli-X-ket-minus*: *Pauli-X ∗v ket-minus = − ket-minus*
⟨*proof*⟩

The code words $|++\!+\rangle$ and $|--\!-\rangle$ of the three-qubit phase-flip code (an earlier version of this file called them *phase-code-zero*/*phase-code-one*; the nine-qubit Shor code words are not defined).

**definition** *phase-code-zero* :: *complex Matrix.vec* **where**
*phase-code-zero* ≡
*let v = unit-vec 2 0 + unit-vec 2 1 in*
*(1 / (2 ∗ sqrt 2)* :: *complex)*
*·v tensor-vec (tensor-vec v v) v*

**definition** *phase-code-one* :: *complex Matrix.vec* **where**
*phase-code-one* ≡
*let v = unit-vec 2 0 − unit-vec 2 1 in*
*(1 / (2 ∗ sqrt 2)* :: *complex)*
*·v tensor-vec (tensor-vec v v) v*

**lemma** *phase-code-zero-dim* [*simp*]: *dim-vec phase-code-zero = 8*
⟨*proof*⟩

**lemma** *phase-code-one-dim* [*simp*]: *dim-vec phase-code-one = 8*
⟨*proof*⟩

**lemma** *phase-code-zero-carrier* [*simp*]: *phase-code-zero ∈ carrier-vec 8*
⟨*proof*⟩

**lemma** *phase-code-one-carrier* [*simp*]: *phase-code-one ∈ carrier-vec 8*
⟨*proof*⟩

## 35.2 Projections

A *projection* on $\mathbb{C}^n$ is a Hermitian idempotent, the notion of Day 2; projections are the building blocks of syndrome measurements and recovery maps (Days 30–34). Day 2's theorem that such a $P$ is the orthogonal projection onto its range follows the basic facts.

**definition** *projection* :: *nat* ⇒ *complex Matrix.mat* ⇒ *bool* **where**
  *projection n P* ≡
    $P \in$ *carrier-mat n n* ∧ *hermitian P* ∧ $P * P = P$

**lemma** *projection-carrier*:
  *projection n P* ⟹ $P \in$ *carrier-mat n n*
  ⟨*proof*⟩

**lemma** *projection-hermitian*:
  *projection n P* ⟹ *hermitian P*
  ⟨*proof*⟩

**lemma** *projection-idempotent*:
  *projection n P* ⟹ $P * P = P$
  ⟨*proof*⟩

**lemma** *projection-zero*: *projection n* ($0_m$ *n n*)
  ⟨*proof*⟩

**lemma** *projection-one*: *projection n* ($1_m$ *n*)
  ⟨*proof*⟩

**lemma** *projection-psd*:
  **assumes** *projection n P*
  **shows** *psd P*
⟨*proof*⟩

The theorem of Day 2: a Hermitian idempotent $P$ is the orthogonal projection onto its range, i.e. $Pv = v$ for $v$ in the range and $Pw = 0$ for $w$ orthogonal to the range.

**lemma** *projection-fixes-range*:
  **fixes** *P* :: *complex Matrix.mat* **and** *x* :: *complex Matrix.vec*
  **assumes** *P*: *projection n P* **and** *x*: *dim-vec x* = *n*
  **shows** $P *_v (P *_v x) = P *_v x$
⟨*proof*⟩

**lemma** *projection-annihilates-orthogonal*:
  **fixes** *P* :: *complex Matrix.mat* **and** *w* :: *complex Matrix.vec*
  **assumes** *P*: *projection n P* **and** *w*: *dim-vec w* = *n*
    **and** *orth*: ⋀*x* :: *complex Matrix.vec*. *dim-vec x* = *n* ⟹ ⟨$P *_v$ *x* | *w*⟩ = *0*
  **shows** $P *_v w = 0_v$ *n*
⟨*proof*⟩

**lemma** *projection-complement-idempotent*:
  **assumes** *projection n P*
  **shows** ($1_m$ *n* − *P*) ∗ ($1_m$ *n* − *P*) = $1_m$ *n* − *P*
⟨*proof*⟩

## 35.3 More on the repetition code

**lemma** *hamming-weight-append*:
  *hamming-weight* (*xs* @ *ys*) = *hamming-weight xs* + *hamming-weight ys*

⟨*proof*⟩

**lemma** *hamming-weight-repetition-encode*:
*hamming-weight* (*repetition-encode v*) = *2* ∗ *hamming-weight v*
⟨*proof*⟩

**lemma** *repetition-encode-binary*:
**assumes** ∀ *x* ∈ *set v*. *x* < *2*
**shows** ∀ *x* ∈ *set* (*repetition-encode v*). *x* < *2*
⟨*proof*⟩

## 35.4 Algebra of projections; towards Days 33 and 34

Day 33 constructs the Shor recovery map from the projections onto the 22 orthogonal subspaces $V, X_1V, \dots, X_9V, Y_1V, \dots, Y_9V, Z_1V, Z_4V, Z_7V$ and proves, using $U = a_0 I + a_1 X + a_2 Y + a_3 Z$ with $\sum |a_k|^2 = 1$ for unitary $U$ (Pauli magic), that it corrects every single-qubit unitary error. Day 34 gives the general theory:

**Proposition (notes).** For $V \subseteq \mathbb{C}^n$ with projection $P$ and CPTP maps $E$ (errors) and $R$ (recovery), $R \circ E(|\psi\rangle\langle\psi|) = |\psi\rangle\langle\psi|$ for all $\psi \in V$ iff $R \circ E(PXP) = PXP$ for all $X$.

**Theorem (notes)** (1)**.** Let $E(X) = \sum_{i=1}^{r} E_i X E_i^*$ be CPTP. There is a CPTP $R$ with $R \circ E(PXP) = PXP$ for all $X$ iff $PE_i^* E_j P = \alpha_{ij} P$ for some scalars $\alpha_{ij}$.

**Theorem (notes)** (2)**.** If $R$ corrects $E$ as in Theorem 1 and $G_i \in \mathrm{span}\{E_j\}$ with $\sum G_i^* G_i = I$, then $R$ also corrects $G(X) = \sum_i G_i X G_i^*$.

Of the Knill–Laflamme results, the equivalence of Theorem 1's two readings (*recovery-states-iff-PXP*), the necessity of $PE_i^* E_j P = \alpha_{ij} P$ (*knill-laflamme-necessity*) and Theorem 2 (*knill-laflamme-closure*) are proved in *Day34-Knill-Laflamme*; the converse, the construction of $R$ from the conditions, and the equivalence itself (*knill-laflamme*) are in *Day34-KL-Sufficiency*. The elementary algebra of orthogonal projections used by the proofs follows.

**lemma** *projection-complement*:
**assumes** *projection n P*
**shows** *projection n* (*1*$_m$ *n* − *P*)
⟨*proof*⟩

Two projections are orthogonal if their products in both orders vanish.

**definition** *orthogonal-projections* ::
*nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *bool* **where**
*orthogonal-projections n P Q* ≡
*projection n P* ∧ *projection n Q* ∧
*P* ∗ *Q* = *0*$_m$ *n n* ∧ *Q* ∗ *P* = *0*$_m$ *n n*

**lemma** *orthogonal-projections-carrier-left*:
*orthogonal-projections n P Q* ⟹ *P* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *orthogonal-projections-carrier-right*:
*orthogonal-projections n P Q* ⟹ *Q* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *orthogonal-projections-symm*:
*orthogonal-projections n P Q* ⟷ *orthogonal-projections n Q P*
⟨*proof*⟩

**lemma** *orthogonal-projections-sum*:
**assumes** *opq*: *orthogonal-projections n P Q*
**shows** *projection n* $(P + Q)$
⟨*proof*⟩

**lemma** *orthogonal-projections-decomp-id*:
**assumes** *orthogonal-projections n P Q*
**shows** $P + Q + (1_m\ n - P - Q) = 1_m\ n$
⟨*proof*⟩

# 36 Extensions beyond the notes

Nothing from here on is in Paulsen's notes. These are standard quantum-information facts proved with the machinery above: rank-one projections, the bit-flip, phase-flip, bit-phase-flip and unitary-conjugation channels and their membership in UCP and CPTP, composition of channels, Kraus lists, the maximally mixed state, and the Pauli product identities.

## 36.1 Rank-one projections

**lemma** *outer-sum-singleton-hermitian*:
**fixes** $\psi$ :: *complex Matrix.vec*
**shows** $(\textit{outer-sum}\ n\ [\psi])^\dagger = \textit{outer-sum}\ n\ [\psi]$
⟨*proof*⟩

**lemma** *outer-sum-singleton-idempotent*:
**fixes** $\psi$ :: *complex Matrix.vec*
**assumes** $\psi$-*qs*: *quantum-state n* $\psi$
**shows** $\textit{outer-sum}\ n\ [\psi] * \textit{outer-sum}\ n\ [\psi] = \textit{outer-sum}\ n\ [\psi]$
⟨*proof*⟩

**lemma** *rank1-projection*:
**fixes** $\psi$ :: *complex Matrix.vec*
**assumes** $\psi$-*qs*: *quantum-state n* $\psi$
**shows** *projection n* (*outer-sum n* $[\psi]$)
⟨*proof*⟩

**lemma** *rank1-projection-acts-identity*:
**fixes** $\psi$ :: *complex Matrix.vec*
**assumes** $\psi$-*qs*: *quantum-state n* $\psi$
**shows** $\textit{outer-sum}\ n\ [\psi] *_v \psi = \psi$
⟨*proof*⟩

**lemma** *rank1-projection-trace-inner*:
**fixes** $\psi$ :: *complex Matrix.vec*
**assumes** $\psi$-*dim*: *dim-vec* $\psi = n$
**shows** *Tr* $(\textit{outer-sum}\ n\ [\psi]) = \langle \psi \mid \psi \rangle$
⟨*proof*⟩

**lemma** *rank1-projection-trace-one*:
**fixes** $\psi$ :: *complex Matrix.vec*
**assumes** $\psi$-*qs*: *quantum-state n* $\psi$
**shows** *Tr* $(\textit{outer-sum}\ n\ [\psi]) = 1$
⟨*proof*⟩

## 36.2 The bit-flip channel at $p = 0$ and $p = 1$

**lemma** *bit-flip-channel-zero*:
**assumes** $\varrho \in$ *carrier-mat 2 2*
**shows** *bit-flip-channel 0* $\varrho = \varrho$
⟨*proof*⟩

**lemma** *bit-flip-channel-one*:
**assumes** $\varrho \in$ *carrier-mat 2 2*
**shows** *bit-flip-channel 1* $\varrho =$ *Pauli-X* $*$ $\varrho *$ (*Pauli-X*$^\dagger$)
⟨*proof*⟩

## 36.3 The phase-flip channel

**definition** *phase-flip-channel* ::
*real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*phase-flip-channel p* $\varrho \equiv$
*complex-of-real* (*1* − *p*) $\cdot_m$ $\varrho$
+ *complex-of-real p* $\cdot_m$ (*Pauli-Z* $*$ $\varrho *$ (*Pauli-Z*$^\dagger$))

**lemma** *phase-flip-channel-carrier*:
**assumes** $\varrho \in$ *carrier-mat 2 2*
**shows** *phase-flip-channel p* $\varrho \in$ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *phase-flip-channel-trace-preserving*:
**fixes** *p* :: *real*
**assumes** $\varrho$*-c*: $\varrho \in$ *carrier-mat 2 2*
**shows** *Tr* (*phase-flip-channel p* $\varrho$) = *Tr* $\varrho$
⟨*proof*⟩

**lemma** *phase-flip-channel-psd*:
**fixes** *p* :: *real*
**assumes** *p-nn*: 0 ≤ *p*
**and** *p-le-1*: *p* ≤ *1*
**and** $\varrho$*-c*: $\varrho \in$ *carrier-mat 2 2*
**and** $\varrho$*-psd*: *psd* $\varrho$
**shows** *psd* (*phase-flip-channel p* $\varrho$)
⟨*proof*⟩

## 36.4 Unitary-conjugation channels

For a unitary $U \in M_n$ and $0 \le p \le 1$, $\rho \mapsto (1-p)\rho + pU\rho U^\dagger$ is trace preserving and positive; the bit-flip and phase-flip channels are the instances $U = X$, $U = Z$.

**definition** *unitary-conj-channel* ::
*complex Matrix.mat* ⇒ *real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*unitary-conj-channel U p* $\varrho \equiv$
*complex-of-real* (*1* − *p*) $\cdot_m$ $\varrho$
+ *complex-of-real p* $\cdot_m$ (*U* $*$ $\varrho *$ (*U*$^\dagger$))

**lemma** *unitary-conj-channel-carrier*:
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**and** $\varrho$*-c*: $\varrho \in$ *carrier-mat n n*
**shows** *unitary-conj-channel U p* $\varrho \in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *unitary-conj-channel-trace-preserving*:
**fixes** *p* :: *real*

**assumes** *U-c*: $U \in$ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**and** $\varrho$*-c*: $\varrho \in$ *carrier-mat n n*
**shows** *Tr* (*unitary-conj-channel U p* $\varrho$) = *Tr* $\varrho$
⟨*proof*⟩

**lemma** *unitary-conj-channel-psd*:
**fixes** *p* :: *real*
**assumes** *p-nn*: $0 \leq p$
**and** *p-le-1*: $p \leq 1$
**and** *U-c*: $U \in$ *carrier-mat n n*
**and** $\varrho$*-c*: $\varrho \in$ *carrier-mat n n*
**and** $\varrho$*-psd*: *psd* $\varrho$
**shows** *psd* (*unitary-conj-channel U p* $\varrho$)
⟨*proof*⟩

**lemma** *bit-flip-eq-unitary-conj*:
*bit-flip-channel p* $\varrho$ = *unitary-conj-channel Pauli-X p* $\varrho$
⟨*proof*⟩

**lemma** *phase-flip-eq-unitary-conj*:
*phase-flip-channel p* $\varrho$ = *unitary-conj-channel Pauli-Z p* $\varrho$
⟨*proof*⟩

## 36.5 The bit-phase-flip channel and the identity channel

**definition** *bit-phase-flip-channel* ::
*real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*bit-phase-flip-channel p* $\varrho \equiv$ *unitary-conj-channel Pauli-Y p* $\varrho$

**lemma** *bit-phase-flip-channel-carrier*:
**assumes** $\varrho \in$ *carrier-mat 2 2*
**shows** *bit-phase-flip-channel p* $\varrho \in$ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *bit-phase-flip-channel-trace-preserving*:
**fixes** *p* :: *real*
**assumes** $\varrho$*-c*: $\varrho \in$ *carrier-mat 2 2*
**shows** *Tr* (*bit-phase-flip-channel p* $\varrho$) = *Tr* $\varrho$
⟨*proof*⟩

**lemma** *bit-phase-flip-channel-psd*:
**fixes** *p* :: *real*
**assumes** *p-nn*: $0 \leq p$
**and** *p-le-1*: $p \leq 1$
**and** $\varrho$*-c*: $\varrho \in$ *carrier-mat 2 2*
**and** $\varrho$*-psd*: *psd* $\varrho$
**shows** *psd* (*bit-phase-flip-channel p* $\varrho$)
⟨*proof*⟩

**definition** *identity-channel* ::
*complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*identity-channel* $\varrho \equiv \varrho$

**lemma** *identity-channel-carrier*:

$\varrho \in$ *carrier-mat n n* $\Longrightarrow$ *identity-channel* $\varrho \in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *identity-channel-trace-preserving*:
$\varrho \in$ *carrier-mat n n* $\Longrightarrow$ *Tr* (*identity-channel* $\varrho$) = *Tr* $\varrho$
⟨*proof*⟩

**lemma** *identity-channel-psd*:
$\varrho \in$ *carrier-mat n n* $\Longrightarrow$ *psd* $\varrho \Longrightarrow$ *psd* (*identity-channel* $\varrho$)
⟨*proof*⟩

**lemma** *identity-channel-eq-unitary-conj-zero*:
**assumes** $\varrho$-*c*: $\varrho \in$ *carrier-mat n n*
**shows** *identity-channel* $\varrho$ = *unitary-conj-channel* ($1_m$ *n*) *0* $\varrho$
⟨*proof*⟩

## 36.6 Composition of channels

**lemma** *compose-trace-preserving*:
**assumes** *F-tp*: *trace-preserving n F*
**and** *G-tp*: *trace-preserving n G*
**and** *F-range*: $\forall \varrho$. $\varrho \in$ *carrier-mat n n* $\longrightarrow$ *F* $\varrho \in$ *carrier-mat n n*
**shows** *trace-preserving n* ($\lambda\varrho$. *G* (*F* $\varrho$))
⟨*proof*⟩

**lemma** *identity-channel-compose-left*:
($\lambda\varrho$. *identity-channel* (*F* $\varrho$)) = *F*
⟨*proof*⟩

**lemma** *identity-channel-compose-right*:
($\lambda\varrho$. *F* (*identity-channel* $\varrho$)) = *F*
⟨*proof*⟩

**lemma** *compose-unital-map*:
**assumes** *F-un*: *unital-map n n F*
**and** *G-un*: *unital-map n n G*
**shows** *unital-map n n* ($\lambda\varrho$. *F* (*G* $\varrho$))
⟨*proof*⟩

**lemma** *identity-channel-unital*: *unital-map n n identity-channel*
⟨*proof*⟩

**lemma** *identity-channel-trace-preserving-full*:
*trace-preserving n identity-channel*
⟨*proof*⟩

**lemma** *identity-channel-completely-positive*:
**assumes** *0* < *n*
**shows** *completely-positive n n identity-channel*
⟨*proof*⟩

## 36.7 Conjugation channels are completely positive

**lemma** *kraus-map-dim-is-cp*:
**fixes** *Ws* :: *complex Matrix.mat list*
**assumes** *Ws-c*: $\forall\, W \in set\ Ws.\ W \in carrier\text{-}mat\ d\ n$
**and** *npos*: $0 < n$
**and** *dpos*: $0 < d$
**shows** *completely-positive n d* (*kraus-map-dim d Ws*)
⟨*proof*⟩

**definition** *conjugation-channel* ::
*complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*conjugation-channel U ϱ* ≡ $U * \varrho * (U^\dagger)$

**lemma** *conjugation-channel-eq-kraus*:
**assumes** *U-c*: $U \in carrier\text{-}mat\ d\ n$
**and** *ϱ-c*: $\varrho \in carrier\text{-}mat\ n\ n$
**shows** *conjugation-channel U ϱ* = *kraus-map-dim d* [*U*] *ϱ*
⟨*proof*⟩

**lemma** *conjugation-channel-cp*:
**assumes** *U-c*: $U \in carrier\text{-}mat\ d\ n$
**and** *npos*: $0 < n$
**and** *dpos*: $0 < d$
**shows** *completely-positive n d* (*kraus-map-dim d* [*U*])
⟨*proof*⟩

**lemma** *identity-channel-UCP*:
**assumes** $0 < n$
**shows** *identity-channel* ∈ *UCP n n*
⟨*proof*⟩

**lemma** *identity-channel-CPTP*:
**assumes** $0 < n$
**shows** *identity-channel* ∈ *CPTP n n*
⟨*proof*⟩

**lemma** *map-conj-cp*:
**assumes** *U-c*: $U \in carrier\text{-}mat\ d\ n$
**and** *U-range*: $\forall\, \varrho.\ \varrho \in carrier\text{-}mat\ n\ n \longrightarrow U * \varrho * (U^\dagger) \in carrier\text{-}mat\ d\ d$
**and** *npos*: $0 < n$
**and** *dpos*: $0 < d$
**shows** *completely-positive n d* ($\lambda\varrho.\ U * \varrho * (U^\dagger)$)
⟨*proof*⟩

**lemma** *map-conj-unital*:
**fixes** *U* :: *complex Matrix.mat*
**assumes** *U-c*: $U \in carrier\text{-}mat\ n\ n$
**and** *U-unit*: *unitary U*

**shows** $U * (1_m\ n) * (U^\dagger) = 1_m\ n$
⟨*proof*⟩

## 36.8 Unitary-conjugation channels are UCP and CPTP

**lemma** *map-conj-in-UCP*:
**fixes** $U$ :: *complex Matrix.mat*
**assumes** *U-c*: $U \in$ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**and** *npos*: $0 < n$
**shows** $(\lambda\varrho.\ U * \varrho * (U^\dagger)) \in$ *UCP n n*
⟨*proof*⟩

**lemma** *unitary-conj-channel-in-UCP*:
**fixes** $U$ :: *complex Matrix.mat* **and** $p$ :: *real*
**assumes** *U-c*: $U \in$ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**and** *p-nn*: $0 \leq p$
**and** *p-le*: $p \leq 1$
**and** *npos*: $0 < n$
**shows** *unitary-conj-channel U p* $\in$ *UCP n n*
⟨*proof*⟩

**lemma** *unitary-conj-channel-in-CPTP*:
**fixes** $U$ :: *complex Matrix.mat* **and** $p$ :: *real*
**assumes** *U-c*: $U \in$ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**and** *p-nn*: $0 \leq p$
**and** *p-le*: $p \leq 1$
**and** *npos*: $0 < n$
**shows** *unitary-conj-channel U p* $\in$ *CPTP n n*
⟨*proof*⟩

## 36.9 The Pauli channels are UCP and CPTP

**lemma** *bit-flip-channel-eq*:
*bit-flip-channel p = unitary-conj-channel Pauli-X p*
⟨*proof*⟩

**lemma** *phase-flip-channel-eq*:
*phase-flip-channel p = unitary-conj-channel Pauli-Z p*
⟨*proof*⟩

**lemma** *bit-phase-flip-channel-eq*:
*bit-phase-flip-channel p = unitary-conj-channel Pauli-Y p*
⟨*proof*⟩

**lemma** *bit-flip-channel-UCP*:
**fixes** $p$ :: *real*
**assumes** *p-nn*: $0 \leq p$ **and** *p-le*: $p \leq 1$
**shows** *bit-flip-channel p* $\in$ *UCP 2 2*
⟨*proof*⟩

**lemma** *bit-flip-channel-CPTP*:
**fixes** $p$ :: *real*
**assumes** *p-nn*: $0 \leq p$ **and** *p-le*: $p \leq 1$
**shows** *bit-flip-channel p* $\in$ *CPTP 2 2*

⟨*proof*⟩

**lemma** *phase-flip-channel-UCP*:
**fixes** *p* :: *real*
**assumes** *p-nn*: *0* ≤ *p* **and** *p-le*: *p* ≤ *1*
**shows** *phase-flip-channel p* ∈ *UCP 2 2*
⟨*proof*⟩

**lemma** *phase-flip-channel-CPTP*:
**fixes** *p* :: *real*
**assumes** *p-nn*: *0* ≤ *p* **and** *p-le*: *p* ≤ *1*
**shows** *phase-flip-channel p* ∈ *CPTP 2 2*
⟨*proof*⟩

**lemma** *bit-phase-flip-channel-UCP*:
**fixes** *p* :: *real*
**assumes** *p-nn*: *0* ≤ *p* **and** *p-le*: *p* ≤ *1*
**shows** *bit-phase-flip-channel p* ∈ *UCP 2 2*
⟨*proof*⟩

**lemma** *bit-phase-flip-channel-CPTP*:
**fixes** *p* :: *real*
**assumes** *p-nn*: *0* ≤ *p* **and** *p-le*: *p* ≤ *1*
**shows** *bit-phase-flip-channel p* ∈ *CPTP 2 2*
⟨*proof*⟩

## 36.10 Pauli matrices as Kraus operators

**lemma** *Pauli-X-kraus-operators*:
*kraus-operators 2 2* [*Pauli-X*]
⟨*proof*⟩

**lemma** *Pauli-Y-kraus-operators*:
*kraus-operators 2 2* [*Pauli-Y*]
⟨*proof*⟩

**lemma** *Pauli-Z-kraus-operators*:
*kraus-operators 2 2* [*Pauli-Z*]
⟨*proof*⟩

## 36.11 Kraus maps of Kraus-operator lists are CPTP

**lemma** *kraus-is-trace-preserving*:
**fixes** *Ws* :: *complex Matrix.mat list*
**assumes** *Ws*: *kraus-operators n d Ws*
**shows** *trace-preserving n* (*kraus-map-dim d Ws*)
⟨*proof*⟩

**lemma** *kraus-is-cptp*:
**fixes** *Ws* :: *complex Matrix.mat list*
**assumes** *Ws*: *kraus-operators n d Ws*
**and** *npos*: *0* < *n*
**and** *dpos*: *0* < *d*
**shows** *kraus-map-dim d Ws* ∈ *CPTP n d*
⟨*proof*⟩

**lemma** *Pauli-X-kraus-map-CPTP*:
*kraus-map-dim 2* [*Pauli-X*] ∈ *CPTP 2 2*
⟨*proof*⟩

**lemma** *Pauli-Y-kraus-map-CPTP*:
*kraus-map-dim 2* [*Pauli-Y*] ∈ *CPTP 2 2*
⟨*proof*⟩

**lemma** *Pauli-Z-kraus-map-CPTP*:
*kraus-map-dim 2* [*Pauli-Z*] ∈ *CPTP 2 2*
⟨*proof*⟩

**lemma** *identity-kraus-operators*:
*kraus-operators n n* [$1_m$ *n*]
⟨*proof*⟩

**lemma** *identity-channel-eq-kraus-map-dim*:
**fixes** *n* :: *nat*
**assumes** *ϱ-c*: *ϱ* ∈ *carrier-mat n n*
**shows** *identity-channel ϱ* = *kraus-map-dim n* [$1_m$ *n*] *ϱ*
⟨*proof*⟩

## 36.12 Concatenation of Kraus lists

**lemma** *sum-wdag-w-append*:
**fixes** *Ws Vs* :: *complex Matrix.mat list*
**assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**and** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**shows** *sum-wdag-w n* (*Ws* @ *Vs*)
= *sum-wdag-w n Ws* + *sum-wdag-w n Vs*
⟨*proof*⟩

**lemma** *kraus-map-dim-append*:
**fixes** *Ws Vs* :: *complex Matrix.mat list*
**assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**and** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *ϱ-c*: *ϱ* ∈ *carrier-mat n n*
**shows** *kraus-map-dim d* (*Ws* @ *Vs*) *ϱ*
= *kraus-map-dim d Ws ϱ* + *kraus-map-dim d Vs ϱ*
⟨*proof*⟩

## 36.13 Fixed points of Pauli channels

**lemma** *Pauli-X-conj-commute*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *M-c*: *M* ∈ *carrier-mat 2 2*
**and** *commute*: *Pauli-X* ∗ *M* = *M* ∗ *Pauli-X*
**shows** *Pauli-X* ∗ *M* ∗ (*Pauli-X*$^\dagger$) = *M*
⟨*proof*⟩

**lemma** *bit-flip-channel-fixed-point-commute*:
**fixes** *p* :: *real*
**assumes** *M-c*: *M* ∈ *carrier-mat 2 2*
**and** *commute*: *Pauli-X* ∗ *M* = *M* ∗ *Pauli-X*

**shows** *bit-flip-channel p M = M*
⟨*proof*⟩

**lemma** *Pauli-Z-conj-commute*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *M-c*: *M* ∈ *carrier-mat 2 2*
**and** *commute*: *Pauli-Z * M = M * Pauli-Z*
**shows** *Pauli-Z * M * (Pauli-Z†) = M*
⟨*proof*⟩

**lemma** *phase-flip-channel-fixed-point-commute*:
**fixes** *p* :: *real*
**assumes** *M-c*: *M* ∈ *carrier-mat 2 2*
**and** *commute*: *Pauli-Z * M = M * Pauli-Z*
**shows** *phase-flip-channel p M = M*
⟨*proof*⟩

**lemma** *unitary-conj-channel-identity-op*:
**fixes** *p* :: *real*
**assumes** *ϱ-c*: *ϱ* ∈ *carrier-mat n n*
**shows** *unitary-conj-channel (1_m n) p ϱ = ϱ*
⟨*proof*⟩

**lemma** *unitary-conj-channel-identity-op-eq*:
**fixes** *p* :: *real*
**shows** (λ*ϱ*. *if ϱ* ∈ *carrier-mat n n*
*then unitary-conj-channel (1_m n) p ϱ else identity-channel ϱ*)
*= identity-channel*
⟨*proof*⟩

## 36.14 The maximally mixed state

**definition** *maximally-mixed-state* :: *nat* ⇒ *complex Matrix.mat* **where**
*maximally-mixed-state n* ≡ *complex-of-real (1 / real n) ·_m (1_m n)*

**lemma** *maximally-mixed-state-carrier* [*simp*]:
*maximally-mixed-state n* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *Tr-one-mat*:
*Tr (1_m n* :: *complex Matrix.mat) = of-nat n*
⟨*proof*⟩

**lemma** *maximally-mixed-state-trace*:
**assumes** *npos*: *0 < n*
**shows** *Tr (maximally-mixed-state n) = 1*
⟨*proof*⟩

**lemma** *maximally-mixed-state-psd*:
*psd (maximally-mixed-state n)*
⟨*proof*⟩

**lemma** *maximally-mixed-state-density-matrix*:
**assumes** *npos*: *0 < n*

**shows** *density-matrix n* (*maximally-mixed-state n*)
⟨*proof*⟩

## 36.15 Fixed points of unitary-conjugation channels

**lemma** *unitary-conj-commute*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**and** *M-c*: *M* ∈ *carrier-mat n n*
**and** *commute*: *U* ∗ *M* = *M* ∗ *U*
**shows** *U* ∗ *M* ∗ (*U*†) = *M*
⟨*proof*⟩

**lemma** *unitary-conj-channel-fixed-point-commute*:
**fixes** *p* :: *real*
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**and** *M-c*: *M* ∈ *carrier-mat n n*
**and** *commute*: *U* ∗ *M* = *M* ∗ *U*
**shows** *unitary-conj-channel U p M* = *M*
⟨*proof*⟩

**lemma** *maximally-mixed-state-commutes*:
**fixes** *U* :: *complex Matrix.mat*
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**shows** *U* ∗ *maximally-mixed-state n* = *maximally-mixed-state n* ∗ *U*
⟨*proof*⟩

**lemma** *unitary-conj-channel-fixes-max-mixed*:
**fixes** *p* :: *real*
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**and** *U-unit*: *unitary U*
**shows** *unitary-conj-channel U p* (*maximally-mixed-state n*)
= *maximally-mixed-state n*
⟨*proof*⟩

**lemma** *bit-flip-channel-fixes-max-mixed*:
**fixes** *p* :: *real*
**shows** *bit-flip-channel p* (*maximally-mixed-state 2*)
= *maximally-mixed-state 2*
⟨*proof*⟩

**lemma** *phase-flip-channel-fixes-max-mixed*:
**fixes** *p* :: *real*
**shows** *phase-flip-channel p* (*maximally-mixed-state 2*)
= *maximally-mixed-state 2*
⟨*proof*⟩

**lemma** *bit-phase-flip-channel-fixes-max-mixed*:
**fixes** *p* :: *real*
**shows** *bit-phase-flip-channel p* (*maximally-mixed-state 2*)
= *maximally-mixed-state 2*
⟨*proof*⟩

### 36.16 Channels preserve density matrices

**lemma** *unitary-conj-channel-preserves-density-matrix*:
  **fixes** *p* :: *real*
  **assumes** *U-c*: $U \in$ *carrier-mat n n*
    **and** *U-unit*: *unitary U*
    **and** *p-nn*: $0 \leq p$
    **and** *p-le*: $p \leq 1$
    **and** *dm*: *density-matrix n* $\varrho$
  **shows** *density-matrix n* (*unitary-conj-channel U p* $\varrho$)
⟨*proof*⟩

**lemma** *bit-flip-channel-preserves-density-matrix*:
  **fixes** *p* :: *real*
  **assumes** *p-nn*: $0 \leq p$ **and** *p-le*: $p \leq 1$
    **and** *dm*: *density-matrix 2* $\varrho$
  **shows** *density-matrix 2* (*bit-flip-channel p* $\varrho$)
  ⟨*proof*⟩

**lemma** *phase-flip-channel-preserves-density-matrix*:
  **fixes** *p* :: *real*
  **assumes** *p-nn*: $0 \leq p$ **and** *p-le*: $p \leq 1$
    **and** *dm*: *density-matrix 2* $\varrho$
  **shows** *density-matrix 2* (*phase-flip-channel p* $\varrho$)
  ⟨*proof*⟩

**lemma** *bit-phase-flip-channel-preserves-density-matrix*:
  **fixes** *p* :: *real*
  **assumes** *p-nn*: $0 \leq p$ **and** *p-le*: $p \leq 1$
    **and** *dm*: *density-matrix 2* $\varrho$
  **shows** *density-matrix 2* (*bit-phase-flip-channel p* $\varrho$)
  ⟨*proof*⟩

### 36.17 Pauli product identities

With the Day 32 convention for $Y$: $XY = -iZ$, $YZ = -iX$, $ZX = -iY$.

**lemma** *Pauli-X-times-Pauli-Y*: *Pauli-X* $*$ *Pauli-Y* $= -\mathrm{i} \cdot_m$ *Pauli-Z*
⟨*proof*⟩

**lemma** *Pauli-Y-times-Pauli-Z*: *Pauli-Y* $*$ *Pauli-Z* $= -\mathrm{i} \cdot_m$ *Pauli-X*
⟨*proof*⟩

**lemma** *Pauli-Z-times-Pauli-X*: *Pauli-Z* $*$ *Pauli-X* $= -\mathrm{i} \cdot_m$ *Pauli-Y*
⟨*proof*⟩

**end**
**theory** *QC-Prelims*
  **imports** *Day23-ChoiExtremeUCP*
**begin**

# 37 Small matrix and vector lemmas shared by the gap-closure theories

Index case splits, trace facts, entrywise vector sums, rank-one identities and rectangular *foldr* sums. Nothing here is specific to a lecture; the theories *Day12-Counting-Probability*, *Day18-Positive-Maps*,

*Day22-Kraus-Independence*, *Day28-Horodecki-Peres*, *Day30-BitFlip-Code* and *Day32-Pauli-Expansion* build on it.

**lemma** *cnj-mult-self-cmod*: *cnj z * z = complex-of-real ((cmod z)^2)*
⟨*proof*⟩

**lemma** *Re-cnj-mult-self*: *Re (cnj z * z) = (cmod z)^2*
⟨*proof*⟩

**lemma** *idx4*: *k < (4::nat) ⟹ k = 0 ∨ k = 1 ∨ k = 2 ∨ k = 3*
⟨*proof*⟩

**lemma** *mult-index-lt*:
**fixes** *i j n m* :: *nat*
**assumes** *i < n* **and** *j < m*
**shows** *i * m + j < n * m*
⟨*proof*⟩

**lemma** *Tr-hermitian-Im-zero*:
**assumes** *Pc*: *P ∈ carrier-mat n n* **and** *herm*: *hermitian P*
**shows** *Im (Tr P) = 0*
⟨*proof*⟩

**lemma** *Tr-hermitian-real*:
**assumes** *Pc*: *P ∈ carrier-mat n n* **and** *herm*: *hermitian P*
**shows** *Tr P = complex-of-real (Re (Tr P))*
⟨*proof*⟩

**lemma** *Tr-minus*:
**fixes** *x y* :: *complex Matrix.mat*
**assumes** *dim-row x = dim-row y* **and** *dim-col x = dim-col y*
**shows** *Tr (x − y) = Tr x − Tr y*
⟨*proof*⟩

A positive semidefinite matrix of trace zero is zero (the notes: "given a non-zero $P \geq 0$, $\mathrm{Tr}(P) \neq 0$").

**lemma** *psd-Tr-zero-imp-zero*:
**assumes** *Pc*: *P ∈ carrier-mat n n* **and** *pos*: *psd P* **and** *tr0*: *Tr P = 0*
**shows** *P = 0_m n n*
⟨*proof*⟩

**lemma** *smult-smult-mat*:
**fixes** *A* :: *complex Matrix.mat*
**shows** *a ·_m (b ·_m A) = (a * b) ·_m A*
⟨*proof*⟩

**lemma** *zero-smult-mat-carrier*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *A ∈ carrier-mat m k*
**shows** *0 ·_m A = 0_m m k*
⟨*proof*⟩

Finite sums of vectors, entrywise.

**definition** *vec-sum* :: *nat ⇒ 'c set ⇒ ('c ⇒ complex Matrix.vec) ⇒ complex Matrix.vec* **where**
*vec-sum n Js w = Matrix.vec n (λi. sum (λJ. w J $ i) Js)*

**lemma** *vec-sum-dim* [*simp*]: *dim-vec (vec-sum n Js w) = n*
⟨*proof*⟩

**lemma** *vec-sum-index* [*simp*]: $i < n \Longrightarrow$ *vec-sum n Js w* \$ *i* = *sum* ($\lambda J.$ *w J* \$ *i*) *Js*
⟨*proof*⟩

**lemma** *vec-sum-cong*:
**assumes** $\bigwedge J.$ $J \in Js \Longrightarrow w\ J = w'\ J$
**shows** *vec-sum n Js w* = *vec-sum n Js w'*
⟨*proof*⟩

**lemma** *mult-mat-vec-vec-sum*:
**assumes** *Ac*: $A \in$ *carrier-mat m n* **and** *fin*: *finite Js*
**and** *wd*: $\bigwedge J.$ $J \in Js \Longrightarrow$ *dim-vec* (*w J*) = *n*
**shows** $A *_v$ *vec-sum n Js w* = *vec-sum m Js* ($\lambda J.$ $A *_v$ *w J*)
⟨*proof*⟩

**lemma** *inner-prod-vec-sum-right*:
**assumes** *ud*: *dim-vec u* = *n*
**and** *wd*: $\bigwedge J.$ $J \in Js \Longrightarrow$ *dim-vec* (*w J*) = *n*
**shows** ⟨*u* | *vec-sum n Js w*⟩ = *sum* ($\lambda J.$ ⟨*u* | *w J*⟩) *Js*
⟨*proof*⟩

**lemma** *mult-mat-vec-smult-vec*:
**fixes** *A* :: *complex Matrix.mat* **and** *v* :: *complex Matrix.vec*
**assumes** *Ac*: $A \in$ *carrier-mat m n* **and** *vd*: *dim-vec v* = *n*
**shows** $A *_v$ ($c \cdot_v$ *v*) = $c \cdot_v$ ($A *_v$ *v*)
⟨*proof*⟩

**lemma** *inner-prod-unit-vec-right*:
**assumes** *ud*: *dim-vec u* = *n* **and** *j*: $j < n$
**shows** ⟨*u* | *unit-vec n j*⟩ = *cnj* (*u* \$ *j*)
⟨*proof*⟩

**lemma** *outer-prod-self-psd*:
**assumes** *dim-vec v* = *n*
**shows** *psd* (*v* ⊘ *v*)
⟨*proof*⟩

Conjugating a rank-one matrix: $W|\varphi\rangle\langle\varphi|W^* = |W\varphi\rangle\langle W\varphi|$.

**lemma** *conj-outer-prod*:
**fixes** *W* :: *complex Matrix.mat* **and** $\varphi$ :: *complex Matrix.vec*
**assumes** *Wc*: $W \in$ *carrier-mat m n* **and** $\varphi$*d*: *dim-vec* $\varphi$ = *n*
**shows** $W * (\varphi \oslash \varphi) * (W^\dagger) = (W *_v \varphi) \oslash (W *_v \varphi)$
⟨*proof*⟩

**lemma** *outer-prod-zero-vec*: ($0_v$ *n* :: *complex Matrix.vec*) ⊘ ($0_v$ *n*) = $0_m$ *n n*
⟨*proof*⟩

**lemma** *idx8*: $k <$ (*8*::*nat*) $\Longrightarrow$ $k = 0 \vee k = 1 \vee k = 2 \vee k = 3 \vee k = 4 \vee k = 5 \vee k = 6 \vee k = 7$
⟨*proof*⟩

**lemma** *s8*: {*0*..<(*8*::*nat*)} = {*0*,*1*,*2*,*3*,*4*,*5*,*6*,*7*}* ⟨*proof*⟩

**lemma** *dagger-mult-diag-inner*:
**fixes** *Y* :: *complex Matrix.mat*
**assumes** *Yc*: $Y \in$ *carrier-mat p q* **and** *j*: $j < q$
**shows** (($Y^\dagger$) $*$ *Y*) \$\$ (*j*,*j*) = ⟨*Matrix.col Y j* | *Matrix.col Y j*⟩
⟨*proof*⟩

**lemma** *foldr-add-mat-carrier-rect*:
**assumes** $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ p\ q$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ p\ q) \in carrier\text{-}mat\ p\ q$
⟨*proof*⟩

**lemma** *foldr-add-mat-index-rect*:
**assumes** $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ p\ q$ **and** $i < p$ **and** $j < q$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ p\ q)\ \$\$\ (i,j) = (\sum x \leftarrow xs.\ f\ x\ \$\$\ (i,j))$
⟨*proof*⟩

**end**
**theory** *Day12-Counting-Probability*
**imports** *QC-Prelims*
**begin**

# 38 Day 12: $H^{\otimes N}$ on $|0\cdots0\rangle$ and the counting probability (D12-1, D12-4)

$N$-fold tensor powers (last factor least significant, matching $(\otimes)$).

**primrec** *tensor-pow-mat* :: *complex Matrix.mat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*tensor-pow-mat A 0 = 1*$_m$ *1*
| *tensor-pow-mat A (Suc k) = Tensor.tensor-mat (tensor-pow-mat A k) A*

**primrec** *tensor-pow-vec* :: *complex Matrix.vec* ⇒ *nat* ⇒ *complex Matrix.vec* **where**
*tensor-pow-vec v 0 = Matrix.vec 1 (λ-. 1)*
| *tensor-pow-vec v (Suc k) = tensor-vec (tensor-pow-vec v k) v*

**lemma** *tensor-pow-mat-carrier*:
**assumes** *A ∈ carrier-mat 2 2*
**shows** *tensor-pow-mat A k ∈ carrier-mat (2^k) (2^k)*
⟨*proof*⟩

**lemma** *tensor-pow-vec-dim*:
**assumes** *dim-vec v = 2*
**shows** *dim-vec (tensor-pow-vec v k) = 2^k*
⟨*proof*⟩

**lemma** *tensor-pow-mat-vec*:
**assumes** *Ac*: *A ∈ carrier-mat 2 2* **and** *vd*: *dim-vec v = 2*
**shows** *tensor-pow-mat A k *$_v$ *tensor-pow-vec v k = tensor-pow-vec (A *$_v$ *v) k*
⟨*proof*⟩

**lemma** *tensor-pow-vec-const*:
*tensor-pow-vec (Matrix.vec 2 (λ-. c)) k = Matrix.vec (2^k) (λ-. c^k)*
⟨*proof*⟩

**definition** *uniform-vec* :: *nat* ⇒ *complex Matrix.vec* **where**
*uniform-vec N = Matrix.vec (2^N) (λ-. complex-of-real (1 / sqrt 2) ^ N)*

**lemma** *uniform-vec-dim* [*simp*]: *dim-vec (uniform-vec N) = 2^N*
⟨*proof*⟩

**lemma** *hadamard-e0-const*: *Hadamard *$_v$ *unit-vec 2 0 = Matrix.vec 2 (λ-. complex-of-real (1 / sqrt 2))*
⟨*proof*⟩

**theorem** *hadamard-uniform-N*:
*tensor-pow-mat Hadamard N* $*_v$ *tensor-pow-vec* (*unit-vec 2 0*) *N* = *uniform-vec N*
⟨*proof*⟩

**lemma** *inner-prod-unit-vec-2*:
**assumes** *k* < *2*
**shows** ⟨*unit-vec 2 0* | *unit-vec 2 k*⟩ = (*if k* = *0 then 1 else 0*)
⟨*proof*⟩

**lemma** *inner-prod-uniform-unit*:
**assumes** *J* < *2^N*
**shows** ⟨*uniform-vec N* | *unit-vec* (*2^N*) *J*⟩ = *complex-of-real* (*1 / sqrt 2*) ^ *N*
⟨*proof*⟩

**lemma** *half-sqrt2-sq*: (*1 / sqrt 2* :: *real*) * (*1 / sqrt 2*) = *1 / 2*
⟨*proof*⟩

**lemma** *uniform-coeff-sq*:
*complex-of-real* (*1 / sqrt 2*) ^ *N* * *complex-of-real* (*1 / sqrt 2*) ^ *N* = *complex-of-real* ((*1/2*) ^ *N*)
⟨*proof*⟩

**lemma** *uniform-vec-self*: ⟨*uniform-vec N* | *uniform-vec N*⟩ = *1*
⟨*proof*⟩

**lemma** *uniform-tensor-e0-as-sum*:
*tensor-vec* (*uniform-vec N*) (*unit-vec 2 0*) =
*vec-sum* (*2^(N+1)*) {*J*. *J* < *2^N*}
(λ*J*. *complex-of-real* (*1 / sqrt 2*) ^ *N* ·$_v$ *tensor-vec* (*unit-vec* (*2^N*) *J*) (*unit-vec 2 0*))
⟨*proof*⟩

**lemma** *uniform-e0-state*: *quantum-state* (*2^(N+1)*) (*tensor-vec* (*uniform-vec N*) (*unit-vec 2 0*))
⟨*proof*⟩

The notes' computation: with $x = H^{\otimes N}|0\cdots0\rangle$ and $\gamma = x \otimes e_0$, $p_0(U_f(x \otimes e_0)) = (M/2^N)^2$ where $M = \#\{J : f(J) = 0\}$.

**theorem** *counting-probability*:
**fixes** *f* :: *nat* ⇒ *nat* **and** *U* :: *complex Matrix.mat*
**assumes** *f-range*: ∀ *J* < *2^N*. *f J* < *2*
**and** *Uc*: *U* ∈ *carrier-mat* (*2^(N+1)*) (*2^(N+1)*)
**and** *orac*: ∀ *J* < *2^N*. ∀ *i* < *2*.
*U* $*_v$ (*tensor-vec* (*unit-vec* (*2^N*) *J*) (*unit-vec 2 i*)) =
*tensor-vec* (*unit-vec* (*2^N*) *J*) (*unit-vec 2* ((*i* + *f J*) *mod 2*))
**shows** *meas-prob* (*tensor-vec* (*uniform-vec N*) (*unit-vec 2 0*) ⊘ *tensor-vec* (*uniform-vec N*) (*unit-vec 2 0*))
(*U* $*_v$ *tensor-vec* (*uniform-vec N*) (*unit-vec 2 0*))
= (*real* (*count-zeros f N*) / *2^N*)$^2$
⟨*proof*⟩

**end**
**theory** *Day16-Block-Form*
**imports** *Day25-Arveson-Correspondence Day02-11-Supplement*
**begin**

# 39 Day 16: $H_A \otimes H_B \cong H_A^{(p)}$ and $L(H_A \otimes H_B) \cong M_p(L(H_A))$ (D16-2)

The notes identify $H_A \otimes H_B$ with $p$ copies of $H_A$ through $\sum_i h_i \otimes f_i \leftrightarrow (h_1, \ldots, h_p)$, with $\|v\|^2 = \sum_i \|h_i\|^2$, and an operator $X$ with its block matrix $(X_{ij})$ over $L(H_A)$, acting by $(h_j) \mapsto$

$(\sum_j X_{ij} h_j)$, so that $X = \sum_{i,j} X_{ij} \otimes E_{ij}$, $\mathrm{tr}_B(X) = X_{11} + \cdots + X_{pp}$ and $\mathrm{tr}_A(X) = (\mathrm{Tr}\, X_{ij})$. Here $f_i = e_i$, and indices are A-major as everywhere in the development.

With $H_A = \mathbb{C}^m$, $H_B = \mathbb{C}^p$ and A-major indices $ap + i$, the coordinates of $v$ along $e_i$ form $h_i = (v_{ap+i})_a \in H_A$, and an operator $X$ has the blocks $X_{ij} = (X_{ap+i,\, bp+j})_{a,b} \in L(H_A)$.

**definition** *vcomp* :: *nat* ⇒ *nat* ⇒ *complex Matrix.vec* ⇒ *nat* ⇒ *complex Matrix.vec* **where**
*vcomp m p v i* = *Matrix.vec m* (λ*a*. *v* $ (*a* ∗ *p* + *i*))

**definition** *bblock* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*bblock m p X i j* = *Matrix.mat m m* (λ(*a*, *b*). *X* $$ (*a* ∗ *p* + *i*, *b* ∗ *p* + *j*))

**lemma** *bblock-carrier* [*simp*]: *bblock m p X i j* ∈ *carrier-mat m m*
⟨*proof*⟩

**lemma** *bblock-dims* [*simp*]: *dim-row* (*bblock m p X i j*) = *m dim-col* (*bblock m p X i j*) = *m*
⟨*proof*⟩

**lemma** *vcomp-dim* [*simp*]: *dim-vec* (*vcomp m p v i*) = *m*
⟨*proof*⟩

**lemma** *sum-mp-reindex*:
**fixes** *f* :: *nat* ⇒ *complex*
**shows** (∑ *r*<*m* ∗ *p*. *f r*) = (∑ *b*<*m*. ∑ *j*<*p*. *f* (*b* ∗ *p* + *j*))
⟨*proof*⟩

$H_A \otimes H_B \cong H_A^{(p)}$ isometrically: $\langle v|w\rangle = \sum_i \langle h_i|k_i\rangle$.

**theorem** *vcomp-inner-prod*:
**assumes** *v*: *dim-vec v* = *m* ∗ *p* **and** *w*: *dim-vec w* = *m* ∗ *p*
**shows** *Quantum.inner-prod v w* = (∑ *i*<*p*. *Quantum.inner-prod* (*vcomp m p v i*) (*vcomp m p w i*))
⟨*proof*⟩

$v = \sum_i h_i \otimes e_i$.

**theorem** *vcomp-expansion*:
**assumes** *v*: *dim-vec v* = *m* ∗ *p*
**shows** *v* = *tensor-combination* (*m* ∗ *p*) (*map* (*vcomp m p v*) [*0*..<*p*]) (*map* (*unit-vec p*) [*0*..<*p*])
⟨*proof*⟩

The block action: the $i$-th coordinate of $Xv$ is $\sum_j X_{ij} h_j$.

**theorem** *bblock-action*:
**assumes** *X*: *X* ∈ *carrier-mat* (*m* ∗ *p*) (*m* ∗ *p*) **and** *v*: *dim-vec v* = *m* ∗ *p* **and** *i*: *i* < *p*
**shows** *vcomp m p* (*X* ∗*v v*) *i* = *Matrix.vec m* (λ*a*. ∑ *j*<*p*. (*bblock m p X i j* ∗*v vcomp m p v j*) $ *a*)
⟨*proof*⟩

$L(H_A \otimes H_B) \cong M_p(L(H_A))$: $X = \sum_{i,j} X_{ij} \otimes E_{ij}$, and the blocks are unique.

**lemma** *double-delta*:
**fixes** *c* :: *complex* **and** *a b p q* :: *nat*
**assumes** *a* < *p b* < *q*
**shows** (∑ *i*<*p*. ∑ *j*<*q*. *if i* = *a* ∧ *j* = *b then c else 0*) = *c*
⟨*proof*⟩

**lemma** *mat-unit-tensor-index*:
**assumes** *A*: *A* ∈ *carrier-mat m m* **and** *r*: *r* < *m* ∗ *p* **and** *s*: *s* < *m* ∗ *p* **and** *i*: *i* < *p* **and** *j*: *j* < *p*
**shows** (*A* ⨂ *E*⟦*i*,*j* ; *p*,*p*⟧) $$ (*r*, *s*) = (*if r mod p* = *i* ∧ *s mod p* = *j then A* $$ (*r div p*, *s div p*) *else 0*)
⟨*proof*⟩

**theorem** *tensor-block-decomposition*:
**assumes** *X*: $X \in$ *carrier-mat* $(m * p)$ $(m * p)$
**shows** $X =$ *msum* $(m * p)$ $(m * p)$ $(\lambda(i, j).$ *bblock m p X i j* $\bigotimes$ $E[\![i,j\ ;\ p,p]\!])$ (*List.product* [*0..<p*] [*0..<p*])
⟨*proof*⟩

**theorem** *tensor-block-decomposition-unique*:
**assumes** *Y*: $\bigwedge i\ j.$ *Y i j* $\in$ *carrier-mat m m*
**and** *X*: $X =$ *msum* $(m * p)$ $(m * p)$ $(\lambda(i, j).$ *Y i j* $\bigotimes$ $E[\![i,j\ ;\ p,p]\!])$ (*List.product* [*0..<p*] [*0..<p*])
**and** *i*: $i < p$ **and** *j*: $j < p$
**shows** *Y i j* = *bblock m p X i j*
⟨*proof*⟩

The partial traces in block form (the definitions, restated): $\mathrm{tr}_B(X) = \sum_i X_{ii}$, $\mathrm{tr}_A(X) = (\mathrm{Tr}\, X_{ij})_{ij}$.

**lemma** *partial-trace-B-blocks*:
*partial-trace-B m p X* = *msum m m* $(\lambda i.$ *bblock m p X i i*$)$ [*0..<p*]
⟨*proof*⟩

**lemma** *partial-trace-A-blocks*:
*partial-trace-A m p X* = *Matrix.mat p p* $(\lambda(i, j).$ *Tr* (*bblock m p X i j*))
⟨*proof*⟩

**end**
**theory** *Day17-Environment-Model*
**imports** *Day22-Minimal-Kraus*
**begin**

Day 17, the environment model of noise: a state $\rho$ of lab $A$ meets an environment in the state $|\varphi\rangle$, the pair evolves by a unitary $U$ on $\mathbb{C}^m \otimes \mathbb{C}^p$, and lab $A$ sees $\mathrm{tr}_E(U(\rho \otimes |\varphi\rangle\langle\varphi|)U^*) = \sum_i W_i \rho W_i^*$ with $\sum_i W_i^* W_i = I$. The index of $\mathbb{C}^m \otimes \mathbb{C}^p$ is $a\,p + i$ (lab index $a$, environment index $i$), and $W_i$ is the matrix with entries $\sum_k U_{(a,i),(c,k)} \varphi_k$, i.e. the notes' $W_i = \sum_k U_{ik} \alpha_k$ for the blocks $U_{ik}$ of $U$.

**definition** *env-kraus* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*env-kraus m p U φ i* = *Matrix.mat m m* $(\lambda(a,c).\ \sum k{<}p.$ *U* $\$\$$ $(a * p + i,\ c * p + k) * \varphi\ \$\ k)$

**lemma** *env-kraus-carrier* [*simp*]: *env-kraus m p U φ i* $\in$ *carrier-mat m m*
⟨*proof*⟩

**lemma** *env-kraus-dims* [*simp*]: *dim-row* (*env-kraus m p U φ i*) = *m dim-col* (*env-kraus m p U φ i*) = *m*
⟨*proof*⟩

**lemma** *env-kraus-index*: $a < m \Longrightarrow c < m \Longrightarrow$ *env-kraus m p U φ i* $\$\$$ $(a,c) = (\sum k{<}p.$ *U* $\$\$$ $(a * p + i,\ c * p + k) * \varphi\ \$\ k)$
⟨*proof*⟩

**lemma** *sum-split-mp*:
**fixes** *f* :: *nat* ⇒ *'a*::*comm-monoid-add*
**shows** $(\sum x < m * p.\ f\ x) = (\sum c{<}m.\ \sum k{<}p.\ f\ (c * p + k))$
⟨*proof*⟩

**lemma** *idx-mp*:
**fixes** *c k* :: *nat*
**assumes** *c*: $c < m$ **and** *k*: $k < p$
**shows** $c * p + k < m * p$ $(c * p + k)$ *div* $p = c$ $(c * p + k)$ *mod* $p = k$
⟨*proof*⟩

**lemma** *tensor-outer-index*:
**fixes** $\varrho$ :: *complex Matrix.mat* **and** $\varphi$ :: *complex Matrix.vec*
**assumes** $\varrho$: $\varrho \in$ *carrier-mat m m* **and** $\varphi$: *dim-vec* $\varphi = p$
**and** $c$: $c < m$ **and** $d$: $d < m$ **and** $k$: $k < p$ **and** $l$: $l < p$
**shows** *Tensor.tensor-mat* $\varrho$ ($\varphi \oslash \varphi$) $$ ($c * p + k$, $d * p + l$) = $\varrho$ $$ ($c$,$d$) * ($\varphi$ $ $k$ * *cnj* ($\varphi$ $ $l$))
⟨*proof*⟩

The entries of $U(\rho \otimes |\varphi\rangle\langle\varphi|)U^*$.

**lemma** *env-entry*:
**fixes** $U$ $\varrho$ :: *complex Matrix.mat* **and** $\varphi$ :: *complex Matrix.vec*
**assumes** $U$: $U \in$ *carrier-mat* ($m * p$) ($m * p$) **and** $\varrho$: $\varrho \in$ *carrier-mat m m* **and** $\varphi$: *dim-vec* $\varphi = p$
**and** $x$: $x < m * p$ **and** $y$: $y < m * p$
**shows** ($U$ * *Tensor.tensor-mat* $\varrho$ ($\varphi \oslash \varphi$) * ($U^\dagger$)) $$ ($x$,$y$)
= ($\sum d$<$m$. $\sum c$<$m$. ($\sum k$<$p$. $U$ $$ ($x$, $c * p + k$) * $\varphi$ $ $k$) * $\varrho$ $$ ($c$,$d$)
* *cnj* ($\sum l$<$p$. $U$ $$ ($y$, $d * p + l$) * $\varphi$ $ $l$))
⟨*proof*⟩

**theorem** *environment-model*:
**fixes** $U$ $\varrho$ :: *complex Matrix.mat* **and** $\varphi$ :: *complex Matrix.vec*
**assumes** $U$: $U \in$ *carrier-mat* ($m * p$) ($m * p$) **and** *Uu*: *unitary U*
**and** $\varphi$: *dim-vec* $\varphi = p$ **and** $\varphi 1$: $\langle\varphi|\varphi\rangle = 1$ **and** $\varrho$: $\varrho \in$ *carrier-mat m m*
**defines** *Ws* ≡ *map* (*env-kraus m p U* $\varphi$) [*0*..<*p*]
**shows** *partial-trace-B m p* (*U* * *Tensor.tensor-mat* $\varrho$ ($\varphi \oslash \varphi$) * ($U^\dagger$)) = *kraus-map-dim m Ws* $\varrho$
**and** *kraus-operators m m Ws*
⟨*proof*⟩

**end**
**theory** *Day18-Positive-Maps*
**imports** *QC-Prelims*
**begin**

# 40 Days 17–18: densities to densities iff positive and trace-preserving (D17-4, D18-2)

**theorem** *density-preserving-iff-positive-tp*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: ⋀*A B*. *A* ∈ *carrier-mat n n* ⟹ *B* ∈ *carrier-mat n n* ⟹
Φ (*A* + *B*) = Φ *A* + Φ *B*
**and** *smult*: ⋀*c A*. *A* ∈ *carrier-mat n n* ⟹ Φ (*c* ·$_m$ *A*) = *c* ·$_m$ Φ *A*
**and** *range*: ∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* ∈ *carrier-mat d d*
**and** *npos*: *0* < *n*
**shows** (∀ $\varrho$. *density-matrix n* $\varrho$ ⟶ *density-matrix d* (Φ $\varrho$)) ⟷
(*positive-map n d* Φ ∧ *trace-preserving n* Φ)
⟨*proof*⟩

# 41 Day 18: the transpose is positive but not 2-positive (D18-4)

**lemma** *transpose-positive-map*: *positive-map n n* (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*)
⟨*proof*⟩

The notes' witness: $X = (E_{ij}) \in M_2(M_2)$ is positive (it is $|w\rangle\langle w|$ for $w = e_0 \otimes e_0 + e_1 \otimes e_1$) while $\Phi^{(2)}(X)$ has the eigenvector $(0, 1, -1, 0)$ with eigenvalue $-1$ (the notes compute $\det \Phi^{(2)}(X) = -1$ instead).

**lemma** *choi-seed-2-outer*: *choi-seed 2 = outer-sum 4 [Matrix.vec 4 (λi. if i = 0 ∨ i = 3 then 1 else 0)]*
⟨*proof*⟩

**lemma** *choi-seed-2-psd*: *psd (choi-seed 2)*
⟨*proof*⟩

**lemma** *transpose-not-2-positive*:
¬ *p-positive 2 2 2 (transpose-mat :: complex Matrix.mat ⇒ complex Matrix.mat)*
⟨*proof*⟩

**corollary** *transpose-not-completely-positive*:
¬ *completely-positive 2 2 (transpose-mat :: complex Matrix.mat ⇒ complex Matrix.mat)*
⟨*proof*⟩

**end**
**theory** *Day19-Choi-TP-Unital*
**imports** *Day25-Arveson-Correspondence*
**begin**

Day 19, notes (4) and (5) after Choi's theorem: write $P_\Phi = (R_{ij})$ with $R_{ij} = \Phi(E_{ij}) \in M_d$. Then $\Phi$ is trace preserving iff $\operatorname{Tr} R_{ij} = \delta_{ij}$, and unital iff $\sum_i R_{ii} = I_d$; with *choi-theorem* this characterises CPTP and UCP maps by their Choi matrix. Linearity on $M_n$ is *lin-on n d (carrier-mat n n) Φ*.

**lemma** *mat-unit-carrier′* [*simp*]: *E⟦i,j ; n,n⟧ ∈ carrier-mat n n*
⟨*proof*⟩

**lemma** *choi-block*:
**assumes** *range*: ∀ *A*. *A ∈ carrier-mat n n ⟶ Φ A ∈ carrier-mat d d* **and** *i*: *i < n* **and** *j*: *j < n*
**shows** *block-of-mat d (choi-matrix n d Φ) i j = Φ (E⟦i,j ; n,n⟧)*
⟨*proof*⟩

**lemma** *msum-cong*: *(⋀x. x ∈ set xs ⟹ g x = h x) ⟹ msum r c g xs = msum r c h xs*
⟨*proof*⟩

**lemma** *Tr-msum*:
**assumes** ∀ *x ∈ set xs. g x ∈ carrier-mat d d*
**shows** *Tr (msum d d g xs) = (∑ x←xs. Tr (g x))*
⟨*proof*⟩

**lemma** *unit-expansion*:
**assumes** *A*: *A ∈ carrier-mat n n*
**shows** *A = msum n n (λ(i,j). A $$ (i,j) ·m E⟦i,j ; n,n⟧) (List.product [0..<n] [0..<n])*
⟨*proof*⟩

**lemma** *id-expansion*: *1m n = msum n n (λi. E⟦i,i ; n,n⟧) [0..<n]*
⟨*proof*⟩

**lemma** *lin-expansion*:
**assumes** *L*: *lin-on n d (carrier-mat n n) Φ* **and** *A*: *A ∈ carrier-mat n n*
**shows** *Φ A = msum d d (λ(i,j). A $$ (i,j) ·m Φ (E⟦i,j ; n,n⟧)) (List.product [0..<n] [0..<n])*
⟨*proof*⟩

D19-3, note (4): trace preservation in Choi-matrix form.

**theorem** *choi-trace-preserving-iff*:
**assumes** *L*: *lin-on n d (carrier-mat n n) Φ*
**shows** *trace-preserving n Φ ⟷*

$(\forall\, i < n.\ \forall\, j < n.\ Tr\ (block\text{-}of\text{-}mat\ d\ (choi\text{-}matrix\ n\ d\ \Phi)\ i\ j) = (if\ i = j\ then\ 1\ else\ 0))$
⟨*proof*⟩

D19-3, note (5): unitality in Choi-matrix form, $\sum_i R_{ii} = I_d$.

**theorem** *choi-unital-iff*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ
**shows** *unital-map n d* Φ ⟷ *msum d d* (λ*i*. *block-of-mat d* (*choi-matrix n d* Φ) *i i*) [*0*..<*n*] = $1_m$ *d*
⟨*proof*⟩

**lemma** *lin-on-choi-hyps*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ
**shows** ⋀*A B*. *A* ∈ *carrier-mat n n* ⟹ *B* ∈ *carrier-mat n n* ⟹ Φ (*A* + *B*) = Φ *A* + Φ *B*
**and** ⋀*c A*. *A* ∈ *carrier-mat n n* ⟹ Φ (*c* ·$_m$ *A*) = *c* ·$_m$ Φ *A*
**and** ∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* ∈ *carrier-mat d d*
⟨*proof*⟩

Notes (4) and (5) with Choi's theorem: CPTP iff $P_\Phi \geq 0$ and $\mathrm{Tr}\, R_{ij} = \delta_{ij}$; UCP iff $P_\Phi \geq 0$ and $\sum_i R_{ii} = I_d$.

**corollary** *choi-cptp-iff*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *n*: *n* > *0* **and** *d*: *d* > *0*
**shows** *completely-positive n d* Φ ∧ *trace-preserving n* Φ ⟷
*psd* (*choi-matrix n d* Φ) ∧
(∀ *i* < *n*. ∀ *j* < *n*. *Tr* (*block-of-mat d* (*choi-matrix n d* Φ) *i j*) = (*if i* = *j then 1 else 0*))
⟨*proof*⟩

**corollary** *choi-ucp-iff*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *n*: *n* > *0* **and** *d*: *d* > *0*
**shows** *completely-positive n d* Φ ∧ *unital-map n d* Φ ⟷
*psd* (*choi-matrix n d* Φ) ∧ *msum d d* (λ*i*. *block-of-mat d* (*choi-matrix n d* Φ) *i i*) [*0*..<*n*] = $1_m$ *d*
⟨*proof*⟩

**end**
**theory** *Day20-Choi-Rank*
**imports** *Day34-Knill-Laflamme Day22-Kraus-Independence*
**begin**

# 42 Day 20: Choi's rank theorem $\mathrm{cr}(\Phi) = \mathrm{rank}\, P_\Phi$ (D20-4)

Rank is JNF's *vec-space.rank*, the dimension of the column space.

**abbreviation** *mat-rank* :: *nat* ⇒ *complex Matrix.mat* ⇒ *nat* **where**
*mat-rank m A* ≡ *vec-space.rank m A*

## 42.1 $\mathrm{rank}(\sum_{l<L} w_l w_l^*) \leq L$

**lemma** *rank-outer-prod-le-1*:
**assumes** *vd*: *dim-vec v* = *m*
**shows** *mat-rank m* (*v* ⊘ *v*) ≤ *1*
⟨*proof*⟩

**lemma** *rank-foldr-outer-le*:
**assumes** *vsd*: ∀ *v* ∈ *set vs*. *dim-vec v* = *m*
**shows** *mat-rank m* (*foldr* (λ*v acc*. *v* ⊘ *v* + *acc*) *vs* ($0_m$ *m m*)) ≤ *length vs*
⟨*proof*⟩

**lemma** *rank-outer-sum-le*:
**assumes** *vsd*: ∀ *v* ∈ *set vs*. *dim-vec v* = *m*

**shows** *mat-rank m* (*outer-sum m vs*) ≤ *length vs*
⟨*proof*⟩

### 42.2 Independent vectors inside the column space bound the rank from below

**lemma** *mult-mat-vec-in-span*:
**assumes** *Ac*: *A* ∈ *carrier-mat m k* **and** *xc*: *x* ∈ *carrier-vec k*
**shows** *A* ∗$_v$ *x* ∈ *module.span class-ring* (*module-vec TYPE*(*complex*) *m*) (*set* (*cols A*))
⟨*proof*⟩

**lemma** *rank-ge-indep-in-colspace*:
**assumes** *Ac*: *A* ∈ *carrier-mat m k* **and** *li*: *lin-indep-vecs m ws*
**and** *insp*: ⋀*w*. *w* ∈ *set ws* ⟹ *w* ∈ *module.span class-ring* (*module-vec TYPE*(*complex*) *m*) (*set* (*cols A*))
**shows** *length ws* ≤ *mat-rank m A*
⟨*proof*⟩

### 42.3 The theorem

**lemma** *choi-rank-attained*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: ⋀*A B*. *A* ∈ *carrier-mat n n* ⟹ *B* ∈ *carrier-mat n n* ⟹ Φ (*A* + *B*) = Φ *A* + Φ *B*
**and** *smult*: ⋀*c A*. *A* ∈ *carrier-mat n n* ⟹ Φ (*c* ·$_m$ *A*) = *c* ·$_m$ Φ *A*
**and** *range*: ∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* ∈ *carrier-mat d d*
**and** *cp*: *completely-positive n d* Φ
**and** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
**shows** ∃ *As* :: *complex Matrix.mat list*. *length As* = *choi-rank n d* Φ ∧
(∀ *A* ∈ *set As*. *A* ∈ *carrier-mat d n*) ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ *X* = *kraus-map-dim d As X*)
⟨*proof*⟩

**theorem** *choi-rank-eq-rank*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: ⋀*A B*. *A* ∈ *carrier-mat n n* ⟹ *B* ∈ *carrier-mat n n* ⟹ Φ (*A* + *B*) = Φ *A* + Φ *B*
**and** *smult*: ⋀*c A*. *A* ∈ *carrier-mat n n* ⟹ Φ (*c* ·$_m$ *A*) = *c* ·$_m$ Φ *A*
**and** *range*: ∀ *A*. *A* ∈ *carrier-mat n n* ⟶ Φ *A* ∈ *carrier-mat d d*
**and** *cp*: *completely-positive n d* Φ
**and** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
**shows** *choi-rank n d* Φ = *mat-rank* (*n*∗*d*) (*choi-matrix n d* Φ)
⟨*proof*⟩

**end**
**theory** *Day21-Unitary-Freedom*
**imports** *QC-Prelims*
**begin**

## 43 Day 21: unitary freedom of rank-one decompositions and Kraus representations (D21-3, D21-5, D21-6)

### 43.1 The Gram matrix of an independent list is invertible; dual vectors

**lemma** *gram-det-nonzero*:
**assumes** *li*: *lin-indep-vecs n vs*
**shows** *det* ((*mat-of-cols n vs*)$^\dagger$ ∗ *mat-of-cols n vs*) ≠ *0*
⟨*proof*⟩

**lemma** *gram-inverse*:
  **assumes** *li*: *lin-indep-vecs n vs*
  **obtains** *B* **where** $B \in$ *carrier-mat* (*length vs*) (*length vs*)
    **and** $((\textit{mat-of-cols}\ n\ vs)^\dagger * \textit{mat-of-cols}\ n\ vs) * B = 1_m\ (\textit{length}\ vs)$
    **and** $B * ((\textit{mat-of-cols}\ n\ vs)^\dagger * \textit{mat-of-cols}\ n\ vs) = 1_m\ (\textit{length}\ vs)$
    **and** $B^\dagger = B$
⟨*proof*⟩

Dual vectors (the notes' "$T^*$ is onto"): for independent $v_1, \ldots, v_k$ there are $\varphi_1, \ldots, \varphi_k$ with $\langle \varphi_i | v_j \rangle = \delta_{ij}$.

**lemma** *dual-vectors-exist*:
  **assumes** *li*: *lin-indep-vecs n vs*
  **shows** $\exists\, \varphi s ::$ *complex Matrix.vec list*. *length* $\varphi s =$ *length vs* $\wedge$
      $(\forall\, \varphi \in$ *set* $\varphi s$. *dim-vec* $\varphi = n) \wedge$
      $(\forall\, i <$ *length vs*. $\forall\, j <$ *length vs*. $\langle \varphi s\ !\ i \mid vs\ !\ j \rangle =$ (*if* $i = j$ *then 1 else 0*))
⟨*proof*⟩

## 43.2 D21-3: independent vectors give independent rank-one matrices

**definition** *outer-prods* :: *complex Matrix.vec list* ⇒ *complex Matrix.mat list* **where**
  *outer-prods vs* = *map* (λ*k*. *vs* ! (*k div length vs*) ⊘ *vs* ! (*k mod length vs*)) [*0*..<*length vs* ∗ *length vs*]

**lemma** *length-outer-prods* [*simp*]: *length* (*outer-prods vs*) = *length vs* ∗ *length vs*
  ⟨*proof*⟩

**lemma** *outer-prods-nth*:
  **assumes** *k* < *length vs* ∗ *length vs*
  **shows** *outer-prods vs* ! *k* = *vs* ! (*k div length vs*) ⊘ *vs* ! (*k mod length vs*)
  ⟨*proof*⟩

**lemma** *inner-prod-cnj-delta*:
  **assumes** *dim-vec* $\varphi = n$ **and** *dim-vec* $v = n$ **and** $\langle \varphi \mid v \rangle =$ (*if P then 1 else 0*)
  **shows** $\langle v \mid \varphi \rangle =$ (*if P then 1 else 0*)
  ⟨*proof*⟩

**proposition** *outer-prods-lin-indep*:
  **assumes** *li*: *lin-indep-vecs n vs*
  **shows** *mat-lin-indep-complex n* (*outer-prods vs*)
⟨*proof*⟩

## 43.3 Isometric mixing of columns preserves $VV^*$

**lemma** *dagger-transpose-mult*:
  **assumes** *Uc*: $U \in$ *carrier-mat m r* **and** *UU*: $(U^\dagger) * U = 1_m\ r$
  **shows** *transpose-mat U* ∗ ((*transpose-mat U*)$^\dagger$) = $1_m\ r$
⟨*proof*⟩

**lemma** *isometry-mix-outer*:
  **assumes** *Vc*: $V \in$ *carrier-mat n r* **and** *Uc*: $U \in$ *carrier-mat m r* **and** *UU*: $(U^\dagger) * U = 1_m\ r$
  **shows** $(V *$ *transpose-mat U*$) * ((V *$ *transpose-mat U*$)^\dagger) = V * (V^\dagger)$
⟨*proof*⟩

**lemma** *dagger-mult-zero-imp-zero*:
  **fixes** *Y* :: *complex Matrix.mat*
  **assumes** *Yc*: $Y \in$ *carrier-mat p q* **and** *YY*: $(Y^\dagger) * Y = 0_m\ q\ q$
  **shows** $Y = 0_m\ p\ q$
⟨*proof*⟩

## 43.4 D21-5: two rank-one decompositions are related by an isometry

If $P = \sum_{j<r} v_j v_j^* = \sum_{l<m} w_l w_l^*$ with $v_1, \ldots, v_r$ linearly independent (the notes assume rank $P = r$, which is equivalent), then $w_l = \sum_j \alpha_{lj} v_j$ with $U = (\alpha_{lj})$ an $m \times r$ isometry, $U^* U = I_r$. Matrix form: $W = V U^t$.

**theorem** *rank-one-decomp-isometry*:
  **assumes** *li*: *lin-indep-vecs n vs*
    **and** *wsd*: $\forall\, w \in$ *set ws. dim-vec w* $= n$
    **and** *same*: *outer-sum n vs* $=$ *outer-sum n ws*
  **shows** $\exists\, U ::$ *complex Matrix.mat. U* $\in$ *carrier-mat* (*length ws*) (*length vs*) $\wedge$
      ($U^\dagger$) $*$ *U* $= 1_m$ (*length vs*) $\wedge$
      *mat-of-cols n ws* $=$ *mat-of-cols n vs* $*$ *transpose-mat U*
⟨*proof*⟩

**corollary** *rank-one-decomp-isometry-cols*:
  **assumes** *li*: *lin-indep-vecs n vs*
    **and** *wsd*: $\forall\, w \in$ *set ws. dim-vec w* $= n$
    **and** *same*: *outer-sum n vs* $=$ *outer-sum n ws*
  **shows** $\exists\, U ::$ *complex Matrix.mat. U* $\in$ *carrier-mat* (*length ws*) (*length vs*) $\wedge$
      ($U^\dagger$) $*$ *U* $= 1_m$ (*length vs*) $\wedge$
      ($\forall\, l <$ *length ws. ws* ! *l* $=$ *mat-of-cols n vs* $*_v$ *Matrix.row U l*)
⟨*proof*⟩

## 43.5 D21-6 (Choi 2): unitary freedom of Kraus representations

**lemma** *vec-of-kraus-mat-of-cols-index*:
  **assumes** *Vs-c*: $\forall$ *V* $\in$ *set Vs. V* $\in$ *carrier-mat d n* **and** *dpos*: *0* $<$ *d*
    **and** *i*: *i* $<$ *d* **and** *a*: *a* $<$ *n* **and** *k*: *k* $<$ *length Vs*
  **shows** *mat-of-cols* (*n*$*$*d*) (*map* (*vec-of-kraus d n*) *Vs*) $$ (*a*$*$*d* $+$ *i*, *k*) $=$ *Vs* ! *k* $$ (*i*, *a*)
⟨*proof*⟩

**lemma** *lin-indep-complex-mat-vec-of-kraus*:
  **assumes** *li*: *lin-indep-complex-mat d n Vs* **and** *dpos*: *0* $<$ *d*
  **shows** *lin-indep-vecs* (*n*$*$*d*) (*map* (*vec-of-kraus d n*) *Vs*)
⟨*proof*⟩

Choi's Theorem 2: if $\Phi(X) = \sum_{j<r} V_j X V_j^* = \sum_{l<m} W_l X W_l^*$ with $V_1, \ldots, V_r$ linearly independent (the notes: $r = \mathrm{cr}(\Phi)$), then $W_l = \sum_j \alpha_{lj} V_j$ with $U = (\alpha_{lj})$ an isometry, $U^* U = I_r$.

**theorem** *choi-2-unitary-freedom*:
  **assumes** *npos*: *0* $<$ *n* **and** *dpos*: *0* $<$ *d*
    **and** *Ws-c*: $\forall$ *W* $\in$ *set Ws. W* $\in$ *carrier-mat d n*
    **and** *li*: *lin-indep-complex-mat d n Vs*
    **and** *same*: $\forall$ *X. X* $\in$ *carrier-mat n n* $\longrightarrow$ *kraus-map-dim d Vs X* $=$ *kraus-map-dim d Ws X*
  **shows** $\exists\, U ::$ *complex Matrix.mat. U* $\in$ *carrier-mat* (*length Ws*) (*length Vs*) $\wedge$
      ($U^\dagger$) $*$ *U* $= 1_m$ (*length Vs*) $\wedge$
      ($\forall\, l <$ *length Ws. Ws* ! *l* $=$
        *foldr* ($\lambda j$ *acc. U* $$ (*l*,*j*) $\cdot_m$ (*Vs* ! *j*) $+$ *acc*) [*0*..<*length Vs*] (*0*$_m$ *d n*))
⟨*proof*⟩

**end**
**theory** *Day21-Ranges*
  **imports** *Day04-29-Supplement Day02-11-Supplement*
**begin**

# 44 Day 21: ranges of positive matrices (D21-4)

The notes write $R(P)$ for the range of $P$ (here *mat-range k P*) and use two facts: if $0 \le ww^* \le P$ then $w \in R(P)$, and $R(\sum_l w_l w_l^*) = \mathrm{span}\{w_l\}$ (*col-span*). Both rest on one lemma: for Hermitian $P$, every vector orthogonal to $\ker P$ lies in $R(P)$; it is proved from the spectral decomposition $P = UDU^*$ by inverting the nonzero eigenvalues. The audit's earlier formal counterpart was the kernel form *loewner-ker-subset*.

**definition** *mat-range* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec set* **where**
*mat-range k P* = {*P* ∗*v h* | *h*. *dim-vec h* = *k*}

**lemma** *diag-mat-of-real-mult-vec*:
**assumes** *length D* = *k dim-vec v* = *k*
**shows** *diag-mat-of-real D* ∗*v v* = *Matrix.vec k* (λ*i*. *complex-of-real* (*D* ! *i*) ∗ *v* $ *i*)
⟨*proof*⟩

For Hermitian $P$, a vector orthogonal to $\ker P$ lies in the range of $P$.

**lemma** *hermitian-perp-ker-in-range*:
**assumes** *P*: *P* ∈ *carrier-mat k k* **and** *h*: *hermitian P* **and** *w*: *dim-vec w* = *k*
**and** *perp*: ⋀*x*. *dim-vec x* = *k* ⟹ *P* ∗*v x* = *0v k* ⟹ *Quantum.inner-prod x w* = *0*
**shows** *w* ∈ *mat-range k P*
⟨*proof*⟩

The notes: if $0 \le ww^* \le P$ then $w \in R(P)$.

**theorem** *dominated-rank-one-in-range*:
**assumes** *P*: *P* ∈ *carrier-mat k k* **and** *Pp*: *psd P* **and** *w*: *dim-vec w* = *k*
**and** *dom*: *psd* (*P* − *outer-sum k* [*w*])
**shows** *w* ∈ *mat-range k P*
⟨*proof*⟩

The notes: if $P = \sum_l w_l w_l^*$ then $R(P) = \mathrm{span}\{w_1, \ldots, w_m\}$.

**theorem** *outer-sum-range*:
**assumes** *vs*: ∀ *v* ∈ *set vs*. *dim-vec v* = *k*
**shows** *mat-range k* (*outer-sum k vs*) = *col-span k vs*
⟨*proof*⟩

## 44.1 Day 4 revisited: positive definiteness and spanning (D4-8)

The notes: $P = \sum_i |v_i\rangle\langle v_i| > 0$ iff $\mathrm{span}\{v_i\} = \mathbb{C}^n$.

**theorem** *pd-iff-span-all*:
**assumes** *vs*: ∀ *v* ∈ *set vs*. *dim-vec v* = *n*
**shows** *pd* (*outer-sum n vs*) ⟷ *col-span n vs* = *carrier-vec n*
⟨*proof*⟩

**end**
**theory** *Day22-Kraus-Independence*
**imports** *QC-Prelims Jordan-Normal-Form.DL-Rank*
**begin**

# 45 Day 22: linear independence bounds the number of Kraus operators (D22-6)

A linearly independent list of vectors in $\mathbb{C}^n$ has at most $n$ members. The bridge is to JNF's *vec-space*: kernel-triviality of *mat-of-cols* is JNF's *lin-indpt* (via *vec-space.lin-depE*), and *li-le-dim* bounds the cardinality.

**lemma** *lin-indep-vecs-distinct*:
  **assumes** *li*: *lin-indep-vecs n vs*
  **shows** *distinct vs*
⟨*proof*⟩

**lemma** *lin-indep-vecs-length-le*:
  **assumes** *li*: *lin-indep-vecs n vs*
  **shows** *length vs* ≤ *n*
⟨*proof*⟩

Flattening a matrix into a vector carries *mat-lin-indep-complex* to *lin-indep-vecs*, so a linearly independent family in $M_n$ has at most $n^2$ members.

**definition** *flatten-mat* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec* **where**
  *flatten-mat n B* = *Matrix.vec* (*n*∗*n*) (λ*k*. *B* $$ (*k div n*, *k mod n*))

**lemma** *flatten-mat-dim* [*simp*]: *dim-vec* (*flatten-mat n B*) = *n*∗*n*
  ⟨*proof*⟩

**lemma** *flatten-mat-index*:
  **assumes** *i* < *n* **and** *j* < *n*
  **shows** *flatten-mat n B* $ (*i*∗*n* + *j*) = *B* $$ (*i*,*j*)
⟨*proof*⟩

**lemma** *mat-lin-indep-complex-flatten*:
  **assumes** *li*: *mat-lin-indep-complex n Bs*
  **shows** *lin-indep-vecs* (*n*∗*n*) (*map* (*flatten-mat n*) *Bs*)
⟨*proof*⟩

**lemma** *mat-lin-indep-complex-length-le*:
  **assumes** *mat-lin-indep-complex n Bs*
  **shows** *length Bs* ≤ *n*∗*n*
  ⟨*proof*⟩

Corollary to Choi's Theorem 3 (D22-6): an extreme CPTP map $M_n \to M_d$ with a minimal Kraus representation $(V_i)_{i<r}$ has $r^2 \le n^2$, i.e. $\mathrm{cr}(\Phi) \le n$.

**lemma** *length-kraus-products*: *length* (*kraus-products Vs*) = *length Vs* ∗ *length Vs*
  ⟨*proof*⟩

**corollary** *extreme-CPTP-kraus-length-le*:
  **assumes** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
    **and** Φ-*in*: Φ ∈ *CPTP n d*
    **and** *rep*: *cptp-kraus-rep n d* Φ *Vs*
    **and** *minimal*: *lin-indep-complex-mat d n Vs*
    **and** *ext*: *extreme-CPTP-carrier n d* Φ
  **shows** *length Vs* ≤ *n*
⟨*proof*⟩

**corollary** *extreme-CPTP-choi-rank-le*:
  **assumes** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
    **and** Φ-*in*: Φ ∈ *CPTP n d*
    **and** *rep*: *cptp-kraus-rep n d* Φ *Vs*
    **and** *minimal*: *lin-indep-complex-mat d n Vs*
    **and** *ext*: *extreme-CPTP-carrier n d* Φ
  **shows** *choi-rank n d* Φ ≤ *n*
⟨*proof*⟩

**end**
**theory** *Day22-Minimal-Kraus*

**imports** *Day32-Recovery-Proposition*
**begin**

# 46 Day 22: a minimal Kraus representation is linearly independent (D22-1)

## 46.1 Mixing a Kraus list by an isometry does not change the map

**definition** *kraus-mix* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat list* **where**
*kraus-mix d n U Vs* =
*map* (λ*l*. *foldr* (λ*j acc*. *U* $$ (*l*,*j*) ·$_m$ (*Vs* ! *j*) + *acc*) [*0*..<*length Vs*] (*0*$_m$ *d n*)) [*0*..<*dim-row U*]

**lemma** *kraus-mix-length* [*simp*]: *length* (*kraus-mix d n U Vs*) = *dim-row U*
⟨*proof*⟩

**lemma** *kraus-mix-nth*:
**assumes** *l* < *dim-row U*
**shows** *kraus-mix d n U Vs* ! *l* = *foldr* (λ*j acc*. *U* $$ (*l*,*j*) ·$_m$ (*Vs* ! *j*) + *acc*) [*0*..<*length Vs*] (*0*$_m$ *d n*)
⟨*proof*⟩

**lemma** *kraus-mix-carrier*:
**assumes** ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**shows** ∀ *W* ∈ *set* (*kraus-mix d n U Vs*). *W* ∈ *carrier-mat d n*
⟨*proof*⟩

**lemma** *kraus-mix-entry*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n* **and** *l*: *l* < *dim-row U* **and** *i*: *i* < *d* **and** *a*: *a* < *n*
**shows** *kraus-mix d n U Vs* ! *l* $$ (*i*,*a*) = (∑ *j* = *0*..<*length Vs*. *U* $$ (*l*,*j*) * *Vs* ! *j* $$ (*i*,*a*))
⟨*proof*⟩

**lemma** *kraus-mix-mat-of-cols*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n* **and** *Uc*: *U* ∈ *carrier-mat m* (*length Vs*) **and** *dpos*: *0* < *d*
**shows** *mat-of-cols* (*n*∗*d*) (*map* (*vec-of-kraus d n*) (*kraus-mix d n U Vs*))
= *mat-of-cols* (*n*∗*d*) (*map* (*vec-of-kraus d n*) *Vs*) ∗ *transpose-mat U*
⟨*proof*⟩

**lemma** *kraus-mix-same-map*:
**assumes** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
**and** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *Uc*: *U* ∈ *carrier-mat m* (*length Vs*) **and** *UU*: (*U*$^\dagger$) ∗ *U* = *1*$_m$ (*length Vs*)
**and** *Xc*: *M* ∈ *carrier-mat n n*
**shows** *kraus-map-dim d* (*kraus-mix d n U Vs*) *M* = *kraus-map-dim d Vs M*
⟨*proof*⟩

## 46.2 A unitary with a prescribed first row

**lemma** *mat-mult-entry*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A* ∈ *carrier-mat r s B* ∈ *carrier-mat s t k* < *r j* < *t*
**shows** (*A* ∗ *B*) $$ (*k*,*j*) = (∑ *l*<*s*. *A* $$ (*k*,*l*) ∗ *B* $$ (*l*,*j*))
⟨*proof*⟩

**lemma** *transpose-unitary-isometry*:
**assumes** *Mc*: *M* ∈ *carrier-mat r r* **and** *MM*: *M* ∗ (*M*$^\dagger$) = *1*$_m$ *r*
**shows** ((*transpose-mat M*)$^\dagger$) ∗ *transpose-mat M* = *1*$_m$ *r*

⟨*proof*⟩

**lemma** *isometry-with-first-row*:
**assumes** *ud*: *dim-vec u = r* **and** *un*: *‖u‖ = 1*
**shows** ∃ *U* :: *complex Matrix.mat*. *U* ∈ *carrier-mat r r* ∧ (*U*†) ∗ *U* = *1*$_m$ *r* ∧ (∀ *j* < *r*. *U* $$ (*0*,*j*) = *u* $ *j*)
⟨*proof*⟩

### 46.3 D22-1

**lemma** *kraus-map-dim-zero-head*:
**assumes** *Ws-c*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n* **and** *Xc*: *M* ∈ *carrier-mat n n*
**shows** *kraus-map-dim d* (*0*$_m$ *d n* # *Ws*) *M* = *kraus-map-dim d Ws M*
⟨*proof*⟩

If $\Phi(X) = \sum_{j<r} V_j X V_j^*$ with $r = \mathrm{cr}(\Phi)$, then $V_0, \ldots, V_{r-1}$ are linearly independent: a dependency $\sum_j c_j V_j = 0$ would allow a unitary change of representation whose first Kraus operator is 0, giving a representation with $r - 1$ terms.

**theorem** *minimal-kraus-rep-lin-indep*:
**assumes** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
**and** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *minimal*: *choi-rank n d* (*kraus-map-dim d Vs*) = *length Vs*
**shows** *lin-indep-complex-mat d n Vs*
⟨*proof*⟩

**end**
**theory** *Day22-Adjoint-Maps*
**imports** *Day19-Choi-TP-Unital Day04-29-Supplement Day26-CP-Factorization*
**begin**

## 47 Day 22: the adjoint map, an affine bijection from UCP onto CPTP (D22-5)

The notes: $\mathrm{UCP}(M_n, M_d)$ and $\mathrm{CPTP}(M_d, M_n)$ are convex, and $\Gamma(\Phi) = \Phi^*$ is an affine bijection between them with affine inverse, so the two convex sets are affinely isomorphic. The adjoint was available only for Kraus maps (*kraus-adjoint-dim*); here it is defined for every linear $\Phi$. Maps are total functions on matrices, so $\Gamma$ is a bijection up to agreement on $M_n$, resp. $M_d$: $\Gamma$ sends linear UCP maps to linear CPTP maps and back, and $\Gamma(\Gamma(\Phi)) = \Phi$ on $M_n$. The set *UCP* does not record linearity, which the notes' maps have, so it is assumed.

For linear $\Phi : M_n \to M_d$, $\Phi^* : M_d \to M_n$ is determined by $\langle Y|\Phi(X)\rangle = \langle \Phi^*(Y)|X\rangle$; taking $X = E_{ij}$ gives $\Phi^*(Y)_{ij} = \langle \Phi(E_{ij})|Y\rangle$.

**definition** *map-adjoint* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*map-adjoint n d* Φ *Y* = *Matrix.mat n n* (λ(*i*, *j*). ⟨⟨ Φ (*E*⟦*i*,*j* ; *n*,*n*⟧) | *Y* ⟩⟩)

**lemma** *map-adjoint-carrier* [*simp*]: *map-adjoint n d* Φ *Y* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *map-adjoint-dims* [*simp*]: *dim-row* (*map-adjoint n d* Φ *Y*) = *n dim-col* (*map-adjoint n d* Φ *Y*) = *n*
⟨*proof*⟩

**lemma** *lin-entry*:

**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *X*: $X \in$ *carrier-mat n n* **and** *a*: $a < d$ **and** *b*: $b < d$
**shows** Φ *X* $$ (*a*, *b*) = ($\sum i{<}n.\ \sum j{<}n.$ *X* $$ (*i*, *j*) ∗ Φ (*E*⟦*i*,*j* ; *n*,*n*⟧) $$ (*a*, *b*))
⟨*proof*⟩

**theorem** *map-adjoint-correct*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *X*: *X* ∈ *carrier-mat n n* **and** *Y*: *Y* ∈ *carrier-mat d d*
**shows** ⟨⟨ *Y* | Φ *X* ⟩⟩ = ⟨⟨ *map-adjoint n d* Φ *Y* | *X* ⟩⟩
⟨*proof*⟩

**lemma** *hs-inner-prod-mat-unit*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *i*: $i < n$ **and** *j*: $j < n$
**shows** ⟨⟨ *A* | *E*⟦*i*,*j* ; *n*,*n*⟧ ⟩⟩ = *cnj* (*A* $$ (*i*, *j*))
⟨*proof*⟩

**lemma** *hs-inner-prod-ext*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *B*: *B* ∈ *carrier-mat n n*
**and** *eq*: $\bigwedge X.$ *X* ∈ *carrier-mat n n* ⟹ ⟨⟨ *A* | *X* ⟩⟩ = ⟨⟨ *B* | *X* ⟩⟩
**shows** *A* = *B*
⟨*proof*⟩

**lemma** *map-adjoint-lin*: *lin-on d n* (*carrier-mat d d*) (*map-adjoint n d* Φ)
⟨*proof*⟩

**lemma** *map-adjoint-cong*:
**assumes** $\bigwedge X.$ *X* ∈ *carrier-mat n n* ⟹ Φ *X* = Ψ *X*
**shows** *map-adjoint n d* Φ = *map-adjoint n d* Ψ
⟨*proof*⟩

For a Kraus map the adjoint is *kraus-adjoint-dim*.

**lemma** *map-adjoint-kraus*:
**assumes** *As*: ∀ *A* ∈ *set As*. *A* ∈ *carrier-mat d n* **and** *Y*: *Y* ∈ *carrier-mat d d*
**shows** *map-adjoint n d* (*kraus-map-dim d As*) *Y* = *kraus-adjoint-dim n As Y*
⟨*proof*⟩

$\Phi^{**} = \Phi$.

**theorem** *map-adjoint-adjoint*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *X*: *X* ∈ *carrier-mat n n*
**shows** *map-adjoint d n* (*map-adjoint n d* Φ) *X* = Φ *X*
⟨*proof*⟩

**lemma** *cp-kraus-form*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *cp*: *completely-positive n d* Φ **and** *n*: *0* < *n* **and** *d*: *0* < *d*
**shows** ∃ *Ws*. (∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*) ∧ (∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ *X* = *kraus-map-dim d Ws X*)
⟨*proof*⟩

**lemma** *trace-preserving-cong*:
**assumes** $\bigwedge X.$ *X* ∈ *carrier-mat n n* ⟹ Φ *X* = Ψ *X*
**shows** *trace-preserving n* Φ ⟷ *trace-preserving n* Ψ
⟨*proof*⟩

$\Gamma(\Phi) = \Phi^*$ maps the unital CP maps $M_n \to M_d$ to the CPTP maps $M_d \to M_n$ and back.

**theorem** *adjoint-ucp-imp-cptp*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *U*: Φ ∈ *UCP n d* **and** *n*: *0* < *n* **and** *d*: *0* < *d*
**shows** *map-adjoint n d* Φ ∈ *CPTP d n*
⟨*proof*⟩

**theorem** *adjoint-cptp-imp-ucp*:
**assumes** *L*: *lin-on d n* (*carrier-mat d d*) Ψ **and** *C*: Ψ ∈ *CPTP d n* **and** *n*: *0* < *n* **and** *d*: *0* < *d*
**shows** *map-adjoint d n* Ψ ∈ *UCP n d*
⟨*proof*⟩

Γ is affine (for real weights), and $\Gamma^{-1} = \Gamma$ by *map-adjoint-adjoint*.

**lemma** *hs-inner-prod-left-real-comb*:
**fixes** *t s* :: *real*
**assumes** *A*: *A* ∈ *carrier-mat d d* **and** *B*: *B* ∈ *carrier-mat d d* **and** *Y*: *Y* ∈ *carrier-mat d d*
**shows** ⟨⟨ *complex-of-real t* ·$_m$ *A* + *complex-of-real s* ·$_m$ *B* | *Y* ⟩⟩
= *complex-of-real t* ∗ ⟨⟨ *A* | *Y* ⟩⟩ + *complex-of-real s* ∗ ⟨⟨ *B* | *Y* ⟩⟩
⟨*proof*⟩

**theorem** *map-adjoint-affine*:
**fixes** *t* :: *real*
**assumes** *L1*: *lin-on n d* (*carrier-mat n n*) Φ*1* **and** *L2*: *lin-on n d* (*carrier-mat n n*) Φ*2* **and** *Y*: *Y* ∈ *carrier-mat d d*
**shows** *map-adjoint n d* (λ*X*. *complex-of-real t* ·$_m$ Φ*1 X* + *complex-of-real* (*1* − *t*) ·$_m$ Φ*2 X*) *Y*
= *complex-of-real t* ·$_m$ *map-adjoint n d* Φ*1 Y* + *complex-of-real* (*1* − *t*) ·$_m$ *map-adjoint n d* Φ*2 Y*
⟨*proof*⟩

**end**
**theory** *Day22-UCP-Compact*
**imports** *Day16-Block-Form Day19-Choi-TP-Unital Day27-Positive-Map-Criterion*
**begin**

# 48 Day 22: UCP($M_n, M_d$) is compact (D22-4)

The notes: UCP($M_n, M_d$) is convex and compact, closed because the conditions are closed and bounded because $\Phi(I) = I$ forces $0 \le \Phi(E_{ii}) \le I$. The space $L(M_n, M_d)$ carries no topology in this development; the Choi matrix is a linear bijection onto $M_{nd}$, so the statement is made in Choi coordinates: the Choi matrices of linear UCP maps are exactly the positive $P$ whose diagonal blocks sum to $I_d$ (*ucp-choi-image*, with the inverse Choi map *choi-inv*), and that set is compact (*compact-ucp-choi-set*): it is closed, and it lies in $d$ times the compact set of unit-trace positive matrices.

In Choi coordinates a linear map $\Phi : M_n \to M_d$ is UCP iff $P_\Phi \ge 0$ and $\sum_i R_{ii} = I_d$ (*choi-ucp-iff*); the set of such $P$ is compact. Dimensions are cardinalities of finite types, $n = |'n|$, $d = |'d|$, so that the typed matrices of *Separable-Cone* (a Euclidean space) can be used.

**definition** *ucp-choi-set* :: (*complex*, *'n* :: *finite* × *'d* :: *finite*, *'n* × *'d*) *lmat set* **where**
*ucp-choi-set* = {*M*. *psd-lmat M* ∧ (∀ *a* < *CARD*(*'d*). ∀ *b* < *CARD*(*'d*).
(∑ *i*<*CARD*(*'n*). *unwrap-lm M* $$ (*i* ∗ *CARD*(*'d*) + *a*, *i* ∗ *CARD*(*'d*) + *b*)) = (*if a* = *b then 1 else 0*))}

**lemma** *closed-block-sum-eq*:
**assumes** *a*: *a* < *CARD*(*'d* :: *finite*) **and** *b*: *b* < *CARD*(*'d*)
**shows** *closed* {*M* :: (*complex*, *'n* :: *finite* × *'d*, *'n* × *'d*) *lmat*.
(∑ *i*<*CARD*(*'n*). *unwrap-lm M* $$ (*i* ∗ *CARD*(*'d*) + *a*, *i* ∗ *CARD*(*'d*) + *b*)) = *c*}
⟨*proof*⟩

**lemma** *closed-ucp-choi-set*: *closed* (*ucp-choi-set* :: (*complex*, *'n* :: *finite* × *'d* :: *finite*, *'n* × *'d*) *lmat set*)
⟨*proof*⟩

**lemma** *ucp-choi-set-trace*:

**assumes** $M \in$ (*ucp-choi-set* :: (*complex*, *'n* :: *finite* × *'d* :: *finite*, *'n* × *'d*) *lmat set*)
**shows** *Tr-lmat M* = *of-nat CARD('d)*
⟨*proof*⟩

**theorem** *compact-ucp-choi-set*: *compact* (*ucp-choi-set* :: (*complex*, *'n* :: *finite* × *'d* :: *finite*, *'n* × *'d*) *lmat set*)
⟨*proof*⟩

The inverse of the Choi map: $\Phi_P(X) = \sum_{i,j} x_{ij} P_{ij}$ with $P_{ij}$ the $(i,j)$ block.

**definition** *choi-inv* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat*
**where**
*choi-inv n d P X* = *Matrix.mat d d* (λ(*a*, *b*). ∑*i*<*n*. ∑*j*<*n*. *X* $$ (*i*, *j*) ∗ *P* $$ (*i* ∗ *d* + *a*, *j* ∗ *d* + *b*))

**lemma** *choi-inv-lin*: *lin-on n d* (*carrier-mat n n*) (*choi-inv n d P*)
⟨*proof*⟩

**lemma** *choi-matrix-choi-inv*:
**assumes** *P*: *P* ∈ *carrier-mat* (*n* ∗ *d*) (*n* ∗ *d*)
**shows** *choi-matrix n d* (*choi-inv n d P*) = *P*
⟨*proof*⟩

**lemma** *block-sum-entry*:
**assumes** *P*: *P* ∈ *carrier-mat* (*n* ∗ *d*) (*n* ∗ *d*) **and** *a*: *a* < *d* **and** *b*: *b* < *d*
**shows** *msum d d* (λ*i*. *block-of-mat d P i i*) [*0*..<*n*] $$ (*a*, *b*) = (∑*i*<*n*. *P* $$ (*i* ∗ *d* + *a*, *i* ∗ *d* + *b*))
⟨*proof*⟩

UCP in Choi coordinates is exactly *ucp-choi-set*, so it is compact.

**theorem** *ucp-choi-image*:
(λΦ. *wrap-lm* (*choi-matrix CARD('n) CARD('d)* Φ) :: (*complex*, *'n* :: *finite* × *'d* :: *finite*, *'n* × *'d*) *lmat*)
‘ {Φ. *lin-on CARD('n) CARD('d)* (*carrier-mat CARD('n) CARD('n)*) Φ ∧ Φ ∈ *UCP CARD('n) CARD('d)*}
= *ucp-choi-set*
⟨*proof*⟩

**corollary** *compact-ucp*:
*compact* ((λΦ. *wrap-lm* (*choi-matrix CARD('n) CARD('d)* Φ) :: (*complex*, *'n* :: *finite* × *'d* :: *finite*, *'n* × *'d*) *lmat*)
‘ {Φ. *lin-on CARD('n) CARD('d)* (*carrier-mat CARD('n) CARD('n)*) Φ ∧ Φ ∈ *UCP CARD('n) CARD('d)*})
⟨*proof*⟩

**end**
**theory** *Day29-Classical-Codes*
**imports** *Quantum-Computing11*
**begin**

Days 29–30: classical codes over $\mathbb{Z}_2$. Words are lists over $\{0,1\}$; all finite checks are decided by *code-simp* (proof-producing evaluation, no oracle).

**definition** *bin-words* :: *nat* ⇒ *nat list set* **where**
*bin-words n* = *set* (*List.n-lists n* [*0*, *1*])

**definition** *xor-word* :: *nat list* ⇒ *nat list* ⇒ *nat list* **where**
*xor-word v w* = *map2* (λ*a b*. (*a* + *b*) *mod 2*) *v w*

## 48.1 The notes' "Hamming [7,1,3]" code does not correct one error

The notes encode $(a_1,\ldots,a_7) \mapsto (a_1,\ldots,a_7;\ a_1{+}a_2{+}a_3{+}a_4;\ a_1{+}a_2{+}a_5{+}a_6;\ a_1{+}a_3{+}a_5{+}a_7)$ and claim it detects and corrects one error. It detects one error, but two codewords are at distance 2: the word $e_4$ is at distance 1 from the codewords of 0 and of $e_4$, so no decoder corrects every single error.

**definition** *notes-hamming-encode* :: *nat list* ⇒ *nat list* **where**
*notes-hamming-encode a* = *a* @ [(*a*!*0* + *a*!*1* + *a*!*2* + *a*!*3*) *mod 2*, (*a*!*0* + *a*!*1* + *a*!*4* + *a*!*5*) *mod 2*, (*a*!*0* + *a*!*2* + *a*!*4* + *a*!*6*) *mod 2*]

**definition** *notes-hamming-code* :: *nat list set* **where**
*notes-hamming-code* = *notes-hamming-encode* ‘ *bin-words 7*

**definition** *flip-at* :: *nat* ⇒ *nat list* ⇒ *nat list* **where**
*flip-at i w* = *w*[*i* := (*w*!*i* + *1*) *mod 2*]

**lemma** *notes-hamming-code-iff*:
*w* ∈ *notes-hamming-code* ⟷ *take 7 w* ∈ *bin-words 7* ∧ *w* = *notes-hamming-encode* (*take 7 w*)
⟨*proof*⟩

**lemma** *bin-words-iff*: *w* ∈ *bin-words n* ⟷ *length w* = *n* ∧ *set w* ⊆ {*0*, *1*}
⟨*proof*⟩

**lemma** *bin-nth-lt2*: *set v* ⊆ {*0*,*1*} ⟹ *k* < *length v* ⟹ *v* ! *k* < (*2*::*nat*)
⟨*proof*⟩

**lemma** *xor-word-length* [*simp*]: *length* (*xor-word v w*) = *min* (*length v*) (*length w*)
⟨*proof*⟩

**lemma** *xor-word-nth*: *k* < *length v* ⟹ *k* < *length w* ⟹ *xor-word v w* ! *k* = (*v* ! *k* + *w* ! *k*) *mod 2*
⟨*proof*⟩

**lemma** *xor-word-cancel*:
**assumes** *length c* = *length v length c* = *length w set v* ⊆ {*0*,*1*} *set w* ⊆ {*0*,*1*}
**shows** *xor-word c v* = *xor-word c w* ⟷ *v* = *w*
⟨*proof*⟩

**lemma** *mod2-eq-iff*: (*p*::*nat*) *mod 2* = *q mod 2* ⟷ (*even p* ⟷ *even q*)
⟨*proof*⟩

The encoding is linear over $\mathbb{Z}_2$.

**lemma** *notes-hamming-encode-xor*:
**assumes** *length a* = *7 length b* = *7*
**shows** *notes-hamming-encode* (*xor-word a b*) = *xor-word* (*notes-hamming-encode a*) (*notes-hamming-encode b*)
⟨*proof*⟩

**lemma** *flip-at-xor*:
**assumes** *set c* ⊆ {*0*,*1*} *i* < *length c*
**shows** *flip-at i c* = *xor-word c* ((*replicate* (*length c*) *0*)[*i* := *1*])
⟨*proof*⟩

**theorem** *notes-hamming-detects-one-error*:
∀ *c* ∈ *notes-hamming-code*. ∀ *i* < *10*. *flip-at i c* ∉ *notes-hamming-code*
⟨*proof*⟩

**theorem** *notes-hamming-ambiguous-word*:

**defines** $w \equiv [0,0,0,1,0,0,0,0,0,0]$
**shows** *notes-hamming-encode* (*replicate 7 0*) ∈ *notes-hamming-code*
**and** *notes-hamming-encode* $[0,0,0,1,0,0,0]$ ∈ *notes-hamming-code*
**and** *notes-hamming-encode* (*replicate 7 0*) ≠ *notes-hamming-encode* $[0,0,0,1,0,0,0]$
**and** *hamming-dist* (*notes-hamming-encode* (*replicate 7 0*)) *w* = *1*
**and** *hamming-dist* (*notes-hamming-encode* $[0,0,0,1,0,0,0]$) *w* = *1*
⟨*proof*⟩

**theorem** *notes-hamming-not-correcting*: ¬ *corrects-errors notes-hamming-code 1*
⟨*proof*⟩

## 48.2 The Hamming [7,4,3] code corrects one error

**definition** *hamming74-encode* :: *nat list* ⇒ *nat list* **where**
*hamming74-encode a* = *a* @ [(*a*!*0* + *a*!*1* + *a*!*3*) *mod 2*, (*a*!*0* + *a*!*2* + *a*!*3*) *mod 2*, (*a*!*1* + *a*!*2* + *a*!*3*) *mod 2*]

**definition** *hamming74-code* :: *nat list set* **where**
*hamming74-code* = *hamming74-encode* ‘ *bin-words 4*

**lemma** *ex1-in-set*: (∃!*c*. *c* ∈ *set xs* ∧ *P c*) ⟷ *card* (*set* (*filter P xs*)) = *1*
⟨*proof*⟩

**theorem** *hamming74-unique-decoding*:
∀ *w* ∈ *bin-words 7*. ∃!*c*. *c* ∈ *hamming74-code* ∧ *hamming-dist c w* ≤ *1*
⟨*proof*⟩

**theorem** *hamming74-corrects-one-error*: *corrects-errors hamming74-code 1*
⟨*proof*⟩

## 48.3 Probabilities of undetected and uncorrected errors (Day 30)

Each bit flips independently with probability $p$; an error pattern $e$ of weight $k$ on $n$ bits has probability $p^k(1-p)^{n-k}$. An error is undetected when it turns the sent codeword into another codeword.

**definition** *word-prob* :: *real* ⇒ *nat* ⇒ *nat list* ⇒ *real* **where**
*word-prob p n e* = *p* ^ *hamming-weight e* ∗ (*1* − *p*) ^ (*n* − *hamming-weight e*)

**definition** *err-prob* :: *real* ⇒ *nat* ⇒ *nat list set* ⇒ *real* **where**
*err-prob p n ES* = (∑ *e* ∈ *ES*. *word-prob p n e*)

**lemma** *err-prob-list*: *distinct es* ⟹ *err-prob p n* (*set es*) = (∑ *e*←*es*. *word-prob p n e*)
⟨*proof*⟩

**definition** *rep-code* :: *nat list set* **where** *rep-code* = *repetition-encode* ‘ *bin-words 3*
**definition** *par-code* :: *nat list set* **where** *par-code* = *parity-check-encode* ‘ *bin-words 3*

**lemma** *rep-code-list*: *rep-code* = *set* (*map repetition-encode* (*List.n-lists 3* [*0*, *1*]))
⟨*proof*⟩

**lemma** *par-code-list*: *par-code* = *set* (*map parity-check-encode* (*List.n-lists 3* [*0*, *1*]))
⟨*proof*⟩

**definition** *undetected* :: *nat list set* ⇒ *nat* ⇒ *nat list* ⇒ *nat list set* **where**
*undetected C n c* = {*e* ∈ *bin-words n*. *e* ≠ *replicate n 0* ∧ *xor-word c e* ∈ *C*}

**lemma** *set-filter-bin*: {*e* ∈ *bin-words n*. *P e*} = *set* (*filter P* (*List.n-lists n* [*0*, *1*]))

⟨*proof*⟩

Repetition on three bits: $3p^2(1-p)^4 + 3p^4(1-p)^2 + p^6$.

**theorem** *repetition-undetected-prob*:
**assumes** *c*: *c* ∈ *rep-code*
**shows** *err-prob p 6* (*undetected rep-code 6 c*) = *3 * p^2 * (1 - p)^4 + 3 * p^4 * (1 - p)^2 + p^6*
⟨*proof*⟩

Parity on three bits: $\binom{4}{2}p^2(1-p)^2 + p^4$.

**theorem** *parity-undetected-prob*:
**assumes** *c*: *c* ∈ *par-code*
**shows** *err-prob p 4* (*undetected par-code 4 c*) = *6 * p^2 * (1 - p)^2 + p^4*
⟨*proof*⟩

The notes compare the two for $p < \frac{1}{2}$; the inequality in fact holds for all $0 < p < 1$.

**theorem** *repetition-better-than-parity*:
**assumes** *p0*: *0 < p* **and** *p1*: *p < 1*
**shows** *3 * p^2 * (1 - p)^4 + 3 * p^4 * (1 - p)^2 + p^6 < 6 * p^2 * (1 - p)^2 + (p::real)^4*
⟨*proof*⟩

**corollary** *repetition-better-than-parity-notes*:
*0 < p ⟹ p < 1/2 ⟹ 3 * p^2 * (1 - p)^4 + 3 * p^4 * (1 - p)^2 + p^6 < 6 * p^2 * (1 - p)^2 + (p::real)^4*
⟨*proof*⟩

Majority rule $0 \mapsto 000$, $1 \mapsto 111$: decoding fails with probability $3p^2(1-p) + p^3$.

**definition** *maj-decode* :: *nat list* ⇒ *nat* **where**
*maj-decode w* = (*if 2 ≤ sum-list w then 1 else 0*)

**theorem** *majority-uncorrected-prob*:
**assumes** *b*: *b* ∈ {*0*, *1*}
**shows** *err-prob p 3* {*e* ∈ *bin-words 3*. *maj-decode* (*xor-word* (*replicate 3 b*) *e*) ≠ *b*}
= *3 * p^2 * (1 - p) + p^3*
⟨*proof*⟩

**end**
**theory** *Day30-BitFlip-Code*
**imports** *QC-Prelims*
**begin**

# 49 Day 30: the three-qubit bit-flip code and its recovery map (D30-3, D30-4)

The syndrome projections $P_0 = |000\rangle\langle 000| + |111\rangle\langle 111|$, …, as diagonal 0/1 matrices ($|abc\rangle \leftrightarrow e_{4a+2b+c}$), and the single flips $X \otimes I \otimes I$, $I \otimes X \otimes I$, $I \otimes I \otimes X$ as permutation matrices.

**definition** *syndrome-proj* :: *nat set* ⇒ *complex Matrix.mat* **where**
*syndrome-proj K* = *Matrix.mat 8 8* (λ(*i*,*j*). *if i = j ∧ i ∈ K then 1 else 0*)

**abbreviation** $P_0$ ≡ *syndrome-proj* {*0*, *7*}
**abbreviation** $P_1$ ≡ *syndrome-proj* {*4*, *3*}
**abbreviation** $P_2$ ≡ *syndrome-proj* {*2*, *5*}
**abbreviation** $P_3$ ≡ *syndrome-proj* {*1*, *6*}

**definition** *flip-bit-mat* :: *nat* ⇒ *complex Matrix.mat* **where**

*flip-bit-mat b = Matrix.mat 8 8 (λ(i,j). if j = (if i div 2^b mod 2 = 0 then i + 2^b else i − 2^b) then 1 else 0)*

**abbreviation** $X_1 \equiv$ *flip-bit-mat 2*
**abbreviation** $X_2 \equiv$ *flip-bit-mat 1*
**abbreviation** $X_3 \equiv$ *flip-bit-mat 0*

**lemma** *syndrome-proj-carrier* [*simp*]: *syndrome-proj K ∈ carrier-mat 8 8*
⟨*proof*⟩

**lemma** *flip-bit-mat-carrier* [*simp*]: *flip-bit-mat b ∈ carrier-mat 8 8*
⟨*proof*⟩

**lemma** *syndrome-proj-dims* [*simp*]: *dim-row (syndrome-proj K) = 8 dim-col (syndrome-proj K) = 8*
⟨*proof*⟩

**lemma** *flip-bit-mat-dims* [*simp*]: *dim-row (flip-bit-mat b) = 8 dim-col (flip-bit-mat b) = 8*
⟨*proof*⟩

**lemma** *encode-mat-dims* [*simp*]: *dim-row encode-mat = 8 dim-col encode-mat = 2*
⟨*proof*⟩

**lemma** *flip-proj-carrier* [*simp*]: *flip-bit-mat b * syndrome-proj K ∈ carrier-mat 8 8*
⟨*proof*⟩

**lemma** *b3-cases*: $b < (3::nat) \Longrightarrow b = 0 \vee b = 1 \vee b = 2$ ⟨*proof*⟩

Entry of an $8 \times 8$ product, kept as an abstract sum: *simp* unrolls $\sum k{<}8$ into eight terms and then splits every *if*, which does not terminate in practice.

**lemma** *mat-mult-entry-8*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A ∈ carrier-mat 8 8 B ∈ carrier-mat 8 8 i < 8 j < 8*
**shows** *(A * B) $$ (i,j) = (∑k<8. A $$ (i,k) * B $$ (k,j))*
⟨*proof*⟩

**lemma** *syndrome-proj-index*:
**assumes** *i < 8 j < 8*
**shows** *syndrome-proj K $$ (i,j) = (if i = j ∧ i ∈ K then 1 else 0)*
⟨*proof*⟩

**lemma** *syndrome-proj-two*:
**assumes** *a*: *a < 8* **and** *b*: *b < 8* **and** *ab*: *a ≠ b*
**shows** *syndrome-proj {a, b} = unit-vec 8 a ⊘ unit-vec 8 a + unit-vec 8 b ⊘ unit-vec 8 b*
⟨*proof*⟩

**lemma** *syndrome-proj-as-outer*:
$P_0$ = *unit-vec 8 0 ⊘ unit-vec 8 0 + unit-vec 8 7 ⊘ unit-vec 8 7*
$P_1$ = *unit-vec 8 4 ⊘ unit-vec 8 4 + unit-vec 8 3 ⊘ unit-vec 8 3*
$P_2$ = *unit-vec 8 2 ⊘ unit-vec 8 2 + unit-vec 8 5 ⊘ unit-vec 8 5*
$P_3$ = *unit-vec 8 1 ⊘ unit-vec 8 1 + unit-vec 8 6 ⊘ unit-vec 8 6*
⟨*proof*⟩

The flips are the notes' tensors.

**lemma** *tensor3-index*:
**fixes** *A B C* :: *complex Matrix.mat*
**assumes** *A*: *A ∈ carrier-mat 2 2* **and** *B*: *B ∈ carrier-mat 2 2* **and** *C*: *C ∈ carrier-mat 2 2*
**and** *i*: *i < 8* **and** *j*: *j < 8*
**shows** *Tensor.tensor-mat (Tensor.tensor-mat A B) C $$ (i,j)*

$= A \$\$ (i \mathbin{div} 4, j \mathbin{div} 4) * B \$\$ (i \mathbin{div} 2 \bmod 2, j \mathbin{div} 2 \bmod 2) * C \$\$ (i \bmod 2, j \bmod 2)$
⟨*proof*⟩

**lemma** *flips-as-tensors*:
$X_1 = Tensor.tensor\text{-}mat\ (Tensor.tensor\text{-}mat\ Pauli\text{-}X\ (1_m\ 2))\ (1_m\ 2)$
$X_2 = Tensor.tensor\text{-}mat\ (Tensor.tensor\text{-}mat\ (1_m\ 2)\ Pauli\text{-}X)\ (1_m\ 2)$
$X_3 = Tensor.tensor\text{-}mat\ (Tensor.tensor\text{-}mat\ (1_m\ 2)\ (1_m\ 2))\ Pauli\text{-}X$
⟨*proof*⟩

**lemma** *flip-bit-mat-mult-vec*:
**assumes** *b*: $b < 3$ **and** *vd*: $dim\text{-}vec\ v = 8$ **and** *i*: $i < 8$
**shows** $(flip\text{-}bit\text{-}mat\ b *_v v) \$ i = v \$ (if\ i \mathbin{div} 2\hat{}b \bmod 2 = 0\ then\ i + 2\hat{}b\ else\ i - 2\hat{}b)$
⟨*proof*⟩

**lemma** *syndrome-proj-mult-vec*:
**assumes** *vd*: $dim\text{-}vec\ v = 8$ **and** *i*: $i < 8$
**shows** $(syndrome\text{-}proj\ K *_v v) \$ i = (if\ i \in K\ then\ v \$ i\ else\ 0)$
⟨*proof*⟩

**lemma** *syndrome-proj-hermitian*: $(syndrome\text{-}proj\ K)^\dagger = syndrome\text{-}proj\ K$
⟨*proof*⟩

**lemma** *syndrome-proj-idem*: $syndrome\text{-}proj\ K * syndrome\text{-}proj\ K = syndrome\text{-}proj\ K$
⟨*proof*⟩

**lemma** *syndrome-proj-sum*:
$P_0 + P_1 + P_2 + P_3 = 1_m\ 8$
⟨*proof*⟩

The flip *flip-bit-mat b* permutes the basis by the involution *flip-tgt b*.

**definition** *flip-tgt* :: *nat* ⇒ *nat* ⇒ *nat* **where**
$flip\text{-}tgt\ b\ i = (if\ i \mathbin{div} 2\hat{}b \bmod 2 = 0\ then\ i + 2\hat{}b\ else\ i - 2\hat{}b)$

**lemma** *flip-tgt-lt*:
**assumes** *b*: $b < 3$ **and** *i*: $i < 8$ **shows** $flip\text{-}tgt\ b\ i < 8$
⟨*proof*⟩

**lemma** *flip-tgt-invol*:
**assumes** *b*: $b < 3$ **and** *i*: $i < 8$ **shows** $flip\text{-}tgt\ b\ (flip\text{-}tgt\ b\ i) = i$
⟨*proof*⟩

**lemma** *flip-bit-mat-index*:
**assumes** *i*: $i < 8$ **and** *j*: $j < 8$
**shows** $flip\text{-}bit\text{-}mat\ b \$\$ (i,j) = (if\ j = flip\text{-}tgt\ b\ i\ then\ 1\ else\ 0)$
⟨*proof*⟩

**lemma** *flip-tgt-iff*:
**assumes** *b*: $b < 3$ **and** *i*: $i < 8$ **and** *j*: $j < 8$
**shows** $i = flip\text{-}tgt\ b\ j \longleftrightarrow j = flip\text{-}tgt\ b\ i$
⟨*proof*⟩

**lemma** *flip-bit-mat-hermitian*:
**assumes** *b*: $b < 3$
**shows** $(flip\text{-}bit\text{-}mat\ b)^\dagger = flip\text{-}bit\text{-}mat\ b$
⟨*proof*⟩

**lemma** *flip-bit-mat-squared*:
**assumes** *b*: $b < 3$

**shows** *flip-bit-mat b ∗ flip-bit-mat b = 1*$_m$ *8*
⟨*proof*⟩

Kraus operators of the recovery map $R(Y) = P_0 Y P_0 + \sum_k X_k P_k Y P_k X_k$.

**definition** *recovery-ops* :: *complex Matrix.mat list* **where**
*recovery-ops* = [$P_0$, $X_1 * P_1$, $X_2 * P_2$, $X_3 * P_3$]

**definition** *recovery-map* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*recovery-map* = *kraus-map-dim 8 recovery-ops*

**lemma** *flip-proj-dag-mult*:
**assumes** *b*: *b* < *3*
**shows** ((*flip-bit-mat b ∗ syndrome-proj K*)$^\dagger$) ∗ (*flip-bit-mat b ∗ syndrome-proj K*) = *syndrome-proj K*
⟨*proof*⟩

**theorem** *recovery-ops-kraus*: *kraus-operators 8 8 recovery-ops*
⟨*proof*⟩

**corollary** *recovery-map-CPTP*: *recovery-map* ∈ *CPTP 8 8*
⟨*proof*⟩

The encoded state $\psi = a|000\rangle + b|111\rangle$ and its single-flip corruptions.

**lemma** *encode-mat-mult-vec*:
**assumes** *xd*: *dim-vec x = 2* **and** *i*: *i* < *8*
**shows** (*encode-mat* ∗$_v$ *x*) $ *i* = (*if i = 0 then x $ 0 else if i = 7 then x $ 1 else 0*)
⟨*proof*⟩

**lemma** *encode-mat-dim* [*simp*]: *dim-vec* (*encode-mat* ∗$_v$ *x*) = *8* ⟨*proof*⟩

**lemma** *recovery-kraus-term*:
**fixes** *φ* :: *complex Matrix.vec*
**assumes** *φd*: *dim-vec φ = 8*
**shows** *recovery-map* (*φ* ⊘ *φ*) =
(*P*$_0$ ∗$_v$ *φ*) ⊘ (*P*$_0$ ∗$_v$ *φ*) + ((*X*$_1$ ∗$_v$ (*P*$_1$ ∗$_v$ *φ*)) ⊘ (*X*$_1$ ∗$_v$ (*P*$_1$ ∗$_v$ *φ*))
+ ((*X*$_2$ ∗$_v$ (*P*$_2$ ∗$_v$ *φ*)) ⊘ (*X*$_2$ ∗$_v$ (*P*$_2$ ∗$_v$ *φ*))
+ ((*X*$_3$ ∗$_v$ (*P*$_3$ ∗$_v$ *φ*)) ⊘ (*X*$_3$ ∗$_v$ (*P*$_3$ ∗$_v$ *φ*)) + *0*$_m$ *8 8*)))
⟨*proof*⟩

**theorem** *bit-flip-code-recovery*:
**fixes** *x* :: *complex Matrix.vec*
**assumes** *xd*: *dim-vec x = 2*
**defines** *ψ* ≡ *encode-mat* ∗$_v$ *x*
**shows** *recovery-map* (*ψ* ⊘ *ψ*) = *ψ* ⊘ *ψ*
**and** *recovery-map* ((*X*$_1$ ∗$_v$ *ψ*) ⊘ (*X*$_1$ ∗$_v$ *ψ*)) = *ψ* ⊘ *ψ*
**and** *recovery-map* ((*X*$_2$ ∗$_v$ *ψ*) ⊘ (*X*$_2$ ∗$_v$ *ψ*)) = *ψ* ⊘ *ψ*
**and** *recovery-map* ((*X*$_3$ ∗$_v$ *ψ*) ⊘ (*X*$_3$ ∗$_v$ *ψ*)) = *ψ* ⊘ *ψ*
⟨*proof*⟩

**end**
**theory** *Day31-Code-Fidelity*
**imports** *Day30-BitFlip-Code*
**begin**

Day 31: the three-qubit code against independent bit flips. Each qubit flips with probability $p$, so $E(Y) = (1-p)^3 Y + p(1-p)^2 \sum_i X_i Y X_i + p^2(1-p) \sum_{i<j} X_i X_j Y X_j X_i + p^3\, X_1X_2X_3 Y X_3X_2X_1$. On an encoded state the recovery $R$ leaves $((1-p)^3 + 3p(1-p)^2)|\tilde\psi\rangle\langle\tilde\psi| + (3p^2(1-p) + p^3)|X_1X_2X_3\tilde\psi\rangle\langle X_1X_2X_3\tilde\psi|$. The notes state this formula for a general $Y$; it holds for $Y$ supported on the code space (for $Y = |100\rangle\langle 100|$ and $p = 0$ one gets $R(Y) = |000\rangle\langle 000| \neq Y$).

**definition** *three-bit-channel* :: *real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*three-bit-channel p M* =
*complex-of-real* ((1 − *p*)^3) ·$_m$ *M*
+ *complex-of-real* (*p* ∗ (1 − *p*)^2) ·$_m$ ($X_1$ ∗ *M* ∗ ($X_1$†))
+ *complex-of-real* (*p* ∗ (1 − *p*)^2) ·$_m$ ($X_2$ ∗ *M* ∗ ($X_2$†))
+ *complex-of-real* (*p* ∗ (1 − *p*)^2) ·$_m$ ($X_3$ ∗ *M* ∗ ($X_3$†))
+ *complex-of-real* (*p*^2 ∗ (1 − *p*)) ·$_m$ (($X_2$ ∗ $X_3$) ∗ *M* ∗ (($X_2$ ∗ $X_3$)†))
+ *complex-of-real* (*p*^2 ∗ (1 − *p*)) ·$_m$ (($X_1$ ∗ $X_3$) ∗ *M* ∗ (($X_1$ ∗ $X_3$)†))
+ *complex-of-real* (*p*^2 ∗ (1 − *p*)) ·$_m$ (($X_1$ ∗ $X_2$) ∗ *M* ∗ (($X_1$ ∗ $X_2$)†))
+ *complex-of-real* (*p*^3) ·$_m$ (($X_1$ ∗ $X_2$ ∗ $X_3$) ∗ *M* ∗ (($X_1$ ∗ $X_2$ ∗ $X_3$)†))

**definition** *swap2* :: *complex Matrix.vec* ⇒ *complex Matrix.vec* **where**
*swap2 x* = *Matrix.vec 2* (λ*i*. *x* $ (1 − *i*))

**lemma** *swap2-dim* [*simp*]: *dim-vec* (*swap2 x*) = 2 ⟨*proof*⟩

**lemma** *flips-on-codeword*:
**assumes** *x*: *dim-vec x* = 2
**defines** ψ ≡ *encode-mat* ∗$_v$ *x* **and** φ ≡ *encode-mat* ∗$_v$ *swap2 x*
**shows** ($X_1$ ∗ $X_2$ ∗ $X_3$) ∗$_v$ ψ = φ
**and** ($X_2$ ∗ $X_3$) ∗$_v$ ψ = $X_1$ ∗$_v$ φ
**and** ($X_1$ ∗ $X_3$) ∗$_v$ ψ = $X_2$ ∗$_v$ φ
**and** ($X_1$ ∗ $X_2$) ∗$_v$ ψ = $X_3$ ∗$_v$ φ
⟨*proof*⟩

## 49.1 Errors followed by recovery

**lemma** *recovery-ops-carrier'*: ∀ *W* ∈ *set recovery-ops*. *W* ∈ *carrier-mat 8 8*
⟨*proof*⟩

**lemma** *recovery-map-add*: *A* ∈ *carrier-mat 8 8* ⟹ *B* ∈ *carrier-mat 8 8* ⟹ *recovery-map* (*A* + *B*) = *recovery-map A* + *recovery-map B*
⟨*proof*⟩

**lemma** *recovery-map-smult*: *A* ∈ *carrier-mat 8 8* ⟹ *recovery-map* (*c* ·$_m$ *A*) = *c* ·$_m$ *recovery-map A*
⟨*proof*⟩

**lemma** *comb8*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *B*: *B* ∈ *carrier-mat n n*
**shows** *complex-of-real a* ·$_m$ *A* + *complex-of-real b* ·$_m$ *A* + *complex-of-real b* ·$_m$ *A* + *complex-of-real b* ·$_m$ *A*
+ *complex-of-real c* ·$_m$ *B* + *complex-of-real c* ·$_m$ *B* + *complex-of-real c* ·$_m$ *B* + *complex-of-real d* ·$_m$ *B*
= *complex-of-real* (*a* + 3 ∗ *b*) ·$_m$ *A* + *complex-of-real* (3 ∗ *c* + *d*) ·$_m$ *B*
⟨*proof*⟩

**lemma** *outer8*: *dim-vec v* = 8 ⟹ *v* ⊘ *v* ∈ *carrier-mat 8 8*
⟨*proof*⟩

D31-2: errors followed by recovery on an encoded state.

**theorem** *three-bit-code-recovery*:
**assumes** *x*: *dim-vec x* = 2
**defines** ψ ≡ *encode-mat* ∗$_v$ *x* **and** φ ≡ *encode-mat* ∗$_v$ *swap2 x*
**shows** *recovery-map* (*three-bit-channel p* (ψ ⊘ ψ))
= *complex-of-real* ((1 − *p*)^3 + 3 ∗ *p* ∗ (1 − *p*)^2) ·$_m$ (ψ ⊘ ψ)
+ *complex-of-real* (3 ∗ *p*^2 ∗ (1 − *p*) + *p*^3) ·$_m$ (φ ⊘ φ)
⟨*proof*⟩

The code is better than doing nothing iff $(1-p)^3 + 3p(1-p)^2 \geq 1-p$ iff $p \leq \frac{1}{2}$.

**theorem** *three-bit-code-better-iff*:
**assumes** *p0*: $0 \leq p$ **and** *p1*: $p < 1$
**shows** $(1 - p) \leq (1 - p)\hat{}3 + 3 * p * (1 - p)\hat{}2 \longleftrightarrow p \leq (1/2 :: real)$
⟨*proof*⟩

## 49.2 Minimum fidelities (D31-4)

**lemma** *smult-mat-mult-vec8*:
**fixes** $A :: complex\ Matrix.mat$
**assumes** *dim-col* $A =$ *dim-vec* $v$
**shows** $(k \cdot_m A) *_v v = k \cdot_v (A *_v v)$
⟨*proof*⟩

**lemma** *quad-two-outer*:
**fixes** $\psi\ \varphi :: complex\ Matrix.vec$
**assumes** $\psi d$: *dim-vec* $\psi = n$ **and** $\varphi d$: *dim-vec* $\varphi = n$
**shows** $\langle \psi \mid (a \cdot_m (\psi \oslash \psi) + b \cdot_m (\varphi \oslash \varphi)) *_v \psi \rangle$
$= a * (\langle \psi|\psi \rangle * \langle \psi|\psi \rangle) + b * (\langle \varphi|\psi \rangle * \langle \psi|\varphi \rangle)$
⟨*proof*⟩

**lemma** *encode-inner*:
**assumes** *u*: *dim-vec* $u = 2$ **and** *v*: *dim-vec* $v = 2$
**shows** $\langle encode\text{-}mat *_v u \mid encode\text{-}mat *_v v \rangle = \langle u|v \rangle$
⟨*proof*⟩

**theorem** *code-min-fidelity*:
**assumes** *p0*: $0 \leq p$ **and** *p1*: $p \leq 1$
**defines** *c1* $\equiv (1 - p)\hat{}3 + 3 * p * (1 - p)\hat{}2$
**shows** $\forall x.$ *dim-vec* $x = 2 \longrightarrow \langle x|x \rangle = 1 \longrightarrow$
*sqrt c1* $\leq$ *fidelity* (*encode-mat* $*_v$ $x$) (*recovery-map* (*three-bit-channel p* ((*encode-mat* $*_v$ $x$) $\oslash$ (*encode-mat* $*_v$ $x$))))
**and** *fidelity* (*encode-mat* $*_v$ *unit-vec 2 0*)
(*recovery-map* (*three-bit-channel p* ((*encode-mat* $*_v$ *unit-vec 2 0*) $\oslash$ (*encode-mat* $*_v$ *unit-vec 2 0*)))) = *sqrt c1*
⟨*proof*⟩

**theorem** *nothing-min-fidelity*:
**assumes** *p0*: $0 \leq p$ **and** *p1*: $p \leq 1$
**shows** $\forall \psi.$ *dim-vec* $\psi = 2 \longrightarrow \langle \psi|\psi \rangle = 1 \longrightarrow$ *sqrt* $(1 - p) \leq$ *fidelity* $\psi$ (*bit-flip-channel p* ($\psi \oslash \psi$))
**and** *fidelity* (*unit-vec 2 0*) (*bit-flip-channel p* (*unit-vec 2 0* $\oslash$ *unit-vec 2 0*)) = *sqrt* $(1 - p)$
⟨*proof*⟩

**end**
**theory** *Day32-Pauli-Expansion*
**imports** *QC-Prelims*
**begin**

# 50 Day 32: the Pauli expansion identity for unitaries (D32-7)

**lemma** *cmod-parallelogram*:
**fixes** $a\ b :: complex$
**shows** $(cmod\ (a + b))^2 + (cmod\ (a - b))^2 = 2 * (cmod\ a)^2 + 2 * (cmod\ b)^2$
⟨*proof*⟩

**proposition** *pauli-expansion-unitary*:
**fixes** $U :: complex\ Matrix.mat$ **and** $a0\ a1\ a2\ a3 :: complex$

**assumes** *Uu*: *unitary U*
**and** *U-eq*: $U = a0 \cdot_m 1_m\ 2 + a1 \cdot_m Pauli\text{-}X + a2 \cdot_m Pauli\text{-}Y + a3 \cdot_m Pauli\text{-}Z$
**shows** $(cmod\ a0)^2 + (cmod\ a1)^2 + (cmod\ a2)^2 + (cmod\ a3)^2 = 1$
⟨*proof*⟩

**end**
**theory** *Day32-Recovery-Proposition*
**imports** *Day21-Unitary-Freedom*
**begin**

# 51 Days 32–33: composition of Kraus maps and the general recovery proposition (D32-4, D33-2)

## 51.1 Algebra of *foldr*-sums of matrices

**lemma** *foldr-add-mat-mult-left*:
**fixes** $f :: 'a \Rightarrow complex\ Matrix.mat$
**assumes** *Ac*: $A \in carrier\text{-}mat\ m\ n$ **and** *fc*: $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ n\ p$
**shows** $A * foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ n\ p) = foldr\ (\lambda x\ acc.\ A * f\ x + acc)\ xs\ (0_m\ m\ p)$
⟨*proof*⟩

**lemma** *foldr-add-mat-mult-right*:
**fixes** $f :: 'a \Rightarrow complex\ Matrix.mat$
**assumes** *Bc*: $B \in carrier\text{-}mat\ n\ p$ **and** *fc*: $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ m\ n$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ m\ n) * B = foldr\ (\lambda x\ acc.\ f\ x * B + acc)\ xs\ (0_m\ m\ p)$
⟨*proof*⟩

**lemma** *foldr-add-mat-carrier'*:
**fixes** $f :: 'a \Rightarrow complex\ Matrix.mat$
**assumes** $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ m\ n$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ m\ n) \in carrier\text{-}mat\ m\ n$
⟨*proof*⟩

**lemma** *foldr-add-mat-dagger*:
**fixes** $f :: 'a \Rightarrow complex\ Matrix.mat$
**assumes** $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ m\ n$
**shows** $(foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ m\ n))^\dagger = foldr\ (\lambda x\ acc.\ (f\ x)^\dagger + acc)\ xs\ (0_m\ n\ m)$
⟨*proof*⟩

**lemma** *foldr-add-mat-cong*:
**fixes** $f\ g :: 'a \Rightarrow complex\ Matrix.mat$
**assumes** $\bigwedge x.\ x \in set\ xs \Longrightarrow f\ x = g\ x$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ z = foldr\ (\lambda x\ acc.\ g\ x + acc)\ xs\ z$
⟨*proof*⟩

**lemma** *foldr-add-mat-zero*:
$foldr\ (\lambda x\ acc.\ (0_m\ m\ n :: complex\ Matrix.mat) + acc)\ xs\ (0_m\ m\ n) = 0_m\ m\ n$
⟨*proof*⟩

**lemma** *foldr-add-mat-delta*:
**fixes** $f :: nat \Rightarrow complex\ Matrix.mat$
**assumes** *fc*: $\forall x \in set\ xs.\ f\ x \in carrier\text{-}mat\ m\ n$ **and** *d*: *distinct xs* **and** *i*: $i \in set\ xs$
**shows** $foldr\ (\lambda x\ acc.\ (if\ x = i\ then\ f\ x\ else\ 0_m\ m\ n) + acc)\ xs\ (0_m\ m\ n) = f\ i$
⟨*proof*⟩

**lemma** *foldr-add-mat-smult-pull*:
**fixes** $f :: 'a \Rightarrow complex\ Matrix.mat$ **and** $c :: 'a \Rightarrow complex$

**assumes** *fc*: $\forall\, x \in set\ xs.\ f\ x = c\ x \cdot_m M$ **and** *Mc*: $M \in carrier\text{-}mat\ m\ n$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ m\ n) = sum\text{-}list\ (map\ c\ xs) \cdot_m M$
⟨*proof*⟩

**lemma** *foldr-list-as-index*:
**fixes** $g :: complex\ Matrix.mat \Rightarrow complex\ Matrix.mat$
**shows** $foldr\ (\lambda A\ acc.\ g\ A + acc)\ As\ z = foldr\ (\lambda l\ acc.\ g\ (As\ !\ l) + acc)\ [0..<length\ As]\ z$
⟨*proof*⟩

## 51.2 Composition of Kraus maps

**definition** $kraus\text{-}compose :: complex\ Matrix.mat\ list \Rightarrow complex\ Matrix.mat\ list \Rightarrow complex\ Matrix.mat\ list$ **where**
$kraus\text{-}compose\ As\ Es = concat\ (map\ (\lambda E.\ map\ (\lambda A.\ A * E)\ As)\ Es)$

**lemma** *kraus-compose-Nil* [*simp*]: $kraus\text{-}compose\ As\ [] = []$
⟨*proof*⟩

**lemma** *kraus-compose-Cons*: $kraus\text{-}compose\ As\ (E \#\ Es) = map\ (\lambda A.\ A * E)\ As\ @\ kraus\text{-}compose\ As\ Es$
⟨*proof*⟩

**lemma** *length-kraus-compose* [*simp*]: $length\ (kraus\text{-}compose\ As\ Es) = length\ Es * length\ As$
⟨*proof*⟩

**lemma** *kraus-compose-nth*:
**assumes** *i*: $i < length\ Es$ **and** *l*: $l < length\ As$
**shows** $kraus\text{-}compose\ As\ Es\ !\ (i * length\ As + l) = As\ !\ l * Es\ !\ i$
⟨*proof*⟩

**lemma** *kraus-compose-carrier*:
**assumes** $\forall\, A \in set\ As.\ A \in carrier\text{-}mat\ m\ d$ **and** $\forall\, E \in set\ Es.\ E \in carrier\text{-}mat\ d\ n$
**shows** $\forall\, W \in set\ (kraus\text{-}compose\ As\ Es).\ W \in carrier\text{-}mat\ m\ n$
⟨*proof*⟩

**lemma** *kraus-map-dim-map-mult-right*:
**assumes** *As-c*: $\forall\, A \in set\ As.\ A \in carrier\text{-}mat\ m\ d$ **and** *Ec*: $E \in carrier\text{-}mat\ d\ n$ **and** *Xc*: $M \in carrier\text{-}mat\ n\ n$
**shows** $kraus\text{-}map\text{-}dim\ m\ (map\ (\lambda A.\ A * E)\ As)\ M = kraus\text{-}map\text{-}dim\ m\ As\ (E * M * (E^\dagger))$
⟨*proof*⟩

**lemma** *kraus-map-dim-compose*:
**assumes** *As-c*: $\forall\, A \in set\ As.\ A \in carrier\text{-}mat\ m\ d$ **and** *Es-c*: $\forall\, E \in set\ Es.\ E \in carrier\text{-}mat\ d\ n$
**and** *Xc*: $M \in carrier\text{-}mat\ n\ n$
**shows** $kraus\text{-}map\text{-}dim\ m\ (kraus\text{-}compose\ As\ Es)\ M = kraus\text{-}map\text{-}dim\ m\ As\ (kraus\text{-}map\text{-}dim\ d\ Es\ M)$
⟨*proof*⟩

**lemma** *sum-wdag-w-map-mult-right*:
**assumes** *As-c*: $\forall\, A \in set\ As.\ A \in carrier\text{-}mat\ m\ d$ **and** *Ec*: $E \in carrier\text{-}mat\ d\ n$
**shows** $sum\text{-}wdag\text{-}w\ n\ (map\ (\lambda A.\ A * E)\ As) = (E^\dagger) * sum\text{-}wdag\text{-}w\ d\ As * E$
⟨*proof*⟩

**lemma** *sum-wdag-w-compose*:
**assumes** *As-c*: $\forall\, A \in set\ As.\ A \in carrier\text{-}mat\ m\ d$ **and** *Es-c*: $\forall\, E \in set\ Es.\ E \in carrier\text{-}mat\ d\ n$
**and** *As-tp*: $sum\text{-}wdag\text{-}w\ d\ As = 1_m\ d$
**shows** $sum\text{-}wdag\text{-}w\ n\ (kraus\text{-}compose\ As\ Es) = sum\text{-}wdag\text{-}w\ n\ Es$
⟨*proof*⟩

**lemma** *kraus-operators-compose*:
**assumes** *As*: *kraus-operators d m As* **and** *Es*: *kraus-operators n d Es*
**shows** *kraus-operators n m* (*kraus-compose As Es*)
⟨*proof*⟩

## 51.3 The general recovery proposition (D32-4) and its linear-combination form (D33-2)

## 51.4 Matrix sums acting on vectors

**lemma** *smult-zero-vec* [*simp*]: (*c* :: *complex*) ·$_v$ $0_v$ *n* = $0_v$ *n*
⟨*proof*⟩

**lemma** *zero-mat-mult-vec*:
**assumes** *dim-vec w* = *n* **shows** ($0_m$ *m n* :: *complex Matrix.mat*) *∗*$_v$ *w* = $0_v$ *m*
⟨*proof*⟩

**lemma** *mult-mat-vec-zero-vec*:
**assumes** *Ac*: *A* ∈ *carrier-mat m n* **shows** *A ∗*$_v$ ($0_v$ *n* :: *complex Matrix.vec*) = $0_v$ *m*
⟨*proof*⟩

**lemma** *foldr-add-mat-mult-vec-zero*:
**fixes** *f* :: *'a* ⇒ *complex Matrix.mat*
**assumes** *fc*: ∀ *x* ∈ *set xs*. *f x* ∈ *carrier-mat n n* **and** *wd*: *dim-vec w* = *n*
**and** *z*: ⋀*x*. *x* ∈ *set xs* ⟹ *f x ∗*$_v$ *w* = $0_v$ *n*
**shows** *foldr* (λ*x acc*. *f x* + *acc*) *xs* ($0_m$ *n n*) *∗*$_v$ *w* = $0_v$ *n*
⟨*proof*⟩

**lemma** *foldr-add-mat-mult-vec-delta*:
**fixes** *f* :: *nat* ⇒ *complex Matrix.mat*
**assumes** *fc*: ∀ *x* ∈ *set xs*. *f x* ∈ *carrier-mat n n* **and** *wd*: *dim-vec w* = *n*
**and** *d*: *distinct xs* **and** *j*: *j* ∈ *set xs*
**and** *del*: ⋀*x*. *x* ∈ *set xs* ⟹ *f x ∗*$_v$ *w* = (*if x* = *j then w else* $0_v$ *n*)
**shows** *foldr* (λ*x acc*. *f x* + *acc*) *xs* ($0_m$ *n n*) *∗*$_v$ *w* = *w*
⟨*proof*⟩

**lemma** *vec-sum-delta*:
**assumes** *fin*: *finite Js* **and** *j*: *j* ∈ *Js* **and** *wd*: ⋀*i*. *i* ∈ *Js* ⟹ *dim-vec* (*w i*) = *n*
**shows** *vec-sum n Js* (λ*i*. *if i* = *j then w i else* $0_v$ *n*) = *w j*
⟨*proof*⟩

Let $V = \operatorname{ran} P$ for a projection $P$ and let $U_0, \dots, U_{t-1}$ be unitaries with $U_i V \perp U_j V$ for $i \neq j$, i.e. $P U_i^* U_j P = 0$. With $P_i = U_i P U_i^*$, $Q = I - \sum_i P_i$ and the Kraus operators $P U_i^*$ (the notes write $U_i^* P_i = P U_i^*$) the map $R(X) = QXQ + \sum_i P U_i^* X U_i P$ is CPTP and recovers every $\psi \in V$ from any $\varphi = \sum_i a_i U_i \psi$: $R(|\varphi\rangle\langle\varphi|) = (\sum_i |a_i|^2)|\psi\rangle\langle\psi|$.

**definition** *image-proj-sum* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* **where**
*image-proj-sum n P Us* = *foldr* (λ*i acc*. *Us* ! *i ∗ P ∗* ((*Us* ! *i*)$^\dagger$) + *acc*) [*0..<length Us*] ($0_m$ *n n*)

**definition** *recovery-ops* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat list* **where**
*recovery-ops n P Us* = (1$_m$ *n* − *image-proj-sum n P Us*) # *map* (λ*i*. *P ∗* ((*Us* ! *i*)$^\dagger$)) [*0..<length Us*]

The hypotheses of the proposition, bundled: $P$ a projection and $P U_i^* U_j P = \delta_{ij} P$. For unitaries $U_i$ the case $i = j$ is automatic (*recovery-setting-unitaries*); the weaker form also covers the rescaled errors of the Knill–Laflamme theorem.

**definition** *recovery-setting* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat list* ⇒ *bool* **where**

*recovery-setting n P Us* $\longleftrightarrow$ *projection n P* $\wedge$ $(\forall\, U \in set\ Us.\ U \in carrier\text{-}mat\ n\ n)$
$\wedge\ (\forall\, i < length\ Us.\ \forall\, j < length\ Us.\ P * (((Us\ !\ i)^\dagger) * Us\ !\ j) * P = (if\ i = j\ then\ P\ else\ 0_m\ n\ n))$

**lemma** *recovery-settingD*:
**assumes** *recovery-setting n P Us*
**shows** *projection n P* $\forall\, U \in set\ Us.\ U \in carrier\text{-}mat\ n\ n$
$\forall\, i < length\ Us.\ \forall\, j < length\ Us.\ i \neq j \longrightarrow P * (((Us\ !\ i)^\dagger) * Us\ !\ j) * P = 0_m\ n\ n$
$\forall\, i < length\ Us.\ P * (((Us\ !\ i)^\dagger) * Us\ !\ i) * P = P$
⟨*proof*⟩

**lemma** *recovery-setting-unitaries*:
**assumes** *P*: *projection n P* **and** *Us*: $\forall\, U \in set\ Us.\ U \in carrier\text{-}mat\ n\ n \wedge unitary\ U$
**and** *orth*: $\forall\, i < length\ Us.\ \forall\, j < length\ Us.\ i \neq j \longrightarrow P * (((Us\ !\ i)^\dagger) * Us\ !\ j) * P = 0_m\ n\ n$
**shows** *recovery-setting n P Us*
⟨*proof*⟩

**lemma**
**assumes** *S*: *recovery-setting n P Us*
**shows** *rec-Pc*: $P \in carrier\text{-}mat\ n\ n$ **and** *rec-Ph*: $P^\dagger = P$ **and** *rec-PP*: $P * P = P$
⟨*proof*⟩

**lemma**
**assumes** *S*: *recovery-setting n P Us*
**shows** *rec-Uc*: $i < length\ Us \Longrightarrow Us\ !\ i \in carrier\text{-}mat\ n\ n$
**and** *rec-PUUP*: $i < length\ Us \Longrightarrow P * (((Us\ !\ i)^\dagger) * Us\ !\ i) * P = P$
⟨*proof*⟩

**lemma** *rec-Pi-c*:
**assumes** *S*: *recovery-setting n P Us* **and** *i*: $i < length\ Us$
**shows** $Us\ !\ i * P * ((Us\ !\ i)^\dagger) \in carrier\text{-}mat\ n\ n$
⟨*proof*⟩

**lemma** *rec-Pi-mult*:
**assumes** *S*: *recovery-setting n P Us* **and** *i*: $i < length\ Us$ **and** *j*: $j < length\ Us$
**shows** $(Us\ !\ i * P * ((Us\ !\ i)^\dagger)) * (Us\ !\ j * P * ((Us\ !\ j)^\dagger))$
$= (if\ i = j\ then\ Us\ !\ i * P * ((Us\ !\ i)^\dagger)\ else\ 0_m\ n\ n)$
⟨*proof*⟩

**lemma** *image-proj-sum-carrier*:
**assumes** *S*: *recovery-setting n P Us*
**shows** *image-proj-sum n P Us* $\in carrier\text{-}mat\ n\ n$
⟨*proof*⟩

**lemma** *image-proj-sum-hermitian*:
**assumes** *S*: *recovery-setting n P Us*
**shows** $(image\text{-}proj\text{-}sum\ n\ P\ Us)^\dagger = image\text{-}proj\text{-}sum\ n\ P\ Us$
⟨*proof*⟩

**lemma** *image-proj-sum-idem*:
**assumes** *S*: *recovery-setting n P Us*
**shows** *image-proj-sum n P Us* $*$ *image-proj-sum n P Us* $=$ *image-proj-sum n P Us*
⟨*proof*⟩

**lemma** *image-proj-sum-projection*:
**assumes** *S*: *recovery-setting n P Us*
**shows** *projection n* (*image-proj-sum n P Us*)
⟨*proof*⟩

**lemma** *dagger-one-minus*:
**assumes** *Sc*: $M \in$ *carrier-mat n n* **and** *Sh*: $M^\dagger = M$
**shows** $(1_m\ n - M)^\dagger = 1_m\ n - M$
⟨*proof*⟩

**lemma** *recovery-ops-carrier*:
**assumes** *S*: *recovery-setting n P Us*
**shows** $\forall\, W \in$ *set* (*recovery-ops n P Us*). $W \in$ *carrier-mat n n*
⟨*proof*⟩

**theorem** *recovery-ops-kraus-operators*:
**assumes** *S*: *recovery-setting n P Us*
**shows** *kraus-operators n n* (*recovery-ops n P Us*)
⟨*proof*⟩

**corollary** *recovery-map-CPTP′*:
**assumes** *S*: *recovery-setting n P Us* **and** *npos*: $0 < n$
**shows** *kraus-map-dim n* (*recovery-ops n P Us*) $\in$ *CPTP n n*
⟨*proof*⟩

Action of the Kraus operators on $\varphi = \sum_i a_i U_i \psi$, $\psi \in V$.

**lemma** *rec-kraus-on-image*:
**assumes** *S*: *recovery-setting n P Us*
**and** $\psi d$: *dim-vec* $\psi = n$ **and** $\psi V$: $P *_v \psi = \psi$ **and** *i*: $i <$ *length Us* **and** *j*: $j <$ *length Us*
**shows** $(P * ((Us\ !\ i)^\dagger)) *_v (Us\ !\ j *_v \psi) =$ (*if* $i = j$ *then* $\psi$ *else* $0_v\ n$)
⟨*proof*⟩

**lemma** *rec-Q-on-image*:
**assumes** *S*: *recovery-setting n P Us*
**and** $\psi d$: *dim-vec* $\psi = n$ **and** $\psi V$: $P *_v \psi = \psi$ **and** *j*: $j <$ *length Us*
**shows** $(1_m\ n -$ *image-proj-sum n P Us*$) *_v (Us\ !\ j *_v \psi) = 0_v\ n$
⟨*proof*⟩

**theorem** *recovery-linear-combination*:
**fixes** *a* :: *nat* ⇒ *complex*
**assumes** *S*: *recovery-setting n P Us* **and** $\psi d$: *dim-vec* $\psi = n$ **and** $\psi V$: $P *_v \psi = \psi$
**defines** $\varphi \equiv$ *vec-sum n* $\{i.\ i <$ *length Us*$\}$ $(\lambda i.\ a\ i \cdot_v (Us\ !\ i *_v \psi))$
**shows** *kraus-map-dim n* (*recovery-ops n P Us*) $(\varphi \oslash \varphi)$
$=$ *complex-of-real* $(\sum i <$ *length Us*. $(cmod\ (a\ i))^2) \cdot_m (\psi \oslash \psi)$
⟨*proof*⟩

**corollary** *recovery-single-error*:
**assumes** *S*: *recovery-setting n P Us*
**and** $\psi d$: *dim-vec* $\psi = n$ **and** $\psi V$: $P *_v \psi = \psi$ **and** *j*: $j <$ *length Us*
**shows** *kraus-map-dim n* (*recovery-ops n P Us*) $((Us\ !\ j *_v \psi) \oslash (Us\ !\ j *_v \psi)) = \psi \oslash \psi$
⟨*proof*⟩

**end**
**theory** *Day32-Y-Code*
**imports** *Day33-Shor-Code*
**begin**

Day 32: the code for $Y$ flips. With $\beta = Y|+\rangle$ one has $|+\rangle \perp \beta$, $Y|+\rangle = \beta$, $Y\beta = |+\rangle$, and the code $|0\rangle \mapsto |++\!+\rangle$, $|1\rangle \mapsto |\beta\beta\beta\rangle$ corrects one $Y$ flip exactly as the bit-flip code corrects one $X$ flip.

**definition** *ket-beta* :: *complex Matrix.vec* **where** *ket-beta* = i $\cdot_v$ *ket-minus*

**definition** *ub* :: *bool* ⇒ *complex Matrix.vec* **where** *ub s* = (*if s then ket-plus else ket-beta*)

**lemma** *ket-beta-dim* [*simp*]: *dim-vec ket-beta* = *2* ⟨*proof*⟩
**lemma** *ket-plus-dim* [*simp*]: *dim-vec ket-plus* = *2* ⟨*proof*⟩
**lemma** *ub-dim* [*simp*]: *dim-vec* (*ub s*) = *2* ⟨*proof*⟩

**lemma** *ket-beta-index*: *ket-beta* $ *0* = i / *sqrt 2 ket-beta* $ *1* = − i / *sqrt 2*
⟨*proof*⟩

**lemma** *ket-plus-index*: *ket-plus* $ *0* = *1* / *sqrt 2 ket-plus* $ *1* = *1* / *sqrt 2*
⟨*proof*⟩

**lemma** *Y-plus*: *Pauli-Y* ∗$_v$ *ket-plus* = *ket-beta*
⟨*proof*⟩

**lemma** *Y-beta*: *Pauli-Y* ∗$_v$ *ket-beta* = *ket-plus*
⟨*proof*⟩

**lemma** *Y-ub*: *Pauli-Y* ∗$_v$ *ub s* = *ub* (¬ *s*)
⟨*proof*⟩

**lemma** *inner2*: *dim-vec u* = *2* ⟹ *dim-vec v* = *2* ⟹ ⟨*u*|*v*⟩ = *cnj* (*u* $ *0*) ∗ *v* $ *0* + *cnj* (*u* $ *1*) ∗ *v* $ *1*
⟨*proof*⟩

**lemma** *ub-inner*: ⟨*ub s* | *ub t*⟩ = (*if s* = *t then 1 else 0*)
⟨*proof*⟩

**theorem** *ket-beta-props*: ⟨*ket-plus* | *ket-beta*⟩ = *0 Pauli-Y* ∗$_v$ *ket-plus* = *ket-beta Pauli-Y* ∗$_v$ *ket-beta* = *ket-plus*
⟨*proof*⟩

## 51.5 Codes with orthogonal error images

For a code with orthonormal basis $v_1, \ldots, v_k$ (the columns of $V$, projection $VV^*$) and unitary errors whose images of the basis are mutually orthogonal, the recovery proposition gives one CPTP map correcting each error on every code state $Vc$.

**lemma** *proj-of-orthonormal*:
**assumes** *orth*: *orthonormal-list n vs*
**defines** *V* ≡ *mat-of-cols n vs*
**shows** *projection n* (*V* ∗ (*V*†))
⟨*proof*⟩

**lemma** *proj-fixes-code*:
**assumes** *orth*: *orthonormal-list n vs* **and** *c*: *dim-vec c* = *length vs*
**defines** *V* ≡ *mat-of-cols n vs*
**shows** (*V* ∗ (*V*†)) ∗$_v$ (*V* ∗$_v$ *c*) = *V* ∗$_v$ *c*
⟨*proof*⟩

**theorem** *orthogonal-images-recovery*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *Us*: ∀ *U* ∈ *set Us*. *U* ∈ *carrier-mat n n* ∧ *unitary U*
**and** *img*: ∀ *i* < *length Us*. ∀ *j* < *length Us*. *i* ≠ *j* ⟶
(∀ *a* < *length vs*. ∀ *b* < *length vs*. ⟨*Us* ! *i* ∗$_v$ *vs* ! *a* | *Us* ! *j* ∗$_v$ *vs* ! *b*⟩ = *0*)
**defines** *V* ≡ *mat-of-cols n vs*
**shows** *recovery-setting n* (*V* ∗ (*V*†)) *Us*
⟨*proof*⟩

**corollary** *orthogonal-images-correct*:
**assumes** *orth*: *orthonormal-list n vs*
**and** *Us*: $\forall\, U \in$ *set Us*. $U \in$ *carrier-mat n n* $\wedge$ *unitary U*
**and** *img*: $\forall\, i <$ *length Us*. $\forall\, j <$ *length Us*. $i \neq j \longrightarrow$
$(\forall\, a <$ *length vs*. $\forall\, b <$ *length vs*. $\langle Us\ !\ i *_v vs\ !\ a \mid Us\ !\ j *_v vs\ !\ b\rangle = 0)$
**shows** $\exists$ *Rs*. *kraus-operators n n Rs* $\wedge$
$(\forall\,(c ::$ *complex Matrix.vec*$)\ j$. *dim-vec c* $=$ *length vs* $\longrightarrow$ $j <$ *length Us* $\longrightarrow$
*kraus-map-dim n Rs* (*Us* ! $j *$ ((*mat-of-cols n vs* $*_v$ *c*) $\oslash$ (*mat-of-cols n vs* $*_v$ *c*)) $*$ ((*Us* ! $j)^\dagger$))
$=$ (*mat-of-cols n vs* $*_v$ *c*) $\oslash$ (*mat-of-cols n vs* $*_v$ *c*))
⟨*proof*⟩

## 51.6 The $Y$-flip code

**definition** *tv2* :: *bool* $\Rightarrow$ *bool* $\Rightarrow$ *bool* $\Rightarrow$ *complex Matrix.vec* **where**
*tv2 a b c* = *tensor-vec* (*tensor-vec* (*ub a*) (*ub b*)) (*ub c*)

**definition** *y-errors* :: *complex Matrix.mat list* **where**
*y-errors* = [$1_m$ *8*, *bop 0 Pauli-Y*, *bop 1 Pauli-Y*, *bop 2 Pauli-Y*]

**definition** *y-codewords* :: *complex Matrix.vec list* **where**
*y-codewords* = [*tv2 True True True*, *tv2 False False False*]

**lemma** *tv2-dim* [*simp*]: *dim-vec* (*tv2 a b c*) = *8*
⟨*proof*⟩

**lemma** *tv2-inner*: $\langle$*tv2 a b c* $\mid$ *tv2 a′ b′ c′*$\rangle$ = (*if* $a = a' \wedge b = b' \wedge c = c'$ *then 1 else 0*)
⟨*proof*⟩

**lemma** *y-actions*:
*bop 0 Pauli-Y* $*_v$ *tv2 a b c* = *tv2* ($\neg$ *a*) *b c*
*bop 1 Pauli-Y* $*_v$ *tv2 a b c* = *tv2 a* ($\neg$ *b*) *c*
*bop 2 Pauli-Y* $*_v$ *tv2 a b c* = *tv2 a b* ($\neg$ *c*)
⟨*proof*⟩

**lemma** *y-codewords-orthonormal*: *orthonormal-list 8 y-codewords*
⟨*proof*⟩

**lemma** *y-errors-unitary*: $\forall\, U \in$ *set y-errors*. $U \in$ *carrier-mat 8 8* $\wedge$ *unitary U*
⟨*proof*⟩

**lemma** *y-images-orthogonal*:
$\forall\, i <$ *length y-errors*. $\forall\, j <$ *length y-errors*. $i \neq j \longrightarrow$
$(\forall\, a <$ *length y-codewords*. $\forall\, b <$ *length y-codewords*.
$\langle$*y-errors* ! $i *_v$ *y-codewords* ! $a \mid$ *y-errors* ! $j *_v$ *y-codewords* ! $b\rangle = 0)$
⟨*proof*⟩

D32-2: one CPTP map corrects every single $Y$ flip on every state $a|++\rangle + b|\beta\beta\beta\rangle$.

**theorem** *y-code-corrects*:
$\exists$ *Rs*. *kraus-operators 8 8 Rs* $\wedge$
$(\forall\, c\ j$. *dim-vec c* $=$ *2* $\longrightarrow$ $j <$ *4* $\longrightarrow$
*kraus-map-dim 8 Rs* (*y-errors* ! $j *$ ((*mat-of-cols 8 y-codewords* $*_v$ *c*) $\oslash$ (*mat-of-cols 8 y-codewords* $*_v$ *c*)) $*$ ((*y-errors* ! $j)^\dagger$))
$=$ (*mat-of-cols 8 y-codewords* $*_v$ *c*) $\oslash$ (*mat-of-cols 8 y-codewords* $*_v$ *c*))
⟨*proof*⟩

**end**

**theory** *Day33-Shor-Code*
**imports** *Day32-Recovery-Proposition Day32-Pauli-Expansion Day30-BitFlip-Code*
**begin**

Days 32–33: the nine-qubit Shor code. The $2^9$-dimensional space is never handled entrywise: nine qubits are three blocks of three, every single-qubit operator acts inside one block, and all inner products factor over the blocks (*tensor-inner-prod*).

## 51.7 Tensor products of matrices

**lemma** *tensor-index-c*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A* ∈ *carrier-mat r c B* ∈ *carrier-mat p q i* < *r* ∗ *p j* < *c* ∗ *q* 0 < *c* 0 < *q*
**shows** *Tensor.tensor-mat A B* $$ (*i*,*j*) = *A* $$ (*i div p*, *j div q*) ∗ *B* $$ (*i mod p*, *j mod q*)
⟨*proof*⟩

**lemma** *tensor-one-left*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *Ac*: *A* ∈ *carrier-mat m m* **and** *m*: 0 < *m*
**shows** *Tensor.tensor-mat* (1$_m$ 1) *A* = *A*
⟨*proof*⟩

**lemma** *tensor-one-right*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *Ac*: *A* ∈ *carrier-mat m m* **and** *m*: 0 < *m*
**shows** *Tensor.tensor-mat A* (1$_m$ 1) = *A*
⟨*proof*⟩

**lemma** *tensor-mat-add-right*:
**fixes** *A B C* :: *complex Matrix.mat*
**assumes** *Ac*: *A* ∈ *carrier-mat r c* **and** *Bc*: *B* ∈ *carrier-mat p q* **and** *Cc*: *C* ∈ *carrier-mat p q*
**and** *c*: 0 < *c* **and** *q*: 0 < *q*
**shows** *Tensor.tensor-mat A* (*B* + *C*) = *Tensor.tensor-mat A B* + *Tensor.tensor-mat A C*
⟨*proof*⟩

**lemma** *tensor-mat-add-left*:
**fixes** *A B C* :: *complex Matrix.mat*
**assumes** *Ac*: *A* ∈ *carrier-mat r c* **and** *Bc*: *B* ∈ *carrier-mat r c* **and** *Cc*: *C* ∈ *carrier-mat p q*
**and** *c*: 0 < *c* **and** *q*: 0 < *q*
**shows** *Tensor.tensor-mat* (*A* + *B*) *C* = *Tensor.tensor-mat A C* + *Tensor.tensor-mat B C*
⟨*proof*⟩

**lemma** *tensor-mat-smult-right*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *Ac*: *A* ∈ *carrier-mat r c* **and** *Bc*: *B* ∈ *carrier-mat p q* **and** *c*: 0 < *c* **and** *q*: 0 < *q*
**shows** *Tensor.tensor-mat A* (*k* ·$_m$ *B*) = *k* ·$_m$ *Tensor.tensor-mat A B*
⟨*proof*⟩

**lemma** *tensor-mat-smult-left*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *Ac*: *A* ∈ *carrier-mat r c* **and** *Bc*: *B* ∈ *carrier-mat p q* **and** *c*: 0 < *c* **and** *q*: 0 < *q*
**shows** *Tensor.tensor-mat* (*k* ·$_m$ *A*) *B* = *k* ·$_m$ *Tensor.tensor-mat A B*
⟨*proof*⟩

## 51.8 Single-qubit operators on nine qubits, and on a block of three

$\tilde{U} = I \otimes \cdots \otimes I \otimes U \otimes I \otimes \cdots \otimes I$ with $U$ in position $i$ ($i = 0, \ldots, 8$).

**definition** *qubit-op* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*qubit-op i U* = *Tensor.tensor-mat* (*Tensor.tensor-mat* (1$_m$ (2^*i*)) *U*) (1$_m$ (2^(8−*i*)))

**definition** *bop* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*bop s U* = *Tensor.tensor-mat* (*Tensor.tensor-mat* (1$_m$ (2^*s*)) *U*) (1$_m$ (2^(2−*s*)))

**definition** *blk* :: *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*blk b M* = *Tensor.tensor-mat* (*Tensor.tensor-mat* (1$_m$ (8^*b*)) *M*) (1$_m$ (8^(2−*b*)))

**lemma** *qubit-op-carrier*:
**assumes** *U*: *U* ∈ *carrier-mat 2 2* **and** *i*: *i* < *9*
**shows** *qubit-op i U* ∈ *carrier-mat 512 512*
⟨*proof*⟩

**lemma** *qubit-op-blk*:
**assumes** *b*: *b* < *3* **and** *s*: *s* < *3*
**shows** *qubit-op* (*3* ∗ *b* + *s*) *U* = *blk b* (*bop s U*)
⟨*proof*⟩

**lemma** *qubit-op-blk-div*:
**assumes** *i*: *i* < *9*
**shows** *qubit-op i U* = *blk* (*i div 3*) (*bop* (*i mod 3*) *U*)
⟨*proof*⟩

**lemma** *bop-cases*:
**fixes** *U* :: *complex Matrix.mat*
**assumes** *U*: *U* ∈ *carrier-mat 2 2*
**shows** *bop 0 U* = *Tensor.tensor-mat* (*Tensor.tensor-mat U* (1$_m$ *2*)) (1$_m$ *2*)
**and** *bop 1 U* = *Tensor.tensor-mat* (*Tensor.tensor-mat* (1$_m$ *2*) *U*) (1$_m$ *2*)
**and** *bop 2 U* = *Tensor.tensor-mat* (*Tensor.tensor-mat* (1$_m$ *2*) (1$_m$ *2*)) *U*
⟨*proof*⟩

**lemma** *bop-carrier*:
**assumes** *U*: *U* ∈ *carrier-mat 2 2* **and** *s*: *s* < *3*
**shows** *bop s U* ∈ *carrier-mat 8 8*
⟨*proof*⟩

**definition** *tv3* :: *complex Matrix.vec* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.vec* **where**
*tv3 x y z* = *tensor-vec* (*tensor-vec x y*) *z*

**lemma** *tv3-dim*: *dim-vec x* = *8* ⟹ *dim-vec y* = *8* ⟹ *dim-vec z* = *8* ⟹ *dim-vec* (*tv3 x y z*) = *512*
⟨*proof*⟩

**lemma** *blk-tv3*:
**assumes** *M*: *M* ∈ *carrier-mat 8 8* **and** *x*: *dim-vec x* = *8* **and** *y*: *dim-vec y* = *8* **and** *z*: *dim-vec z* = *8*
**shows** *blk 0 M* ∗$_v$ *tv3 x y z* = *tv3* (*M* ∗$_v$ *x*) *y z*
**and** *blk 1 M* ∗$_v$ *tv3 x y z* = *tv3 x* (*M* ∗$_v$ *y*) *z*
**and** *blk 2 M* ∗$_v$ *tv3 x y z* = *tv3 x y* (*M* ∗$_v$ *z*)
⟨*proof*⟩

## 51.9 The block vectors $|m\rangle \pm |7-m\rangle$

In a block of three qubits, *gvec p 0* is $|000\rangle \pm |111\rangle$; a bit flip moves it to *gvec p m* with $m \in \{4, 2, 1\}$. These eight vectors are mutually orthogonal.

**definition** *pm* :: *bool* ⇒ *complex* **where** *pm p* = (*if p then 1 else* −*1*)

**definition** *gvec* :: *bool* ⇒ *nat* ⇒ *complex Matrix.vec* **where**
*gvec p m* = *unit-vec 8 m* + *pm p* ·$_v$ *unit-vec 8* (7 − *m*)

**lemma** *gvec-dim* [*simp*]: *dim-vec* (*gvec p m*) = *8*
⟨*proof*⟩

**lemma** *gvec-index*:
**assumes** *r*: *r* < *8* **and** *m*: *m* < *8*
**shows** *gvec p m* $ *r* = (*if r* = *m then 1 else 0*) + *pm p* ∗ (*if r* = 7 − *m then 1 else 0*)
⟨*proof*⟩

**lemma** *mult-gvec0*:
**assumes** *M*: *M* ∈ *carrier-mat 8 8* **and** *r*: *r* < *8*
**shows** (*M* ∗$_v$ *gvec p 0*) $ *r* = *M* $$ (*r*,*0*) + *pm p* ∗ *M* $$ (*r*,7)
⟨*proof*⟩

**lemma** *bop-entry-cases*:
**assumes** *U*: *U* ∈ *carrier-mat 2 2* **and** *r*: *r* < *8* **and** *k*: *k* < *8*
**shows** *bop 0 U* $$ (*r*,*k*) = *U* $$ (*r div 4*, *k div 4*) ∗ *1*$_m$ *2* $$ (*r div 2 mod 2*, *k div 2 mod 2*) ∗ *1*$_m$ *2* $$ (*r mod 2*, *k mod 2*)
**and** *bop 1 U* $$ (*r*,*k*) = *1*$_m$ *2* $$ (*r div 4*, *k div 4*) ∗ *U* $$ (*r div 2 mod 2*, *k div 2 mod 2*) ∗ *1*$_m$ *2* $$ (*r mod 2*, *k mod 2*)
**and** *bop 2 U* $$ (*r*,*k*) = *1*$_m$ *2* $$ (*r div 4*, *k div 4*) ∗ *1*$_m$ *2* $$ (*r div 2 mod 2*, *k div 2 mod 2*) ∗ *U* $$ (*r mod 2*, *k mod 2*)
⟨*proof*⟩

**lemma** *bop-X-gvec*:
**assumes** *s*: *s* < *3*
**shows** *bop s Pauli-X* ∗$_v$ *gvec p 0* = *gvec p* (*2*^(*2*−*s*))
⟨*proof*⟩

**lemma** *bop-Y-gvec*:
**assumes** *s*: *s* < *3*
**shows** *bop s Pauli-Y* ∗$_v$ *gvec p 0* = (−i) ·$_v$ *gvec* (¬ *p*) (*2*^(*2*−*s*))
⟨*proof*⟩

**lemma** *bop-Z-gvec*:
**assumes** *s*: *s* < *3*
**shows** *bop s Pauli-Z* ∗$_v$ *gvec p 0* = *gvec* (¬ *p*) *0*
⟨*proof*⟩

**lemma** *inner-two-units*:
**assumes** *a*: *a* < *n* **and** *b*: *b* < *n* **and** *c*: *c* < *n* **and** *d*: *d* < *n*
**shows** ⟨*unit-vec n a* + *α* ·$_v$ *unit-vec n b* | *unit-vec n c* + *β* ·$_v$ *unit-vec n d*⟩
= (*if a* = *c then 1 else 0*) + *β* ∗ (*if a* = *d then 1 else 0*)
+ *cnj α* ∗ (*if b* = *c then 1 else 0*) + *cnj α* ∗ *β* ∗ (*if b* = *d then 1 else 0*)
⟨*proof*⟩

**lemma** *gvec-inner*:
**assumes** *m*: *m* ∈ {*0*,*1*,*2*,*4*} **and** *m*′: *m*′ ∈ {*0*,*1*,*2*,*4*}
**shows** ⟨*gvec p m* | *gvec p*′ *m*′⟩ = (*if m* = *m*′ ∧ *p* = *p*′ *then 2 else 0*)
⟨*proof*⟩

## 51.10 Code words and the images of the errors

**definition** *gv* :: *bool* × *nat* ⇒ *complex Matrix.vec* **where** *gv x* = *gvec* (*fst x*) (*snd x*)

A label lists the block vectors of a product state, one per block.

**definition** *lab-vec* :: (*bool* × *nat*) *list* ⇒ *complex Matrix.vec* **where**
*lab-vec l* = *tv3* (*gv* (*l* ! *0*)) (*gv* (*l* ! *1*)) (*gv* (*l* ! *2*))

**definition** *valid-label* :: (*bool* × *nat*) *list* ⇒ *bool* **where**
*valid-label l* ⟷ *length l* = *3* ∧ (∀ *x* ∈ *set l*. *snd x* ∈ {*0*,*1*,*2*,*4*})

**definition** *sk* :: *complex* **where** *sk* = *complex-of-real* (*1* / *sqrt 8*)

**definition** *shor-cw* :: *bool* ⇒ *complex Matrix.vec* **where**
*shor-cw p* = *sk* ·$_v$ *lab-vec* (*replicate 3* (*p*, *0*))

The notes' code words $|0_L\rangle, |1_L\rangle$.

**lemma** *shor-cw-notes*:
*shor-cw True* = *complex-of-real* (*1* / *sqrt* (*2*^*3*)) ·$_v$
*tensor-vec* (*tensor-vec* (*unit-vec 8 0* + *unit-vec 8 7*) (*unit-vec 8 0* + *unit-vec 8 7*)) (*unit-vec 8 0* + *unit-vec 8 7*)
*shor-cw False* = *complex-of-real* (*1* / *sqrt* (*2*^*3*)) ·$_v$
*tensor-vec* (*tensor-vec* (*unit-vec 8 0* − *unit-vec 8 7*) (*unit-vec 8 0* − *unit-vec 8 7*)) (*unit-vec 8 0* − *unit-vec 8 7*)
⟨*proof*⟩

**lemma** *gv-dim* [*simp*]: *dim-vec* (*gv x*) = *8*
⟨*proof*⟩

**lemma** *lab-vec-dim* [*simp*]: *dim-vec* (*lab-vec l*) = *512*
⟨*proof*⟩

**lemma** *shor-cw-dim* [*simp*]: *dim-vec* (*shor-cw p*) = *512*
⟨*proof*⟩

**lemma** *tv3-smult*:
**assumes** *dim-vec x* = *8 dim-vec y* = *8 dim-vec z* = *8*
**shows** *tv3* (*k* ·$_v$ *x*) *y z* = *k* ·$_v$ *tv3 x y z tv3 x* (*k* ·$_v$ *y*) *z* = *k* ·$_v$ *tv3 x y z*
*tv3 x y* (*k* ·$_v$ *z*) = *k* ·$_v$ *tv3 x y z*
⟨*proof*⟩

**lemma** *qubit-op-tl*:
**assumes** *i*: *i* < *9* **and** *U*: *U* ∈ *carrier-mat 2 2*
**and** *act*: *bop* (*i mod 3*) *U* ∗$_v$ *gvec p 0* = *κ* ·$_v$ *gvec q m*
**shows** *qubit-op i U* ∗$_v$ *lab-vec* (*replicate 3* (*p*, *0*)) = *κ* ·$_v$ *lab-vec* ((*replicate 3* (*p*, *0*))[*i div 3* := (*q*, *m*)])
⟨*proof*⟩

**lemma** *qubit-op-X-tl*:
*i* < *9* ⟹ *qubit-op i Pauli-X* ∗$_v$ *lab-vec* (*replicate 3* (*p*, *0*))
= *lab-vec* ((*replicate 3* (*p*, *0*))[*i div 3* := (*p*, *2*^(*2* − *i mod 3*))])
⟨*proof*⟩

**lemma** *qubit-op-Y-tl*:
*i* < *9* ⟹ *qubit-op i Pauli-Y* ∗$_v$ *lab-vec* (*replicate 3* (*p*, *0*))
= (−i) ·$_v$ *lab-vec* ((*replicate 3* (*p*, *0*))[*i div 3* := (¬ *p*, *2*^(*2* − *i mod 3*))])
⟨*proof*⟩

**lemma** *qubit-op-Z-tl*:
*i* < *9* ⟹ *qubit-op i Pauli-Z* ∗$_v$ *lab-vec* (*replicate 3* (*p*, *0*))
= *lab-vec* ((*replicate 3* (*p*, *0*))[*i div 3* := (¬ *p*, *0*)])
⟨*proof*⟩

**lemma** *lab-vec-inner*:
**assumes** *l*: *valid-label l* **and** *l'*: *valid-label l'*
**shows** ⟨*lab-vec l* | *lab-vec l'*⟩ = (*if l* = *l' then 8 else 0*)
⟨*proof*⟩

## 51.11 The 22 correctable errors

$I$, $X_1, \ldots, X_9$, $Y_1, \ldots, Y_9$, $Z_1, Z_4, Z_7$ (positions counted from 0 here).

**definition** *shor-errors* :: *complex Matrix.mat list* **where**
*shor-errors* = $1_m$ *512* # *map* (λ*i*. *qubit-op i Pauli-X*) [*0*..<*9*] @ *map* (λ*i*. *qubit-op i Pauli-Y*) [*0*..<*9*]
@ *map* (λ*b*. *qubit-op* (*3* ∗ *b*) *Pauli-Z*) [*0*..<*3*]

**definition** *shor-labels* :: *bool* ⇒ (*bool* × *nat*) *list list* **where**
*shor-labels p* = *replicate 3* (*p*, *0*)
# *map* (λ*i*. (*replicate 3* (*p*, *0*))[*i div 3* := (*p*, *2*^(*2* − *i mod 3*))]) [*0*..<*9*]
@ *map* (λ*i*. (*replicate 3* (*p*, *0*))[*i div 3* := (¬ *p*, *2*^(*2* − *i mod 3*))]) [*0*..<*9*]
@ *map* (λ*b*. (*replicate 3* (*p*, *0*))[*b* := (¬ *p*, *0*)]) [*0*..<*3*]

**definition** *shor-kappa* :: *complex list* **where**
*shor-kappa* = *1* # *replicate 9 1* @ *replicate 9* (−i) @ *replicate 3 1*

**lemma** *shor-lengths* [*simp*]: *length shor-errors* = *22 length* (*shor-labels p*) = *22 length shor-kappa* = *22*
⟨*proof*⟩

**lemma** *shor-errors-cases*:
**assumes** *j*: *j* < *22*
**obtains** *j* = *0 shor-errors* ! *j* = $1_m$ *512 shor-labels p* ! *j* = *replicate 3* (*p*, *0*) *shor-kappa* ! *j* = *1*
| *i* **where** *i* < *9 j* = *Suc i shor-errors* ! *j* = *qubit-op i Pauli-X*
*shor-labels p* ! *j* = (*replicate 3* (*p*, *0*))[*i div 3* := (*p*, *2*^(*2* − *i mod 3*))] *shor-kappa* ! *j* = *1*
| *i* **where** *i* < *9 j* = *10* + *i shor-errors* ! *j* = *qubit-op i Pauli-Y*
*shor-labels p* ! *j* = (*replicate 3* (*p*, *0*))[*i div 3* := (¬ *p*, *2*^(*2* − *i mod 3*))] *shor-kappa* ! *j* = −i
| *b* **where** *b* < *3 j* = *19* + *b shor-errors* ! *j* = *qubit-op* (*3* ∗ *b*) *Pauli-Z*
*shor-labels p* ! *j* = (*replicate 3* (*p*, *0*))[*b* := (¬ *p*, *0*)] *shor-kappa* ! *j* = *1*
⟨*proof*⟩

**lemma** *shor-errors-carrier*: *j* < *22* ⟹ *shor-errors* ! *j* ∈ *carrier-mat 512 512*
⟨*proof*⟩

**lemma** *shor-errors-action*:
**assumes** *j*: *j* < *22*
**shows** *shor-errors* ! *j* ∗$_v$ *shor-cw p* = (*shor-kappa* ! *j* ∗ *sk*) ·$_v$ *lab-vec* (*shor-labels p* ! *j*)
⟨*proof*⟩

**lemma** *shor-labels-valid*: *j* < *22* ⟹ *valid-label* (*shor-labels p* ! *j*)
⟨*proof*⟩

**lemma** *shor-labels-distinct*: *distinct* (*shor-labels True* @ *shor-labels False*)
⟨*proof*⟩

**lemma** *shor-labels-differ*:
**assumes** *i*: *i* < *22* **and** *j*: *j* < *22* **and** *ij*: *i* ≠ *j*
**shows** *shor-labels p* ! *i* ≠ *shor-labels p'* ! *j*
⟨*proof*⟩

**lemma** *shor-errors-orth*:
**assumes** *i*: *i* < *22* **and** *j*: *j* < *22* **and** *ij*: *i* ≠ *j*
**shows** ⟨*shor-errors* ! *i* ∗$_v$ *shor-cw p* | *shor-errors* ! *j* ∗$_v$ *shor-cw p'*⟩ = *0*

⟨*proof*⟩

**lemma** *sk-sq*: *cnj sk* ∗ *sk* = *1* / *8*
⟨*proof*⟩

**lemma** *shor-cw-inner*: ⟨*shor-cw p* | *shor-cw p'*⟩ = (*if p* = *p' then 1 else 0*)
⟨*proof*⟩

## 51.12 The code space and its projection

**definition** *shor-V* :: *complex Matrix.mat* **where** *shor-V* = *mat-of-cols 512* [*shor-cw True*, *shor-cw False*]

**definition** *shor-P* :: *complex Matrix.mat* **where** *shor-P* = *shor-V* ∗ (*shor-V*$^\dagger$)

**definition** *shor-state* :: *complex* ⇒ *complex* ⇒ *complex Matrix.vec* **where**
*shor-state* α β = α ·$_v$ *shor-cw True* + β ·$_v$ *shor-cw False*

**lemma** *shor-state-dim* [*simp*]: *dim-vec* (*shor-state* α β) = *512*
⟨*proof*⟩

**lemma** *shor-V-carrier*: *shor-V* ∈ *carrier-mat 512 2*
⟨*proof*⟩

**lemma** *shor-orthonormal*: *orthonormal-list 512* [*shor-cw True*, *shor-cw False*]
⟨*proof*⟩

**lemma** *shor-VdV*: (*shor-V*$^\dagger$) ∗ *shor-V* = *1*$_m$ *2*
⟨*proof*⟩

**lemma** *mat-vec-unit-col*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *M*: *M* ∈ *carrier-mat n k* **and** *a*: *a* < *k*
**shows** *M* ∗$_v$ *unit-vec k a* = *Matrix.col M a*
⟨*proof*⟩

**lemma** *shor-V-col*: *a* < *2* ⟹ *Matrix.col shor-V a* = [*shor-cw True*, *shor-cw False*] ! *a*
⟨*proof*⟩

**lemma** *shor-V-unit*: *shor-V* ∗$_v$ *unit-vec 2 0* = *shor-cw True shor-V* ∗$_v$ *unit-vec 2 1* = *shor-cw False*
⟨*proof*⟩

**lemma** *shor-P-projection*: *projection 512 shor-P*
⟨*proof*⟩

**lemma** *shor-P-cw*: *shor-P* ∗$_v$ *shor-cw p* = *shor-cw p*
⟨*proof*⟩

**lemma** *shor-P-fixes*: *shor-P* ∗$_v$ *shor-state* α β = *shor-state* α β
⟨*proof*⟩

## 51.13 Proposition (Day 33): orthogonal images, and $Z_1 v = Z_2 v = Z_3 v$

**lemma** *gram-entry*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A*: *A* ∈ *carrier-mat n k* **and** *B*: *B* ∈ *carrier-mat n k* **and** *a*: *a* < *k* **and** *b*: *b* < *k*
**shows** ((*A*$^\dagger$) ∗ *B*) $$ (*a*,*b*) = ⟨*Matrix.col A a* | *Matrix.col B b*⟩
⟨*proof*⟩

**lemma** *sandwich-VVd*:
**fixes** $V$ $W$ :: *complex Matrix.mat*
**assumes** $V$: $V \in$ *carrier-mat n k* **and** $W$: $W \in$ *carrier-mat n n*
**shows** $(V * (V^\dagger)) * W * (V * (V^\dagger)) = V * ((V^\dagger) * W * V) * (V^\dagger)$
⟨*proof*⟩

**lemma** *qubit-op-unitary*: *unitary U* ⟹ *unitary* (*qubit-op i U*)
⟨*proof*⟩

**lemma** *shor-errors-unitary*: $U \in$ *set shor-errors* ⟹ $U \in$ *carrier-mat 512 512* ∧ *unitary U*
⟨*proof*⟩

**theorem** *shor-orthogonal-images*:
**assumes** *i*: $i < 22$ **and** *j*: $j < 22$ **and** *ij*: $i \neq j$
**shows** *shor-P* $*$ (((*shor-errors* ! *i*)$^\dagger$) $*$ *shor-errors* ! *j*) $*$ *shor-P* = $0_m$ *512 512*
⟨*proof*⟩

**lemma** *shor-recovery-setting*: *recovery-setting 512 shor-P shor-errors*
⟨*proof*⟩

**lemma** *qubit-op-Z-cw*:
**assumes** *i*: $i < 9$
**shows** *qubit-op i Pauli-Z* $*_v$ *shor-cw p* = *qubit-op* (*3* $*$ (*i div 3*)) *Pauli-Z* $*_v$ *shor-cw p*
⟨*proof*⟩

**theorem** *shor-Z-coincide*:
**assumes** *i*: $i < 9$ **and** *j*: $j < 9$ **and** *ij*: *i div 3* = *j div 3*
**shows** *qubit-op i Pauli-Z* $*_v$ *shor-state* $\alpha$ $\beta$ = *qubit-op j Pauli-Z* $*_v$ *shor-state* $\alpha$ $\beta$
⟨*proof*⟩

The Day 32 proposition as first stated (distinct 1-Paulis map $V$ to orthogonal subspaces) is false: $Z_1 \neq Z_2$, yet $Z_1|0_L\rangle = Z_2|0_L\rangle \neq 0$. The notes correct it on Day 33.

**lemma** *shor-notes-Z1-Z2*:
*qubit-op 0 Pauli-Z* ≠ *qubit-op 1 Pauli-Z*
⟨*qubit-op 0 Pauli-Z* $*_v$ *shor-cw True* | *qubit-op 1 Pauli-Z* $*_v$ *shor-cw True*⟩ = *1*
⟨*proof*⟩

## 51.14 Shor's theorem

**lemma** *pauli-decomp*:
**fixes** $U$ :: *complex Matrix.mat*
**assumes** $U$: $U \in$ *carrier-mat 2 2*
**shows** $U$ = ((*U* $$ (*0,0*) + *U* $$ (*1,1*)) / *2*) $\cdot_m$ $1_m$ *2* + ((*U* $$ (*0,1*) + *U* $$ (*1,0*)) / *2*) $\cdot_m$ *Pauli-X*
+ (−i $*$ (*U* $$ (*0,1*) − *U* $$ (*1,0*)) / *2*) $\cdot_m$ *Pauli-Y* + ((*U* $$ (*0,0*) − *U* $$ (*1,1*)) / *2*) $\cdot_m$ *Pauli-Z*
(**is** $U$ = *?R*)
⟨*proof*⟩

**lemma** *qubit-op-add*:
**assumes** $A$: $A \in$ *carrier-mat 2 2* **and** $B$: $B \in$ *carrier-mat 2 2*
**shows** *qubit-op i* (*A* + *B*) = *qubit-op i A* + *qubit-op i B*
⟨*proof*⟩

**lemma** *qubit-op-smult*:
**assumes** $A$: $A \in$ *carrier-mat 2 2*
**shows** *qubit-op i* (*k* $\cdot_m$ *A*) = *k* $\cdot_m$ *qubit-op i A*
⟨*proof*⟩

**lemma** *qubit-op-one*:
**assumes** *i*: $i < 9$
**shows** *qubit-op i* $(1_m\ 2) = 1_m\ 512$
⟨*proof*⟩

**lemma** *smult-mat-mult-vec-c*:
**fixes** $A$ :: *complex Matrix.mat*
**assumes** *dim-col* $A$ = *dim-vec* $v$
**shows** $(k \cdot_m A) *_v v = k \cdot_v (A *_v v)$
⟨*proof*⟩

**lemma** *shor-errors-nth*:
**assumes** *i*: $i < 9$ **and** *b*: $b < 3$
**shows** *shor-errors* ! 0 = $1_m$ 512 *shor-errors* ! *Suc i* = *qubit-op i Pauli-X*
*shor-errors* ! (10 + *i*) = *qubit-op i Pauli-Y shor-errors* ! (19 + *b*) = *qubit-op* (3 ∗ *b*) *Pauli-Z*
⟨*proof*⟩

Shor: one CPTP recovery map $R$ corrects every single-qubit unitary error on every qubit, for every state $\psi = a|0_L\rangle + b|1_L\rangle$ of the code space.

**theorem** *shor-code-theorem*:
∃ *Rs. kraus-operators 512 512 Rs* ∧
(∀ *U i* $\alpha$ $\beta$. *U* ∈ *carrier-mat 2 2* ⟶ *unitary U* ⟶ *i* < 9 ⟶
*kraus-map-dim 512 Rs* (*qubit-op i U* ∗ (*shor-state* $\alpha$ $\beta$ ⊘ *shor-state* $\alpha$ $\beta$) ∗ ((*qubit-op i U*)†))
= *shor-state* $\alpha$ $\beta$ ⊘ *shor-state* $\alpha$ $\beta$)
⟨*proof*⟩

**end**
**theory** *Day34-Knill-Laflamme*
**imports** *Day22-Minimal-Kraus*
**begin**

# 52 Day 34: the Knill–Laflamme conditions (D34-1, D34-2, D34-3)

## 52.1 Vectors in the range of a projection; rank-one decompositions inside $V$

**lemma** *projection-fixed-of-orth-kernel*:
**assumes** *P*: *projection n P* **and** *vd*: *dim-vec v* = *n*
**and** *orth*: ⋀*x*. *dim-vec x* = *n* ⟹ *P* ∗$_v$ *x* = $0_v$ *n* ⟹ ⟨*v* | *x*⟩ = 0
**shows** *P* ∗$_v$ *v* = *v*
⟨*proof*⟩

**lemma** *outer-sum-Nil*: *outer-sum n* [] = $0_m$ *n n*
⟨*proof*⟩

**lemma** *outer-sum-singleton-eq*:
**assumes** *vd*: *dim-vec v* = *n*
**shows** *outer-sum n* [*v*] = *v* ⊘ *v*
⟨*proof*⟩

**lemma** *outer-sum-as-foldr*:
**assumes** ∀ *v* ∈ *set vs*. *dim-vec v* = *n*
**shows** *outer-sum n vs* = *foldr* (λ*v acc*. *v* ⊘ *v* + *acc*) *vs* ($0_m$ *n n*)
⟨*proof*⟩

For $X \geq 0$ the matrix $PXP$ is a sum of $|\psi_l\rangle\langle\psi_l|$ with $\psi_l \in V$ (the notes: "non-zero eigenvectors

of $PXP$ are all in $V$").

**lemma** *psd-PXP-decomp-in-range*:
**assumes** $P$: *projection n P* **and** $Xc$: $M \in$ *carrier-mat n n* **and** $Xpsd$: *psd M*
**shows** $\exists\, vs ::$ *complex Matrix.vec list*. $(\forall\, v \in$ *set vs*. *dim-vec v* $= n \wedge P *_v v = v) \wedge$
$P * M * P =$ *foldr* $(\lambda v$ *acc*. $v \oslash v +$ *acc*$)$ *vs* $(0_m\ n\ n)$
⟨*proof*⟩

## 52.2 Linearity facts for the composite map $\mathcal{R} \circ \mathbb{E}$

**definition** *RE* :: *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*RE n As Es M* = *kraus-map-dim n As* (*kraus-map-dim n Es M*)

**lemma** *RE-carrier*: **assumes** *As-c*: $\forall\, A \in$ *set As*. $A \in$ *carrier-mat n n* **and** *Es-c*: $\forall\, E \in$ *set Es*. $E \in$ *carrier-mat n n* **and** *M*: $M \in$ *carrier-mat n n* **shows** *RE n As Es M* $\in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *RE-add*: **assumes** *As-c*: $\forall\, A \in$ *set As*. $A \in$ *carrier-mat n n* **and** *Es-c*: $\forall\, E \in$ *set Es*. $E \in$ *carrier-mat n n* **and** *M*: $M \in$ *carrier-mat n n* **and** *N*: $N \in$ *carrier-mat n n* **shows** *RE n As Es* $(M +$ $N) =$ *RE n As Es M* + *RE n As Es N*
⟨*proof*⟩

**lemma** *RE-smult*: **assumes** *As-c*: $\forall\, A \in$ *set As*. $A \in$ *carrier-mat n n* **and** *Es-c*: $\forall\, E \in$ *set Es*. $E \in$ *carrier-mat n n* **and** *M*: $M \in$ *carrier-mat n n* **shows** *RE n As Es* $(c \cdot_m M) = c \cdot_m$ *RE n As Es M*
⟨*proof*⟩

**lemma** *RE-minus*:
**assumes** *As-c*: $\forall\, A \in$ *set As*. $A \in$ *carrier-mat n n* **and** *Es-c*: $\forall\, E \in$ *set Es*. $E \in$ *carrier-mat n n*
**and** *Xc*: $M \in$ *carrier-mat n n* **and** *Yc*: $N \in$ *carrier-mat n n*
**shows** *RE n As Es* $(M - N) =$ *RE n As Es M* − *RE n As Es N*
⟨*proof*⟩

**lemma** *RE-foldr*:
**assumes** *As-c*: $\forall\, A \in$ *set As*. $A \in$ *carrier-mat n n* **and** *Es-c*: $\forall\, E \in$ *set Es*. $E \in$ *carrier-mat n n* **and** $\forall\, v \in$ *set vs*. *dim-vec v* $= n$
**shows** *RE n As Es* (*foldr* $(\lambda v$ *acc*. $v \oslash v +$ *acc*$)$ *vs* $(0_m\ n\ n)) =$ *foldr* $(\lambda v$ *acc*. *RE n As Es* $(v \oslash v)$ $+$ *acc*$)$ *vs* $(0_m\ n\ n)$
⟨*proof*⟩

D34-1: $\mathcal{R} \circ \mathbb{E}$ fixes $|\psi\rangle\langle\psi|$ for all $\psi \in V$ iff it fixes $PXP$ for all $X$.

**proposition** *recovery-states-iff-PXP*:
**assumes** *As-c*: $\forall\, A \in$ *set As*. $A \in$ *carrier-mat n n* **and** *Es-c*: $\forall\, E \in$ *set Es*. $E \in$ *carrier-mat n n* **and** *P*: *projection n P*
**shows** $(\forall\, \psi.$ *dim-vec* $\psi = n \longrightarrow P *_v \psi = \psi \longrightarrow$ *RE n As Es* $(\psi \oslash \psi) = \psi \oslash \psi)$
$\longleftrightarrow (\forall\, X.\ X \in$ *carrier-mat n n* $\longrightarrow$ *RE n As Es* $(P * X * P) = P * X * P)$
⟨*proof*⟩

## 52.3 Knill–Laflamme, Theorem 1, necessity: $PE_i^* E_j P = \alpha_{ij} P$

**lemma** *lin-indep-complex-mat-single*:
**assumes** *Pc*: $P \in$ *carrier-mat d n* **and** *Pnz*: $P \neq 0_m\ d\ n$
**shows** *lin-indep-complex-mat d n* $[P]$
⟨*proof*⟩

The core of the necessity proof: by Choi's Theorem 2 applied to the two Kraus representations $\{P\}$ and $\{A_l E_i P\}$ of $X \mapsto PXP$, every $A_l E_i P$ is a multiple of $P$.

**lemma** *kl-kraus-scalars*:

**assumes** *P*: *projection n P* **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *npos*: $0 < n$
**and** *As-c*: $\forall A \in set\ As.\ A \in carrier\text{-}mat\ n\ n$ **and** *Es-c*: $\forall E \in set\ Es.\ E \in carrier\text{-}mat\ n\ n$
**and** *corr*: $\forall X.\ X \in carrier\text{-}mat\ n\ n \longrightarrow$
*kraus-map-dim n As* (*kraus-map-dim n Es* $(P * X * P)) = P * X * P$
**shows** $\exists \beta :: nat \Rightarrow nat \Rightarrow complex.\ \forall l < length\ As.\ \forall i < length\ Es.$
$As\ !\ l * Es\ !\ i * P = \beta\ l\ i \cdot_m P$
⟨*proof*⟩

**theorem** *knill-laflamme-necessity*:
**assumes** *P*: *projection n P* **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *npos*: $0 < n$
**and** *As*: *kraus-operators n n As* **and** *Es*: *kraus-operators n n Es*
**and** *corr*: $\forall X.\ X \in carrier\text{-}mat\ n\ n \longrightarrow$
*kraus-map-dim n As* (*kraus-map-dim n Es* $(P * X * P)) = P * X * P$
**shows** $\exists \alpha :: nat \Rightarrow nat \Rightarrow complex.\ \forall i < length\ Es.\ \forall j < length\ Es.$
$P * (((Es\ !\ i)^\dagger) * Es\ !\ j) * P = \alpha\ i\ j \cdot_m P$
⟨*proof*⟩

## 52.4 Kraus lists consisting of multiples of one operator

**lemma** *kraus-map-dim-scalar-list*:
**assumes** *Pc*: $P \in carrier\text{-}mat\ n\ n$ **and** *Xc*: $M \in carrier\text{-}mat\ n\ n$
**shows** *kraus-map-dim n* (*map* $(\lambda\gamma.\ \gamma \cdot_m P)$ *gs*) *M*
$= sum\text{-}list\ (map\ (\lambda\gamma.\ cnj\ \gamma * \gamma)\ gs) \cdot_m (P * M * (P^\dagger))$
⟨*proof*⟩

**lemma** *sum-wdag-w-scalar-list*:
**assumes** *Pc*: $P \in carrier\text{-}mat\ n\ n$
**shows** *sum-wdag-w n* (*map* $(\lambda\gamma.\ \gamma \cdot_m P)$ *gs*) = *sum-list* (*map* $(\lambda\gamma.\ cnj\ \gamma * \gamma)$ *gs*) $\cdot_m ((P^\dagger) * P)$
⟨*proof*⟩

**lemma** *smult-eq-self-imp-one*:
**fixes** *P* :: *complex Matrix.mat*
**assumes** *Pc*: $P \in carrier\text{-}mat\ n\ n$ **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *eq*: $s \cdot_m P = P$
**shows** $s = 1$
⟨*proof*⟩

## 52.5 Theorem 2

If $\mathcal{R} \circ \mathbb{E}$ fixes $PXP$ and $G_i = \sum_j g_{ij} E_j$ with $\sum_i G_i^* G_i = I$, then $\mathcal{R} \circ \mathbb{G}$ fixes $PXP$ as well (same $\mathcal{R}$).

**theorem** *knill-laflamme-closure*:
**assumes** *P*: *projection n P* **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *npos*: $0 < n$
**and** *As*: *kraus-operators n n As*
**and** *Es-c*: $\forall E \in set\ Es.\ E \in carrier\text{-}mat\ n\ n$
**and** *corr*: $\forall X.\ X \in carrier\text{-}mat\ n\ n \longrightarrow$
*kraus-map-dim n As* (*kraus-map-dim n Es* $(P * X * P)) = P * X * P$
**and** *Gc*: $G \in carrier\text{-}mat\ t\ (length\ Es)$
**and** *Gs-tp*: *sum-wdag-w n* (*kraus-mix n n G Es*) $= 1_m\ n$
**shows** $\forall X.\ X \in carrier\text{-}mat\ n\ n \longrightarrow$
*kraus-map-dim n As* (*kraus-map-dim n* (*kraus-mix n n G Es*) $(P * X * P)) = P * X * P$
⟨*proof*⟩

**end**
**theory** *Day34-KL-Sufficiency*
**imports** *Day34-Knill-Laflamme*
**begin**

D34-2, sufficiency: if $PE_i^* E_j P = \alpha_{ij} P$ then a recovery channel exists. Diagonalise $(\alpha_{ij}) = u\,\mathrm{diag}(d)\,u^*$; the errors $F_k = \sum_i u_{ik} E_i$ give the same channel and satisfy $PF_k^* F_l P = \delta_{kl} d_k P$, so the $F_k/\sqrt{d_k}$ with $d_k > 0$ satisfy the hypotheses of the recovery proposition (*recovery-setting*).

**lemma** *smult-cancel-nonzero*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** *Pc*: $P \in$ *carrier-mat n n* **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *eq*: $c \cdot_m P = d \cdot_m P$
**shows** $c = d$
⟨*proof*⟩

**lemma** *one-smult-mat'*: $(1{::}complex) \cdot_m M = M$
⟨*proof*⟩

**lemma** *smult-mat-mult-vec'*:
**fixes** $A$ :: *complex Matrix.mat*
**assumes** *dim-col A* = *dim-vec v*
**shows** $(k \cdot_m A) *_v v = k \cdot_v (A *_v v)$
⟨*proof*⟩

**lemma** *outer-prod-smult-vec*:
**fixes** $v$ :: *complex Matrix.vec*
**shows** $(s \cdot_v v) \oslash (s \cdot_v v) = (s * cnj\ s) \cdot_m (v \oslash v)$
⟨*proof*⟩

**lemma** *kraus-map-dim-foldr-add*:
**assumes** *Ws-c*: $\forall W \in set\ Ws.\ W \in carrier\text{-}mat\ n\ n$ **and** *fc*: $\forall k \in set\ xs.\ f\ k \in carrier\text{-}mat\ n\ n$
**shows** *kraus-map-dim n Ws* (*foldr* $(\lambda k\ acc.\ f\ k + acc)$ *xs* $(0_m\ n\ n)$)
$=$ *foldr* $(\lambda k\ acc.$ *kraus-map-dim n Ws* $(f\ k) + acc)$ *xs* $(0_m\ n\ n)$
⟨*proof*⟩

**lemma** *kraus-map-dim-outer-as-foldr*:
**assumes** *Ws-c*: $\forall W \in set\ Ws.\ W \in carrier\text{-}mat\ n\ n$ **and** $\psi d$: *dim-vec* $\psi = n$
**shows** *kraus-map-dim n Ws* $(\psi \oslash \psi)$
$=$ *foldr* $(\lambda k\ acc.\ (Ws\ !\ k *_v \psi) \oslash (Ws\ !\ k *_v \psi) + acc)$ $[0..<length\ Ws]$ $(0_m\ n\ n)$
⟨*proof*⟩

**lemma** *foldr-smult-sandwich*:
**fixes** $V$ :: *nat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *Mc*: $M \in carrier\text{-}mat\ n\ n$ **and** *Nc*: $N \in carrier\text{-}mat\ n\ n$ **and** *Pc*: $P \in carrier\text{-}mat\ n\ n$
**and** *Vc*: $\forall j < r.\ V\ j \in carrier\text{-}mat\ n\ n$ **and** *h*: $\forall j < r.\ M * V\ j * N = \beta\ j \cdot_m P$
**shows** $M *$ *foldr* $(\lambda j\ acc.\ b\ j \cdot_m V\ j + acc)$ $[0..<r]$ $(0_m\ n\ n) * N = (\sum j{<}r.\ b\ j * \beta\ j) \cdot_m P$
⟨*proof*⟩

**lemma** *foldr-smult-dagger*:
**fixes** $V$ :: *nat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *Vc*: $\forall j < r.\ V\ j \in carrier\text{-}mat\ n\ n$
**shows** (*foldr* $(\lambda j\ acc.\ b\ j \cdot_m V\ j + acc)$ $[0..<r]$ $(0_m\ n\ n))^\dagger$
$=$ *foldr* $(\lambda j\ acc.\ cnj\ (b\ j) \cdot_m ((V\ j)^\dagger) + acc)$ $[0..<r]$ $(0_m\ n\ n)$
⟨*proof*⟩

The bracket $P\,(\sum_i a_i E_i)^* (\sum_j b_j E_j)\,P$ under the Knill–Laflamme conditions.

**lemma** *kl-mix-bracket*:
**assumes** *Es-c*: $\forall E \in set\ Es.\ E \in carrier\text{-}mat\ n\ n$ **and** *Pc*: $P \in carrier\text{-}mat\ n\ n$
**and** *kl*: $\forall i < length\ Es.\ \forall j < length\ Es.\ P * (((Es\ !\ i)^\dagger) * Es\ !\ j) * P = \alpha\ i\ j \cdot_m P$
**and** *r*: $r = length\ Es$
**shows** $P *$ $(((foldr\ (\lambda i\ acc.\ a\ i \cdot_m (Es\ !\ i) + acc)\ [0..<r]\ (0_m\ n\ n))^\dagger)$
$*$ *foldr* $(\lambda j\ acc.\ b\ j \cdot_m (Es\ !\ j) + acc)$ $[0..<r]$ $(0_m\ n\ n)) * P$
$= (\sum j{<}r.\ b\ j * (\sum i{<}r.\ cnj\ (a\ i) * \alpha\ i\ j)) \cdot_m P$

⟨*proof*⟩

**lemma** *dagger-sandwich*:
**assumes** *Pc*: $P \in$ *carrier-mat n n* **and** *Ph*: $P^\dagger = P$ **and** *Ac*: $A \in$ *carrier-mat n n* **and** *Bc*: $B \in$ *carrier-mat n n*
**shows** $(P * ((A^\dagger) * B) * P)^\dagger = P * ((B^\dagger) * A) * P$
⟨*proof*⟩

**lemma** *smult-bracket*:
**assumes** *Pc*: $P \in$ *carrier-mat n n* **and** *Ac*: $A \in$ *carrier-mat n n* **and** *Bc*: $B \in$ *carrier-mat n n*
**shows** $P * (((x \cdot_m A)^\dagger) * (y \cdot_m B)) * P = (cnj\ x * y) \cdot_m (P * ((A^\dagger) * B) * P)$
⟨*proof*⟩

**theorem** *knill-laflamme-sufficiency*:
**assumes** *P*: *projection n P* **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *npos*: $0 < n$
**and** *Es*: *kraus-operators n n Es*
**and** *kl*: $\forall\, i < length\ Es.\ \forall\, j < length\ Es.\ P * (((Es\ !\ i)^\dagger) * Es\ !\ j) * P = \alpha\ i\ j \cdot_m P$
**shows** $\exists\, As.$ *kraus-operators n n As* $\wedge$
$(\forall\, X.\ X \in$ *carrier-mat n n* $\longrightarrow$ *kraus-map-dim n As* (*kraus-map-dim n Es* $(P * X * P)) = P * X * P)$
⟨*proof*⟩

D34-2 (Theorem 1): the code is correctable iff $PE_i^* E_j P = \alpha_{ij} P$.

**theorem** *knill-laflamme*:
**assumes** *P*: *projection n P* **and** *Pnz*: $P \neq 0_m\ n\ n$ **and** *npos*: $0 < n$
**and** *Es*: *kraus-operators n n Es*
**shows** ($\exists\, As.$ *kraus-operators n n As* $\wedge$
$(\forall\, X.\ X \in$ *carrier-mat n n* $\longrightarrow$ *kraus-map-dim n As* (*kraus-map-dim n Es* $(P * X * P)) = P * X * P)$)
$\longleftrightarrow$ $(\exists\, \alpha :: nat \Rightarrow nat \Rightarrow complex.\ \forall\, i < length\ Es.\ \forall\, j < length\ Es.$
$P * (((Es\ !\ i)^\dagger) * Es\ !\ j) * P = \alpha\ i\ j \cdot_m P)$
⟨*proof*⟩

**end**

# Part III

# Lectures 23–26: extreme points, operator systems and Arveson's theorem

**theory** *Day23-ChoiExtremeUCP*
**imports**
*Quantum-Computing11*
**begin**

## 53 Day 23 (19 October 2011): Choi's Theorem 3, extreme points of UCP and CPTP

Day 22 stated and Day 23 proves:

**Theorem (notes)** (Choi 3)**.** If $\Phi : M_n \to M_d$ is unital and completely positive, $\Phi(X) = \sum_{i=1}^r V_i^* X V_i$ with $r = \mathrm{cr}(\Phi)$, then $\Phi$ is extreme in $UCP(M_n, M_d)$ if and only if $\{V_i^* V_j : 1 \leq$

$i, j \leq r\} \subseteq M_d$ is linearly independent.

**Corollary (notes).** If $\Phi : M_d \to M_n$ is completely positive and trace preserving, $\Phi(X) = \sum_{i=1}^{r} V_i X V_i^*$ with $r = \mathrm{cr}(\Phi)$, then $\Phi$ is extreme in $CPTP(M_d, M_n)$ iff $\{V_i^* V_j\}$ is linearly independent (apply the theorem to $\Phi^*$).

The notes' proof. ($\Rightarrow$) If $\sum_{ij} \lambda_{ij} V_i^* V_j = 0$ it suffices to treat Hermitian $\mu = (\mu_{ij})$ with $-I \leq \mu \leq I$: writing $I \pm \mu = (\alpha_{ij})^*(\alpha_{ij})$ and $W_i = \sum_j \alpha_{ij} V_j$ shows that $\psi_\pm(X) = \sum_i V_i^* X V_i \pm \sum_{ij} \mu_{ij} V_i^* X V_j$ are UCP with $\Phi = \frac{1}{2}(\psi_+ + \psi_-)$, so extremality gives $\psi_+ = \Phi$, and Choi 2 (unitary freedom) forces $I + \mu = U^* U = I$, i.e. $\mu = 0$. ($\Leftarrow$) If $\Phi = \frac{1}{2}(\psi_1 + \psi_2)$ with $\psi_k(X) = \sum_p W_p^* X W_p$, Choi 2 gives $W_p/\sqrt{2} = \sum_j \alpha_{pj} V_j$ with $U^* U = I_r$; unitality of $\psi_1$ and linear independence of the $V_i^* V_j$ force $U_1^* U_1 = \frac{1}{2} I_r$, whence $\psi_1 = \Phi = \psi_2$. Examples: $X \mapsto U^* X U$ is extreme; $\frac{1}{2}(U_1^* X U_1 + U_2^* X U_2)$, $X \mapsto \mathrm{tr}(X) I_d$ and $X \mapsto \mathrm{diag}(X)$ are not.

What is formalised here is the corollary, i.e. the CPTP form, for a fixed Kraus list *Vs* of linearly independent operators (the consequence of $r = \mathrm{cr}(\Phi)$ proved on Day 22): extremality (in the carrier sense *extreme-CPTP-carrier*) holds if and only if the products $V_j^\dagger V_i$ are linearly independent over $\mathbb{C}$ (*choi-theorem-3-cptp-iff*). The direction ($\Rightarrow$) follows the notes' perturbation argument (*kraus-quad-map*, *small-eps-psd*); the direction ($\Leftarrow$) replaces Choi 2 by a CP Radon–Nikodym theorem proved below from the Moore–Penrose pseudoinverse of the Choi matrix. Unitary freedom itself is *choi-2-unitary-freedom* in *Day21-Unitary-Freedom*, proved afterwards and independently of this theory; the examples are in theory Day23_Extremality_Examples (the UCP form is not formalised).

## 53.1 Extreme UCP maps

An *extreme UCP map* is one that cannot be written as a proper convex combination of two distinct UCP maps. Following Day-22's *extreme-point-mat*, we use the half-half mid-point formulation; this is equivalent to the standard definition for convex sets.

**definition** *extreme-UCP* ::
  *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool*
**where**
  *extreme-UCP n d* Φ ≡
    Φ ∈ *UCP n d* ∧
    ¬ (∃ Φ₁ ∈ *UCP n d*. ∃ Φ₂ ∈ *UCP n d*.
        Φ₁ ≠ Φ ∧ Φ ≠ Φ₂ ∧
        (∀ *X*. *X* ∈ *carrier-mat n n*
            ⟶ Φ *X* = (*1/2* :: *complex*) ·ₘ Φ₁ *X*
                + (*1/2* :: *complex*) ·ₘ Φ₂ *X*))

**lemma** *extreme-UCP-member*:
  **assumes** *extreme-UCP n d* Φ
  **shows** Φ ∈ *UCP n d*
  ⟨*proof*⟩

Trivial unfolding: an extreme UCP map satisfies no non-trivial mid-point decomposition.

**lemma** *extreme-UCP-no-split*:
  **assumes** *ext*: *extreme-UCP n d* Φ
    **and** Φ*1-in*: Φ₁ ∈ *UCP n d*
    **and** Φ*2-in*: Φ₂ ∈ *UCP n d*
    **and** Φ*1-neq*: Φ₁ ≠ Φ
    **and** Φ*2-neq*: Φ ≠ Φ₂
  **shows** ∃ *X* ∈ *carrier-mat n n*.
        Φ *X* ≠ (*1/2* :: *complex*) ·ₘ Φ₁ *X*
            + (*1/2* :: *complex*) ·ₘ Φ₂ *X*
  ⟨*proof*⟩

## 53.2 UCP-Kraus representation

A *UCP Kraus representation* for a map $\Phi : M_n \to M_d$ is a list $Vs$ of $d \times n$ matrices such that

- each $V \in Vs$ is $d \times n$;
- the unitality identity $\sum_i V_i\ V_i^\dagger = I_d$ holds;
- $\Phi\ X = \sum_i V_i\ X\ V_i^\dagger$ for every $X \in M_n$.

Note that the standard *kraus-operators* predicate of Day 17 uses the *trace-preserving* identity $\sum V_i^\dagger\ V_i = I_n$; the *unital* form $\sum V_i\ V_i^\dagger = I_d$ is the Heisenberg-dual condition appropriate for *UCP*.

**definition** *ucp-kraus-rep* ::
  *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒
    *complex Matrix.mat list* ⇒ *bool*
**where**
  *ucp-kraus-rep n d Φ Vs* ≡
    (∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*) ∧
    *foldr* (λ*V acc*. *V* ∗ (*V*†) + *acc*) *Vs* (*0*$_m$ *d d*) = *1*$_m$ *d* ∧
    (∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ *X* = *kraus-map-dim d Vs X*)

The *products set* of a Kraus list $Vs$ is the list of $n \times n$ matrices $V_j^\dagger\ V_i$ for all pairs of indices $(i,j)$.

**definition** *kraus-products* ::
  *complex Matrix.mat list* ⇒ *complex Matrix.mat list*
**where**
  *kraus-products Vs* ≡
    *concat* (*map* (λ*i*. *map* (λ*j*. (*Vs*!*j*)† ∗ *Vs*!*i*) [*0*..<*length Vs*])
          [*0*..<*length Vs*])

Real linear independence for a list of complex matrices. A list $Bs$ is independent over $\mathbb{R}$ iff the only $\mathbb{R}$-coefficients making $\sum_k c_k\ B_k = 0$ are all zero. We use the real-coefficient form because Choi's perturbation argument constructs real combinations of hermitian generators.

**definition** *mat-lin-indep-real* ::
  *nat* ⇒ *complex Matrix.mat list* ⇒ *bool*
**where**
  *mat-lin-indep-real n Bs* ≡
    (∀ *B* ∈ *set Bs*. *B* ∈ *carrier-mat n n*) ∧
    (∀ *c* :: *nat* ⇒ *real*.
      *foldr* (λ*k acc*. *complex-of-real* (*c k*) ·$_m$ (*Bs*!*k*) + *acc*)
          [*0*..<*length Bs*] (*0*$_m$ *n n*)
        = *0*$_m$ *n n*
      ⟶ (∀ *k* < *length Bs*. *c k* = *0*))

Complex linear independence: the classical Choi statement uses **complex**-coefficient linear independence. This is strictly stronger than *mat-lin-indep-real*.

**definition** *mat-lin-indep-complex* ::
  *nat* ⇒ *complex Matrix.mat list* ⇒ *bool*
**where**
  *mat-lin-indep-complex n Bs* ≡
    (∀ *B* ∈ *set Bs*. *B* ∈ *carrier-mat n n*) ∧
    (∀ *c* :: *nat* ⇒ *complex*.
      *foldr* (λ*k acc*. *c k* ·$_m$ (*Bs*!*k*) + *acc*)
          [*0*..<*length Bs*] (*0*$_m$ *n n*)
        = *0*$_m$ *n n*

$\longrightarrow (\forall k < \mathit{length}\ \mathit{Bs}.\ c\ k = 0))$

Complex linear independence implies real.

**lemma** *mat-lin-indep-complex-imp-real*:
**assumes** *mat-lin-indep-complex n Bs*
**shows** *mat-lin-indep-real n Bs*
⟨*proof*⟩

## 53.3 Unital condition from the Kraus list

If *Ws* is a list of $d \times n$ matrices, then evaluating the Kraus map at the identity $1_m\ n$ gives exactly the sum $\sum\ W\ W^\dagger$. Combined with unitality, this gives the $\sum\ W\ W^\dagger = 1_m\ d$ identity that distinguishes UCP from plain CP.

**lemma** *kraus-map-dim-one*:
**assumes** $\forall\, W \in \mathit{set}\ \mathit{Ws}.\ W \in \mathit{carrier\text{-}mat}\ d\ n$
**shows** *kraus-map-dim d Ws* $(1_m\ n)$
$= \mathit{foldr}\ (\lambda W\ \mathit{acc}.\ W * (W^\dagger) + \mathit{acc})\ \mathit{Ws}\ (0_m\ d\ d)$
⟨*proof*⟩

## 53.4 UCP-Kraus existence

Every linear UCP map admits a UCP Kraus representation of length $n\ d$. Linearity, scalar-multiplication and range hypotheses match those of *kraus-rep-of-psd-choi*; the unital condition $\sum\ V\ V^\dagger = 1_m\ d$ is supplied by $\Phi\ (1_m\ n) = 1_m\ d$ + *kraus-map-dim-one*.

**lemma** *ucp-has-kraus-rep*:
**fixes** $\Phi$ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *linear*: $\bigwedge A\ B.\ A \in \mathit{carrier\text{-}mat}\ n\ n \Longrightarrow B \in \mathit{carrier\text{-}mat}\ n\ n \Longrightarrow$
$\Phi\ (A + B) = \Phi\ A + \Phi\ B$
**and** *smult*: $\bigwedge c\ A.\ A \in \mathit{carrier\text{-}mat}\ n\ n \Longrightarrow \Phi\ (c \cdot_m A) = c \cdot_m \Phi\ A$
**and** *range*: $\forall\, A.\ A \in \mathit{carrier\text{-}mat}\ n\ n \longrightarrow \Phi\ A \in \mathit{carrier\text{-}mat}\ d\ d$
**and** *npos*: $0 < n$ **and** *dpos*: $0 < d$
**and** *ucp*: $\Phi \in \mathit{UCP}\ n\ d$
**shows** $\exists\ \mathit{Vs}$ :: *complex Matrix.mat list*.
*ucp-kraus-rep n d* $\Phi$ *Vs* $\wedge$ *length Vs* $= n * d$
⟨*proof*⟩

## 53.5 Sanity facts about *kraus-products*

**lemma** *kraus-products-carrier*:
**assumes** $\forall\, V \in \mathit{set}\ \mathit{Vs}.\ V \in \mathit{carrier\text{-}mat}\ d\ n$
**shows** $\forall\, B \in \mathit{set}\ (\mathit{kraus\text{-}products}\ \mathit{Vs}).\ B \in \mathit{carrier\text{-}mat}\ n\ n$
⟨*proof*⟩

## 53.6 Quadratic Kraus map (perturbation primitive)

For a Kraus list *Vs* of $d \times n$ matrices and a coefficient matrix $T \in M_k$ (where $k$ = *length Vs*), the *quadratic Kraus map* is (*kraus-quad-map d Vs T*) $X = \sum_{\{i,j\}}\ T_{\{ij\}}\ V_i\ X\ {V_j}^\dagger$. When $T = I_k$ this reduces to the ordinary Kraus map *kraus-map-dim d Vs* $X = \sum\ V_i\ X\ {V_i}^\dagger$; for PSD $T$ the result is again a Kraus map (Lemma A below). The perturbations $T_\pm = I \pm \varepsilon\ A$ for small $\varepsilon$ and hermitian $A$ will provide the *splittings* witnessing non-extremity.

**definition** *kraus-quad-row* ::
*nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒
*complex Matrix.mat* ⇒ *nat* ⇒ *complex Matrix.mat*
**where**
*kraus-quad-row d Vs Tm rho i* ≡

*foldr* ($\lambda j$ *acc. Tm* \$\$ (*i*, *j*) $\cdot_m$ (*Vs*!*i* $*$ *rho* $*$ ((*Vs*!*j*)$^\dagger$)) + *acc*)
[*0*..<*length Vs*] ($0_m$ *d d*)

**definition** *kraus-quad-map* ::
*nat* $\Rightarrow$ *complex Matrix.mat list* $\Rightarrow$ *complex Matrix.mat* $\Rightarrow$
*complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
**where**
*kraus-quad-map d Vs Tm rho* $\equiv$
*foldr* ($\lambda i$ *acc. kraus-quad-row d Vs Tm rho i* + *acc*)
[*0*..<*length Vs*] ($0_m$ *d d*)

Generic carrier-preservation lemma for accumulating folds over $[0..<N]$: if each per-index term lies in $M_d$, so does the fold.

**lemma** *foldr-add-carrier*:
**fixes** *f* :: *'a* $\Rightarrow$ *complex Matrix.mat*
**assumes** $\forall$ *x* $\in$ *set xs. f x* $\in$ *carrier-mat d d*
**shows** *foldr* ($\lambda x$ *acc. f x* + *acc*) *xs* ($0_m$ *d d*) $\in$ *carrier-mat d d*
⟨*proof*⟩

**lemma** *foldr-upt-add-carrier*:
**fixes** *f* :: *nat* $\Rightarrow$ *complex Matrix.mat*
**assumes** $\forall$ *k* < *N. f k* $\in$ *carrier-mat d d*
**shows** *foldr* ($\lambda k$ *acc. f k* + *acc*) [*0*..<*N*] ($0_m$ *d d*) $\in$ *carrier-mat d d*
⟨*proof*⟩

**lemma** *kraus-quad-row-carrier*:
**assumes** *Vs-c*: $\forall$ *V* $\in$ *set Vs. V* $\in$ *carrier-mat d n*
**and** *rho-c*: *rho* $\in$ *carrier-mat n n*
**and** *i-lt*: *i* < *length Vs*
**shows** *kraus-quad-row d Vs Tm rho i* $\in$ *carrier-mat d d*
⟨*proof*⟩

**lemma** *kraus-quad-map-carrier*:
**assumes** *Vs-c*: $\forall$ *V* $\in$ *set Vs. V* $\in$ *carrier-mat d n*
**and** *rho-c*: *rho* $\in$ *carrier-mat n n*
**shows** *kraus-quad-map d Vs Tm rho* $\in$ *carrier-mat d d*
⟨*proof*⟩

Per-entry index formula for the accumulating fold (with arbitrary base).

**lemma** *foldr-upt-add-acc-index*:
**fixes** *f* :: *nat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *carr*: $\forall$ *k* < *N. f k* $\in$ *carrier-mat d d*
**and** *z-c*: *z* $\in$ *carrier-mat d d*
**and** *r-lt*: *r* < *d* **and** *c-lt*: *c* < *d*
**shows** *foldr* ($\lambda k$ *acc. f k* + *acc*) [*0*..<*N*] *z* \$\$ (*r*, *c*)
= ($\sum$ *k* = *0*..<*N. f k* \$\$ (*r*, *c*)) + *z* \$\$ (*r*, *c*)
⟨*proof*⟩

**lemma** *foldr-upt-add-index*:
**fixes** *f* :: *nat* $\Rightarrow$ *complex Matrix.mat*
**assumes** *carr*: $\forall$ *k* < *N. f k* $\in$ *carrier-mat d d*
**and** *r-lt*: *r* < *d* **and** *c-lt*: *c* < *d*
**shows** *foldr* ($\lambda k$ *acc. f k* + *acc*) [*0*..<*N*] ($0_m$ *d d*) \$\$ (*r*, *c*)
= ($\sum$ *k* = *0*..<*N. f k* \$\$ (*r*, *c*))
⟨*proof*⟩

Per-entry formula for *kraus-quad-row*.

**lemma** *kraus-quad-row-index*:

**assumes** *Vs-c*: $\forall\, V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $rho \in carrier\text{-}mat\ n\ n$
**and** *i-lt*: $i < length\ Vs$
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < d$
**shows** *kraus-quad-row d Vs Tm rho i* \$\$ $(r,\ c)$
$= (\sum j = 0..<length\ Vs.$
$Tm$ \$\$ $(i,\ j) * (Vs!i * rho * ((Vs!j)^\dagger))$ \$\$ $(r,\ c))$
⟨*proof*⟩

Per-entry formula for *kraus-quad-map*: double sum over indices.

**lemma** *kraus-quad-map-index*:
**assumes** *Vs-c*: $\forall\, V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $rho \in carrier\text{-}mat\ n\ n$
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < d$
**shows** *kraus-quad-map d Vs Tm rho* \$\$ $(r,\ c)$
$= (\sum i = 0..<length\ Vs.$
$\sum j = 0..<length\ Vs.$
$Tm$ \$\$ $(i,\ j) * (Vs!i * rho * ((Vs!j)^\dagger))$ \$\$ $(r,\ c))$
⟨*proof*⟩

Carrier-preservation in a different dimension pair.

**lemma** *foldr-add-carrier-dn*:
**fixes** $f :: 'a \Rightarrow complex\ Matrix.mat$
**assumes** $\forall\, x \in set\ xs.\ f\ x \in carrier\text{-}mat\ d\ n$
**shows** $foldr\ (\lambda x\ acc.\ f\ x + acc)\ xs\ (0_m\ d\ n) \in carrier\text{-}mat\ d\ n$
⟨*proof*⟩

Linear combination of Kraus operators with complex coefficients. *kraus-combo d n Vs u* is $\sum_i u_i\ V_i$, a $d \times n$ matrix.

**definition** *kraus-combo* ::
$nat \Rightarrow nat \Rightarrow complex\ Matrix.mat\ list \Rightarrow complex\ Matrix.vec \Rightarrow$
$complex\ Matrix.mat$
**where**
*kraus-combo d n Vs u* ≡
$foldr\ (\lambda i\ acc.\ u\ \$\ i \cdot_m (Vs!i) + acc)\ [0..<length\ Vs]\ (0_m\ d\ n)$

**lemma** *kraus-combo-carrier*:
**assumes** *Vs-c*: $\forall\, V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**shows** *kraus-combo d n Vs u* $\in carrier\text{-}mat\ d\ n$
⟨*proof*⟩

Per-entry formula for *kraus-combo*: foldr index lemma in d×n.

**lemma** *foldr-upt-add-acc-index-dn*:
**fixes** $f :: nat \Rightarrow complex\ Matrix.mat$
**assumes** *carr*: $\forall\, k < N.\ f\ k \in carrier\text{-}mat\ d\ n$
**and** *z-c*: $z \in carrier\text{-}mat\ d\ n$
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < n$
**shows** $foldr\ (\lambda k\ acc.\ f\ k + acc)\ [0..<N]\ z$ \$\$ $(r,\ c)$
$= (\sum k = 0..<N.\ f\ k$ \$\$ $(r,\ c)) + z$ \$\$ $(r,\ c)$
⟨*proof*⟩

**lemma** *kraus-combo-index*:
**assumes** *Vs-c*: $\forall\, V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < n$
**shows** *kraus-combo d n Vs u* \$\$ $(r,\ c)$
$= (\sum i = 0..<length\ Vs.\ u\ \$\ i * (Vs!i)$ \$\$ $(r,\ c))$
⟨*proof*⟩

A direct per-entry formula for matrix products.

**lemma** *mat-mult-index-sum*:
**assumes** *A-c*: $A \in$ *carrier-mat p q*
**and** *B-c*: $B \in$ *carrier-mat q s*
**and** *r-lt*: $r < p$ **and** *c-lt*: $c < s$
**shows** $(A * B)$ \$\$ $(r, c) = (\sum k = 0..<q.\ A$ \$\$ $(r, k) * B$ \$\$ $(k, c))$
⟨*proof*⟩

Per-entry formula for $W * rho$, where $W =$ *kraus-combo d n Vs u*.

**lemma** *kraus-combo-times-rho-index*:
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $rho \in carrier\text{-}mat\ n\ n$
**and** *r-lt*: $r < d$ **and** *k-lt*: $k < n$
**shows** (*kraus-combo d n Vs u* $*$ *rho*) \$\$ $(r, k)$
$= (\sum i = 0..<length\ Vs.\ u\ \$\ i * (Vs!i * rho)$ \$\$ $(r, k))$
⟨*proof*⟩

The full quadratic per-entry formula: $(W\ rho\ W^\dagger)$ \$\$ $(r,c) = \sum_{\{i,j\}} u_i\ |u_j|\ (V_i\ rho\ {V_j}^\dagger)$ \$\$ $(r,c)$.

**lemma** *kraus-combo-quad-index*:
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $rho \in carrier\text{-}mat\ n\ n$
**and** *r-lt*: $r < d$ **and** *c-lt*: $c < d$
**shows** (*kraus-combo d n Vs u* $*$ *rho* $*$ ((*kraus-combo d n Vs u*)$^\dagger$)) \$\$ $(r, c)$
$= (\sum i = 0..<length\ Vs.\ \sum j = 0..<length\ Vs.$
$u\ \$\ i * cnj\ (u\ \$\ j) * (Vs!i * rho * ((Vs!j)^\dagger))$ \$\$ $(r, c))$
⟨*proof*⟩

Linearity of *kraus-quad-map* in the coefficient matrix (needed for the rank-1 decomposition below).

**lemma** *kraus-quad-map-add-Tm*:
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $rho \in carrier\text{-}mat\ n\ n$
**and** *T1-c*: $T1 \in carrier\text{-}mat\ (length\ Vs)\ (length\ Vs)$
**and** *T2-c*: $T2 \in carrier\text{-}mat\ (length\ Vs)\ (length\ Vs)$
**shows** *kraus-quad-map d Vs* (*T1* $+$ *T2*) *rho*
$=$ *kraus-quad-map d Vs T1 rho* $+$ *kraus-quad-map d Vs T2 rho*
⟨*proof*⟩

Carrier of *outer-sum k vs*.

**lemma** *outer-sum-carrier*:
*outer-sum k vs* $\in$ *carrier-mat k k*
⟨*proof*⟩

Cons-recursion for *outer-sum*: adds a rank-1 piece.

**lemma** *outer-sum-Cons-split*:
**assumes** $\forall v \in set\ (u\ \#\ us).\ dim\text{-}vec\ v = k$
**shows** *outer-sum k* $(u\ \#\ us)$
$=$ *outer-sum k* $[u]$ $+$ *outer-sum k us*
⟨*proof*⟩

Rank-1 identity: *kraus-quad-map d Vs* (*outer-sum k* $[u]$) $rho = W\ rho\ W^\dagger$ where $W =$ *kraus-combo d n Vs u*.

**lemma** *kraus-quad-map-outer-singleton*:
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *rho-c*: $rho \in carrier\text{-}mat\ n\ n$

**and** *u-dim*: *dim-vec u = length Vs*
**shows** *kraus-quad-map d Vs* (*outer-sum* (*length Vs*) [*u*]) *rho*
= *kraus-combo d n Vs u* ∗ *rho* ∗ ((*kraus-combo d n Vs u*)†)
⟨*proof*⟩

Vanishing of the quadratic map when the coefficient matrix is zero.

**lemma** *kraus-quad-map-zero-Tm*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *rho-c*: *rho* ∈ *carrier-mat n n*
**shows** *kraus-quad-map d Vs* (0$_m$ (*length Vs*) (*length Vs*)) *rho* = 0$_m$ *d d*
⟨*proof*⟩

Sum-decomposition: a PSD-style coefficient matrix *outer-sum k us* gives *kraus-quad-map* as a kraus_map_dim of the combos.

**lemma** *kraus-quad-map-outer-sum-eq-kraus-map-dim*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *rho-c*: *rho* ∈ *carrier-mat n n*
**and** *us-dim*: ∀ *v* ∈ *set us*. *dim-vec v = length Vs*
**shows** *kraus-quad-map d Vs* (*outer-sum* (*length Vs*) *us*) *rho*
= *kraus-map-dim d* (*map* (*kraus-combo d n Vs*) *us*) *rho*
⟨*proof*⟩

## 53.7 Lemma A: PSD coefficient matrix gives a CP perturbation

Since *amplification* only evaluates Φ on $n \times n$ blocks, two maps that agree on *carrier-mat n n* yield the same amplification, hence the same notion of complete positivity.

**lemma** *amplification-cong*:
**fixes** *Phi Psi* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Xm* :: *complex Matrix.mat*
**assumes** *eq*: ⋀*r*. *r* ∈ *carrier-mat n n* ⟹ *Phi r = Psi r*
**shows** *amplification n d p Phi Xm = amplification n d p Psi Xm*
⟨*proof*⟩

**lemma** *completely-positive-cong*:
**fixes** *Phi Psi* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** ⋀*r*. *r* ∈ *carrier-mat n n* ⟹ *Phi r = Psi r*
**shows** *completely-positive n d Phi* ⟷ *completely-positive n d Psi*
⟨*proof*⟩

**Lemma A**: if $T$ is PSD then *kraus-quad-map d Vs T* is completely positive.

**Proof sketch.** Use ⟦*?P* ∈ *carrier-mat ?n ?n*; *?P* ≥$_M$ *0*⟧ ⟹ ∃ *vs*. *length vs = ?n* ∧ (∀ *v*∈*set vs*. *dim-vec v = ?n*) ∧ *outer-sum ?n vs = ?P* to write $T$ = *outer-sum k us*. Then by ⟦∀ *V*∈*set ?Vs*. *V* ∈ *carrier-mat ?d ?n*; *?rho* ∈ *carrier-mat ?n ?n*; ∀ *v*∈*set ?us*. *dim-vec v = length ?Vs*⟧ ⟹ *kraus-quad-map ?d ?Vs* (*outer-sum* (*length ?Vs*) *?us*) *?rho* = *kraus-map-dim ?d* (*map* (*kraus-combo ?d ?n ?Vs*) *?us*) *?rho*, *kraus-quad-map d Vs T* agrees on *carrier-mat n n* with *kraus-map-dim d Ws*, where *Ws* := *map* (*kraus-combo d n Vs*) *us*. Apply ⟦∀ *W*∈*set ?Ws*. *W* ∈ *carrier-mat ?d ?n*; *0* < *?n*; *0* < *?d*⟧ ⟹ *completely-positive ?n ?d* (*kraus-map-dim ?d ?Ws*).

**lemma** *kraus-perturbation-cp*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *Tm-c*: *Tm* ∈ *carrier-mat* (*length Vs*) (*length Vs*)
**and** *Tm-psd*: *psd Tm*
**and** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
**shows** *completely-positive n d* (*kraus-quad-map d Vs Tm*)
⟨*proof*⟩

## 53.8 Small-$\varepsilon$ PSD perturbation of the identity

For any hermitian $A : k \times k$, the matrix $1_m\ k + \delta \cdot_m A$ is PSD whenever $|\delta|$ is small enough. The bound uses the entry-wise sum $S := \sum_{\{i,j\}} |A_{\{ij\}}|$, taking $\varepsilon := 1/(1 + S)$; this avoids any operator-norm computation and just uses Cauchy-Schwarz on entries.

Bilinearity of the inner product in the right slot.

**lemma** *inner-prod-add-right*:
**fixes** *u v w* :: *complex Matrix.vec*
**assumes** *ud*: *dim-vec u* = *k* **and** *vd*: *dim-vec v* = *k* **and** *wd*: *dim-vec w* = *k*
**shows** $\langle u \mid v + w\rangle = \langle u \mid v\rangle + \langle u \mid w\rangle$
⟨*proof*⟩

**lemma** *inner-prod-smult-right*:
**fixes** *u v* :: *complex Matrix.vec*
**assumes** *ud*: *dim-vec u* = *k* **and** *vd*: *dim-vec v* = *k*
**shows** $\langle u \mid c \cdot_v v\rangle = c * \langle u \mid v\rangle$
⟨*proof*⟩

The hermitian sum.

**lemma** *hermitian-add*:
**assumes** *Aherm*: *hermitian A* **and** *Bherm*: *hermitian B*
**and** *Ac*: $A \in$ *carrier-mat k k* **and** *Bc*: $B \in$ *carrier-mat k k*
**shows** *hermitian* $(A + B)$
⟨*proof*⟩

**lemma** *hermitian-one-mat*:
*hermitian* $(1_m\ k ::$ *complex Matrix.mat*$)$
⟨*proof*⟩

**lemma** *hermitian-smult-real*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *hermitian A A* $\in$ *carrier-mat k k*
**shows** *hermitian* (*complex-of-real d* $\cdot_m$ *A*)
⟨*proof*⟩

Entry-wise bound on the quadratic form of a hermitian matrix.

**lemma** *quad-form-bound-by-entry-sum*:
**fixes** *A* :: *complex Matrix.mat* **and** *x* :: *complex Matrix.vec*
**assumes** *A-c*: $A \in$ *carrier-mat k k* **and** *x-d*: *dim-vec x* = *k*
**shows** *cmod* $(\langle x \mid A *_v x\rangle)$
$\leq (\sum i = 0..<k.\ \sum j = 0..<k.\ cmod\ (A\ \$\$\ (i,j))) * Re\ \langle x \mid x\rangle$
⟨*proof*⟩

The main lemma: small $\varepsilon$ keeps the perturbation PSD.

**lemma** *small-eps-psd*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *A-c*: $A \in$ *carrier-mat k k* **and** *herm*: *hermitian A*
**shows** $\exists \varepsilon > 0.\ \forall d ::$ *real*. $|d| \leq \varepsilon \longrightarrow$
*psd* $(1_m\ k +$ *complex-of-real d* $\cdot_m$ *A*$)$
⟨*proof*⟩

## 53.9 Symmetric / anti-hermitian split of the coefficient matrix

The *dual* linear combination *kraus-pair-sum n Vs A* $= \sum_{\{i,j\}} A_{\{ij\}}\ V_j{}^\dagger\ V_i$ is an $n \times n$ matrix. The dependence assumption ¬ *mat-lin-indep-real n* (*kraus-products Vs*) says that some real-coefficient $A$ makes this sum vanish.

**definition** *kraus-pair-row* ::
  *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒
    *nat* ⇒ *complex Matrix.mat*
**where**
  *kraus-pair-row n Vs Amat i* ≡
    *foldr* (λ*j acc*. *Amat* $$ (*i*, *j*) ·$_m$ ((*Vs*!*j*)† ∗ *Vs*!*i*) + *acc*)
      [*0*..<*length Vs*] (*0*$_m$ *n n*)

**definition** *kraus-pair-sum* ::
  *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* ⇒
    *complex Matrix.mat*
**where**
  *kraus-pair-sum n Vs Amat* ≡
    *foldr* (λ*i acc*. *kraus-pair-row n Vs Amat i* + *acc*)
      [*0*..<*length Vs*] (*0*$_m$ *n n*)

**lemma** *kraus-pair-row-carrier*:
  **assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
    **and** *i-lt*: *i* < *length Vs*
  **shows** *kraus-pair-row n Vs Amat i* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *kraus-pair-sum-carrier*:
  **assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
  **shows** *kraus-pair-sum n Vs Amat* ∈ *carrier-mat n n*
⟨*proof*⟩

Per-entry index for the dual sum.

**lemma** *kraus-pair-row-index*:
  **assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
    **and** *i-lt*: *i* < *length Vs*
    **and** *r-lt*: *r* < *n* **and** *c-lt*: *c* < *n*
  **shows** *kraus-pair-row n Vs Amat i* $$ (*r*, *c*)
      = (∑ *j* = *0*..<*length Vs*.
          *Amat* $$ (*i*, *j*) ∗ ((*Vs*!*j*)† ∗ *Vs*!*i*) $$ (*r*, *c*))
⟨*proof*⟩

**lemma** *kraus-pair-sum-index*:
  **assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
    **and** *r-lt*: *r* < *n* **and** *c-lt*: *c* < *n*
  **shows** *kraus-pair-sum n Vs Amat* $$ (*r*, *c*)
      = (∑ *i* = *0*..<*length Vs*. ∑ *j* = *0*..<*length Vs*.
          *Amat* $$ (*i*, *j*) ∗ ((*Vs*!*j*)† ∗ *Vs*!*i*) $$ (*r*, *c*))
⟨*proof*⟩

Linearity in the coefficient matrix.

**lemma** *kraus-pair-sum-add-A*:
  **assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
    **and** *A1-c*: *A1* ∈ *carrier-mat* (*length Vs*) (*length Vs*)
    **and** *A2-c*: *A2* ∈ *carrier-mat* (*length Vs*) (*length Vs*)
  **shows** *kraus-pair-sum n Vs* (*A1* + *A2*)
      = *kraus-pair-sum n Vs A1* + *kraus-pair-sum n Vs A2*
⟨*proof*⟩

**lemma** *kraus-pair-sum-smult-A*:
  **assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
    **and** *Amat-c*: *Amat* ∈ *carrier-mat* (*length Vs*) (*length Vs*)
  **shows** *kraus-pair-sum n Vs* (*c* ·$_m$ *Amat*)

$= c \cdot_m$ *kraus-pair-sum n Vs Amat*
⟨*proof*⟩

Dagger identity: taking the dagger of *kraus-pair-sum n Vs A* swaps and conjugates the coefficient indices.

**lemma** *kraus-pair-sum-dagger*:
**assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
**and** *Amat-c*: *Amat* $\in$ *carrier-mat* (*length Vs*) (*length Vs*)
**shows** (*kraus-pair-sum n Vs Amat*)$^\dagger$
$=$ *kraus-pair-sum n Vs*
(*Matrix.mat* (*length Vs*) (*length Vs*)
(λ(*i*,*j*). *cnj* (*Amat* $$ (*j*, *i*))))
⟨*proof*⟩

Existence of a hermitian witness: if *A-orig* (with real entries) gives *kraus-pair-sum n Vs A-orig* $=$ *0* and *A-orig* $\neq$ *0*, then either the symmetric or the anti-hermitian part yields a non-zero hermitian matrix with the same property.

**lemma** *sym-or-antisym-witness*:
**fixes** *A-orig* :: *complex Matrix.mat*
**assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
**and** *A-c*: *A-orig* $\in$ *carrier-mat* (*length Vs*) (*length Vs*)
**and** *A-real*: $\bigwedge i\ j$. $i <$ *length Vs* $\Longrightarrow$ $j <$ *length Vs* $\Longrightarrow$
*cnj* (*A-orig* $$ (*i*, *j*)) = *A-orig* $$ (*i*, *j*)
**and** *A-zero*: *kraus-pair-sum n Vs A-orig* $= 0_m$ *n n*
**and** *A-nontriv*: *A-orig* $\neq 0_m$ (*length Vs*) (*length Vs*)
**shows** $\exists\, H$. $H \in$ *carrier-mat* (*length Vs*) (*length Vs*)
$\wedge$ *hermitian H*
$\wedge$ $H \neq 0_m$ (*length Vs*) (*length Vs*)
$\wedge$ *kraus-pair-sum n Vs H* $= 0_m$ *n n*
⟨*proof*⟩

## 53.10 Bridge: ¬ *mat-lin-indep-real* gives a real *A-orig* witness

Generic concat-nth for uniform-length rows.

**lemma** *concat-uniform-length*:
**assumes** $\forall\, xs \in$ *set L*. *length xs* $=$ *m*
**shows** *length* (*concat L*) $=$ *length L* $*$ *m*
⟨*proof*⟩

**lemma** *concat-uniform-nth*:
**assumes** *uniform*: $\forall\, xs \in$ *set L*. *length xs* $=$ *m*
**and** *i-lt*: $i <$ *length L*
**and** *j-lt*: $j < m$
**shows** *concat L* ! ($i * m + j$) $=$ *L* ! *i* ! *j*
⟨*proof*⟩

*kraus-products Vs* has length $k^2$ with *kraus-products Vs* ! ($i$*$k + j$) $= (V_j)^\dagger\ V_i$.

**lemma** *length-kraus-products*:
*length* (*kraus-products Vs*) $=$ *length Vs* $*$ *length Vs*
⟨*proof*⟩

**lemma** *kraus-products-nth*:
**assumes** *i-lt*: $i <$ *length Vs* **and** *j-lt*: $j <$ *length Vs*
**shows** *kraus-products Vs* ! ($i *$ *length Vs* $+ j$) $= (Vs!j)^\dagger *$ *Vs*!*i*
⟨*proof*⟩

Per-entry of the foldr that mat_lin_indep_real builds.

**lemma** *foldr-smult-nth-index*:
**fixes** $c :: nat \Rightarrow real$ **and** $Bs :: complex\ Matrix.mat\ list$
**assumes** *Bs-c*: $\forall B \in set\ Bs.\ B \in carrier\text{-}mat\ n\ n$
**and** *r-lt*: $r < n$ **and** *c-lt*: $c1 < n$
**shows** $foldr\ (\lambda k\ acc.\ complex\text{-}of\text{-}real\ (c\ k) \cdot_m (Bs\ !\ k) + acc)$
$[0..<length\ Bs]\ (0_m\ n\ n)\ \$\$\ (r,\ c1)$
$= (\sum k = 0..<length\ Bs.$
$complex\text{-}of\text{-}real\ (c\ k) * (Bs\ !\ k)\ \$\$\ (r,\ c1))$
⟨*proof*⟩

Reindex $\sum_{\{}\kappa = 0..<n{\cdot}k\}$ via $\kappa = i{*}k + j$.

**lemma** *sum-split-prod-atLeastLessThan*:
**fixes** $f :: nat \Rightarrow 'a{::}comm\text{-}monoid\text{-}add$
**shows** $(\sum \kappa = 0..<N * k.\ f\ \kappa) = (\sum i = 0..<N.\ \sum j = 0..<k.\ f\ (i * k + j))$
⟨*proof*⟩

The main bridge: a non-trivial real dependence yields a real, non-zero matrix *A-orig* for which *kraus-pair-sum* vanishes.

**lemma** *mat-lin-indep-real-neg-witness*:
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *npos*: $0 < n$
**and** *dep*: $\neg\ mat\text{-}lin\text{-}indep\text{-}real\ n\ (kraus\text{-}products\ Vs)$
**shows** $\exists A\text{-}orig :: complex\ Matrix.mat.$
$A\text{-}orig \in carrier\text{-}mat\ (length\ Vs)\ (length\ Vs)$
$\wedge\ (\forall i\ j.\ i < length\ Vs \longrightarrow j < length\ Vs \longrightarrow$
$cnj\ (A\text{-}orig\ \$\$\ (i,\ j)) = A\text{-}orig\ \$\$\ (i,\ j))$
$\wedge\ A\text{-}orig \neq 0_m\ (length\ Vs)\ (length\ Vs)$
$\wedge\ kraus\text{-}pair\text{-}sum\ n\ Vs\ A\text{-}orig = 0_m\ n\ n$
⟨*proof*⟩

The same with complex coefficients: a non-trivial complex relation among the products $V_j^\dagger V_i$ gives a non-zero complex coefficient matrix with vanishing pair sum, and then (closure of the family under adjoints) a non-zero *Hermitian* one.

**lemma** *mat-lin-indep-complex-neg-witness*:
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *npos*: $0 < n$
**and** *dep*: $\neg\ mat\text{-}lin\text{-}indep\text{-}complex\ n\ (kraus\text{-}products\ Vs)$
**shows** $\exists A\text{-}orig :: complex\ Matrix.mat.$
$A\text{-}orig \in carrier\text{-}mat\ (length\ Vs)\ (length\ Vs)$
$\wedge\ A\text{-}orig \neq 0_m\ (length\ Vs)\ (length\ Vs)$
$\wedge\ kraus\text{-}pair\text{-}sum\ n\ Vs\ A\text{-}orig = 0_m\ n\ n$
⟨*proof*⟩

**lemma** *herm-witness-of-complex*:
**fixes** *A-orig* :: *complex Matrix.mat*
**assumes** *Vs-c*: $\forall V \in set\ Vs.\ V \in carrier\text{-}mat\ d\ n$
**and** *A-c*: $A\text{-}orig \in carrier\text{-}mat\ (length\ Vs)\ (length\ Vs)$
**and** *A-zero*: $kraus\text{-}pair\text{-}sum\ n\ Vs\ A\text{-}orig = 0_m\ n\ n$
**and** *A-nontriv*: $A\text{-}orig \neq 0_m\ (length\ Vs)\ (length\ Vs)$
**shows** $\exists H.\ H \in carrier\text{-}mat\ (length\ Vs)\ (length\ Vs)$
$\wedge\ hermitian\ H$
$\wedge\ H \neq 0_m\ (length\ Vs)\ (length\ Vs)$
$\wedge\ kraus\text{-}pair\text{-}sum\ n\ Vs\ H = 0_m\ n\ n$
⟨*proof*⟩

## 53.11 Auxiliary lemmas for the assembly

Scalar-multiplication linearity of *kraus-quad-map* in $T$.

**lemma** *kraus-quad-map-smult-Tm*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *rho-c*: *rho* $\in$ *carrier-mat n n*
    **and** *Tm-c*: *Tm* $\in$ *carrier-mat* (*length Vs*) (*length Vs*)
  **shows** *kraus-quad-map d Vs* (*c* $\cdot_m$ *Tm*) *rho*
      = *c* $\cdot_m$ *kraus-quad-map d Vs Tm rho*
⟨*proof*⟩

The identity-coefficient case: *kraus-quad-map* with $1_m\ k$ reduces to the ordinary *kraus-map-dim*.

**lemma** *kraus-quad-map-one-eq-kraus-map-dim*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *rho-c*: *rho* $\in$ *carrier-mat n n*
  **shows** *kraus-quad-map d Vs* ($1_m$ (*length Vs*)) *rho* = *kraus-map-dim d Vs rho*
⟨*proof*⟩

The pair-sum at the identity coefficient is exactly *sum-wdag-w*.

**lemma** *kraus-pair-sum-one-eq-sum-wdag-w*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
  **shows** *kraus-pair-sum n Vs* ($1_m$ (*length Vs*)) = *sum-wdag-w n Vs*
⟨*proof*⟩

## 53.12 Minimality and the distinctness lemma (Lemma B)

Linear independence of a complex-matrix list (over $\mathbb{C}$). This is the *minimality* assumption needed to make the reverse direction of Choi's Theorem 3 go through: without it, perturbations of a non-minimal Kraus list can produce the same map.

**definition** *lin-indep-complex-mat* ::
  *nat* ⇒ *nat* ⇒ *complex Matrix.mat list* ⇒ *bool*
**where**
  *lin-indep-complex-mat d n Ws* ≡
    (∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*) ∧
    (∀ *c* :: *nat* ⇒ *complex*.
      *foldr* (λ*k acc*. *c k* $\cdot_m$ (*Ws*!*k*) + *acc*) [*0*..<*length Ws*] (*0*$_m$ *d n*)
        = *0*$_m$ *d n*
      ⟶ (∀ *k* < *length Ws*. *c k* = *0*))

**Lemma B**: under minimality of *Vs*, a non-zero coefficient matrix gives a non-vanishing quadratic Kraus map for some $\varrho$.

The witness $\varrho$ := *mat-unit p q n n* (a single 1 at $(p,q)$) makes $(V_i\ \varrho\ V_j{}^\dagger)$ \$\$ $(r,c) = V_i$ \$\$ $(r,p)\ |V_j$ \$\$ $(c,q)|$, so universal vanishing of *kraus-quad-map* on all $(p,q,r,c)$ combined with linear independence of *Vs* (applied twice, separated by conjugation) forces $A = 0$.

A convenient form of minimality: if the per-entry sum vanishes for all $(r,p)$, then all coefficients are zero.

**lemma** *lin-indep-zero-combo*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *minimal*: *lin-indep-complex-mat d n Vs*
    **and** *coeff-kills*: $\bigwedge r\ p.\ r < d \Longrightarrow p < n \Longrightarrow$
                  $(\sum i = 0..<$*length Vs*. *c i* ∗ (*Vs*!*i*) \$\$ (*r*, *p*)) = *0*
  **shows** ∀ *i* < *length Vs*. *c i* = *0*
⟨*proof*⟩

The per-entry computation for the rank-1 $\varrho$ = *mat-unit p q n n*.

**lemma** *kraus-quad-map-mat-unit-index*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *i-lt*: $i <$ *length Vs* **and** *j-lt*: $j <$ *length Vs*
    **and** *p-lt*: $p < n$ **and** *q-lt*: $q < n$
    **and** *r-lt*: $r < d$ **and** *c-lt*: $c < d$
  **shows** (*Vs*!$i$ $*$ *mat-unit p q n n* $*$ ((*Vs*!$j$)$^\dagger$)) $$ $(r,\ c)$
      $=$ (*Vs*!$i$) $$ $(r,\ p)$ $*$ *cnj* ((*Vs*!$j$) $$ $(c,\ q)$)
⟨*proof*⟩

**Lemma B.** Under minimality, a non-zero coefficient matrix gives a non-vanishing quadratic Kraus map for some rank-1 $\varrho$.

**lemma** *kraus-quad-map-nonzero-witness*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *minimal*: *lin-indep-complex-mat d n Vs*
    **and** *A-c*: $A \in$ *carrier-mat* (*length Vs*) (*length Vs*)
    **and** *A-nz*: $A \neq 0_m$ (*length Vs*) (*length Vs*)
    **and** *npos*: $0 < n$ **and** *dpos*: $0 < d$
  **shows** $\exists$ *rho* $\in$ *carrier-mat n n*. *kraus-quad-map d Vs A rho* $\neq 0_m$ *d d*
⟨*proof*⟩

Trace-preservation lemma for *kraus-quad-map*.

The proof shows *Tr* (*kraus-quad-map d Vs T* $\varrho$) $=$ *Tr* (*kraus-pair-sum n Vs T* $*$ $\varrho$) by per-entry sum manipulation and one application of $[\![$*dim-col ?A* $=$ *dim-row ?B*; *dim-row ?A* $=$ *dim-col ?B*$]\!] \Longrightarrow$ *Tr* (*?A* $*$ *?B*) $=$ *Tr* (*?B* $*$ *?A*) to flip $V_i\ \varrho\ {V_j}^\dagger$ into ${V_j}^\dagger\ V_i\ \varrho$; the TP conclusion follows when the pair-sum equals $1_m$ *n*.

**lemma** *Tr-kraus-quad-map-eq*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *Tmat-c*: *Tmat* $\in$ *carrier-mat* (*length Vs*) (*length Vs*)
    **and** *rho-c*: *rho* $\in$ *carrier-mat n n*
  **shows** *Tr* (*kraus-quad-map d Vs Tmat rho*)
      $=$ *Tr* (*kraus-pair-sum n Vs Tmat* $*$ *rho*)
⟨*proof*⟩

**lemma** *kraus-quad-map-is-TP*:
  **assumes** *Vs-c*: $\forall\, V \in$ *set Vs*. $V \in$ *carrier-mat d n*
    **and** *Tmat-c*: *Tmat* $\in$ *carrier-mat* (*length Vs*) (*length Vs*)
    **and** *Tmat-pair-sum*: *kraus-pair-sum n Vs Tmat* $= 1_m$ *n*
    **and** *npos*: $0 < n$
  **shows** *trace-preserving n* (*kraus-quad-map d Vs Tmat*)
  ⟨*proof*⟩

## 53.13 CPTP-Kraus representation and extreme CPTP maps

A *cptp-kraus-rep n d* $\Phi$ *Vs* says *Vs* is a Kraus operator list (trace-preserving) and $\Phi$ equals the corresponding *kraus-map-dim* on every $X \in$ *carrier-mat n n*.

**definition** *cptp-kraus-rep* ::
  *nat* $\Rightarrow$ *nat* $\Rightarrow$ (*complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*) $\Rightarrow$
    *complex Matrix.mat list* $\Rightarrow$ *bool*
**where**
  *cptp-kraus-rep n d* $\Phi$ *Vs* $\equiv$
    *kraus-operators n d Vs* $\wedge$
    $(\forall\, X.\ X \in$ *carrier-mat n n* $\longrightarrow\ \Phi\ X =$ *kraus-map-dim d Vs X*$)$

Extreme points of $CPTP(n,d)$: analog of *extreme-UCP*.

**definition** *extreme-CPTP* ::
  *nat* $\Rightarrow$ *nat* $\Rightarrow$ (*complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*) $\Rightarrow$ *bool*

**where**
*extreme-CPTP n d Φ ≡*
*Φ ∈ CPTP n d ∧*
*¬ (∃ Φ₁ ∈ CPTP n d. ∃ Φ₂ ∈ CPTP n d.*
*Φ₁ ≠ Φ ∧ Φ ≠ Φ₂ ∧*
*(∀ X. X ∈ carrier-mat n n*
*⟶ Φ X = (1/2 :: complex) ·ₘ Φ₁ X*
*+ (1/2 :: complex) ·ₘ Φ₂ X))*

## 53.14 Vectorisation of Kraus operators

The *vectorisation* of a $d \times n$ matrix $V$ is the length-$n{*}d$ vector with $r$-th entry $V(r \bmod d,\ r \operatorname{div} d)$. This is the column-stacking isomorphism $M_{\{d \times n\}} \cong \mathbb{C}^{\{nd\}}$ used in the Choi/Stinespring picture.

**definition** *vec-of-kraus :: nat ⇒ nat ⇒ complex Matrix.mat ⇒ complex Matrix.vec*
**where**
*vec-of-kraus d n V = Matrix.vec (n∗d) (λr. V $$ (r mod d, r div d))*

**lemma** *vec-of-kraus-dim* [*simp*]: *dim-vec (vec-of-kraus d n V) = n∗d*
⟨*proof*⟩

**lemma** *vec-of-kraus-index*:
**assumes** *r-lt*: *r < n∗d*
**shows** *vec-of-kraus d n V $ r = V $$ (r mod d, r div d)*
⟨*proof*⟩

For $i < d$ and $a < n$, indexing the vectorisation at the *combined* index $a{*}d + i$ recovers the matrix entry $V(i,\ a)$.

**lemma** *vec-of-kraus-combined-index*:
**assumes** *V-c*: *V ∈ carrier-mat d n*
**and** *a-lt*: *a < n* **and** *i-lt*: *i < d* **and** *dpos*: *0 < d*
**shows** *vec-of-kraus d n V $ (a∗d + i) = V $$ (i, a)*
⟨*proof*⟩

## 53.15 Choi matrix of a Kraus map

**Key structural identity**: the Choi matrix of a Kraus map $\Phi(X) = \sum_i V_i\ X\ {V_i}^\dagger$ equals the outer sum $\sum_i v_i\ {v_i}^\dagger$ where $v_i =$ *vec-of-kraus*$(V_i)$. In particular, its range is the linear span of the vectorisations of *Vs*.

**lemma** *kraus-map-dim-mat-unit-entry*:
**assumes** *Vs-c*: *∀ V ∈ set Vs. V ∈ carrier-mat d n*
**and** *a-lt*: *a < n* **and** *b-lt*: *b < n*
**and** *i-lt*: *i < d* **and** *j-lt*: *j < d*
**shows** *(kraus-map-dim d Vs (mat-unit a b n n)) $$ (i, j)*
*= (∑ k<length Vs. Vs!k $$ (i, a) ∗ cnj (Vs!k $$ (j, b)))*
⟨*proof*⟩

**lemma** *choi-matrix-kraus-eq*:
**assumes** *Vs-c*: *∀ V ∈ set Vs. V ∈ carrier-mat d n*
**and** *npos*: *0 < n* **and** *dpos*: *0 < d*
**shows** *choi-matrix n d (kraus-map-dim d Vs)*
*= outer-sum (n∗d) (map (vec-of-kraus d n) Vs)*
⟨*proof*⟩

## 53.16 Choi matrix is linear in the map

The Choi matrix is a linear function of the underlying map (evaluated on the basis *mat-unit*'s). In particular, $P_{\{}\Phi - \Psi\} = P_\Phi - P_\Psi$.

**lemma** *choi-matrix-carrier* [*simp*]:
*choi-matrix n d* $\Phi \in$ *carrier-mat* $(n{*}d)$ $(n{*}d)$
⟨*proof*⟩

**lemma** *choi-matrix-diff*:
**fixes** *Phi Psi* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *Phi-range*: $\bigwedge X.$ $X \in$ *carrier-mat n n* ⟹ *Phi X* ∈ *carrier-mat d d*
**and** *Psi-range*: $\bigwedge X.$ $X \in$ *carrier-mat n n* ⟹ *Psi X* ∈ *carrier-mat d d*
**and** *npos*: $0 < n$ **and** *dpos*: $0 < d$
**shows** *choi-matrix n d* $(\lambda X.$ *Phi X* − *Psi X*$)$
= *choi-matrix n d Phi* − *choi-matrix n d Psi*
⟨*proof*⟩

The Choi matrix agrees on maps that agree on the carrier.

**lemma** *choi-matrix-cong*:
**assumes** *agree*: $\bigwedge X.$ $X \in$ *carrier-mat n n* ⟹ *Phi X* = *Psi X*
**shows** *choi-matrix n d Phi* = *choi-matrix n d Psi*
⟨*proof*⟩

## 53.17 The Choi matrix of a *kraus-quad-map*

The Choi matrix of $X \mapsto \sum_{\{} ij\}$ $T_{\{} ij\}$ $V_i$ $X$ ${V_j}^\dagger$ equals $V\ T\ V^\dagger$ where $V =$ *mat-of-cols* $(n{*}d)$ (*map vec-of-kraus Vs*).

This is the bridge from the classical dominated-PSD parametrisation back to a Kraus quadratic form.

**definition** *kraus-vec-mat* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat*
**where** *kraus-vec-mat d n Vs* = *mat-of-cols* $(n{*}d)$ (*map* (*vec-of-kraus d n*) *Vs*)

**lemma** *kraus-vec-mat-carrier* [*simp*]:
*kraus-vec-mat d n Vs* ∈ *carrier-mat* $(n{*}d)$ (*length Vs*)
⟨*proof*⟩

**lemma** *kraus-vec-mat-index*:
**assumes** *r-lt*: $r < n{*}d$ **and** *k-lt*: $k <$ *length Vs*
**shows** *kraus-vec-mat d n Vs* $\$\$$ $(r, k)$ = *vec-of-kraus d n* (*Vs*!*k*) $\$$ *r*
⟨*proof*⟩

**lemma** *kraus-vec-mat-dagger-index*:
**assumes** *Vs-c*: $\forall$ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *k-lt*: $k <$ *length Vs* **and** *c-lt*: $c < n{*}d$
**shows** (*kraus-vec-mat d n Vs*)$^\dagger$ $\$\$$ $(k, c)$ = *cnj* (*vec-of-kraus d n* (*Vs*!*k*) $\$$ *c*)
⟨*proof*⟩

The Choi matrix of *kraus-quad-map d Vs T* is the sandwich $V\ T\ V^\dagger$, where $V =$ *kraus-vec-mat d n Vs*.

**lemma** *choi-matrix-kraus-quad-eq*:
**fixes** *Tmat* :: *complex Matrix.mat*
**assumes** *Vs-c*: $\forall$ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *T-c*: *Tmat* ∈ *carrier-mat* (*length Vs*) (*length Vs*)
**and** *npos*: $0 < n$ **and** *dpos*: $0 < d$
**shows** *choi-matrix n d* (*kraus-quad-map d Vs Tmat*)
= (*kraus-vec-mat d n Vs*) ∗ *Tmat* ∗ ((*kraus-vec-mat d n Vs*)$^\dagger$)

⟨*proof*⟩

## 53.18 CP Radon–Nikodym — the dominated-PSD kernel

**Dominated PSD parametrisation**. This is the genuinely classical fact behind CP Radon–Nikodym: any PSD matrix $B$ dominated by the outer sum $\sum\ v_i\ {v_i}^\dagger$ can be sandwiched as $V\ T\ V^\dagger$ for some PSD $T$, where $V = \textit{mat-of-cols}\ k\ \textit{vs}$.

Geometric content: $\textit{range}(B) \subseteq \textit{range}(\textit{outer-sum}\ k\ \textit{vs})$ (equivalent to B being Loewner-below outer_sum k vs after PSD), and that range is spanned by the columns of $V$. Pulling B back through that span gives $T$.

Proved below as *dominated-outer-sum-param* through the Moore–Penrose pseudoinverse of the outer sum (no axioms).

**Loewner–kernel inclusion.** If $A \geq B \geq 0$ in Loewner order, then every vector annihilated by $A$ is also annihilated by $B$. Used to derive the range inclusion $\textit{range}(B) \subseteq \textit{range}(A)$.

Proof structure:

1. $\langle x \mid A\ x\rangle = 0$ from $A\ x = 0$.
2. Linearity of the inner product: $\langle x \mid A\ x\rangle = \langle x \mid B\ x\rangle + \langle x \mid (A-B)\ x\rangle$.
3. Both summands have non-negative real part (PSD); the sum is 0, so each is 0.
4. Hermiticity of $B$ makes $\langle x \mid B\ x\rangle$ real, so it equals 0.
5. Decompose $B = N\ N^\dagger$ via $[\![?P \in \textit{carrier-mat}\ ?n\ ?n;\ ?P \geq_M 0]\!] \Longrightarrow \exists\, \textit{vs}.\ \textit{length}\ \textit{vs} = ?n \wedge (\forall\, v{\in}\textit{set}\ \textit{vs}.\ \textit{dim-vec}\ v = ?n) \wedge \textit{outer-sum}\ ?n\ \textit{vs} = ?P$, then $\langle x \mid B\ x\rangle = \langle N^\dagger\ x \mid N^\dagger\ x\rangle = 0$ forces $N^\dagger\ x = 0$, hence $B\ x = N\ (N^\dagger\ x) = 0$.

**lemma** *loewner-ker-subset*:
  **fixes** $A\ B$ :: *complex Matrix.mat* **and** $x$ :: *complex Matrix.vec*
  **assumes** *A-c*: $A \in \textit{carrier-mat}\ k\ k$
    **and** *B-c*: $B \in \textit{carrier-mat}\ k\ k$
    **and** *B-psd*: *psd* $B$
    **and** *diff-psd*: *psd* $(A - B)$
    **and** *x-dim*: $\textit{dim-vec}\ x = k$
    **and** *Ax-zero*: $A *_v x = 0_v\ k$
  **shows** $B *_v x = 0_v\ k$
⟨*proof*⟩

**PSD congruence.** If $P$ is PSD and $M$ is any matrix of compatible shape, then $M^\dagger\ P\ M$ is PSD. This is the standard "sandwich" fact, generalising $?M \in \textit{carrier-mat}\ ?n\ ?m \Longrightarrow ?M^\dagger * ?M \geq_M 0$ (the case $P = 1_m$).

**lemma** *psd-conjugate*:
  **fixes** $P$ :: *complex Matrix.mat* **and** $M$ :: *complex Matrix.mat*
  **assumes** *P-c*: $P \in \textit{carrier-mat}\ k\ k$
    **and** *P-psd*: *psd* $P$
    **and** *M-c*: $M \in \textit{carrier-mat}\ k\ n$
  **shows** *psd* $(M^\dagger * P * M)$
⟨*proof*⟩

### 53.18.1 Diagonal pseudo-inverse list and its identity

**Pseudo-inverse of a real list (interpreted as a diagonal).** Zero entries are kept at zero; non-zero entries are reciprocated. This is the diagonal Moore–Penrose pseudoinverse.

**definition** *pinv-list* :: *real list* ⇒ *real list* **where**
*pinv-list D* = *map* (λ*d*. *if d* = *0 then 0 else 1*/*d*) *D*

**lemma** *pinv-list-length* [*simp*]: *length* (*pinv-list D*) = *length D*
⟨*proof*⟩

**lemma** *pinv-list-index*:
**assumes** *i* < *length D*
**shows** *pinv-list D* ! *i* = (*if D* ! *i* = *0 then 0 else 1* / (*D* ! *i*))
⟨*proof*⟩

**lemma** *pinv-list-nonneg*:
**assumes** ⋀*i*. *i* < *length D* ⟹ *0* ≤ *D* ! *i*
**shows** ⋀*i*. *i* < *length D* ⟹ *0* ≤ *pinv-list D* ! *i*
⟨*proof*⟩

The central identity: $D \cdot D^{+} \cdot D = D$ entrywise on the diagonal. For $D_i = 0$ the left-hand side is $0 \cdot 0 \cdot 0 = 0 = D_i$; for $D_i \neq 0$ it is $D_i \cdot (1/D_i) \cdot D_i = D_i$. This is the core diagonal pseudoinverse identity that lifts to operator level through unitary conjugation.

**lemma** *diag-pinv-diag-eq*:
**fixes** *D* :: *real list*
**shows** *diag-mat-of-real D*
∗ *diag-mat-of-real* (*pinv-list D*)
∗ *diag-mat-of-real D*
= *diag-mat-of-real D*
⟨*proof*⟩

**Pseudoinverse identity at operator level.** Given a hermitian spectral decomposition $A = U\ D\ U^{\dagger}$, define the Moore–Penrose pseudoinverse via $A^{+} = U\ D^{+}\ U^{\dagger}$. Then $A\ A^{+}\ A = A$. This is the unitary-conjugation lift of *diag-mat-of-real ?D* ∗ *diag-mat-of-real* (*pinv-list ?D*) ∗ *diag-mat-of-real ?D* = *diag-mat-of-real ?D*.

**lemma** *A-Aplus-A-eq*:
**fixes** *U* :: *complex Matrix.mat* **and** *D* :: *real list*
**assumes** *Uc*: *U* ∈ *carrier-mat k k*
**and** *UU*: *unitary U*
**and** *Dlen*: *length D* = *k*
**shows** (*U* ∗ *diag-mat-of-real D* ∗ (*U*†))
∗ (*U* ∗ *diag-mat-of-real* (*pinv-list D*) ∗ (*U*†))
∗ (*U* ∗ *diag-mat-of-real D* ∗ (*U*†))
= *U* ∗ *diag-mat-of-real D* ∗ (*U*†)
⟨*proof*⟩

Two square matrices agreeing on every vector are equal.

**lemma** *mat-eq-of-mult-vec-eq*:
**fixes** *M N* :: *complex Matrix.mat*
**assumes** *M-c*: *M* ∈ *carrier-mat k k*
**and** *N-c*: *N* ∈ *carrier-mat k k*
**and** *eq*: ⋀*x*. *x* ∈ *carrier-vec k* ⟹ *M* ∗*v x* = *N* ∗*v x*
**shows** *M* = *N*
⟨*proof*⟩

Loewner-implied pseudoinverse identity, proved at operator level: with $Q = A^{+}\ A$ one has $A\ Q = A$, so $x - Q\ x \in ker\ A \subseteq ker\ B$, whence $B = B\ Q$; taking adjoints gives $B = A\ A^{+}\ B$.

**lemma** *A-Aplus-B-eq-B*:
**fixes** *U* :: *complex Matrix.mat* **and** *D* :: *real list* **and** *B* :: *complex Matrix.mat*
**assumes** *Uc*: *U* ∈ *carrier-mat k k*

**and** *UU*: *unitary U*
**and** *Dlen*: *length D* = *k*
**and** *Dnn*: $\bigwedge i.\ i < k \Longrightarrow 0 \leq D\ !\ i$
**and** *B-c*: *B* ∈ *carrier-mat k k*
**and** *B-psd*: *psd B*
**and** *dom-psd*: *psd* ((*U* ∗ *diag-mat-of-real D* ∗ (*U*†)) − *B*)
**shows** (*U* ∗ *diag-mat-of-real D* ∗ (*U*†))
∗ (*U* ∗ *diag-mat-of-real* (*pinv-list D*) ∗ (*U*†)) ∗ *B* = *B*
⟨*proof*⟩

Spectral decomposition of a psd matrix with non-negative diagonal.

Dominated psd matrices are parametrised through the dominating frame: if $0 \leq B \leq V\ V^\dagger$ then $B = V\ T\ V^\dagger$ with $T = V^\dagger\ A^+\ B\ A^+\ V \geq 0$, $A = V\ V^\dagger$, $A^+$ the Moore–Penrose pseudoinverse.

**lemma** *dominated-outer-sum-param*:
**fixes** *vs* :: *complex Matrix.vec list*
**and** *B* :: *complex Matrix.mat*
**assumes** *vs-dim*: ∀ *v* ∈ *set vs*. *dim-vec v* = *k*
**and** *k-pos*: *0* < *k*
**and** *B-c*: *B* ∈ *carrier-mat k k*
**and** *B-psd*: *psd B*
**and** *B-dom*: *psd* (*outer-sum k vs* − *B*)
**shows** ∃ *T*. *T* ∈ *carrier-mat* (*length vs*) (*length vs*) ∧ *psd T* ∧
*B* = (*mat-of-cols k vs*) ∗ *T* ∗ ((*mat-of-cols k vs*)†)
⟨*proof*⟩

## 53.19 Linearity of *kraus-quad-map* in its matrix argument

Per-entry computation shows *kraus-quad-map d Vs Tm* is linear in its third (matrix) argument: $\varrho \mapsto \sum_{\{ij\}}\ T_{\{ij\}}\ V_i\ \varrho\ {V_j}^\dagger$.

**lemma** *kraus-quad-map-add-rho*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *rho1-c*: *rho1* ∈ *carrier-mat n n*
**and** *rho2-c*: *rho2* ∈ *carrier-mat n n*
**shows** *kraus-quad-map d Vs Tm* (*rho1* + *rho2*)
= *kraus-quad-map d Vs Tm rho1* + *kraus-quad-map d Vs Tm rho2*
⟨*proof*⟩

**lemma** *kraus-quad-map-smult-rho*:
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *rho-c*: *rho* ∈ *carrier-mat n n*
**shows** *kraus-quad-map d Vs Tm* (*c* ·$_m$ *rho*)
= *c* ·$_m$ *kraus-quad-map d Vs Tm rho*
⟨*proof*⟩

## 53.20 Choi matrix determines a linear CP map on carrier

If two maps that are linear on carrier have the same Choi matrix and the same target carrier, they agree pointwise on carrier.

**lemma** *map-eq-from-choi-when-linear*:
**fixes** *Phi Psi* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *Phi-add*: $\bigwedge X\ Y.\ X \in$ *carrier-mat n n* $\Longrightarrow Y \in$ *carrier-mat n n*
$\Longrightarrow$ *Phi* (*X* + *Y*) = *Phi X* + *Phi Y*
**and** *Phi-smult*: $\bigwedge c\ X.\ X \in$ *carrier-mat n n* $\Longrightarrow$ *Phi* (*c* ·$_m$ *X*) = *c* ·$_m$ *Phi X*
**and** *Phi-car*: $\bigwedge X.\ X \in$ *carrier-mat n n* $\Longrightarrow$ *Phi X* ∈ *carrier-mat d d*
**and** *Psi-add*: $\bigwedge X\ Y.\ X \in$ *carrier-mat n n* $\Longrightarrow Y \in$ *carrier-mat n n*

$\Longrightarrow$ *Psi* (*X* + *Y*) = *Psi X* + *Psi Y*
**and** *Psi-smult*: $\bigwedge$*c X*. *X* ∈ *carrier-mat n n* $\Longrightarrow$ *Psi* (*c* ·$_m$ *X*) = *c* ·$_m$ *Psi X*
**and** *Psi-car*: $\bigwedge$*X*. *X* ∈ *carrier-mat n n* $\Longrightarrow$ *Psi X* ∈ *carrier-mat d d*
**and** *choi-eq*: *choi-matrix n d Phi* = *choi-matrix n d Psi*
**and** *npos*: *n* > *0* **and** *dpos*: *d* > *0*
**and** *A-c*: *A* ∈ *carrier-mat n n*
**shows** *Phi A* = *Psi A*
⟨*proof*⟩

## 53.21 CP Radon–Nikodym

**CP Radon–Nikodym (Choi level)**. At the Choi-matrix level, the parametrisation falls out from ⟦∀ *v*∈*set ?vs*. *dim-vec v* = *?k*; *0* < *?k*; *?B* ∈ *carrier-mat ?k ?k*; *?B* ≥$_M$ *0*; *outer-sum ?k ?vs* − *?B* ≥$_M$ *0*⟧ $\Longrightarrow$ ∃ *T*. *T* ∈ *carrier-mat* (*length ?vs*) (*length ?vs*) ∧ *T* ≥$_M$ *0* ∧ *?B* = *mat-of-cols ?k ?vs* ∗ *T* ∗ *mat-of-cols ?k ?vs*$^\dagger$ composed with the structural identities ⟦∀ *V*∈*set ?Vs*. *V* ∈ *carrier-mat ?d ?n*; *0* < *?n*; *0* < *?d*⟧ $\Longrightarrow$ $P_\Phi$ *?n ?d* (*kraus-map-dim ?d ?Vs*) = *outer-sum* (*?n* ∗ *?d*) (*map* (*vec-of-kraus ?d ?n*) *?Vs*) and ⟦∀ *V*∈*set ?Vs*. *V* ∈ *carrier-mat ?d ?n*; *?Tmat* ∈ *carrier-mat* (*length ?Vs*) (*length ?Vs*); *0* < *?n*; *0* < *?d*⟧ $\Longrightarrow$ $P_\Phi$ *?n ?d* (*kraus-quad-map ?d ?Vs ?Tmat*) = *kraus-vec-mat ?d ?n ?Vs* ∗ *?Tmat* ∗ *kraus-vec-mat ?d ?n ?Vs*$^\dagger$. This proves what is needed at the matrix level; passing to the ∀ *X*-quantified form requires linearity of Ψ, which is the project's *completely-positive* does *not* include (see the lemma after).

**lemma** *cp-radon-nikodym-choi*:
**fixes** *Phi Psi* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list*
**assumes** *Vs-c*: ∀ *V* ∈ *set Vs*. *V* ∈ *carrier-mat d n*
**and** *Phi-kraus*: $\bigwedge$*X*. *X* ∈ *carrier-mat n n* $\Longrightarrow$ *Phi X* = *kraus-map-dim d Vs X*
**and** *Psi-cp*: *completely-positive n d Psi*
**and** *Psi-range*: $\bigwedge$*X*. *X* ∈ *carrier-mat n n* $\Longrightarrow$ *Psi X* ∈ *carrier-mat d d*
**and** *diff-cp*: *completely-positive n d* (λ*X*. *Phi X* − *Psi X*)
**and** *npos*: *0* < *n*
**and** *dpos*: *0* < *d*
**shows** ∃ *T*. *T* ∈ *carrier-mat* (*length Vs*) (*length Vs*) ∧ *psd T* ∧
*choi-matrix n d Psi* = *choi-matrix n d* (*kraus-quad-map d Vs T*)
⟨*proof*⟩

**CP Radon–Nikodym (operator level)**. With Ψ linear on *carrier-mat n n*, the Choi-level parametrisation gives the pointwise equality $\Psi(X) = \sum\ T_{\{ij\}}\ V_i\ X\ V_j{}^\dagger$.
Linearity of Ψ is supplied as an explicit hypothesis: the project's *completely-positive* predicate does *not* imply linearity (it constrains only positivity under amplification). This corresponds to the classical fact that "CP map" standardly includes linearity; we make it explicit so the consumer can supply it.

**Operator-level CP Radon–Nikodym, with linearity assumption.** This version is fully discharged via ⟦∀ *V*∈*set ?Vs*. *V* ∈ *carrier-mat ?d ?n*; $\bigwedge$*X*. *X* ∈ *carrier-mat ?n ?n* $\Longrightarrow$ *?Phi X* = *kraus-map-dim ?d ?Vs X*; *completely-positive ?n ?d ?Psi*; $\bigwedge$*X*. *X* ∈ *carrier-mat ?n ?n* $\Longrightarrow$ *?Psi X* ∈ *carrier-mat ?d ?d*; *completely-positive ?n ?d* (λ*X*. *?Phi X* − *?Psi X*); *0* < *?n*; *0* < *?d*⟧ $\Longrightarrow$ ∃ *T*. *T* ∈ *carrier-mat* (*length ?Vs*) (*length ?Vs*) ∧ *T* ≥$_M$ *0* ∧ $P_\Phi$ *?n ?d ?Psi* = $P_\Phi$ *?n ?d* (*kraus-quad-map ?d ?Vs T*) + ⟦$\bigwedge$*X Y*. ⟦*X* ∈ *carrier-mat ?n ?n*; *Y* ∈ *carrier-mat ?n ?n*⟧ $\Longrightarrow$ *?Phi* (*X* + *Y*) = *?Phi X* + *?Phi Y*; $\bigwedge$*c X*. *X* ∈ *carrier-mat ?n ?n* $\Longrightarrow$ *?Phi* (*c* ·$_m$ *X*) = *c* ·$_m$ *?Phi X*; $\bigwedge$*X*. *X* ∈ *carrier-mat ?n ?n* $\Longrightarrow$ *?Phi X* ∈ *carrier-mat ?d ?d*; $\bigwedge$*X Y*. ⟦*X* ∈ *carrier-mat ?n ?n*; *Y* ∈ *carrier-mat ?n ?n*⟧ $\Longrightarrow$ *?Psi* (*X* + *Y*) = *?Psi X* + *?Psi Y*; $\bigwedge$*c X*. *X* ∈ *carrier-mat ?n ?n* $\Longrightarrow$ *?Psi* (*c* ·$_m$ *X*) = *c* ·$_m$ *?Psi X*; $\bigwedge$*X*. *X* ∈ *carrier-mat ?n ?n* $\Longrightarrow$ *?Psi X* ∈ *carrier-mat ?d ?d*; $P_\Phi$ *?n ?d ?Phi* = $P_\Phi$ *?n ?d ?Psi*; *0* < *?n*; *0* < *?d*; *?A* ∈ *carrier-mat ?n ?n*⟧ $\Longrightarrow$ *?Phi ?A* = *?Psi ?A*, which rest on ⟦∀ *v*∈*set ?vs*. *dim-vec v* = *?k*; *0* < *?k*; *?B* ∈

*carrier-mat ?k ?k*; *?B* $\geq_M$ *0*; *outer-sum ?k ?vs − ?B* $\geq_M$ *0*$\rrbracket \Longrightarrow \exists\, T.\ T \in$ *carrier-mat* (*length ?vs*) (*length ?vs*) $\wedge$ *T* $\geq_M$ *0* $\wedge$ *?B = mat-of-cols ?k ?vs* $*$ *T* $*$ *mat-of-cols ?k ?vs*$^\dagger$. It requires *Psi* to be linear on *carrier-mat n n*, which is not implied by the project's *completely-positive* predicate.

**lemma** *cp-radon-nikodym-linear*:
  **fixes** *Phi Psi* :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
    **and** *Vs* :: *complex Matrix.mat list*
  **assumes** *Vs-c*: $\forall$ *V* $\in$ *set Vs. V* $\in$ *carrier-mat d n*
      **and** *Phi-kraus*: $\bigwedge$*X. X* $\in$ *carrier-mat n n* $\Longrightarrow$ *Phi X = kraus-map-dim d Vs X*
      **and** *Psi-cp*: *completely-positive n d Psi*
      **and** *Psi-range*: $\bigwedge$*X. X* $\in$ *carrier-mat n n* $\Longrightarrow$ *Psi X* $\in$ *carrier-mat d d*
      **and** *Psi-add*: $\bigwedge$*X Y. X* $\in$ *carrier-mat n n* $\Longrightarrow$ *Y* $\in$ *carrier-mat n n*
                  $\Longrightarrow$ *Psi* (*X* + *Y*) = *Psi X* + *Psi Y*
      **and** *Psi-smult*: $\bigwedge$*c X. X* $\in$ *carrier-mat n n* $\Longrightarrow$ *Psi* (*c* $\cdot_m$ *X*) = *c* $\cdot_m$ *Psi X*
      **and** *diff-cp*: *completely-positive n d* ($\lambda$*X. Phi X* − *Psi X*)
      **and** *npos*: *0* < *n*
      **and** *dpos*: *0* < *d*
  **shows** $\exists$ *T. T* $\in$ *carrier-mat* (*length Vs*) (*length Vs*) $\wedge$ *psd T* $\wedge$
          ($\forall$ *X. X* $\in$ *carrier-mat n n* $\longrightarrow$ *Psi X = kraus-quad-map d Vs T X*)
⟨*proof*⟩

The use site *choi-theorem-3-cptp-forward-complex* obtains linearity of its $\Psi$ through the strengthened *cptp-pair-mid* predicate (which carries *lin-on-carrier* for both midpoint witnesses) and dispatches to $[\![\forall$ *V*$\in$*set ?Vs. V* $\in$ *carrier-mat ?d ?n*; $\bigwedge$*X. X* $\in$ *carrier-mat ?n ?n* $\Longrightarrow$ *?Phi X = kraus-map-dim ?d ?Vs X*; *completely-positive ?n ?d ?Psi*; $\bigwedge$*X. X* $\in$ *carrier-mat ?n ?n* $\Longrightarrow$ *?Psi X* $\in$ *carrier-mat ?d ?d*; $\bigwedge$*X Y.* $[\![$*X* $\in$ *carrier-mat ?n ?n*; *Y* $\in$ *carrier-mat ?n ?n*$]\!] \Longrightarrow$ *?Psi* (*X* + *Y*) = *?Psi X* + *?Psi Y*; $\bigwedge$*c X. X* $\in$ *carrier-mat ?n ?n* $\Longrightarrow$ *?Psi* (*c* $\cdot_m$ *X*) = *c* $\cdot_m$ *?Psi X*; *completely-positive ?n ?d* ($\lambda$*X. ?Phi X* − *?Psi X*); *0* < *?n*; *0* < *?d*$]\!] \Longrightarrow \exists$ *T. T* $\in$ *carrier-mat* (*length ?Vs*) (*length ?Vs*) $\wedge$ *T* $\geq_M$ *0* $\wedge$ ($\forall$ *X. X* $\in$ *carrier-mat ?n ?n* $\longrightarrow$ *?Psi X = kraus-quad-map ?d ?Vs T X*). No operator-level *cp-radon-nikodym* without linearity is needed any more.

## 53.22 Choi's Theorem 3 — reverse direction

**Reverse direction of Choi's Theorem 3 (CPTP form).** Under minimality of the Kraus list *Vs*, the non-linear-independence of the products $\{V_j^\dagger\ V_i\}$ forces $\Phi$ to fail to be an extreme point of *CPTP n d*.

The witness uses the explicit perturbation $\Phi_\pm(\varrho) := \sum(I_{\{ij\}} \pm \varepsilon\ H_{\{ij\}})\ V_i\ \varrho\ V_j^\dagger$ for a hermitian *H* (from $[\![\forall$ *V*$\in$*set ?Vs. V* $\in$ *carrier-mat ?d ?n*; *?A-orig* $\in$ *carrier-mat* (*length ?Vs*) (*length ?Vs*); $\bigwedge$*i j.* $[\![$*i* < *length ?Vs*; *j* < *length ?Vs*$]\!] \Longrightarrow$ *cnj* (*?A-orig* $$ (*i*, *j*)) = *?A-orig* $$ (*i*, *j*); *kraus-pair-sum ?n ?Vs ?A-orig = 0*$_m$ *?n ?n*; *?A-orig* $\neq$ *0*$_m$ (*length ?Vs*) (*length ?Vs*)$]\!]$ $\Longrightarrow \exists$ *H. H* $\in$ *carrier-mat* (*length ?Vs*) (*length ?Vs*) $\wedge$ *hermitian H* $\wedge$ *H* $\neq$ *0*$_m$ (*length ?Vs*) (*length ?Vs*) $\wedge$ *kraus-pair-sum ?n ?Vs H = 0*$_m$ *?n ?n*) and a small $\varepsilon$ (from $[\![$*?A* $\in$ *carrier-mat ?k ?k*; *hermitian ?A*$]\!] \Longrightarrow \exists\varepsilon{>}0.\ \forall\, d.\ |d| \leq \varepsilon \longrightarrow$ *1*$_m$ *?k* + *complex-of-real d* $\cdot_m$ *?A* $\geq_M$ *0*). Distinctness uses $[\![\forall$ *V*$\in$*set ?Vs. V* $\in$ *carrier-mat ?d ?n*; *lin-indep-complex-mat ?d ?n ?Vs*; *?A* $\in$ *carrier-mat* (*length ?Vs*) (*length ?Vs*); *?A* $\neq$ *0*$_m$ (*length ?Vs*) (*length ?Vs*); *0* < *?n*; *0* < *?d*$]\!]$ $\Longrightarrow \exists$ *rho*$\in$*carrier-mat ?n ?n. kraus-quad-map ?d ?Vs ?A rho* $\neq$ *0*$_m$ *?d ?d* (Lemma B).

**theorem** *choi-theorem-3-cptp-reverse*:
  **fixes** $\Phi$ :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
    **and** *Vs* :: *complex Matrix.mat list*
  **assumes** *npos*: *0* < *n* **and** *dpos*: *0* < *d*
      **and** $\Phi$*-in*: $\Phi \in$ *CPTP n d*
      **and** *rep*: *cptp-kraus-rep n d* $\Phi$ *Vs*

**and** *minimal*: *lin-indep-complex-mat d n Vs*
**and** *dep*: ¬ *mat-lin-indep-real n* (*kraus-products Vs*)
**shows** ¬ *extreme-CPTP n d* Φ
⟨*proof*⟩

**Caveat on** *extreme-CPTP* **(forward shadow).** Because both *completely-positive* and *trace-preserving* only constrain behaviour on *carrier-mat n n*, the raw function inequality $\Phi_1 \neq \Phi$ in *extreme-CPTP* is trivially satisfiable. We therefore introduce a *carrier*-point non-extremity notion below.

To keep *extreme-CPTP-carrier*'s definition manageable for automated proofs, we factor it through a helper predicate *cptp-pair-mid*: the data of a CPTP midpoint pair witnessing non-extremity (range, linearity on carrier, off-equal at some carrier-point, and the midpoint identity). Unfolding only the outer definition keeps the goal small.

**definition** *lin-on-carrier* ::
*nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool*
**where**
*lin-on-carrier n* Φ ≡
(∀ *X Y*. *X* ∈ *carrier-mat n n* ⟶ *Y* ∈ *carrier-mat n n*
⟶ Φ (*X* + *Y*) = Φ *X* + Φ *Y*) ∧
(∀ *c X*. *X* ∈ *carrier-mat n n* ⟶ Φ (*c* ·$_m$ *X*) = *c* ·$_m$ Φ *X*)

**lemma** *lin-on-carrier-add*:
**assumes** *lin-on-carrier n* Φ *X* ∈ *carrier-mat n n Y* ∈ *carrier-mat n n*
**shows** Φ (*X* + *Y*) = Φ *X* + Φ *Y*
⟨*proof*⟩

**lemma** *lin-on-carrier-smult*:
**assumes** *lin-on-carrier n* Φ *X* ∈ *carrier-mat n n*
**shows** Φ (*c* ·$_m$ *X*) = *c* ·$_m$ Φ *X*
⟨*proof*⟩

**lemma** *lin-on-carrierI*:
**assumes** ⋀*X Y*. *X* ∈ *carrier-mat n n* ⟹ *Y* ∈ *carrier-mat n n*
⟹ Φ (*X* + *Y*) = Φ *X* + Φ *Y*
**and** ⋀*c X*. *X* ∈ *carrier-mat n n* ⟹ Φ (*c* ·$_m$ *X*) = *c* ·$_m$ Φ *X*
**shows** *lin-on-carrier n* Φ
⟨*proof*⟩

**definition** *cptp-pair-mid* ::
*nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*)
⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*)
⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool*
**where**
*cptp-pair-mid n d* Φ Φ$_1$ Φ$_2$ ≡
Φ$_1$ ∈ *CPTP n d* ∧ Φ$_2$ ∈ *CPTP n d* ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ$_1$ *X* ∈ *carrier-mat d d*) ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ$_2$ *X* ∈ *carrier-mat d d*) ∧
*lin-on-carrier n* Φ$_1$ ∧ *lin-on-carrier n* Φ$_2$ ∧
(∃ *X*$_1$ ∈ *carrier-mat n n*. Φ$_1$ *X*$_1$ ≠ Φ *X*$_1$) ∧
(∃ *X*$_2$ ∈ *carrier-mat n n*. Φ *X*$_2$ ≠ Φ$_2$ *X*$_2$) ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶
Φ *X* = (*1/2* :: *complex*) ·$_m$ Φ$_1$ *X* + (*1/2* :: *complex*) ·$_m$ Φ$_2$ *X*)

**definition** *extreme-CPTP-carrier* ::
*nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool*
**where**

*extreme-CPTP-carrier n d Φ ≡*
*Φ ∈ CPTP n d ∧ ¬ (∃ Φ₁ Φ₂. cptp-pair-mid n d Φ Φ₁ Φ₂)*

The same construction as in ⟦*0 < ?n*; *0 < ?d*; *?Φ ∈ CPTP ?n ?d*; *cptp-kraus-rep ?n ?d ?Φ ?Vs*; *lin-indep-complex-mat ?d ?n ?Vs*; ¬ *mat-lin-indep-real ?n* (*kraus-products ?Vs*)⟧ ⟹ ¬ *extreme-CPTP ?n ?d ?Φ* yields the stronger fact that Φ fails the *extreme-CPTP-carrier* test: the perturbed maps *Phip*, *Phim* are CPTP, have range *carrier-mat d d*, and disagree with Φ at the carrier-point *rho*.

**lemma** *not-extreme-CPTP-carrier-of-herm-witness*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list* **and** *H* :: *complex Matrix.mat*
**assumes** *npos*: *0 < n* **and** *dpos*: *0 < d*
**and** Φ-*in*: *Φ ∈ CPTP n d*
**and** *rep*: *cptp-kraus-rep n d Φ Vs*
**and** *minimal*: *lin-indep-complex-mat d n Vs*
**and** *H-c*: *H ∈ carrier-mat* (*length Vs*) (*length Vs*)
**and** *H-herm*: *hermitian H*
**and** *H-nz*: *H ≠ 0*$_m$ (*length Vs*) (*length Vs*)
**and** *H-pair*: *kraus-pair-sum n Vs H = 0*$_m$ *n n*
**shows** ¬ *extreme-CPTP-carrier n d Φ*
⟨*proof*⟩

Immediate contrapositive: extreme-in-carrier-sense forces linear independence of the products.

**theorem** *choi-theorem-3-cptp-reverse-carrier*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list*
**assumes** *npos*: *0 < n* **and** *dpos*: *0 < d*
**and** Φ-*in*: *Φ ∈ CPTP n d*
**and** *rep*: *cptp-kraus-rep n d Φ Vs*
**and** *minimal*: *lin-indep-complex-mat d n Vs*
**and** *dep*: ¬ *mat-lin-indep-real n* (*kraus-products Vs*)
**shows** ¬ *extreme-CPTP-carrier n d Φ*
⟨*proof*⟩

**theorem** *choi-theorem-3-cptp-reverse-carrier-complex*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list*
**assumes** *npos*: *0 < n* **and** *dpos*: *0 < d*
**and** Φ-*in*: *Φ ∈ CPTP n d*
**and** *rep*: *cptp-kraus-rep n d Φ Vs*
**and** *minimal*: *lin-indep-complex-mat d n Vs*
**and** *dep*: ¬ *mat-lin-indep-complex n* (*kraus-products Vs*)
**shows** ¬ *extreme-CPTP-carrier n d Φ*
⟨*proof*⟩

**corollary** *choi-theorem-3-cptp-carrier-forward*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list*
**assumes** *npos*: *0 < n* **and** *dpos*: *0 < d*
**and** Φ-*in*: *Φ ∈ CPTP n d*
**and** *rep*: *cptp-kraus-rep n d Φ Vs*
**and** *minimal*: *lin-indep-complex-mat d n Vs*
**and** *ext*: *extreme-CPTP-carrier n d Φ*
**shows** *mat-lin-indep-real n* (*kraus-products Vs*)
⟨*proof*⟩

## 53.23 Trace pairing duality and TP-converse

If *Tr* (*M* ∗ *ϱ*) = *Tr* (*N* ∗ *ϱ*) for every *ϱ* in carrier, then *M* = *N*. Used to invert ⟦∀ *V*∈*set ?Vs*. *V* ∈ *carrier-mat ?d ?n*; *?Tmat* ∈ *carrier-mat* (*length ?Vs*) (*length ?Vs*); *kraus-pair-sum ?n ?Vs ?Tmat* = $1_m$ *?n*; 0 < *?n*⟧ ⟹ *trace-preserving ?n* (*kraus-quad-map ?d ?Vs ?Tmat*).

**lemma** *Tr-mat-unit-mult*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *M-c*: *M* ∈ *carrier-mat n n*
**and** *r-lt*: *r* < *n* **and** *s-lt*: *s* < *n*
**shows** *Tr* (*M* ∗ *mat-unit s r n n*) = *M* $$ (*r*, *s*)
⟨*proof*⟩

**lemma** *mat-trace-pairing-unique*:
**fixes** *M N* :: *complex Matrix.mat*
**assumes** *M-c*: *M* ∈ *carrier-mat n n*
**and** *N-c*: *N* ∈ *carrier-mat n n*
**and** *pair-eq*: ⋀*rho*. *rho* ∈ *carrier-mat n n* ⟹ *Tr* (*M* ∗ *rho*) = *Tr* (*N* ∗ *rho*)
**shows** *M* = *N*
⟨*proof*⟩

## 53.24 Choi's Theorem 3 — forward direction

**Important caveat on** *extreme-CPTP***.** The current definition uses raw function inequality Φ$_1$ ≠ Φ, but both *completely-positive* and *trace-preserving* only constrain behaviour on *carrier-mat n n*. Hence for *any* CPTP Φ one can construct Φ$_1$ that agrees with Φ on the carrier but disagrees on (say) a *1*×*1* matrix; this Φ$_1$ is itself in CPTP, so the negation of *extreme-CPTP* is vacuously satisfied. Consequently *extreme-CPTP* **as currently defined is always false**, and Choi's Theorem 3 as stated (an iff with *mat-lin-indep-real*) does not hold.

The fix is to use a *carrier*-point notion of inequality (defined above as *extreme-CPTP-carrier*).

The forward direction targets the *carrier*-flavoured version: *mat-lin-indep-complex* ⟹ *extreme-CPTP-carrier*. Proof uses ⟦∀ *V*∈*set ?Vs*. *V* ∈ *carrier-mat ?d ?n*; ⋀*X*. *X* ∈ *carrier-mat ?n ?n* ⟹ *?Phi X* = *kraus-map-dim ?d ?Vs X*; *completely-positive ?n ?d ?Psi*; ⋀*X*. *X* ∈ *carrier-mat ?n ?n* ⟹ *?Psi X* ∈ *carrier-mat ?d ?d*; ⋀*X Y*. ⟦*X* ∈ *carrier-mat ?n ?n*; *Y* ∈ *carrier-mat ?n ?n*⟧ ⟹ *?Psi* (*X* + *Y*) = *?Psi X* + *?Psi Y*; ⋀*c X*. *X* ∈ *carrier-mat ?n ?n* ⟹ *?Psi* (*c* ·$_m$ *X*) = *c* ·$_m$ *?Psi X*; *completely-positive ?n ?d* (λ*X*. *?Phi X* − *?Psi X*); 0 < *?n*; 0 < *?d*⟧ ⟹ ∃ *T*. *T* ∈ *carrier-mat* (*length ?Vs*) (*length ?Vs*) ∧ *T* ≥$_M$ 0 ∧ (∀ *X*. *X* ∈ *carrier-mat ?n ?n* ⟶ *?Psi X* = *kraus-quad-map ?d ?Vs T X*), obtaining linearity of Ψ from the *lin-on-carrier* conjuncts carried by *cptp-pair-mid*.

**theorem** *choi-theorem-3-cptp-forward-complex*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list*
**assumes** *npos*: 0 < *n* **and** *dpos*: 0 < *d*
**and** Φ-*in*: Φ ∈ *CPTP n d*
**and** *rep*: *cptp-kraus-rep n d* Φ *Vs*
**and** *lin-indep*: *mat-lin-indep-complex n* (*kraus-products Vs*)
**shows** *extreme-CPTP-carrier n d* Φ
⟨*proof*⟩

## 53.25 Choi's Theorem 3 (statement)

**NOTE on conventions.** Choi (1975, Theorem 3) gives an extreme-point characterisation in terms of *linear independence of certain products of Kraus operators*. Which products appear depends on which of the two natural normalisations the underlying convex set uses:

- **CPTP** ($\sum\ V_i{}^\dagger\ V_i = I_n$): products are $\{V_j{}^\dagger\ V_i\}$ in $M_n$. The perturbation $\Phi_\varepsilon(X) = \sum(\delta_{\{}ij\} + \varepsilon\ a_{\{}ij\})\ V_i\ X\ V_j{}^\dagger$ stays trace-preserving iff $\sum\ a_{\{}ij\}\ V_j{}^\dagger\ V_i = 0$.
- **UCP** ($\sum\ V_i\ V_i{}^\dagger = I_d$): products are $\{V_i\ V_j{}^\dagger\}$ in $M_d$. Same perturbation stays unital iff $\sum\ a_{\{}ij\}\ V_i\ V_j{}^\dagger = 0$.

The two are interchanged by the Hilbert–Schmidt adjoint $\Phi \mapsto \Phi^*$; this provides a UCP $\leftrightarrow$ CPTP bijection.

We state both versions, parametrised by a fixed Kraus representation; the "any/every" character follows from the unitary-freedom of Kraus reps (Day 22 / "Kraus uniqueness"), which is not yet formalised in this development.

**Reverse direction (extreme $\Rightarrow$ lin. indep.)**: contrapositive. For the CPTP version: if $\sum\ a_{\{}ij\}\ V_j{}^\dagger\ V_i = 0$ non-trivially, define $\Phi_\varepsilon$ as above; for small hermitian $a$, both $\Phi_{\pm\varepsilon}$ are CPTP and distinct, giving a non-trivial midpoint of $\Phi$.

**Forward direction (lin. indep. $\Rightarrow$ extreme)**: requires Radon–Nikodym for CP maps (every CP $\Psi \leq_{\{} cp\}\ \lambda\Phi$ has $\Psi(X) = \sum\ T_{\{}ij\}\ V_i\ X\ V_j{}^\dagger$ for a PSD matrix $T$) which is itself a substantial development.

**The CPTP form of Choi's Theorem 3 (carrier-flavoured).** $\Phi$ is extreme in CPTP iff the products $\{V_j{}^\dagger\ V_i\}$ are linearly independent in $M_n$.

The project's *extreme-CPTP* uses a raw function inequality $\Phi_1 \neq \Phi$; since *completely-positive* and *trace-preserving* only constrain behaviour on *carrier-mat n n*, this is technically vacuous (every CPTP map admits a CPTP $\Phi_1$ that agrees with it on carrier but disagrees off carrier). The theorem is therefore stated in terms of *extreme-CPTP-carrier*, the carrier-point variant.

The first biconditional below uses different lin-indep notions on each direction (the historical form); *choi-theorem-3-cptp-iff* states the genuine iff:

- *extreme-CPTP-carrier* $\Longrightarrow$ *mat-lin-indep-real* (contrapositive of ⟦*0 < ?n*; *0 < ?d*; *?Φ ∈ CPTP ?n ?d*; *cptp-kraus-rep ?n ?d ?Φ ?Vs*; *lin-indep-complex-mat ?d ?n ?Vs*; ¬ *mat-lin-indep-real ?n* (*kraus-products ?Vs*)⟧ $\Longrightarrow$ ¬ *extreme-CPTP-carrier ?n ?d ?Φ*);
- *mat-lin-indep-complex* $\Longrightarrow$ *extreme-CPTP-carrier* (the strengthened Choi forward direction ⟦*0 < ?n*; *0 < ?d*; *?Φ ∈ CPTP ?n ?d*; *cptp-kraus-rep ?n ?d ?Φ ?Vs*; *mat-lin-indep-complex ?n* (*kraus-products ?Vs*)⟧ $\Longrightarrow$ *extreme-CPTP-carrier ?n ?d ?Φ*, modulo CP Radon–Nikodym).

  The complex-coefficient witness *herm-witness-of-complex* bridges the two notions, so the direction "extreme $\Rightarrow$ independent" also holds over $\mathbb{C}$.

**theorem** *choi-theorem-3-cptp*:
  **fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
    **and** *Vs* :: *complex Matrix.mat list*
  **assumes** *npos*: *0 < n* **and** *dpos*: *0 < d*
      **and** Φ-*in*: *Φ ∈ CPTP n d*
      **and** *rep*: *cptp-kraus-rep n d Φ Vs*
      **and** *minimal*: *lin-indep-complex-mat d n Vs*
  **shows** (*extreme-CPTP-carrier n d Φ*
          ⟶ *mat-lin-indep-real n* (*kraus-products Vs*))
      ∧ (*mat-lin-indep-complex n* (*kraus-products Vs*)
          ⟶ *extreme-CPTP-carrier n d Φ*)
⟨*proof*⟩

With the complex-coefficient witness, the two directions use the same notion of linear independence: Choi's Theorem 3 in the form of the notes' corollary.

**theorem** *choi-theorem-3-cptp-iff*:

**fixes** $\Phi$ :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat*
**and** *Vs* :: *complex Matrix.mat list*
**assumes** *npos*: $0 < n$ **and** *dpos*: $0 < d$
**and** $\Phi$-*in*: $\Phi \in$ *CPTP n d*
**and** *rep*: *cptp-kraus-rep n d* $\Phi$ *Vs*
**and** *minimal*: *lin-indep-complex-mat d n Vs*
**shows** *extreme-CPTP-carrier n d* $\Phi \longleftrightarrow$ *mat-lin-indep-complex n* (*kraus-products Vs*)
⟨*proof*⟩

**The UCP form of Choi's Theorem 3.** In the UCP / Heisenberg convention, the relevant products live in $M_d$ with the *opposite* order $\{V_i\ {V_j}^\dagger\}$.

The project's *kraus-products* uses the CPTP-side convention $\{{V_j}^\dagger\ V_i\}$, and there is no UCP-side carrier-extreme notion or analogue of ⟦*0 < ?n*; *0 < ?d*; *?*$\Phi \in$ *CPTP ?n ?d*; *cptp-kraus-rep ?n ?d ?*$\Phi$ *?Vs*; *lin-indep-complex-mat ?d ?n ?Vs*; ¬ *mat-lin-indep-real ?n* (*kraus-products ?Vs*)⟧ $\Longrightarrow$ ¬ *extreme-CPTP-carrier ?n ?d ?*$\Phi$ in this development. A proper UCP form of Choi's theorem requires either:

- a parallel *kraus-products-ucp* $(V_i\ {V_j}^\dagger)$, an *extreme-UCP-carrier* predicate, and full UCP-side analogues of the carrier-flavoured lemmas; or
- a Hilbert–Schmidt duality between *extreme-UCP* and *extreme-CPTP* that transports the CPTP theorem above.

  The second route is taken in theory Day23_Choi3_UCP, which states the UCP form.

## 53.26 Sanity / easy facts about extreme UCP

If a UCP map has a non-trivial mid-point representation by two distinct UCP maps, it is not extreme *by definition*.

**lemma** *not-extreme-UCP-if-midpoint*:
**assumes** $\Phi_1 \in$ *UCP n d* $\Phi_2 \in$ *UCP n d*
**and** $\Phi_1 \neq \Phi$ **and** $\Phi \neq \Phi_2$
**and** $\Phi \in$ *UCP n d*
**and** $\bigwedge X.\ X \in$ *carrier-mat n n*
$\Longrightarrow \Phi\ X = (1/2 :: \mathit{complex}) \cdot_m \Phi_1\ X$
$+ (1/2 :: \mathit{complex}) \cdot_m \Phi_2\ X$
**shows** ¬ *extreme-UCP n d* $\Phi$
⟨*proof*⟩

The contrapositive: every extreme UCP map, when written as a mid-point of two UCP maps, must have those maps coincide with itself on every $X \in$ *carrier-mat n n*.

**lemma** *extreme-UCP-midpoint-collapses*:
**assumes** *ext*: *extreme-UCP n d* $\Phi$
**and** $\Phi$*1-in*: $\Phi_1 \in$ *UCP n d*
**and** $\Phi$*2-in*: $\Phi_2 \in$ *UCP n d*
**and** *mid*: $\bigwedge X.\ X \in$ *carrier-mat n n*
$\Longrightarrow \Phi\ X = (1/2 :: \mathit{complex}) \cdot_m \Phi_1\ X$
$+ (1/2 :: \mathit{complex}) \cdot_m \Phi_2\ X$
**shows** $\Phi_1 = \Phi \vee \Phi = \Phi_2$
⟨*proof*⟩

This closes the Day-23 stub. The full content of Choi's Theorem 3 awaits a Kraus-list formalisation; the definitions collected above will plug into that proof once available.

**end**

**theory** *Day23-Choi3-UCP*
**imports** *Day22-Adjoint-Maps*
**begin**

# 54 Day 23: Choi's Theorem 3 for unital maps (D23-1)

The notes state Choi's Theorem 3 for UCP maps; *choi-theorem-3-cptp-iff* proves the CPTP form. The UCP form follows by the duality of Day 22: $\Gamma(\Phi) = \Phi^*$ maps midpoint decompositions of a UCP map to midpoint decompositions of its adjoint and back (*extreme-UCP-iff-adjoint*), and the Kraus operators of $\Phi^*$ are the adjoints $W_i^*$, which are linearly independent with the $W_i$. Extremality is the carrier-point notion used for CPTP maps.

The UCP-side midpoint relation and extremality, mirroring *cptp-pair-mid* and *extreme-CPTP-carrier*.

**definition** *ucp-pair-mid* ::
*nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*)
⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*ucp-pair-mid n d* Φ Φ$_1$ Φ$_2$ ≡
Φ$_1$ ∈ *UCP n d* ∧ Φ$_2$ ∈ *UCP n d* ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ$_1$ *X* ∈ *carrier-mat d d*) ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ$_2$ *X* ∈ *carrier-mat d d*) ∧
*lin-on-carrier n* Φ$_1$ ∧ *lin-on-carrier n* Φ$_2$ ∧
(∃ *X*$_1$ ∈ *carrier-mat n n*. Φ$_1$ *X*$_1$ ≠ Φ *X*$_1$) ∧
(∃ *X*$_2$ ∈ *carrier-mat n n*. Φ *X*$_2$ ≠ Φ$_2$ *X*$_2$) ∧
(∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ *X* = (*1/2* :: *complex*) ·$_m$ Φ$_1$ *X* + (*1/2* :: *complex*) ·$_m$ Φ$_2$ *X*)

**definition** *extreme-UCP-carrier* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*extreme-UCP-carrier n d* Φ ≡ Φ ∈ *UCP n d* ∧ ¬ (∃ Φ$_1$ Φ$_2$. *ucp-pair-mid n d* Φ Φ$_1$ Φ$_2$)

**lemma** *lin-on-from-carrier*:
**assumes** ∀ *X*. *X* ∈ *carrier-mat n n* ⟶ Φ *X* ∈ *carrier-mat d d lin-on-carrier n* Φ
**shows** *lin-on n d* (*carrier-mat n n*) Φ
⟨*proof*⟩

**lemma** *lin-on-carrier-from-lin-on*: *lin-on n d* (*carrier-mat n n*) Φ ⟹ *lin-on-carrier n* Φ
⟨*proof*⟩

**lemma** *map-adjoint-half*:
**assumes** *L1*: *lin-on n d* (*carrier-mat n n*) *Φ1* **and** *L2*: *lin-on n d* (*carrier-mat n n*) *Φ2* **and** *Y*: *Y* ∈ *carrier-mat d d*
**shows** *map-adjoint n d* (λ*X*. (*1/2* :: *complex*) ·$_m$ *Φ1 X* + (*1/2* :: *complex*) ·$_m$ *Φ2 X*) *Y*
= (*1/2* :: *complex*) ·$_m$ *map-adjoint n d Φ1 Y* + (*1/2* :: *complex*) ·$_m$ *map-adjoint n d Φ2 Y*
⟨*proof*⟩

Γ carries midpoint decompositions to midpoint decompositions, in both directions.

**lemma** *ucp-mid-imp-cptp-mid*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *mid*: *ucp-pair-mid n d* Φ *Φ1 Φ2* **and** *n*: *0* < *n* **and** *d*: *0* < *d*
**shows** *cptp-pair-mid d n* (*map-adjoint n d* Φ) (*map-adjoint n d Φ1*) (*map-adjoint n d Φ2*)
⟨*proof*⟩

**lemma** *cptp-mid-imp-ucp-mid*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *mid*: *cptp-pair-mid d n* (*map-adjoint n d* Φ) *Ψ1 Ψ2*
**and** *n*: *0* < *n* **and** *d*: *0* < *d*
**shows** *ucp-pair-mid n d* Φ (*map-adjoint d n Ψ1*) (*map-adjoint d n Ψ2*)

⟨*proof*⟩

**theorem** *extreme-UCP-iff-adjoint*:
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *U*: Φ ∈ *UCP n d* **and** *n*: *0* < *n* **and** *d*: *0* < *d*
**shows** *extreme-UCP-carrier n d* Φ ⟷ *extreme-CPTP-carrier d n* (*map-adjoint n d* Φ)
⟨*proof*⟩

Linear independence passes to the adjoints: $\sum_k c_k W_k^* = 0$ gives $\sum_k \bar{c}_k W_k = 0$.

**lemma** *lin-indep-complex-mat-dagger*:
**assumes** *li*: *lin-indep-complex-mat d n Ws*
**shows** *lin-indep-complex-mat n d* (*map* (λ*W*. *W*†) *Ws*)
⟨*proof*⟩

**lemma** *kraus-operators-dagger-of-unital*:
**assumes** *Wc*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n* **and** *un*: *kraus-map-dim d Ws* (*1*$_m$ *n*) = *1*$_m$ *d*
**shows** *kraus-operators d n* (*map* (λ*W*. *W*†) *Ws*)
⟨*proof*⟩

Choi's Theorem 3: a unital CP map $\Phi(X) = \sum_i W_i X W_i^*$ with linearly independent $W_i$ is extreme iff the products $W_j W_i^*$ are linearly independent (*kraus-products* of the adjoints, whose entries are $(W_j^*)^* W_i^* = W_j W_i^*$). The notes write $\Phi(X) = \sum_i V_i^* X V_i$, i.e. $V_i = W_i^*$, and the products $V_i^* V_j$.

**theorem** *choi-theorem-3-ucp*:
**assumes** *n*: *0* < *n* **and** *d*: *0* < *d* **and** *U*: Φ ∈ *UCP n d*
**and** *Wc*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**and** *rep*: ⋀*X*. *X* ∈ *carrier-mat n n* ⟹ Φ *X* = *kraus-map-dim d Ws X*
**and** *li*: *lin-indep-complex-mat d n Ws*
**shows** *extreme-UCP-carrier n d* Φ ⟷ *mat-lin-indep-complex d* (*kraus-products* (*map* (λ*W*. *W*†) *Ws*))
⟨*proof*⟩

**end**
**theory** *Day23-Extremality-Examples*
**imports** *Day22-Kraus-Independence Day25-Arveson-Correspondence*
**begin**

Day 23, examples after Choi's Theorem 3, in the CPTP form and with the Kraus convention $\Phi(X) = \sum_i W_i X W_i^*$ of the formalisation (the notes write $U^* X U$; replace $U$ by $U^*$).

## 54.1 Linear dependence tests

**lemma** *foldr-smult-index*:
**assumes** *Bs*: ∀ *B* ∈ *set Bs*. *B* ∈ *carrier-mat n m* **and** *r*: *r* < *n* **and** *s*: *s* < *m*
**shows** *foldr* (λ*k acc*. *c k* ·$_m$ (*Bs* ! *k*) + *acc*) [*0*..<*length Bs*] (*0*$_m$ *n m*) $$ (*r*,*s*)
= (∑ *k*<*length Bs*. *c k* ∗ *Bs* ! *k* $$ (*r*,*s*))
⟨*proof*⟩

**lemma** *not-indep-of-dup*:
**assumes** *a*: *a* < *length Bs* **and** *b*: *b* < *length Bs* **and** *ab*: *a* ≠ *b* **and** *eq*: *Bs* ! *a* = *Bs* ! *b*
**shows** ¬ *mat-lin-indep-complex n Bs*
⟨*proof*⟩

**lemma** *not-indep-of-zero*:
**assumes** *a*: *a* < *length Bs* **and** *z*: *Bs* ! *a* = *0*$_m$ *n n*
**shows** ¬ *mat-lin-indep-complex n Bs*
⟨*proof*⟩

**lemma** *single-lin-indep*:
**assumes** *W*: $W \in$ *carrier-mat d n* **and** *r*: $r < d$ **and** *s*: $s < n$ **and** *nz*: *W* $$ $(r,s) \neq 0$
**shows** *lin-indep-complex-mat d n* [*W*]
⟨*proof*⟩

**lemma** *single-mat-lin-indep*:
**assumes** *B*: $B \in$ *carrier-mat n n* **and** *r*: $r < n$ **and** *s*: $s < n$ **and** *nz*: *B* $$ $(r,s) \neq 0$
**shows** *mat-lin-indep-complex n* [*B*]
⟨*proof*⟩

**lemma** *nonzero-entry*:
**fixes** *A* :: *complex Matrix.mat*
**assumes** *A*: $A \in$ *carrier-mat m k* **and** *nz*: $A \neq 0_m\ m\ k$
**obtains** *r s* **where** $r < m$ $s < k$ *A* $$ $(r,s) \neq 0$
⟨*proof*⟩

**lemma** *mat-unit-carrier-sq* [*simp*]: $E⟦i,j\ ;\ n,n⟧ \in$ *carrier-mat n n*
⟨*proof*⟩

**lemma** *one-smult-mat'*: $(1::complex) \cdot_m A = A$
⟨*proof*⟩

**lemma** *dagger-smult-mat*: $(c \cdot_m A)^\dagger = cnj\ c \cdot_m (A^\dagger)$
⟨*proof*⟩

## 54.2 Example 1: unitary conjugation is extreme

**theorem** *unitary-conjugation-extreme*:
**assumes** *U*: $U \in$ *carrier-mat n n* **and** *Uu*: *unitary U* **and** *n*: $0 < n$
**shows** *extreme-CPTP-carrier n n* (*kraus-map-dim n* [*U*])
⟨*proof*⟩

## 54.3 Example 2: an average of two unitary conjugations

**definition** *avg2* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat list* **where**
*avg2 U1 U2* = [(*1 / csqrt 2*) $\cdot_m$ *U1*, (*1 / csqrt 2*) $\cdot_m$ *U2*]

**lemma** *half-sq*: *cnj* (*1 / csqrt 2*) ∗ (*1 / csqrt 2*) = *1/2*
⟨*proof*⟩

**lemma** *smult-prod*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A* ∈ *carrier-mat m k B* ∈ *carrier-mat k l*
**shows** $(a \cdot_m A) * (b \cdot_m B) = (a * b) \cdot_m (A * B)$
⟨*proof*⟩

**lemma** *smult-triple*:
**fixes** *A X B* :: *complex Matrix.mat*
**assumes** *A* ∈ *carrier-mat m k X* ∈ *carrier-mat k k B* ∈ *carrier-mat k l*
**shows** $(a \cdot_m A) * X * (b \cdot_m B) = (a * b) \cdot_m (A * X * B)$
⟨*proof*⟩

**lemma** *avg2-products*:
**assumes** *U1*: *U1* ∈ *carrier-mat n n unitary U1* **and** *U2*: *U2* ∈ *carrier-mat n n unitary U2*
**shows** (*avg2 U1 U2* ! *0*)$^\dagger$ ∗ *avg2 U1 U2* ! *0* = *(1/2)* $\cdot_m$ $1_m$ *n*
**and** (*avg2 U1 U2* ! *1*)$^\dagger$ ∗ *avg2 U1 U2* ! *1* = *(1/2)* $\cdot_m$ $1_m$ *n*
⟨*proof*⟩

**lemma** *avg2-kraus*:
**assumes** *U1*: *U1* ∈ *carrier-mat n n unitary U1* **and** *U2*: *U2* ∈ *carrier-mat n n unitary U2*
**shows** *kraus-operators n n* (*avg2 U1 U2*)
⟨*proof*⟩

**lemma** *avg2-map*:
**assumes** *U1*: *U1* ∈ *carrier-mat n n* **and** *U2*: *U2* ∈ *carrier-mat n n* **and** *M*: *M* ∈ *carrier-mat n n*
**shows** *kraus-map-dim n* (*avg2 U1 U2*) *M* = (*1/2*) ·$_m$ (*U1* ∗ *M* ∗ (*U1*$^\dagger$) + *U2* ∗ *M* ∗ (*U2*$^\dagger$))
⟨*proof*⟩

The notes' claim needs $U_1, U_2$ linearly independent: then the products $V_i^* V_j$ contain $\frac{1}{2}I$ twice and $\Phi$ is not extreme.

**theorem** *avg2-not-extreme*:
**assumes** *U1*: *U1* ∈ *carrier-mat n n unitary U1* **and** *U2*: *U2* ∈ *carrier-mat n n unitary U2*
**and** *n*: *0* < *n* **and** *ind*: *lin-indep-complex-mat n n* (*avg2 U1 U2*)
**shows** ¬ *extreme-CPTP-carrier n n* (*kraus-map-dim n* (*avg2 U1 U2*))
⟨*proof*⟩

Without that hypothesis the claim fails: for $U_2 = \lambda U_1$ with $|\lambda| = 1$ the map is $X \mapsto U_1 X U_1^*$, which is extreme.

**theorem** *avg2-proportional-extreme*:
**assumes** *U1*: *U1* ∈ *carrier-mat n n unitary U1* **and** *lam*: *cmod z* = *1* **and** *n*: *0* < *n*
**shows** *extreme-CPTP-carrier n n* (*kraus-map-dim n* (*avg2 U1* (*z* ·$_m$ *U1*)))
⟨*proof*⟩

## 54.4 Example 4: the diagonal map

**lemma** *unit-mult-left-entry*:
**assumes** *Bm*: *Bm* ∈ *carrier-mat n m* **and** *r*: *r* < *p* **and** *k*: *k* < *m* **and** *j*: *j* < *n*
**shows** (*E*⟦*i*,*j* ; *p*,*n*⟧ ∗ *Bm*) $$ (*r*,*k*) = (*if r* = *i then Bm* $$ (*j*,*k*) *else 0*)
⟨*proof*⟩

**lemma** *unit-mult-right-entry*:
**assumes** *Cm*: *Cm* ∈ *carrier-mat p n* **and** *r*: *r* < *p* **and** *s*: *s* < *m* **and** *i*: *i* < *n*
**shows** (*Cm* ∗ *E*⟦*i*,*j* ; *n*,*m*⟧) $$ (*r*,*s*) = (*if s* = *j then Cm* $$ (*r*,*i*) *else 0*)
⟨*proof*⟩

**definition** *diag-kraus* :: *nat* ⇒ *complex Matrix.mat list* **where**
*diag-kraus n* = *map* (λ*i*. *E*⟦*i*,*i* ; *n*,*n*⟧) [*0*..<*n*]

**lemma** *diag-kraus-carrier*: ∀ *W* ∈ *set* (*diag-kraus n*). *W* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *diag-kraus-map*:
**assumes** *M*: *M* ∈ *carrier-mat n n*
**shows** *kraus-map-dim n* (*diag-kraus n*) *M* = *Matrix.mat n n* (λ(*r*,*s*). *if r* = *s then M* $$ (*r*,*r*) *else 0*)
⟨*proof*⟩

**lemma** *diag-kraus-operators*: *kraus-operators n n* (*diag-kraus n*)
⟨*proof*⟩

**lemma** *diag-kraus-lin-indep*: *lin-indep-complex-mat n n* (*diag-kraus n*)
⟨*proof*⟩

**theorem** *diag-not-extreme*:
**assumes** *n*: *2* ≤ *n*
**shows** ¬ *extreme-CPTP-carrier n n* (*kraus-map-dim n* (*diag-kraus n*))
⟨*proof*⟩

### 54.5 Example 3: $X \mapsto \mathrm{Tr}(X)\, I_d$

As written, $\Phi(X) = \mathrm{Tr}(X) I_d$ is neither unital ($\Phi(I_n) = n I_d$) nor trace preserving ($\mathrm{Tr}\,\Phi(X) = d\,\mathrm{Tr}\,X$) unless $n = 1$ resp. $d = 1$. The CPTP normalisation is $\Phi(X) = \mathrm{Tr}(X) I_d/d$, with the $nd$ Kraus operators $E_{ab}/\sqrt{d}$; these are linearly independent, so $\mathrm{cr}(\Phi) = nd > n$ for $d \geq 2$ and $\Phi$ is not extreme (D22-6).

**definition** *tr-kraus* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat list* **where**
*tr-kraus n d* = *map* (λ(*a*,*b*). *complex-of-real* (*1* / *sqrt* (*real d*)) ·$_m$ *E*⟦*a*,*b* ; *d*,*n*⟧) (*List.product* [*0*..<*d*] [*0*..<*n*])

**lemma** *tr-kraus-carrier*: ∀ *W* ∈ *set* (*tr-kraus n d*). *W* ∈ *carrier-mat d n*
⟨*proof*⟩

**lemma** *tr-kraus-length*: *length* (*tr-kraus n d*) = *d* ∗ *n*
⟨*proof*⟩

**lemma** *tr-scalar*:
**assumes** *d*: *0* < *d*
**shows** *cnj* (*complex-of-real* (*1* / *sqrt* (*real d*))) ∗ *complex-of-real* (*1* / *sqrt* (*real d*)) = *1* / *of-nat d*
**and** *complex-of-real* (*1* / *sqrt* (*real d*)) ∗ *cnj* (*complex-of-real* (*1* / *sqrt* (*real d*))) = *1* / *of-nat d*
⟨*proof*⟩

**lemma** *tr-kraus-operators*:
**assumes** *d*: *0* < *d*
**shows** *kraus-operators n d* (*tr-kraus n d*)
⟨*proof*⟩

**lemma** *tr-kraus-map*:
**assumes** *d*: *0* < *d* **and** *M*: *M* ∈ *carrier-mat n n*
**shows** *kraus-map-dim d* (*tr-kraus n d*) *M* = (*Tr M* / *of-nat d*) ·$_m$ *1*$_m$ *d*
⟨*proof*⟩

**lemma** *tr-kraus-lin-indep*:
**assumes** *d*: *0* < *d* **and** *n*: *0* < *n*
**shows** *lin-indep-complex-mat d n* (*tr-kraus n d*)
⟨*proof*⟩

**theorem** *tr-map-not-extreme*:
**assumes** *n*: *0* < *n* **and** *d*: *2* ≤ *d*
**shows** ¬ *extreme-CPTP-carrier n d* (*kraus-map-dim d* (*tr-kraus n d*))
⟨*proof*⟩

**end**
**theory** *Day24-Operator-Systems*
**imports** *Day17-Environment-Model*
**begin**

Day 24: operator systems $S \subseteq M_n$ (subspaces containing $I$ and closed under adjoints), the matrices $M_p(S)$ over them, and completely positive maps on $S$. An element of $M_p(M_n)$ is a $pn \times pn$ matrix whose $(a,b)$ block is *block-of-mat n X a b*; the ampliation $\Phi^{(p)}$ is *amplification n d p* Φ (Day 19).

**definition** *operator-system* :: *nat* ⇒ *complex Matrix.mat set* ⇒ *bool* **where**
*operator-system n OS* ⟷ *OS* ⊆ *carrier-mat n n* ∧ *1*$_m$ *n* ∈ *OS* ∧ (∀ *A* ∈ *OS*. *A*† ∈ *OS*)
∧ (∀ *A* ∈ *OS*. ∀ *B* ∈ *OS*. *A* + *B* ∈ *OS*) ∧ (∀ *A* ∈ *OS*. ∀ *c*. *c* ·$_m$ *A* ∈ *OS*)

**definition** *Mp* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat set* ⇒ *complex Matrix.mat set* **where**
*Mp p n OS* = {*A* ∈ *carrier-mat* (*p* ∗ *n*) (*p* ∗ *n*). ∀ *a* < *p*. ∀ *b* < *p*. *block-of-mat n A a b* ∈ *OS*}

**definition** *lin-on* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat set* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*lin-on n d OS* Φ ⟷ (∀ *A* ∈ *OS*. Φ *A* ∈ *carrier-mat d d*) ∧ (∀ *A* ∈ *OS*. ∀ *B* ∈ *OS*. Φ (*A* + *B*) = Φ *A* + Φ *B*)
∧ (∀ *A* ∈ *OS*. ∀ *c*. Φ (*c* ·$_m$ *A*) = *c* ·$_m$ Φ *A*)

**definition** *p-positive-on* :: *nat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat set* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*p-positive-on n d p OS* Φ ⟷ (∀ *A* ∈ *Mp p n OS*. *psd A* ⟶ *psd* (*amplification n d p* Φ *A*))

**definition** *cp-on* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat set* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *bool* **where**
*cp-on n d OS* Φ ⟷ (∀ *p* > *0*. *p-positive-on n d p OS* Φ)

**lemma** *block-of-mat-dims* [*simp*]: *dim-row* (*block-of-mat n A a b*) = *n dim-col* (*block-of-mat n A a b*) = *n*
⟨*proof*⟩

**lemma** *block-of-mat-idx*:
**assumes** *r* < *n* **and** *c* < *n*
**shows** *block-of-mat n A a b* $$ (*r*,*c*) = *A* $$ (*a*∗*n* + *r*, *b*∗*n* + *c*)
⟨*proof*⟩

**lemma** *operator-systemD*:
**assumes** *operator-system n OS*
**shows** *OS* ⊆ *carrier-mat n n* $1_m$ *n* ∈ *OS* ⋀*A*. *A* ∈ *OS* ⟹ $A^\dagger$ ∈ *OS*
⋀*A B*. *A* ∈ *OS* ⟹ *B* ∈ *OS* ⟹ *A* + *B* ∈ *OS* ⋀*A c*. *A* ∈ *OS* ⟹ *c* ·$_m$ *A* ∈ *OS*
⟨*proof*⟩

**lemma** *operator-system-zero*:
**assumes** *OS*: *operator-system n OS*
**shows** $0_m$ *n n* ∈ *OS*
⟨*proof*⟩

**lemma** *operator-system-diff*:
**assumes** *OS*: *operator-system n OS* **and** *A*: *A* ∈ *OS* **and** *B*: *B* ∈ *OS*
**shows** *A* − *B* ∈ *OS*
⟨*proof*⟩

**lemma** *operator-system-full*: *operator-system n* (*carrier-mat n n*)
⟨*proof*⟩

**lemma** *Mp-full*: *Mp p n* (*carrier-mat n n*) = *carrier-mat* (*p* ∗ *n*) (*p* ∗ *n*)
⟨*proof*⟩

**lemma** *cp-on-full*: *cp-on n d* (*carrier-mat n n*) Φ ⟷ *completely-positive n d* Φ
⟨*proof*⟩

**lemma** *cp-on-mono*:
**assumes** *cp-on n d OS* Φ **and** *OS*′ ⊆ *OS*
**shows** *cp-on n d OS*′ Φ
⟨*proof*⟩

## 54.6 $M_p(OS)$ is an operator system

**lemma** *block-of-mat-add*:
**assumes** *A* ∈ *carrier-mat* (*p* ∗ *n*) (*p* ∗ *n*) *B* ∈ *carrier-mat* (*p* ∗ *n*) (*p* ∗ *n*) *a* < *p b* < *p*

**shows** *block-of-mat n (A + B) a b = block-of-mat n A a b + block-of-mat n B a b*
⟨*proof*⟩

**lemma** *block-of-mat-smult*:
**assumes** $A \in$ *carrier-mat* $(p * n)$ $(p * n)$ $a < p$ $b < p$
**shows** *block-of-mat n* $(k \cdot_m A)$ *a b* $= k \cdot_m$ *block-of-mat n A a b*
⟨*proof*⟩

**lemma** *block-of-mat-dagger*:
**assumes** $A \in$ *carrier-mat* $(p * n)$ $(p * n)$ $a < p$ $b < p$
**shows** *block-of-mat n* $(A^\dagger)$ *a b* $=$ $(\textit{block-of-mat n A b a})^\dagger$
⟨*proof*⟩

**lemma** *block-of-mat-one*:
**assumes** $a < p$ $b < p$
**shows** *block-of-mat n* $(1_m\ (p * n))$ *a b* $=$ (*if a = b then* $1_m$ *n else* $0_m$ *n n*)
⟨*proof*⟩

The note of Day 24: $M_p(S)$ is again an operator system.

**lemma** *Mp-operator-system*:
**assumes** *OS*: *operator-system n OS*
**shows** *operator-system* $(p * n)$ (*Mp p n OS*)
⟨*proof*⟩

## 54.7 Every element is a combination of four positives

**lemma** *herm-bound*:
**assumes** *Hmc*: *Hm* $\in$ *carrier-mat n n* **and** *Hmh*: *hermitian Hm*
**shows** $\exists\, c$::*real*. $0 \le c \wedge$ *psd* (*complex-of-real c* $\cdot_m$ $1_m$ *n* $-$ *Hm*) $\wedge$ *psd* (*complex-of-real c* $\cdot_m$ $1_m$ *n* $+$ *Hm*)
⟨*proof*⟩

Proposition (Day 24): $X = (P_1 - P_2) + i(P_3 - P_4)$ with $P_k \in S^+$.

**proposition** *operator-system-four-positives*:
**assumes** *OS*: *operator-system n OS* **and** *A*: *A* $\in$ *OS*
**shows** $\exists$ *P1 P2 P3 P4*. *P1* $\in$ *OS* $\wedge$ *P2* $\in$ *OS* $\wedge$ *P3* $\in$ *OS* $\wedge$ *P4* $\in$ *OS* $\wedge$ *psd P1* $\wedge$ *psd P2* $\wedge$ *psd P3* $\wedge$ *psd P4*
$\wedge$ *A* $=$ (*P1* $-$ *P2*) $+$ i $\cdot_m$ (*P3* $-$ *P4*)
⟨*proof*⟩

**end**
**theory** *Day25-Arveson-Correspondence*
**imports** *Day24-Operator-Systems*
**begin**

Days 24–25: Arveson's correspondence between maps $\Phi : S \to M_d$ and functionals $S_\Phi : M_d(S) \to \mathbb{C}$, $S_\Phi((A_{ij})) = \frac{1}{d}\sum_{i,j} \Phi(A_{ij})_{ij}$, with inverse $f \mapsto \Phi_f$, $\Phi_f(M) = d\,(f(E_{ij} \otimes M))_{ij}$.

## 54.8 Finite sums of matrices

**definition** *msum* :: *nat* $\Rightarrow$ *nat* $\Rightarrow$ (*'a* $\Rightarrow$ *complex Matrix.mat*) $\Rightarrow$ *'a list* $\Rightarrow$ *complex Matrix.mat* **where**
*msum r c g xs* $=$ *foldr* $(\lambda x\ acc.\ g\ x + acc)$ *xs* $(0_m\ r\ c)$

**lemma** *msum-Nil* [*simp*]: *msum r c g* [] $=$ $0_m$ *r c*
⟨*proof*⟩

**lemma** *msum-Cons* [*simp*]: *msum r c g* $(x \#\ xs) = g\ x +$ *msum r c g xs*

⟨*proof*⟩

**lemma** *msum-carrier*: ∀ *x* ∈ *set xs*. *g x* ∈ *carrier-mat r c* ⟹ *msum r c g xs* ∈ *carrier-mat r c*
⟨*proof*⟩

**lemma** *msum-index*:
**assumes** ∀ *x* ∈ *set xs*. *g x* ∈ *carrier-mat r c* **and** *i* < *r* **and** *j* < *c*
**shows** *msum r c g xs* $$ (*i*,*j*) = (∑ *x*←*xs*. *g x* $$ (*i*,*j*))
⟨*proof*⟩

**lemma** *msum-in-os*:
**assumes** *OS*: *operator-system n OS* **and** *g*: ∀ *x* ∈ *set xs*. *g x* ∈ *OS*
**shows** *msum n n g xs* ∈ *OS*
⟨*proof*⟩

**lemma** *lin-on-zero*:
**assumes** *OS*: *operator-system n OS* **and** *L*: *lin-on n d OS* Φ
**shows** Φ (*0*$_m$ *n n*) = *0*$_m$ *d d*
⟨*proof*⟩

**lemma** *lin-on-msum*:
**assumes** *OS*: *operator-system n OS* **and** *L*: *lin-on n d OS* Φ **and** *g*: ∀ *x* ∈ *set xs*. *g x* ∈ *OS*
**shows** Φ (*msum n n g xs*) = *msum d d* (λ*x*. Φ (*g x*)) *xs*
⟨*proof*⟩

**lemma** *sum-list-product-upt*:
(∑ *x*←*List.product* [*0*..<*a*] [*0*..<*b*]. *h x*) = (∑ *i*<*a*. ∑ *j*<*b*. *h* (*i*,*j*))
⟨*proof*⟩

**lemma** *sum-delta2*:
**fixes** *a d b e* :: *nat*
**assumes** *a*: *a* < *d* **and** *b*: *b* < *e*
**shows** (∑ *i*<*d*. ∑ *j*<*e*. *if i* = *a* ∧ *j* = *b then v else 0*) = *v*
⟨*proof*⟩

## 54.9 $E_{ij} \otimes M$ and the block decomposition

**definition** *unit-block* :: *nat* ⇒ *nat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*unit-block d n i j M* = *Matrix.mat* (*d* ∗ *n*) (*d* ∗ *n*)
(λ(*r*,*c*). *if r div n* = *i* ∧ *c div n* = *j then M* $$ (*r mod n*, *c mod n*) *else 0*)

**lemma** *unit-block-carrier* [*simp*]: *unit-block d n i j M* ∈ *carrier-mat* (*d* ∗ *n*) (*d* ∗ *n*)
⟨*proof*⟩

**lemma** *unit-block-dims* [*simp*]: *dim-row* (*unit-block d n i j M*) = *d* ∗ *n dim-col* (*unit-block d n i j M*) = *d* ∗ *n*
⟨*proof*⟩

*unit-block d n i j M* is $E_{ij} \otimes M$.

**lemma** *unit-block-tensor*:
**assumes** *M*: *M* ∈ *carrier-mat n n* **and** *i*: *i* < *d* **and** *j*: *j* < *d*
**shows** *unit-block d n i j M* = *Tensor.tensor-mat* (*Matrix.mat d d* (λ(*a*,*b*). *if a* = *i* ∧ *b* = *j then 1 else 0*)) *M*
⟨*proof*⟩

**lemma** *block-unit-block*:
**assumes** *M*: *M* ∈ *carrier-mat n n* **and** *a*: *a* < *d* **and** *b*: *b* < *d*
**shows** *block-of-mat n* (*unit-block d n i j M*) *a b* = (*if a* = *i* ∧ *b* = *j then M else 0*$_m$ *n n*)

⟨*proof*⟩

**lemma** *unit-block-Mp*:
**assumes** *OS*: *operator-system n OS* **and** *M*: $M \in OS$ **and** *i*: $i < d$ **and** *j*: $j < d$
**shows** *unit-block d n i j M* ∈ *Mp d n OS*
⟨*proof*⟩

**lemma** *block-decomposition*:
**assumes** *A*: $A \in$ *carrier-mat* $(d * n)$ $(d * n)$
**shows** $A =$ *msum* $(d * n)$ $(d * n)$ $(\lambda(i,j).$ *unit-block d n i j* (*block-of-mat n A i j*)) (*List.product* [*0..<d*] [*0..<d*])
⟨*proof*⟩

## 54.10 The correspondence

**definition** *arveson-functional* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* ⇒ *complex* **where**
*arveson-functional n d* Φ *A* = $(\sum i{<}d.\ \sum j{<}d.$ Φ (*block-of-mat n A i j*) $$ $(i,j))$ / *of-nat d*

**definition** *arveson-map* :: *nat* ⇒ *nat* ⇒ (*complex Matrix.mat* ⇒ *complex*) ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*arveson-map n d f M* = *Matrix.mat d d* $(\lambda(i,j).$ *of-nat d* $*$ *f* (*unit-block d n i j M*))

**definition** *lin-fun-on* :: *complex Matrix.mat set* ⇒ (*complex Matrix.mat* ⇒ *complex*) ⇒ *bool* **where**
*lin-fun-on TS f* ⟷ $(\forall A \in TS.\ \forall B \in TS.\ f\ (A + B) = f\ A + f\ B) \wedge (\forall A \in TS.\ \forall c.\ f\ (c \cdot_m A) = c * f\ A)$

**definition** *positive-fun-on* :: *complex Matrix.mat set* ⇒ (*complex Matrix.mat* ⇒ *complex*) ⇒ *bool* **where**
*positive-fun-on TS f* ⟷ $(\forall A \in TS.$ *psd A* ⟶ $0 \leq f\ A)$

**lemma** *amplification-carrier'*: *amplification n d p* Φ $A \in$ *carrier-mat* $(p * d)$ $(p * d)$
⟨*proof*⟩

**lemma** *amplification-idx*:
**assumes** *r*: $r < q$ **and** *s*: $s < q$ **and** *i*: $i < d$ **and** *j*: $j < d$
**shows** *amplification n d q* Φ *A* $$ $(r * d + i,\ s * d + j) =$ Φ (*block-of-mat n A r s*) $$ $(i,j)$
⟨*proof*⟩

**lemma** *arveson-map-index*: $i < d \Longrightarrow j < d \Longrightarrow$ *arveson-map n d f M* $$ $(i,j) =$ *of-nat d* $*$ *f* (*unit-block d n i j M*)
⟨*proof*⟩

**lemma** *arveson-map-carrier* [*simp*]: *arveson-map n d f M* ∈ *carrier-mat d d*
⟨*proof*⟩

**lemma** *arveson-map-dims* [*simp*]: *dim-row* (*arveson-map n d f M*) = *d dim-col* (*arveson-map n d f M*) = *d*
⟨*proof*⟩

**lemma** *lin-fun-zero*:
**assumes** *TS*: *operator-system N TS* **and** *f*: *lin-fun-on TS f*
**shows** *f* $(0_m\ N\ N) = 0$
⟨*proof*⟩

**lemma** *lin-fun-msum*:
**assumes** *TS*: *operator-system N TS* **and** *f*: *lin-fun-on TS f* **and** *g*: $\forall x \in$ *set xs. g x* ∈ *TS*
**shows** *f* (*msum N N g xs*) = $(\sum x{\leftarrow}xs.\ f\ (g\ x))$
⟨*proof*⟩

$S_{\Phi_f} = f$ on $M_d(S)$.

**theorem** *arveson-functional-of-map*:
**assumes** *OS*: *operator-system n OS* **and** *f*: *lin-fun-on* (*Mp d n OS*) *f* **and** *A*: $A \in Mp\ d\ n\ OS$ **and** *d*: $0 < d$
**shows** *arveson-functional n d* (*arveson-map n d f*) *A* = *f A*
⟨*proof*⟩

$\Phi_{S_\Phi} = \Phi$ on $S$.

**theorem** *arveson-map-of-functional*:
**assumes** *OS*: *operator-system n OS* **and** *L*: *lin-on n d OS* Φ **and** *M*: $M \in OS$ **and** *d*: $0 < d$
**shows** *arveson-map n d* (*arveson-functional n d* Φ) *M* = Φ *M*
⟨*proof*⟩

**lemma** *unit-block-add*:
**assumes** *M*: $M \in$ *carrier-mat n n* **and** *M'*: $M' \in$ *carrier-mat n n*
**shows** *unit-block d n i j* (*M* + *M'*) = *unit-block d n i j M* + *unit-block d n i j M'*
⟨*proof*⟩

**lemma** *unit-block-smult*:
**assumes** *M*: $M \in$ *carrier-mat n n*
**shows** *unit-block d n i j* (*c* $\cdot_m$ *M*) = *c* $\cdot_m$ *unit-block d n i j M*
⟨*proof*⟩

**lemma** *arveson-map-lin*:
**assumes** *OS*: *operator-system n OS* **and** *f*: *lin-fun-on* (*Mp d n OS*) *f*
**shows** *lin-on n d OS* (*arveson-map n d f*)
⟨*proof*⟩

**lemma** *arveson-functional-lin*:
**assumes** *OS*: *operator-system n OS* **and** *L*: *lin-on n d OS* Φ
**shows** *lin-fun-on* (*Mp d n OS*) (*arveson-functional n d* Φ)
⟨*proof*⟩

**lemma** *quad-split*:
**fixes** *M* :: *complex Matrix.mat* **and** *w* :: *complex Matrix.vec*
**assumes** *M*: $M \in$ *carrier-mat* (*q* ∗ *d*) (*q* ∗ *d*) **and** *w*: *dim-vec w* = *q* ∗ *d*
**shows** ⟨*w* | *M* $*_v$ *w*⟩ = ($\sum r{<}q.\ \sum i{<}d.\ \sum s{<}q.\ \sum j{<}d.$
*cnj* (*w* $ (*r* ∗ *d* + *i*)) ∗ *M* $$ (*r* ∗ *d* + *i*, *s* ∗ *d* + *j*) ∗ *w* $ (*s* ∗ *d* + *j*))
⟨*proof*⟩

## 54.11 Theorem (Arveson): completely positive iff $d$-positive iff $S_\Phi$ positive

**lemma** *if-zero-mult*: (*if P then x else 0*) ∗ (*y*::*complex*) = (*if P then x* ∗ *y else 0*)
⟨*proof*⟩

**lemma** *mult-if-zero*: (*y*::*complex*) ∗ (*if P then x else 0*) = (*if P then y* ∗ *x else 0*)
⟨*proof*⟩

**lemma** *cnj-if-zero*: *cnj* (*if P then x else 0*) = (*if P then cnj x else 0*)
⟨*proof*⟩

**lemma** *nonneg-of-real-quad*:
**fixes** *z* :: *complex*
**assumes** *z* ∈ ℝ **and** 0 ≤ *Re z* **and** *d*: $0 < d$
**shows** 0 ≤ *z* / *of-nat d*
⟨*proof*⟩

**lemma** *diag-quad-amplification*:

**fixes** $d\ n$ :: *nat* **and** $\Phi$ :: *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat* **and** $A$ :: *complex Matrix.mat*
**defines** $v \equiv$ *Matrix.vec* $(d * d)$ $(\lambda x.$ *if* $x$ *div* $d = x$ *mod* $d$ *then 1 else* $(0$::*complex*$))$
**shows** $\langle v \mid$ *amplification* $n\ d\ d\ \Phi\ A *_v v\rangle = (\sum i{<}d.\ \sum j{<}d.\ \Phi$ (*block-of-mat* $n\ A\ i\ j$) $\$\$\ (i,j))$
⟨*proof*⟩

**lemma** *dpos-imp-functional-pos*:
**assumes** *OS*: *operator-system* $n$ *OS* **and** $d$: $0 < d$ **and** *P*: *p-positive-on* $n\ d\ d$ *OS* $\Phi$
**shows** *positive-fun-on* (*Mp* $d\ n$ *OS*) (*arveson-functional* $n\ d\ \Phi$)
⟨*proof*⟩

**lemma** *conj-tensor-index*:
**fixes** $w$ :: *complex Matrix.vec*
**assumes** $s$: $s < q$ **and** $k$: $k < d$ **and** $b'$: $b' < n$ **and** $b$: $b < n$
**defines** $V \equiv$ *Tensor.tensor-mat* (*Matrix.mat* $q\ d$ $(\lambda(r,t).\ w\ \$\ (r * d + t)))$ $(1_m\ n)$
**shows** $V\ \$\$\ (s * n + b',\ k * n + b) =$ (*if* $b' = b$ *then* $w\ \$\ (s * d + k)$ *else* $0$)
⟨*proof*⟩

**lemma** *conj-block*:
**fixes** $A$ :: *complex Matrix.mat* **and** $w$ :: *complex Matrix.vec*
**assumes** $A$: $A \in$ *carrier-mat* $(q * n)$ $(q * n)$ **and** $i$: $i < d$ **and** $j$: $j < d$ **and** $a$: $a < n$ **and** $b$: $b < n$
**defines** $V \equiv$ *Tensor.tensor-mat* (*Matrix.mat* $q\ d$ $(\lambda(r,t).\ w\ \$\ (r * d + t)))$ $(1_m\ n)$
**shows** $((V^\dagger) * A * V)\ \$\$\ (i * n + a,\ j * n + b)$
$= (\sum r{<}q.\ \sum s{<}q.\ cnj\ (w\ \$\ (r * d + i)) * w\ \$\ (s * d + j) * A\ \$\$\ (r * n + a,\ s * n + b))$
⟨*proof*⟩

**lemma** *functional-pos-imp-cp*:
**assumes** *OS*: *operator-system* $n$ *OS* **and** *L*: *lin-on* $n\ d$ *OS* $\Phi$ **and** $d$: $0 < d$
**and** *Pos*: *positive-fun-on* (*Mp* $d\ n$ *OS*) (*arveson-functional* $n\ d\ \Phi$)
**shows** *cp-on* $n\ d$ *OS* $\Phi$
⟨*proof*⟩

Theorem (Arveson, Day 24): for linear $\Phi : S \to M_d$, completely positive $\iff$ $d$-positive $\iff$ $S_\Phi$ positive.

**theorem** *arveson-cp-iff*:
**assumes** *OS*: *operator-system* $n$ *OS* **and** *L*: *lin-on* $n\ d$ *OS* $\Phi$ **and** $d$: $0 < d$
**shows** *cp-on* $n\ d$ *OS* $\Phi \longleftrightarrow$ *p-positive-on* $n\ d\ d$ *OS* $\Phi$
**and** *p-positive-on* $n\ d\ d$ *OS* $\Phi \longleftrightarrow$ *positive-fun-on* (*Mp* $d\ n$ *OS*) (*arveson-functional* $n\ d\ \Phi$)
⟨*proof*⟩

Theorem (Arveson correspondence, Day 25): $\Phi \mapsto S_\Phi$ maps $CP(S, M_d)$ onto the positive functionals on $M_d(S)$, with inverse $f \mapsto \Phi_f$; it is affine and sends unital maps to states.

**theorem** *arveson-correspondence*:
**assumes** *OS*: *operator-system* $n$ *OS* **and** $d$: $0 < d$
**shows** *lin-on* $n\ d$ *OS* $\Phi \Longrightarrow$ *cp-on* $n\ d$ *OS* $\Phi \longleftrightarrow$ *positive-fun-on* (*Mp* $d\ n$ *OS*) (*arveson-functional* $n\ d\ \Phi$)
**and** *lin-fun-on* (*Mp* $d\ n$ *OS*) $f \Longrightarrow$ *cp-on* $n\ d$ *OS* (*arveson-map* $n\ d\ f$) $\longleftrightarrow$ *positive-fun-on* (*Mp* $d\ n$ *OS*) $f$
**and** *lin-on* $n\ d$ *OS* $\Phi \Longrightarrow M \in$ *OS* $\Longrightarrow$ *arveson-map* $n\ d$ (*arveson-functional* $n\ d\ \Phi$) $M = \Phi\ M$
**and** *lin-fun-on* (*Mp* $d\ n$ *OS*) $f \Longrightarrow A \in$ *Mp* $d\ n$ *OS* $\Longrightarrow$ *arveson-functional* $n\ d$ (*arveson-map* $n\ d\ f$) $A = f\ A$
⟨*proof*⟩

**lemma** *arveson-functional-affine*:
**assumes** *OS*: *operator-system* $n$ *OS* **and** *L1*: *lin-on* $n\ d$ *OS* $\Phi$ **and** *L2*: *lin-on* $n\ d$ *OS* $\Psi$ **and** $A$: $A \in$ *Mp* $d\ n$ *OS*
**shows** *arveson-functional* $n\ d$ $(\lambda M.\ t \cdot_m \Phi\ M + (1 - t) \cdot_m \Psi\ M)\ A$
$=$ $t *$ *arveson-functional* $n\ d\ \Phi\ A + (1 - t) *$ *arveson-functional* $n\ d\ \Psi\ A$
⟨*proof*⟩

**lemma** *arveson-functional-unital*:
**assumes** *OS*: *operator-system n OS* **and** *L*: *lin-on n d OS* Φ **and** *u*: Φ (*1*$_m$ *n*) = *1*$_m$ *d* **and** *d*: *0* < *d*
**shows** *arveson-functional n d* Φ (*1*$_m$ (*d* ∗ *n*)) = *1*
⟨*proof*⟩

**end**
**theory** *Day25-Unital-Functional-Norm*
**imports** *Day25-Arveson-Correspondence*
*HOL−Analysis.L2-Norm*
**begin**

Day 25 proposition (D25-4): for an operator system $S \subseteq M_n$ and a linear $f : S \to \mathbb{C}$ with $f(I) = 1$, $f$ is positive iff $\|f\| = 1$. The operator norm is $\|A\| = \sup\{\|Av\| : \|v\| \le 1\}$ and, as in the notes, $\|f\|$ is defined for bounded $f$. The proof differs from the notes in one step: for $P \ge 0$ the bound $\|P - sI + itI\|^2 \le s^2 + t^2$ ($s = \|P\|/2$) comes from $P^2 \le \|P\|\,P$, which uses the square root of $P$, instead of from the spectrum of $P$. Cauchy–Schwarz is *cauchy-schwarz-ineq* of Isabelle_Marries_Dirac.

**definition** *op-norm* :: *nat* ⇒ *complex Matrix.mat* ⇒ *real* **where**
*op-norm n A* = *Sup* {‖*A* ∗$_v$ *v*‖ | *v*. *dim-vec v* = *n* ∧ ‖*v*‖ ≤ *1*}

**definition** *fun-norm* :: *nat* ⇒ *complex Matrix.mat set* ⇒ (*complex Matrix.mat* ⇒ *complex*) ⇒ *real* **where**
*fun-norm n TS f* = *Sup* {*cmod* (*f A*) | *A*. *A* ∈ *TS* ∧ *op-norm n A* ≤ *1*}

## 54.12 Vectors

**lemma** *vnorm-nonneg*: ‖*v*‖ ≥ *0*
⟨*proof*⟩

**lemma** *inner-self-norm*: ⟨*v*|*v*⟩ = *complex-of-real* (‖*v*‖$^2$)
⟨*proof*⟩

**lemma** *cauchy-schwarz-norm*:
**assumes** *dim-vec v* = *dim-vec w*
**shows** *cmod* ⟨*v*|*w*⟩ ≤ ‖*v*‖ ∗ ‖*w*‖
⟨*proof*⟩

**lemma** *inner-add-smult-expand*:
**assumes** *dim-vec u* = *dim-vec x*
**shows** ⟨*u* + *z* ·$_v$ *x* | *u* + *z* ·$_v$ *x*⟩ = ⟨*u*|*u*⟩ + *z* ∗ ⟨*u*|*x*⟩ + *cnj z* ∗ ⟨*x*|*u*⟩ + *cnj z* ∗ *z* ∗ ⟨*x*|*x*⟩
⟨*proof*⟩

**lemma** *smult-mat-vec*: *A* ∈ *carrier-mat n m* ⟹ *dim-vec v* = *m* ⟹ (*c* ·$_m$ *A*) ∗$_v$ *v* = *c* ·$_v$ (*A* ∗$_v$ *v*)
⟨*proof*⟩

**lemma** *entry-le-vnorm*: *i* < *dim-vec v* ⟹ *cmod* (*v* $ *i*) ≤ ‖*v*‖
⟨*proof*⟩

**lemma** *vnorm-le-sum*: ‖*w*‖ ≤ (∑ *i*<*dim-vec w*. *cmod* (*w* $ *i*))
⟨*proof*⟩

**lemma** *vnorm-zero-iff*: *dim-vec v* = *n* ⟹ ‖*v*‖ = *0* ⟷ *v* = *0*$_v$ *n*
⟨*proof*⟩

## 54.13 The operator norm

**lemma** *mult-vec-norm-bound*:

**assumes** *A*: $A \in$ *carrier-mat n n* **and** *v*: *dim-vec v* = *n*
**shows** $\|A *_v v\| \leq (\sum i{<}n.\ \sum k{<}n.\ cmod\ (A\ \$\$\ (i,k))) * \|v\|$
⟨*proof*⟩

**lemma** *op-norm-bdd*:
**assumes** *A*: $A \in$ *carrier-mat n n*
**shows** *bdd-above* $\{\|A *_v v\| \mid v.\ dim\text{-}vec\ v = n \wedge \|v\| \leq 1\}$
⟨*proof*⟩

**lemma** *mult-zero-vec'*: $A \in$ *carrier-mat n n* $\Longrightarrow A *_v 0_v\ n = 0_v\ n$
⟨*proof*⟩

**lemma** *op-norm-ge*:
**assumes** *A*: $A \in$ *carrier-mat n n* **and** *v*: *dim-vec v* = *n* $\|v\| \leq 1$
**shows** $\|A *_v v\| \leq$ *op-norm n A*
⟨*proof*⟩

**lemma** *op-norm-nonneg*:
**assumes** *A*: $A \in$ *carrier-mat n n*
**shows** $0 \leq$ *op-norm n A*
⟨*proof*⟩

**lemma** *op-norm-mult*:
**assumes** *A*: $A \in$ *carrier-mat n n* **and** *v*: *dim-vec v* = *n*
**shows** $\|A *_v v\| \leq$ *op-norm n A* $* \|v\|$
⟨*proof*⟩

**lemma** *op-norm-le*:
**assumes** *A*: $A \in$ *carrier-mat n n* **and** *c*: $0 \leq c$ **and** *b*: $\bigwedge v.$ *dim-vec v* = *n* $\Longrightarrow \|A *_v v\| \leq c * \|v\|$
**shows** *op-norm n A* $\leq c$
⟨*proof*⟩

**lemma** *op-norm-one*:
**assumes** *n*: $0 < n$
**shows** *op-norm n* $(1_m\ n) = 1$
⟨*proof*⟩

**lemma** *op-norm-smult-le*:
**assumes** *A*: $A \in$ *carrier-mat n n*
**shows** *op-norm n* $(c \cdot_m A) \leq$ *cmod c* $*$ *op-norm n A*
⟨*proof*⟩

**lemma** *op-norm-zero-imp*:
**assumes** *A*: $A \in$ *carrier-mat n n* **and** *z*: *op-norm n A* $= 0$
**shows** $A = 0_m\ n\ n$
⟨*proof*⟩

## 54.14 Unital functionals

**context**
**fixes** *n* :: *nat* **and** *TS* :: *complex Matrix.mat set* **and** *f* :: *complex Matrix.mat* $\Rightarrow$ *complex*
**assumes** *OS*: *operator-system n TS* **and** *L*: *lin-fun-on TS f* **and** *f1*: *f* $(1_m\ n) = 1$
**begin**

**lemma** *TSc*: $A \in TS \Longrightarrow A \in$ *carrier-mat n n* ⟨*proof*⟩

**lemma** *lin2*: $A \in TS \Longrightarrow B \in TS \Longrightarrow f\ (a \cdot_m A + b \cdot_m B) = a * f\ A + b * f\ B$
⟨*proof*⟩

**lemma** *comb-in*: $A \in TS \Longrightarrow B \in TS \Longrightarrow a \cdot_m A + b \cdot_m B \in TS$
⟨*proof*⟩

**lemma** *f-zero*: $f\ (0_m\ n\ n) = 0$
⟨*proof*⟩

A Hermitian $A \in S$ with $|\Re\langle x|Ax\rangle| \le c\|x\|^2$ satisfies $cI \pm A \ge 0$, so a positive $f$ has $f(A)$ real with $|f(A)| \le c$.

**lemma** *herm-bound*:
**assumes** *pos*: *positive-fun-on TS f* **and** *A*: $A \in TS$ *hermitian A* **and** *c*: $0 \le c$
**and** *b*: $\bigwedge x.\ dim\text{-}vec\ x = n \Longrightarrow |Re\ \langle x|A *_v x\rangle| \le c * \|x\|^2$
**shows** $Im\ (f\ A) = 0$ **and** $|Re\ (f\ A)| \le c$
⟨*proof*⟩

**lemma** *adjoint-inner*:
**assumes** *A*: $A \in carrier\text{-}mat\ n\ n$ **and** *x*: $dim\text{-}vec\ x = n$
**shows** $\langle x|A^\dagger *_v x\rangle = cnj\ \langle x|A *_v x\rangle$
⟨*proof*⟩

($\Rightarrow$): a positive unital $f$ has $|f(A)| \le \|A\|$.

**lemma** *positive-imp-bound*:
**assumes** *pos*: *positive-fun-on TS f* **and** *A*: $A \in TS$
**shows** $cmod\ (f\ A) \le op\text{-}norm\ n\ A$
⟨*proof*⟩

($\Leftarrow$): if $|f(A)| \le \|A\|$ on $S$ then $f$ is positive.

**lemma** *sq-le-norm-quad*:
**assumes** *P*: $P \in carrier\text{-}mat\ n\ n\ psd\ P$ **and** *x*: $dim\text{-}vec\ x = n$
**shows** $\|P *_v x\|^2 \le op\text{-}norm\ n\ P * Re\ \langle x|P *_v x\rangle$
⟨*proof*⟩

**lemma** *bound-imp-positive*:
**assumes** *n*: $0 < n$ **and** *b*: $\forall A \in TS.\ cmod\ (f\ A) \le op\text{-}norm\ n\ A$
**shows** *positive-fun-on TS f*
⟨*proof*⟩

**lemma** *fun-norm-one-iff*:
**assumes** *n*: $0 < n$ **and** *bdd*: $\exists C.\ \forall A \in TS.\ cmod\ (f\ A) \le C * op\text{-}norm\ n\ A$
**shows** $fun\text{-}norm\ n\ TS\ f = 1 \longleftrightarrow (\forall A \in TS.\ cmod\ (f\ A) \le op\text{-}norm\ n\ A)$
⟨*proof*⟩

D25-4.

**theorem** *unital-positive-iff-norm-one*:
**assumes** *n*: $0 < n$ **and** *bdd*: $\exists C.\ \forall A \in TS.\ cmod\ (f\ A) \le C * op\text{-}norm\ n\ A$
**shows** *positive-fun-on TS f* $\longleftrightarrow$ $fun\text{-}norm\ n\ TS\ f = 1$
⟨*proof*⟩

**end**

**end**
**theory** *Day26-Arveson-Extension*
**imports** *Day25-Arveson-Correspondence Vern-Paulsen-QC-3-10-Lmat.Separable-Cone*
**begin**

Day 26: Arveson's extension theorem. The notes extend the state $S_\Phi$ by Hahn–Banach (norm one) and use that a unital functional of norm one is positive. Here the positive functional

$S_\Phi$ on the operator system $M_d(S)$ is extended to a positive functional on $M_d(M_n)$ directly, by separating the positive definite cone from the kernel of $S_\Phi$ (*separating-hyperplane-set-0* in the Euclidean space of typed matrices); the extension is $X \mapsto \mathrm{Tr}(DX)$ with $D \geq 0$. Dimensions are cardinalities of finite types, as in *Separable-Cone*.

## 54.15 Auxiliary facts

**lemma** *operator-system-herm-parts*:
**assumes** *OS*: *operator-system n OS* **and** *A*: $A \in OS$
**shows** $\exists\, Hp\ Kp.\ Hp \in OS \wedge Kp \in OS \wedge$ *hermitian Hp* $\wedge$ *hermitian Kp* $\wedge\ A = Hp + \mathrm{i} \cdot_m Kp$
⟨*proof*⟩

**lemma** *Tr-dagger*:
**assumes** *A*: $A \in$ *carrier-mat n n*
**shows** *Tr* $(A^\dagger) =$ *cnj* (*Tr A*)
⟨*proof*⟩

**lemma** *Tr-herm-mult-real*:
**assumes** *Wc*: $W \in$ *carrier-mat n n* **and** *Mc*: $M \in$ *carrier-mat n n* **and** *Wh*: *hermitian W* **and** *Mh*: *hermitian M*
**shows** *Tr* $(W * M) \in \mathbb{R}$
⟨*proof*⟩

**lemma** *Tr-mult-outer*:
**fixes** *A* :: *complex Matrix.mat* **and** *v* :: *complex Matrix.vec*
**assumes** *A*: $A \in$ *carrier-mat n n* **and** *v*: *dim-vec v* $= n$
**shows** *Tr* $(A * (v \oslash v)) = \langle v \mid A *_v v \rangle$
⟨*proof*⟩

**lemma** *Tr-psd-mult-nonneg*:
**assumes** *Dc*: $D \in$ *carrier-mat n n* **and** *Pc*: $P \in$ *carrier-mat n n* **and** *Dp*: *psd D* **and** *Pp*: *psd P*
**shows** $0 \leq$ *Tr* $(D * P)$
⟨*proof*⟩

**lemma** *herm-entry*:
**assumes** $M \in$ *carrier-mat n n hermitian M* $i < n\ j < n$
**shows** *M* \$\$ (*i*,*j*) = *cnj* (*M* \$\$ (*j*,*i*))
⟨*proof*⟩

**lemma** *herm-real-comb*:
**assumes** *Ac*: $A \in$ *carrier-mat n n* **and** *Bc*: $B \in$ *carrier-mat n n* **and** *Ah*: *hermitian A* **and** *Bh*: *hermitian B*
**shows** *hermitian* (*complex-of-real u* $\cdot_m$ *A* + *complex-of-real v* $\cdot_m$ *B*)
⟨*proof*⟩

**lemma** *herm-real-smult*:
**assumes** *Ac*: $A \in$ *carrier-mat n n* **and** *Ah*: *hermitian A*
**shows** *hermitian* (*complex-of-real t* $\cdot_m$ *A*)
⟨*proof*⟩

**lemma** *hermitian-one*: *hermitian* (*1*$_m$ *n*)
⟨*proof*⟩

**lemma** *lin-t-zero*:
**fixes** *b c* :: *real*
**assumes** $\forall\, t.\ 0 \leq c + t * b$
**shows** $b = 0$

⟨*proof*⟩

**lemma** *eps-nonneg*:
**fixes** *x y* :: *real*
**assumes** ∀ε>0. *0* ≤ *x* + ε ∗ *y*
**shows** *0* ≤ *x*
⟨*proof*⟩

**lemma** *psd-one-mat*: *psd* (*1*$_m$ *n*)
⟨*proof*⟩

## 54.16 Matrices as points of the Euclidean space *lmat*

**lemma** *unwrap-wrap-lm*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *M* ∈ *carrier-mat CARD('k::finite) CARD('k)*
**shows** *unwrap-lm* (*wrap-lm M* :: (*complex*, *'k*, *'k*) *lmat*) = *M*
⟨*proof*⟩

**lemma** *wrap-lm-add*:
**fixes** *M M'* :: *complex Matrix.mat*
**assumes** *M*: *M* ∈ *carrier-mat CARD('k::finite) CARD('k)* **and** *M'*: *M'* ∈ *carrier-mat CARD('k) CARD('k)*
**shows** (*wrap-lm* (*M* + *M'*) :: (*complex*, *'k*, *'k*) *lmat*) = *wrap-lm M* + *wrap-lm M'*
⟨*proof*⟩

**lemma** *wrap-lm-scale*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *M*: *M* ∈ *carrier-mat CARD('k::finite) CARD('k)*
**shows** (*wrap-lm* (*complex-of-real r* ·$_m$ *M*) :: (*complex*, *'k*, *'k*) *lmat*) = *r* ∗$_R$ *wrap-lm M*
⟨*proof*⟩

**lemma** *wrap-lm-zero*: (*wrap-lm* (*0*$_m$ *CARD('k::finite) CARD('k)*) :: (*complex*, *'k*, *'k*) *lmat*) = *0*
⟨*proof*⟩

## 54.17 Extension of positive functionals

A positive functional on an operator system $T \subseteq M_N$ is $X \mapsto \mathrm{Tr}(DX)$ on $T$ for some $D \geq 0$; hence it extends to a positive functional on $M_N$.

**theorem** *positive-functional-extension*:
**fixes** *TS* :: *complex Matrix.mat set* **and** *f* :: *complex Matrix.mat* ⇒ *complex*
**assumes** *TS*: *operator-system CARD('k::finite) TS* **and** *f*: *lin-fun-on TS f* **and** *fpos*: *positive-fun-on TS f*
**shows** ∃ *D*. *D* ∈ *carrier-mat CARD('k) CARD('k)* ∧ *psd D* ∧ (∀ *A* ∈ *TS*. *f A* = *Tr* (*D* ∗ *A*))
⟨*proof*⟩

## 54.18 Arveson's extension theorem

**theorem** *arveson-extension*:
**fixes** *OS* :: *complex Matrix.mat set* **and** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *OS*: *operator-system CARD('n::finite) OS* **and** *L*: *lin-on CARD('n) CARD('d::finite) OS* Φ **and** *CP*: *cp-on CARD('n) CARD('d) OS* Φ
**shows** ∃ Ψ. *lin-on CARD('n) CARD('d)* (*carrier-mat CARD('n) CARD('n)*) Ψ
∧ *completely-positive CARD('n) CARD('d)* Ψ ∧ (∀ *M* ∈ *OS*. Ψ *M* = Φ *M*)
⟨*proof*⟩

Corollary: a completely positive map on an operator system has a Kraus form.

**corollary** *arveson-kraus*:

**fixes** *OS* :: *complex Matrix.mat set* **and** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *OS*: *operator-system CARD('n::finite) OS* **and** *L*: *lin-on CARD('n) CARD('d::finite) OS* Φ
**and** *CP*: *cp-on CARD('n) CARD('d) OS* Φ
**shows** ∃ *Ws*. (∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat CARD('d) CARD('n)*) ∧ (∀ *M* ∈ *OS*. Φ *M* = *kraus-map-dim CARD('d) Ws M*)
⟨*proof*⟩

**end**
**theory** *Day26-CP-Factorization*
**imports** *Day26-Arveson-Extension Day32-Recovery-Proposition*
**begin**

# 55 Day 26: factoring a CP map through a unital or a trace-preserving one (D26-1)

The $i$-th block embedding $B_i : \mathbb{C}^n \to \mathbb{C}^{kn}$. The $B_i$ are isometries with orthogonal ranges, $\sum_i B_i B_i^* = I$, so a Kraus representation $\Phi(X) = \sum_i W_i X W_i^*$ gives $\Phi(X) = V^* \psi(X) V$ with $\psi(X) = \sum_i B_i X B_i^*$ unital and $V = \sum_i B_i W_i^*$.

**definition** *blk-emb* :: *nat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*blk-emb k n i* = *Matrix.mat* (*k* ∗ *n*) *n* (λ(*x*, *y*). *if x div n* = *i* ∧ *x mod n* = *y then 1 else 0*)

**lemma** *blk-emb-carrier* [*simp*]: *blk-emb k n i* ∈ *carrier-mat* (*k* ∗ *n*) *n*
⟨*proof*⟩

**lemma** *blk-emb-dims* [*simp*]: *dim-row* (*blk-emb k n i*) = *k* ∗ *n dim-col* (*blk-emb k n i*) = *n*
⟨*proof*⟩

**lemma** *blk-cond*:
**fixes** *x y i n* :: *nat*
**assumes** *y*: *y* < *n*
**shows** (*x div n* = *i* ∧ *x mod n* = *y*) ⟷ *x* = *i* ∗ *n* + *y*
⟨*proof*⟩

**lemma** *blk-less*:
**fixes** *i y k n* :: *nat*
**assumes** *i* < *k y* < *n*
**shows** *i* ∗ *n* + *y* < *k* ∗ *n*
⟨*proof*⟩

**lemma** *foldr-add-carrier-rect*:
∀ *x* ∈ *set xs*. *f x* ∈ *carrier-mat a b* ⟹ *foldr* (λ*x acc*. *f x* + *acc*) *xs* (0$_m$ *a b*) ∈ *carrier-mat a b*
⟨*proof*⟩

**lemma** *foldr-add-index*:
**assumes** ∀ *x* ∈ *set xs*. *f x* ∈ *carrier-mat a b i* < *a j* < *b*
**shows** *foldr* (λ*x acc*. *f x* + *acc*) *xs* (0$_m$ *a b*) $$ (*i*, *j*) = (∑ *x*←*xs*. *f x* $$ (*i*, *j*))
⟨*proof*⟩

**lemma** *blk-emb-gram-entry*:
**assumes** *x*: *x* < *k* ∗ *n* **and** *y*: *y* < *k* ∗ *n*
**shows** (*blk-emb k n i* ∗ ((*blk-emb k n i*)$^\dagger$)) $$ (*x*, *y*) =
(*if x div n* = *i* ∧ *y div n* = *i* ∧ *x mod n* = *y mod n then 1 else 0*)
⟨*proof*⟩

**lemma** *blk-emb-gram*:
*foldr* (λ*B acc*. *B* ∗ (*B*$^\dagger$) + *acc*) (*map* (*blk-emb k n*) [0..<*k*]) (0$_m$ (*k* ∗ *n*) (*k* ∗ *n*)) = 1$_m$ (*k* ∗ *n*)

⟨*proof*⟩

**lemma** *kraus-lin-on*:
**assumes** ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n*
**shows** *lin-on n d* (*carrier-mat n n*) (*kraus-map-dim d Ws*)
⟨*proof*⟩

**lemma** *kraus-compose-single*: *kraus-compose* [*A*] *Es* = *map* (λ*E*. *A* ∗ *E*) *Es*
⟨*proof*⟩

$V$ has the blocks $W_i^*$ stacked vertically, so $V^*B_i = W_i$.

**definition** *kraus-stack-dag* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* **where**
*kraus-stack-dag n d Ws* = *Matrix.mat* (*length Ws* ∗ *n*) *d* (λ(*x*, *c*). *cnj* (*Ws* ! (*x div n*) $$ (*c*, *x mod n*)))

**lemma** *kraus-stack-dag-blk*:
**assumes** *Wc*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n* **and** *i*: *i* < *length Ws*
**shows** (*kraus-stack-dag n d Ws*)† ∗ *blk-emb* (*length Ws*) *n i* = *Ws* ! *i*
⟨*proof*⟩

Proposition (Day 26): a CP map $\Phi : S \to M_d$ on an operator system with $\Phi(I) = P$ is $\Phi(X) = V^*\psi(X)V$ with $\psi$ unital and completely positive (here even on all of $M_n$) and $V^*V = P$. The notes compress $\Phi$ to the range of $P$; here the Kraus form of *arveson-kraus* is used.

**theorem** *cp-unital-factor*:
**fixes** *OS* :: *complex Matrix.mat set* **and** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *OS*: *operator-system CARD*('*n*::*finite*) *OS* **and** *L*: *lin-on CARD*('*n*) *CARD*('*d*::*finite*) *OS* Φ
**and** *CP*: *cp-on CARD*('*n*) *CARD*('*d*) *OS* Φ
**shows** ∃ *r V ψ*. *0* < *r* ∧ *V* ∈ *carrier-mat r CARD*('*d*) ∧ *ψ* ∈ *UCP CARD*('*n*) *r*
∧ *lin-on CARD*('*n*) *r* (*carrier-mat CARD*('*n*) *CARD*('*n*)) *ψ*
∧ (∀ *X* ∈ *OS*. Φ *X* = (*V*†) ∗ *ψ X* ∗ *V*) ∧ (*V*†) ∗ *V* = Φ (*1*$_m$ *CARD*('*n*))
⟨*proof*⟩

$V$ with the blocks $W_i$ stacked vertically, so $B_i^*V = W_i$.

**definition** *kraus-stack* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat list* ⇒ *complex Matrix.mat* **where**
*kraus-stack n d Ws* = *Matrix.mat* (*length Ws* ∗ *d*) *n* (λ(*x*, *c*). *Ws* ! (*x div d*) $$ (*x mod d*, *c*))

**lemma** *kraus-stack-blk*:
**assumes** *Wc*: ∀ *W* ∈ *set Ws*. *W* ∈ *carrier-mat d n* **and** *i*: *i* < *length Ws*
**shows** (*blk-emb* (*length Ws*) *d i*)† ∗ *kraus-stack n d Ws* = *Ws* ! *i*
⟨*proof*⟩

Proposition (Day 26): a CP map $\Phi : M_n \to M_d$ is $\Phi(X) = \psi(VXV^*)$ with $\psi : M_r \to M_d$ completely positive and trace preserving. The notes dualise the previous proposition; here $\psi(Y) = \sum_i B_i^*YB_i$ and $V = \sum_i B_iW_i$ directly.

**theorem** *cp-tp-factor*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *L*: *lin-on CARD*('*n*::*finite*) *CARD*('*d*::*finite*) (*carrier-mat CARD*('*n*) *CARD*('*n*)) Φ
**and** *CP*: *completely-positive CARD*('*n*) *CARD*('*d*) Φ
**shows** ∃ *r V ψ*. *0* < *r* ∧ *V* ∈ *carrier-mat r CARD*('*n*) ∧ *ψ* ∈ *CPTP r CARD*('*d*)
∧ *lin-on r CARD*('*d*) (*carrier-mat r r*) *ψ*
∧ (∀ *X* ∈ *carrier-mat CARD*('*n*) *CARD*('*n*). Φ *X* = *ψ* (*V* ∗ *X* ∗ (*V*†)))
⟨*proof*⟩

**end**

# Part IV
# Functional-analytic infrastructure (Lecture 25: Hahn–Banach)

**theory** *Matrix-Typed*
**imports**
*HOL−Hahn-Banach.Hahn-Banach*
*Jordan-Normal-Form.Matrix*
*HOL−Library.Cardinality*
*Vern-Paulsen-QC-3-10.Basic-Math*
**begin**

## 56 Typed-dimension matrix wrapper

The type $('a, 'm :: finite, 'n :: finite)$ *lmat* represents matrices of size $CARD('m) \times CARD('n)$ with entries in $'a$, by typedef of the JNF matrix type at the appropriate dimension.

**typedef** (**overloaded**) $('a, 'm :: finite, 'n :: finite)$ *lmat* =
$\{M :: 'a$ *Matrix.mat*. *dim-row* $M = CARD('m) \wedge$ *dim-col* $M = CARD('n)\}$
**morphisms** *unwrap-lm wrap-lm*
⟨*proof*⟩

**setup-lifting** *type-definition-lmat*

**lemma** *unwrap-lm-dim-row* [*simp*]:
*dim-row* (*unwrap-lm* $(M :: ('a, 'm :: finite, 'n :: finite)$ *lmat*$)) = CARD('m)$
⟨*proof*⟩

**lemma** *unwrap-lm-dim-col* [*simp*]:
*dim-col* (*unwrap-lm* $(M :: ('a, 'm :: finite, 'n :: finite)$ *lmat*$)) = CARD('n)$
⟨*proof*⟩

**lemma** *unwrap-lm-carrier* [*simp*]:
*unwrap-lm* $(M :: ('a, 'm :: finite, 'n :: finite)$ *lmat*$)$
$\in$ *carrier-mat* $(CARD('m))$ $(CARD('n))$
⟨*proof*⟩

## 57 Typeclass instances

The unique zero matrix of the right shape.

**instantiation** *lmat* :: (*zero*, *finite*, *finite*) *zero*
**begin**
**lift-definition** *zero-lmat* :: $('a, 'm :: finite, 'n :: finite)$ *lmat*
**is** $0_m$ $(CARD('m))$ $(CARD('n))$ ⟨*proof*⟩
**instance** ⟨*proof*⟩
**end**

**lemma** *unwrap-lm-zero* [*simp*]:
*unwrap-lm* $(0 :: ('a :: zero, 'm :: finite, 'n :: finite)$ *lmat*$)$
$= 0_m$ $(CARD('m))$ $(CARD('n))$
⟨*proof*⟩

**instantiation** *lmat* :: (*plus*, *finite*, *finite*) *plus*
**begin**
**lift-definition** *plus-lmat* ::
(*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat* ⇒ (*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat* ⇒ (*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
**is** (+) ⟨*proof*⟩
**instance** ⟨*proof*⟩
**end**

**lemma** *unwrap-lm-plus* [*simp*]:
*unwrap-lm* ((*A* :: (*'a* :: *plus*, *'m* :: *finite*, *'n* :: *finite*) *lmat*) + *B*)
= *unwrap-lm A* + *unwrap-lm B*
⟨*proof*⟩

**instantiation** *lmat* :: (*uminus*, *finite*, *finite*) *uminus*
**begin**
**lift-definition** *uminus-lmat* ::
(*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat* ⇒ (*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
**is** *uminus* ⟨*proof*⟩
**instance** ⟨*proof*⟩
**end**

**lemma** *unwrap-lm-uminus* [*simp*]:
*unwrap-lm* (− (*A* :: (*'a* :: *uminus*, *'m* :: *finite*, *'n* :: *finite*) *lmat*))
= − *unwrap-lm A*
⟨*proof*⟩

**instantiation** *lmat* :: (*minus*, *finite*, *finite*) *minus*
**begin**
**lift-definition** *minus-lmat* ::
(*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat* ⇒ (*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat* ⇒ (*'a*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
**is** (−) ⟨*proof*⟩
**instance** ⟨*proof*⟩
**end**

**lemma** *unwrap-lm-minus* [*simp*]:
*unwrap-lm* ((*A* :: (*'a* :: *minus*, *'m* :: *finite*, *'n* :: *finite*) *lmat*) − *B*)
= *unwrap-lm A* − *unwrap-lm B*
⟨*proof*⟩

Lift the additive structure laws via ⟦⋀*i j*. ⟦*i* < *dim-row ?B*; *j* < *dim-col ?B*⟧ ⟹ *?A* $$ (*i*, *j*) = *?B* $$ (*i*, *j*); *dim-row ?A* = *dim-row ?B*; *dim-col ?A* = *dim-col ?B*⟧ ⟹ *?A* = *?B* on the underlying JNF matrices.

**instance** *lmat* :: (*semigroup-add*, *finite*, *finite*) *semigroup-add*
⟨*proof*⟩

**instance** *lmat* :: (*ab-semigroup-add*, *finite*, *finite*) *ab-semigroup-add*
⟨*proof*⟩

**instance** *lmat* :: (*monoid-add*, *finite*, *finite*) *monoid-add*
⟨*proof*⟩

**instance** *lmat* :: (*comm-monoid-add*, *finite*, *finite*) *comm-monoid-add*
⟨*proof*⟩

**instance** *lmat* :: (*group-add*, *finite*, *finite*) *group-add*
⟨*proof*⟩

**instance** *lmat* :: (*ab-group-add*, *finite*, *finite*) *ab-group-add*
⟨*proof*⟩

## 58 Real scalar multiplication and Hahn-Banach overloading

HOL's $(\cdot)$ is the abstract real-scalar multiplication used by the Hahn-Banach locales. We overload it on the type (*complex*, $'m$, $'n$) *lmat* as multiplication by the embedded real, lifted from JNF's $(\cdot_m)$.

**overloading**
*prod-lmat* ≡
*Vector-Space.prod* ::
*real* ⇒ (*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat*
⇒ (*complex*, $'m$, $'n$) *lmat*
**begin**
**definition** *prod-lmat* ::
*real* ⇒ (*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat*
⇒ (*complex*, $'m$, $'n$) *lmat*
**where**
*prod-lmat r M* = *wrap-lm* (*complex-of-real r* $\cdot_m$ *unwrap-lm M*)
**end**

**lemma** *unwrap-lm-prod* [*simp*]:
*unwrap-lm* (*Vector-Space.prod r* (*M* :: (*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat*))
= *complex-of-real r* $\cdot_m$ *unwrap-lm M*
⟨*proof*⟩

## 59 Real-vectorspace locale on the full type

Every (*complex*, $'m$, $'n$) *lmat* with the universe set is a real vector space in HOL/Hahn_Banach's sense.

**lemma** *vectorspace-UNIV-lmat*:
*Vector-Space.vectorspace* (*UNIV* :: (*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat set*)
⟨*proof*⟩

## 60 The Hilbert-Schmidt norm on (*complex*, $'m$, $'n$) *lmat*

The HS / Frobenius norm: $\|M\|_H S = sqrt\ (Re\ \langle\langle M \mid M\rangle\rangle)$. We define it on (*complex*, $'m$, $'n$) *lmat* using the underlying matrix's HS inner product from *Basic-Math*.

**definition** *hs-norm-lmat* ::
(*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat* ⇒ *real*
**where**
*hs-norm-lmat M* = *sqrt* (*Re* (⟨⟨ *unwrap-lm M* | *unwrap-lm M* ⟩⟩))

**lemma** *hs-norm-lmat-nonneg*: *0* ≤ *hs-norm-lmat M*
⟨*proof*⟩

**lemma** *hs-norm-lmat-sq*:
(*hs-norm-lmat M*)$^2$ = *Re* (⟨⟨ *unwrap-lm M* | *unwrap-lm M* ⟩⟩)
⟨*proof*⟩

**lemma** *hs-norm-lmat-smult*:
*hs-norm-lmat* (*Vector-Space.prod a M*) = |*a*| ∗ *hs-norm-lmat M*
⟨*proof*⟩

# 61 Triangle inequality for the HS norm

Cauchy-Schwarz on the HS inner product (real form), used in the triangle inequality proof.

**lemma** *hs-inner-prod-real-self*:
⟨⟨ *A* | *A* ⟩⟩ = *complex-of-real* (*Re* (⟨⟨ *A* | *A* ⟩⟩))
⟨*proof*⟩

**lemma** *hs-inner-prod-left-add*:
**assumes** *dim-row A1* = *dim-row B* **and** *dim-col A1* = *dim-col B*
**and** *dim-row A2* = *dim-row B* **and** *dim-col A2* = *dim-col B*
**shows** ⟨⟨ *A1* + *A2* | *B* ⟩⟩ = ⟨⟨ *A1* | *B* ⟩⟩ + ⟨⟨ *A2* | *B* ⟩⟩
⟨*proof*⟩

**lemma** *hs-cauchy-schwarz-re*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *dr*: *dim-row A* = *dim-row B* **and** *dc*: *dim-col A* = *dim-col B*
**shows** (*Re* (⟨⟨ *A* | *B* ⟩⟩))$^2$ ≤ *Re* (⟨⟨ *A* | *A* ⟩⟩) ∗ *Re* (⟨⟨ *B* | *B* ⟩⟩)
⟨*proof*⟩

**lemma** *hs-norm-lmat-triangle*:
**fixes** *M N* :: (*complex*, ′*m* :: *finite*, ′*n* :: *finite*) *lmat*
**shows** *hs-norm-lmat* (*M* + *N*) ≤ *hs-norm-lmat M* + *hs-norm-lmat N*
⟨*proof*⟩

# 62 Subspace, seminorm, normed-vectorspace instances

HS norm is a seminorm on the whole real vector space of (*complex*, ′*m*, ′*n*) *lmat*.

**lemma** *seminorm-UNIV-hs-norm-lmat*:
*seminorm* (*UNIV* :: (*complex*, ′*m* :: *finite*, ′*n* :: *finite*) *lmat set*) *hs-norm-lmat*
⟨*proof*⟩

HS norm is a (real) norm: it vanishes only on the zero matrix.

**lemma** *hs-norm-lmat-zero-iff*:
**fixes** *M* :: (*complex*, ′*m* :: *finite*, ′*n* :: *finite*) *lmat*
**shows** (*hs-norm-lmat M* = *0*) ⟷ (*M* = *0*)
⟨*proof*⟩

**lemma** *norm-UNIV-hs-norm-lmat*:
*Normed-Space.norm*
(*UNIV* :: (*complex*, ′*m* :: *finite*, ′*n* :: *finite*) *lmat set*) *hs-norm-lmat*
⟨*proof*⟩

**lemma** *normed-vectorspace-UNIV-lmat*:
*normed-vectorspace*
(*UNIV* :: (*complex*, ′*m* :: *finite*, ′*n* :: *finite*) *lmat set*) *hs-norm-lmat*
⟨*proof*⟩

A complex-linear subspace V of (*complex*, ′*m*, ′*n*) *lmat* (i.e. nonempty, closed under addition and complex-scalar multiplication) is in particular a real-linear subspace, hence a *subspace* of *UNIV* in HOL/Hahn_Banach's sense.

**definition** *cmat-subspace-lmat* ::
(*complex*, ′*m* :: *finite*, ′*n* :: *finite*) *lmat set* ⇒ *bool*
**where**
*cmat-subspace-lmat V* ≡
*V* ≠ {} ∧

$(\forall x \in V.\ \forall y \in V.\ x + y \in V) \wedge$
$(\forall x \in V.\ \forall c :: complex.$
$wrap\text{-}lm\ (c \cdot_m unwrap\text{-}lm\ x) \in V)$

**lemma** *subspace-of-cmat-subspace-lmat*:
**fixes** $V :: (complex, 'm :: finite, 'n :: finite)\ lmat\ set$
**assumes** *cmat-subspace-lmat V*
**shows** *Subspace.subspace V* $(UNIV :: (complex, 'm, 'n)\ lmat\ set)$
⟨*proof*⟩

# 63 Complex scalar multiplication on lmat

Complex scalar multiplication on $(complex, 'm, 'n)\ lmat$, extending the *Vector-Space.prod* overload (which is only by reals).

**definition** *cmult-lmat* ::
$complex \Rightarrow (complex, 'm :: finite, 'n :: finite)\ lmat$
$\Rightarrow (complex, 'm, 'n)\ lmat$
**where**
$cmult\text{-}lmat\ c\ M = wrap\text{-}lm\ (c \cdot_m unwrap\text{-}lm\ M)$

**lemma** *unwrap-lm-cmult* [*simp*]:
*unwrap-lm* $(cmult\text{-}lmat\ c\ (M :: (complex, 'm :: finite, 'n :: finite)\ lmat))$
$= c \cdot_m unwrap\text{-}lm\ M$
⟨*proof*⟩

**lemma** *cmult-lmat-real*:
$cmult\text{-}lmat\ (complex\text{-}of\text{-}real\ a)\ M = Vector\text{-}Space.prod\ a\ M$
⟨*proof*⟩

**lemma** *cmult-lmat-add*:
**fixes** $x\ y :: (complex, 'm :: finite, 'n :: finite)\ lmat$
**shows** $cmult\text{-}lmat\ c\ (x + y) = cmult\text{-}lmat\ c\ x + cmult\text{-}lmat\ c\ y$
⟨*proof*⟩

**lemma** *cmult-lmat-scalar-add*:
**fixes** $x :: (complex, 'm :: finite, 'n :: finite)\ lmat$
**shows** $cmult\text{-}lmat\ (c + d)\ x = cmult\text{-}lmat\ c\ x + cmult\text{-}lmat\ d\ x$
⟨*proof*⟩

**lemma** *cmult-lmat-scalar-mult*:
**fixes** $x :: (complex, 'm :: finite, 'n :: finite)\ lmat$
**shows** $cmult\text{-}lmat\ (c * d)\ x = cmult\text{-}lmat\ c\ (cmult\text{-}lmat\ d\ x)$
⟨*proof*⟩

**lemma** *cmult-lmat-one*:
**fixes** $x :: (complex, 'm :: finite, 'n :: finite)\ lmat$
**shows** $cmult\text{-}lmat\ 1\ x = x$
⟨*proof*⟩

**lemma** *cmult-lmat-uminus-one*:
**fixes** $x :: (complex, 'm :: finite, 'n :: finite)\ lmat$
**shows** $cmult\text{-}lmat\ (-\ 1)\ x = -\ x$
⟨*proof*⟩

**lemma** *cmult-lmat-i-sq*:
**fixes** $x :: (complex, 'm :: finite, 'n :: finite)\ lmat$
**shows** $cmult\text{-}lmat\ \mathrm{i}\ (cmult\text{-}lmat\ \mathrm{i}\ x) = -\ x$

⟨*proof*⟩

**lemma** *hs-norm-lmat-cmult*:
**fixes** $M$ :: (*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
**shows** *hs-norm-lmat* (*cmult-lmat c M*) = *cmod c* ∗ *hs-norm-lmat M*
⟨*proof*⟩

**lemma** *hs-norm-lmat-cmult-i* [*simp*]:
**fixes** $M$ :: (*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
**shows** *hs-norm-lmat* (*cmult-lmat* i *M*) = *hs-norm-lmat M*
⟨*proof*⟩

# 64 Day 25 (26 October 2011): the Hahn–Banach theorem

**Definition (notes).** Let $W$ be a normed space. A linear functional $f : W \to \mathbb{C}$ is *bounded* if $|f(w)| \le C\|w\|$ for some constant $C$; the least such $C$ is the norm $\|f\| = \sup\{|f(w)| : \|w\| \le 1\}$.

**Theorem (notes)** (Hahn–Banach)**.** Let $W$ be a normed space, $V \subseteq W$ a subspace and $g : V \to \mathbb{C}$ a bounded linear functional. Then there is a bounded linear functional $f : W \to \mathbb{C}$ with $\|f\| = \|g\|$ and $f(v) = g(v)$ for all $v \in V$.

In the notes the theorem is used (Day 26) to extend the state $S_\Phi$ from $M_d(S)$ to $M_d(M_n)$ in the proof of Arveson's extension theorem, which is not formalised. Here it is proved for $W = (\mathbb{C}$-valued $m \times n$ matrices) with the Hilbert–Schmidt norm: HOL's Hahn–Banach gives a real-linear norm-preserving extension of $\Re g$, and the Bohnenblust–Sobczyk construction $f(x) = F(x) - iF(ix)$ recovers a complex-linear extension with the same bound.

# 65 Bounded complex-linear functionals on lmat

**definition** *clinear-functional-on* ::
(*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat set*
⇒ ((*complex*, *'m*, *'n*) *lmat* ⇒ *complex*) ⇒ *bool*
**where**
*clinear-functional-on V g* ≡
(∀ *x* ∈ *V*. ∀ *y* ∈ *V*. *g* (*x* + *y*) = *g x* + *g y*) ∧
(∀ *x* ∈ *V*. ∀ *c*. *g* (*cmult-lmat c x*) = *c* ∗ *g x*)

**definition** *bounded-clinear-functional-on* ::
(*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat set*
⇒ ((*complex*, *'m*, *'n*) *lmat* ⇒ *complex*) ⇒ *bool*
**where**
*bounded-clinear-functional-on V g* ≡
*clinear-functional-on V g* ∧
(∃ *C* ≥ *0*. ∀ *x* ∈ *V*. *cmod* (*g x*) ≤ *C* ∗ *hs-norm-lmat x*)

# 66 Real-part functional and linearform/continuous instances

**definition** *re-part-lmat* ::
((*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat* ⇒ *complex*)
⇒ ((*complex*, *'m*, *'n*) *lmat* ⇒ *real*)
**where**
*re-part-lmat g x* = *Re* (*g x*)

**lemma** *linearform-re-part-lmat*:
**fixes** $V$ :: (*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat set*

**and** $g :: (complex, 'm, 'n)\ lmat \Rightarrow complex$
**assumes** *V-sub*: *Subspace.subspace V UNIV*
**and** *g-clin*: *clinear-functional-on V g*
**shows** *linearform V* (*re-part-lmat g*)
⟨*proof*⟩

**lemma** *continuous-re-part-lmat*:
**fixes** $V :: (complex, 'm :: finite, 'n :: finite)\ lmat\ set$
**and** $g :: (complex, 'm, 'n)\ lmat \Rightarrow complex$
**assumes** *V-sub*: *Subspace.subspace V UNIV*
**and** *g-bnd*: *bounded-clinear-functional-on V g*
**shows** *continuous V* (*re-part-lmat g*) *hs-norm-lmat*
⟨*proof*⟩

# 67 Application of HOL/Hahn_Banach (norm-preserving extension)

Apply HOL/Hahn_Banach to extend the real-linearform *re-part-lmat g* to all of UNIV preserving the function norm.

**theorem** *hahn-banach-re-part-lmat*:
**fixes** $V :: (complex, 'm :: finite, 'n :: finite)\ lmat\ set$
**and** $g :: (complex, 'm, 'n)\ lmat \Rightarrow complex$
**assumes** *V-sub*: *Subspace.subspace V* ($UNIV :: (complex, 'm, 'n)\ lmat\ set$)
**and** *g-bnd*: *bounded-clinear-functional-on V g*
**shows**
$\exists F :: (complex, 'm, 'n)\ lmat \Rightarrow real.$
*linearform UNIV F*
$\wedge$ *continuous UNIV F hs-norm-lmat*
$\wedge\ (\forall x \in V.\ F\ x = re\text{-}part\text{-}lmat\ g\ x)$
⟨*proof*⟩

# 68 Bohnenblust-Sobczyk: complex extension from real extension

Given a real-linear extension $F$ of *re-part-lmat g* on all of UNIV, define the complex-linear extension $f\ x = F\ x - i \cdot F\ (i \cdot x)$.

**definition** *bs-complex-extension* ::
$((complex, 'm :: finite, 'n :: finite)\ lmat \Rightarrow real)$
$\Rightarrow ((complex, 'm, 'n)\ lmat \Rightarrow complex)$
**where**
*bs-complex-extension F x* =
*complex-of-real* (*F x*) − i ∗ *complex-of-real* (*F* (*cmult-lmat* i *x*))

**lemma** *bs-complex-extension-add*:
**fixes** $F :: (complex, 'm :: finite, 'n :: finite)\ lmat \Rightarrow real$
**assumes** *F-lin*: *linearform UNIV F*
**shows** *bs-complex-extension F* $(x + y)$
$=$ *bs-complex-extension F x* + *bs-complex-extension F y*
⟨*proof*⟩

**lemma** *bs-complex-extension-real-smult*:
**fixes** $F :: (complex, 'm :: finite, 'n :: finite)\ lmat \Rightarrow real$
**assumes** *F-lin*: *linearform UNIV F*
**shows** *bs-complex-extension F* (*Vector-Space.prod a x*)

$= \mathit{complex\text{-}of\text{-}real}\ a * \mathit{bs\text{-}complex\text{-}extension}\ F\ x$
⟨*proof*⟩

**lemma** *bs-complex-extension-i-smult*:
**fixes** $F :: (\mathit{complex}, 'm :: \mathit{finite}, 'n :: \mathit{finite})\ \mathit{lmat} \Rightarrow \mathit{real}$
**assumes** *F-lin*: *linearform UNIV F*
**shows** $\mathit{bs\text{-}complex\text{-}extension}\ F\ (\mathit{cmult\text{-}lmat}\ \mathrm{i}\ x)$
$= \mathrm{i} * \mathit{bs\text{-}complex\text{-}extension}\ F\ x$
⟨*proof*⟩

**lemma** *bs-complex-extension-csmult*:
**fixes** $F :: (\mathit{complex}, 'm :: \mathit{finite}, 'n :: \mathit{finite})\ \mathit{lmat} \Rightarrow \mathit{real}$
**assumes** *F-lin*: *linearform UNIV F*
**shows** $\mathit{bs\text{-}complex\text{-}extension}\ F\ (\mathit{cmult\text{-}lmat}\ c\ x)$
$= c * \mathit{bs\text{-}complex\text{-}extension}\ F\ x$
⟨*proof*⟩

**lemma** *bs-complex-extension-clinear*:
**fixes** $F :: (\mathit{complex}, 'm :: \mathit{finite}, 'n :: \mathit{finite})\ \mathit{lmat} \Rightarrow \mathit{real}$
**assumes** *F-lin*: *linearform UNIV F*
**shows** $\mathit{clinear\text{-}functional\text{-}on}\ \mathit{UNIV}\ (\mathit{bs\text{-}complex\text{-}extension}\ F)$
⟨*proof*⟩

The exact Bohnenblust-Sobczyk bound: any $C$ bound on $F$ lifts verbatim to a bound on *bs-complex-extension F*. This is the standard $e^{-i\vartheta}$-rotation argument used to prove the complex Hahn-Banach theorem from the real one.

**lemma** *bs-complex-extension-tight-bound*:
**fixes** $F :: (\mathit{complex}, 'm :: \mathit{finite}, 'n :: \mathit{finite})\ \mathit{lmat} \Rightarrow \mathit{real}$
**assumes** *F-lin*: *linearform UNIV F*
**and** *F-C*: $\bigwedge x.\ |F\ x| \leq C * \mathit{hs\text{-}norm\text{-}lmat}\ x$
**shows** $\mathit{cmod}\ (\mathit{bs\text{-}complex\text{-}extension}\ F\ x) \leq C * \mathit{hs\text{-}norm\text{-}lmat}\ x$
⟨*proof*⟩

**lemma** *bs-complex-extension-bounded*:
**fixes** $F :: (\mathit{complex}, 'm :: \mathit{finite}, 'n :: \mathit{finite})\ \mathit{lmat} \Rightarrow \mathit{real}$
**assumes** *F-lin*: *linearform UNIV F*
**and** *F-cont*: *continuous UNIV F hs-norm-lmat*
**shows** $\exists\, C \geq 0.\ \forall x.\ \mathit{cmod}\ (\mathit{bs\text{-}complex\text{-}extension}\ F\ x) \leq C * \mathit{hs\text{-}norm\text{-}lmat}\ x$
⟨*proof*⟩

**lemma** *bs-complex-extension-extends*:
**fixes** $V :: (\mathit{complex}, 'm :: \mathit{finite}, 'n :: \mathit{finite})\ \mathit{lmat}\ \mathit{set}$
**and** $g :: (\mathit{complex}, 'm, 'n)\ \mathit{lmat} \Rightarrow \mathit{complex}$
**and** $F :: (\mathit{complex}, 'm, 'n)\ \mathit{lmat} \Rightarrow \mathit{real}$
**assumes** *g-clin*: *clinear-functional-on V g*
**and** *V-clos-i*: $\forall x \in V.\ \mathit{cmult\text{-}lmat}\ \mathrm{i}\ x \in V$
**and** *F-ext*: $\forall x \in V.\ F\ x = \mathit{re\text{-}part\text{-}lmat}\ g\ x$
**shows** $\forall x \in V.\ \mathit{bs\text{-}complex\text{-}extension}\ F\ x = g\ x$
⟨*proof*⟩

# 69 The matrix-space Hahn-Banach theorem

Combine all the steps: a bounded complex-linear functional on a complex-linear subspace V of $(\mathit{complex}, 'm, 'n)\ \mathit{lmat}$ extends to a bounded complex-linear functional on all of $(\mathit{complex}, 'm, 'n)\ \mathit{lmat}$ (without claiming norm preservation; for the application to Arveson Extension this is enough together with the bound chase).

**theorem** *matrix-hahn-banach-lmat*:
**fixes** $V$ :: (*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat set*
**and** $g$ :: (*complex*, $'m$, $'n$) *lmat* $\Rightarrow$ *complex*
**assumes** *V-csub*: *cmat-subspace-lmat V*
**and** *g-bnd*: *bounded-clinear-functional-on V g*
**shows** $\exists f$ :: (*complex*, $'m$, $'n$) *lmat* $\Rightarrow$ *complex*.
*bounded-clinear-functional-on UNIV f*
$\wedge$ ($\forall x \in V$. $f\ x = g\ x$)
⟨*proof*⟩

The explicit-bound form of the matrix Hahn-Banach theorem: any $C$-bound on $|Re \circ g|$ over the subspace $V$ lifts verbatim to a $C$-bound on $cmod \circ f$ over the whole space.

The proof unfolds HOL/Hahn_Banach's locale-local *fn-norm* to its *Sup*-based definition, bounds the supremum on the subspace side by $C$ via *g-re-C*, transports the bound to the extension side via *norm-Hahn-Banach*'s norm-preservation conclusion, then uses *bs-complex-extension-tight-bound* to turn the bound on the real extension into a bound on the complex extension.

**theorem** *matrix-hahn-banach-lmat-with-bound*:
**fixes** $V$ :: (*complex*, $'m$ :: *finite*, $'n$ :: *finite*) *lmat set*
**and** $g$ :: (*complex*, $'m$, $'n$) *lmat* $\Rightarrow$ *complex*
**and** $C$ :: *real*
**assumes** *V-csub*: *cmat-subspace-lmat V*
**and** *g-clin*: *clinear-functional-on V g*
**and** *g-re-C*: $\forall x \in V$. $|Re\ (g\ x)| \leq C * hs\text{-}norm\text{-}lmat\ x$
**and** *C-nn*: $0 \leq C$
**shows** $\exists f$ :: (*complex*, $'m$, $'n$) *lmat* $\Rightarrow$ *complex*.
*clinear-functional-on UNIV f*
$\wedge$ ($\forall x \in V$. $f\ x = g\ x$)
$\wedge$ ($\forall x$. $cmod\ (f\ x) \leq C * hs\text{-}norm\text{-}lmat\ x$)
⟨*proof*⟩

**end**

**theory** *Essential-Functional-Analysis*
**imports**
*Matrix-Typed*
*Quantum-Computing-3-10*
*Jordan-Normal-Form.Schur-Decomposition*
*Jordan-Normal-Form.DL-Rank*
**begin**

# 70 Overview

The notes take the analysis of finite-dimensional spaces for granted ("it is easy to see that it is closed", Day 22; "$C$ is a closed cone", Day 28). This theory supplies those facts for the matrix space $M_n$ with the Hilbert–Schmidt norm $\|A\| = \sqrt{\mathrm{Tr}(A^*A)}$: the norm axioms, the parallelogram law, entrywise bounds, completeness, closedness of the sets of Hermitian and of positive semidefinite matrices, and Carathéodory's theorem in the form needed on Day 28: a non-negative combination of product matrices $\sigma \otimes \tau$ in $M_{mp}$ can be rewritten using at most $2(mp)^2 + 1$ of them.

# 71 HS inner-product algebra

**lemma** *hs-inner-prod-self-eq-of-real*:

$\langle\langle\ A \mid A\ \rangle\rangle = \textit{complex-of-real}\ (\textit{Re}\ (\langle\langle\ A \mid A\ \rangle\rangle))$
⟨*proof*⟩

**lemma** *hs-inner-prod-self-zero-iff*:
**fixes** $A :: \textit{complex Matrix.mat}$
**shows** $\langle\langle\ A \mid A\ \rangle\rangle = 0 \longleftrightarrow A = 0_m\ (\textit{dim-row}\ A)\ (\textit{dim-col}\ A)$
⟨*proof*⟩

**lemma** *Re-hs-inner-prod-right-smult-real*:
**fixes** $A\ B :: \textit{complex Matrix.mat}$ **and** $t :: \textit{real}$
**assumes** $\textit{dim-row}\ A = \textit{dim-row}\ B$ **and** $\textit{dim-col}\ A = \textit{dim-col}\ B$
**shows** $\textit{Re}\ (\langle\langle\ A \mid \textit{complex-of-real}\ t \cdot_m B\ \rangle\rangle) = t * \textit{Re}\ (\langle\langle\ A \mid B\ \rangle\rangle)$
⟨*proof*⟩

**lemma** *Re-hs-inner-prod-swap*:
**fixes** $A\ B :: \textit{complex Matrix.mat}$
**assumes** $\textit{dim-row}\ A = \textit{dim-row}\ B$ **and** $\textit{dim-col}\ A = \textit{dim-col}\ B$
**shows** $\textit{Re}\ (\langle\langle\ A \mid B\ \rangle\rangle) = \textit{Re}\ (\langle\langle\ B \mid A\ \rangle\rangle)$
⟨*proof*⟩

# 72 Trace lemmas

**lemma** *Tr-dagger*:
**fixes** $M :: \textit{complex Matrix.mat}$
**assumes** *M-c*: $M \in \textit{carrier-mat}\ n\ n$
**shows** $\textit{Tr}\ (M^\dagger) = \textit{cnj}\ (\textit{Tr}\ M)$
⟨*proof*⟩

**lemma** *Tr-smult-mat*:
**fixes** $M :: \textit{complex Matrix.mat}$ **and** $c :: \textit{complex}$
**assumes** *M-c*: $M \in \textit{carrier-mat}\ n\ n$
**shows** $\textit{Tr}\ (c \cdot_m M) = c * \textit{Tr}\ M$
⟨*proof*⟩

**lemma** *Re-hs-inner-eq-Re-Tr-dagger*:
**fixes** $A\ B :: \textit{complex Matrix.mat}$
**assumes** *A-c*: $A \in \textit{carrier-mat}\ n\ n$
**and** *B-c*: $B \in \textit{carrier-mat}\ n\ n$
**shows** $\textit{Re}\ (\langle\langle\ A \mid B\ \rangle\rangle) = \textit{Re}\ (\textit{Tr}\ ((A^\dagger) * B))$
⟨*proof*⟩

# 73 HS norm definition and basic facts

HS / Frobenius norm on raw complex matrices. Definition duplicates the one in *Matrix-Typed* for the typed wrapper, but here for the raw matrices.

**definition** $\textit{hs-norm} :: \textit{complex Matrix.mat} \Rightarrow \textit{real}$ **where**
$\textit{hs-norm}\ A \equiv \textit{sqrt}\ (\textit{Re}\ (\langle\langle\ A \mid A\ \rangle\rangle))$

**lemma** *hs-norm-nonneg*: $\textit{hs-norm}\ A \geq 0$
⟨*proof*⟩

**lemma** *hs-norm-sq*:
$(\textit{hs-norm}\ A)^2 = \textit{Re}\ (\langle\langle\ A \mid A\ \rangle\rangle)$
⟨*proof*⟩

**lemma** *hs-norm-zero*: $\textit{hs-norm}\ (0_m\ m\ n) = 0$

⟨*proof*⟩

**lemma** *hs-norm-smult*:
*hs-norm* (*c* ·$_m$ *A*) = *cmod c* ∗ *hs-norm A*
⟨*proof*⟩

Triangle inequality for the HS norm via Cauchy-Schwarz.

**lemma** *hs-norm-triangle*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A-c*: *A* ∈ *carrier-mat n n*
**and** *B-c*: *B* ∈ *carrier-mat n n*
**shows** *hs-norm* (*A* + *B*) ≤ *hs-norm A* + *hs-norm B*
⟨*proof*⟩

HS-norm-squared of $U + (-t) \cdot V$, expanded along the bilinear structure. We extract this into a stand-alone lemma so the parameter names do not clash with let-bound names in callers.

**lemma** *hs-norm-sq-diff-expand*:
**fixes** *U V* :: *complex Matrix.mat* **and** *t* :: *real*
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**and** *V-c*: *V* ∈ *carrier-mat n n*
**shows** (*hs-norm* (*U* + *complex-of-real* (−*t*) ·$_m$ *V*))$^2$
= *Re* (⟨⟨ *U* | *U* ⟩⟩) − *2* ∗ *t* ∗ *Re* (⟨⟨ *U* | *V* ⟩⟩)
+ *t*$^2$ ∗ *Re* (⟨⟨ *V* | *V* ⟩⟩)
⟨*proof*⟩

HS norm satisfies the parallelogram identity.

**lemma** *hs-norm-parallelogram*:
**fixes** *U V* :: *complex Matrix.mat*
**assumes** *U-c*: *U* ∈ *carrier-mat n n*
**and** *V-c*: *V* ∈ *carrier-mat n n*
**shows** (*hs-norm* (*U* + *V*))$^2$ + (*hs-norm* (*U* − *V*))$^2$
= *2* ∗ (*hs-norm U*)$^2$ + *2* ∗ (*hs-norm V*)$^2$
⟨*proof*⟩

**lemma** *hs-norm-zero-iff*:
**fixes** *A* :: *complex Matrix.mat*
**shows** *hs-norm A* = *0* ⟷ *A* = *0*$_m$ (*dim-row A*) (*dim-col A*)
⟨*proof*⟩

Every entry of a matrix is bounded in modulus-squared by the HS-norm squared. Useful for transferring HS-norm convergence to entry-wise convergence.

**lemma** *cmod-entry-sq-le-hs-norm-sq*:
**fixes** *A* :: *complex Matrix.mat* **and** *i j* :: *nat*
**assumes** *A-c*: *A* ∈ *carrier-mat n n*
**and** *i-lt*: *i* < *n* **and** *j-lt*: *j* < *n*
**shows** (*cmod* (*A* $$ (*i*,*j*)))$^2$ ≤ (*hs-norm A*)$^2$
⟨*proof*⟩

Modulus form of the entry bound: $|A_{i,j}| \le \|A\|_H S$.

**lemma** *cmod-entry-le-hs-norm*:
**fixes** *A* :: *complex Matrix.mat* **and** *i j* :: *nat*
**assumes** *A-c*: *A* ∈ *carrier-mat n n*
**and** *i-lt*: *i* < *n* **and** *j-lt*: *j* < *n*
**shows** *cmod* (*A* $$ (*i*,*j*)) ≤ *hs-norm A*
⟨*proof*⟩

# 74 HS-norm convergence vs entry-wise convergence

HS-norm convergence implies entry-wise convergence: if $(M_n)$ converges to $L$ in HS-norm then for every $(i, j)$ the entry sequence $M_n$ \$\$ $(i, j)$ tends to $L$ \$\$ $(i, j)$.

**lemma** *HS-lim-entry-lim*:
  **fixes** *M-seq* :: *nat* ⇒ *complex Matrix.mat* **and** *r* :: *nat*
    **and** *i j* :: *nat* **and** *L* :: *complex Matrix.mat*
  **assumes** *carrier*: ∀ *n*. *M-seq n* ∈ *carrier-mat r r*
    **and** *L-c*: *L* ∈ *carrier-mat r r*
    **and** *lim*: (λ*n*. *hs-norm* (*M-seq n* − *L*)) ⟶ *0*
    **and** *i-lt*: *i* < *r* **and** *j-lt*: *j* < *r*
  **shows** (λ*n*. *M-seq n* \$\$ (*i*,*j*)) ⟶ *L* \$\$ (*i*,*j*)
⟨*proof*⟩

HS-Cauchy implies entry-wise Cauchy: the dual of *HS-lim-entry-lim*. Combined with the completeness of $\mathbb{C}$ it yields a candidate limit for every entry.

**lemma** *HS-Cauchy-entry-Cauchy*:
  **fixes** *M-seq* :: *nat* ⇒ *complex Matrix.mat* **and** *r* :: *nat*
    **and** *i j* :: *nat*
  **assumes** *carrier*: ∀ *n*. *M-seq n* ∈ *carrier-mat r r*
    **and** *hs-cauchy*: ∀ *ε*>*0*. ∃ *N*. ∀ *x* ≥ *N*. ∀ *y* ≥ *N*.
                  *hs-norm* (*M-seq x* − *M-seq y*) < *ε*
    **and** *i-lt*: *i* < *r* **and** *j-lt*: *j* < *r*
  **shows** *Cauchy* (λ*n*. *M-seq n* \$\$ (*i*,*j*))
⟨*proof*⟩

HS-norm-squared as a real double sum of entry-modulus squares. This is the entry-wise expansion of $\|A\|^2 = Re\ \langle\langle\ A\ |\ A\ \rangle\rangle$.

**lemma** *hs-norm-sq-as-entry-sum*:
  **fixes** *A* :: *complex Matrix.mat*
  **assumes** *A-c*: *A* ∈ *carrier-mat r r*
  **shows** (*hs-norm A*)²
      = (∑ *i*∈{*0*..<*r*}. ∑ *j*∈{*0*..<*r*}. (*cmod* (*A* \$\$ (*i*,*j*)))²)
⟨*proof*⟩

Pointwise entry bounds give an HS-norm-squared bound.

**lemma** *hs-norm-sq-le-from-entry-bounds*:
  **fixes** *A* :: *complex Matrix.mat*
  **assumes** *A-c*: *A* ∈ *carrier-mat r r*
    **and** *entries*: ⋀*i j*. *i* < *r* ⟹ *j* < *r* ⟹ (*cmod* (*A* \$\$ (*i*,*j*)))² ≤ *M*
  **shows** (*hs-norm A*)² ≤ *real r* ∗ *real r* ∗ *M*
⟨*proof*⟩

Entry-cmod bound gives an HS-norm bound. Combines the *hs-norm-sq-le-from-entry-bounds* bound with *sqrt*.

**lemma** *hs-norm-le-from-entry-bounds*:
  **fixes** *A* :: *complex Matrix.mat* **and** *δ* :: *real*
  **assumes** *A-c*: *A* ∈ *carrier-mat r r*
    **and** *δ-nn*: *0* ≤ *δ*
    **and** *entries*: ⋀*i j*. *i* < *r* ⟹ *j* < *r* ⟹ *cmod* (*A* \$\$ (*i*,*j*)) ≤ *δ*
  **shows** *hs-norm A* ≤ *real r* ∗ *δ*
⟨*proof*⟩

Entry-wise convergence implies HS-norm convergence. This is the slow half of HS-completeness; we use the entry-bound helper to keep it tractable.

**lemma** *entry-lim-imp-HS-lim*:

**fixes** *M-seq* :: *nat* ⇒ *complex Matrix.mat* **and** *M-lim* :: *complex Matrix.mat*
**and** *r* :: *nat*
**assumes** *carrier*: ∀ *n*. *M-seq n* ∈ *carrier-mat r r*
**and** *M-lim-c*: *M-lim* ∈ *carrier-mat r r*
**and** *entry-lim*: ⋀*i j*. *i* < *r* ⟹ *j* < *r*
⟹ (λ*n*. *M-seq n* $$ (*i*,*j*)) ⟶ *M-lim* $$ (*i*,*j*)
**shows** (λ*n*. *hs-norm* (*M-seq n* − *M-lim*)) ⟶ *0*
⟨*proof*⟩

HS-completeness of the matrix space: every HS-Cauchy sequence in *carrier-mat r r* converges in HS norm to a matrix in *carrier-mat r r*. Combines *HS-Cauchy-entry-Cauchy* + completeness of $\mathbb{C}$ + *entry-lim-imp-HS-lim*.

**lemma** *HS-Cauchy-imp-convergent*:
**fixes** *M-seq* :: *nat* ⇒ *complex Matrix.mat* **and** *r* :: *nat*
**assumes** *carrier*: ∀ *n*. *M-seq n* ∈ *carrier-mat r r*
**and** *hs-cauchy*: ∀ *ε*>*0*. ∃ *N*. ∀ *x* ≥ *N*. ∀ *y* ≥ *N*.
*hs-norm* (*M-seq x* − *M-seq y*) < *ε*
**shows** ∃ *M-lim*. *M-lim* ∈ *carrier-mat r r*
∧ (∀ *ε*>*0*. ∃ *N*. ∀ *n* ≥ *N*. *hs-norm* (*M-seq n* − *M-lim*) < *ε*)
⟨*proof*⟩

Symmetry of HS-distance: $\|A - B\| = \|B - A\|$.

**lemma** *hs-norm-sub-swap*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A-c*: *A* ∈ *carrier-mat n n*
**and** *B-c*: *B* ∈ *carrier-mat n n*
**shows** *hs-norm* (*A* − *B*) = *hs-norm* (*B* − *A*)
⟨*proof*⟩

HS-norm closedness of the Hermitian set.

**lemma** *hermitian-HS-closed*:
**fixes** *M-seq* :: *nat* ⇒ *complex Matrix.mat* **and** *L* :: *complex Matrix.mat*
**and** *r* :: *nat*
**assumes** *carrier*: ∀ *n*. *M-seq n* ∈ *carrier-mat r r*
**and** *L-c*: *L* ∈ *carrier-mat r r*
**and** *herm-n*: ∀ *n*. *hermitian* (*M-seq n*)
**and** *lim*: (λ*n*. *hs-norm* (*M-seq n* − *L*)) ⟶ *0*
**shows** *hermitian L*
⟨*proof*⟩

HS-norm closedness of the PSD set. The proof combines hermitian_HS_closed for the Hermitian part with continuity of the quadratic form $\sigma \mapsto \langle x \mid \sigma *_v x\rangle$ (linear in the entries of $\sigma$, hence HS-continuous via entry convergence).

**lemma** *psd-HS-closed*:
**fixes** *σ-n* :: *nat* ⇒ *complex Matrix.mat* **and** *σ* :: *complex Matrix.mat*
**and** *r* :: *nat*
**assumes** *carrier*: ∀ *n*. *σ-n n* ∈ *carrier-mat r r*
**and** *σ-c*: *σ* ∈ *carrier-mat r r*
**and** *psd-n*: ∀ *n*. *psd* (*σ-n n*)
**and** *lim*: (λ*n*. *hs-norm* (*σ-n n* − *σ*)) ⟶ *0*
**shows** *psd σ*
⟨*proof*⟩

Reverse triangle inequality: $|\|A\| - \|B\|| \le \|A - B\|$.

**lemma** *hs-norm-sub-triangle-ineq*:
**fixes** *A B* :: *complex Matrix.mat*

**assumes** *A-c*: $A \in$ *carrier-mat n n*
**and** *B-c*: $B \in$ *carrier-mat n n*
**shows** $|$*hs-norm A* $-$ *hs-norm B*$| \leq$ *hs-norm* $(A - B)$
⟨*proof*⟩

# 75 Carathéodory's theorem for conic combinations

Carathéodory's theorem for cones: in a real vector space $V$ of dimension $d$, every element of the conic hull of a set $S$ (a finite non-negative combination of elements of $S$) is already a non-negative combination of at most $d$ elements of $S$. We need it for $V$ the real space of Hermitian $(mp)\times(mp)$ matrices, of real dimension $(mp)^2$, and $S$ the product positive matrices $\sigma \otimes \tau$: it bounds the length of the decompositions of elements of the separable cone uniformly, which is what makes the cone closed (Separable_Cone). HOL-Analysis has Carathéodory for *euclidean-space*; using it directly would require a Euclidean-space instance on the Hermitian matrices, so the theorem is proved here directly for lists: any $d+1$ vectors are linearly dependent, and renormalising the weights along a dependence relation removes one term while keeping all weights non-negative.

Below is a fold-based weighted sum of the cone-style list, suitable for stating Carathéodory at the matrix level.

**fun** *cone-list-sum* ::
*nat* ⇒ *nat* ⇒
(*real* × *complex Matrix.mat* × *complex Matrix.mat*) *list* ⇒
*complex Matrix.mat*
**where**
*cone-list-sum m p* [] = $0_m$ (*m*∗*p*) (*m*∗*p*)
| *cone-list-sum m p* ((*w*, *σ*, *τ*) # *ts*) =
*complex-of-real w* ·$_m$ (*σ* ⨂ *τ*) + *cone-list-sum m p ts*

**lemma** *cone-list-sum-carrier*:
*cone-list-sum m p ts* ∈ *carrier-mat* (*m*∗*p*) (*m*∗*p*)
⟨*proof*⟩

*cone-list-sum* is additive over list-append.

**lemma** *cone-list-sum-append*:
*cone-list-sum m p* (*xs* @ *ys*)
= *cone-list-sum m p xs* + *cone-list-sum m p ys*
⟨*proof*⟩

A zero-weight singleton over a properly-sized matrix pair contributes the zero matrix.

**lemma** *cone-list-sum-zero-singleton*:
**assumes** *σ-c*: *σ* ∈ *carrier-mat m m* **and** *τ-c*: *τ* ∈ *carrier-mat p p*
**shows** *cone-list-sum m p* [(*0*, *σ*, *τ*)] = $0_m$ (*m*∗*p*) (*m*∗*p*)
⟨*proof*⟩

Dropping a zero-weight entry from a cone list (with properly typed matrix part) preserves the cone sum.

**lemma** *cone-list-sum-drop-zero*:
**assumes** *i-lt*: *i* < *length xs*
**and** *zero-w*: *fst* (*xs* ! *i*) = *0*
**and** *typing*: ∀(*w*, *σ*, *τ*) ∈ *set xs*.
*σ* ∈ *carrier-mat m m* ∧ *τ* ∈ *carrier-mat p p*
**shows** *cone-list-sum m p* (*take i xs* @ *drop* (*Suc i*) *xs*)
= *cone-list-sum m p xs*
⟨*proof*⟩

Linearity of *cone-list-sum* in the weight. Two lists with matching matrix parts but weights related by $w_k = {w_k}^1 + \mu * r_k$ satisfy the corresponding sum-equation.

**lemma** *cone-list-sum-weight-split*:
  **fixes** *f g h* :: *nat* ⇒ *real*
    **and** $\sigma f\ \tau f$ :: *nat* ⇒ *complex Matrix.mat*
    **and** $\mu$ :: *real* **and** *L* :: *nat*
  **assumes** *typing*: $\bigwedge k.\ k < L \Longrightarrow$
                $\sigma f\ k \in$ *carrier-mat m m* $\wedge\ \tau f\ k \in$ *carrier-mat p p*
      **and** *split*: $\bigwedge k.\ k < L \Longrightarrow f\ k = g\ k + \mu * h\ k$
  **shows** *cone-list-sum m p* (*map* ($\lambda k.$ (*f k*, $\sigma f\ k$, $\tau f\ k$)) [*0*..<*L*])
      = *cone-list-sum m p* (*map* ($\lambda k.$ (*g k*, $\sigma f\ k$, $\tau f\ k$)) [*0*..<*L*])
        + *complex-of-real* $\mu\ \cdot_m$
          *cone-list-sum m p* (*map* ($\lambda k.$ (*h k*, $\sigma f\ k$, $\tau f\ k$)) [*0*..<*L*])
⟨*proof*⟩

Entry-wise expression for *cone-list-sum*: each entry is a HOL sum over the list indices of weight times tensor entry.

**lemma** *cone-list-sum-entry*:
  **fixes** *m p* :: *nat*
      **and** *ts* :: (*real* × *complex Matrix.mat* × *complex Matrix.mat*) *list*
  **assumes** *typing*: $\forall (w, \sigma, \tau) \in$ *set ts*.
                $\sigma \in$ *carrier-mat m m* $\wedge\ \tau \in$ *carrier-mat p p*
      **and** *i-lt*: $i < m * p$ **and** *j-lt*: $j < m * p$
  **shows** *cone-list-sum m p ts* $$ (*i*, *j*)
        = ($\sum k$<*length ts*.
            *complex-of-real* (*fst* (*ts*!*k*))
              * (*fst* (*snd* (*ts*!*k*)) ⨂ *snd* (*snd* (*ts*!*k*))) $$ (*i*, *j*))
⟨*proof*⟩

An entry-wise zero criterion: if every entry is 0, the matrix is $0_m$.

**lemma** *matrix-eq-zero-via-entries*:
  **fixes** *M* :: *complex Matrix.mat*
  **assumes** *M-c*: *M* ∈ *carrier-mat* (*m*∗*p*) (*m*∗*p*)
      **and** *zero*: $\bigwedge i\ j.\ i < m*p \Longrightarrow j < m*p \Longrightarrow M\ \$\$\ (i, j) = 0$
  **shows** $M = 0_m\ (m*p)\ (m*p)$
⟨*proof*⟩

Top-level linear-algebra core. A list of more than $2\ (m\ p)^2$ complex $(m\ p) \times (m\ p)$ matrix pairs admits a non-trivial real linear combination of their tensor products that vanishes.

The proof is the standard JNF *vec-space TYPE*(*real*) ($2\ (m\ p)^2$) *li-le-dim* contrapositive applied to a real-flatten of the tensor products. Admitted here as a single named lemma so it can be discharged independently of the cone Carathéodory reduction machinery (which is fully proved below modulo this one lemma).

**lemma** *cone-list-real-lin-dep*:
  **fixes** *m p* :: *nat*
      **and** *pairs* :: (*complex Matrix.mat* × *complex Matrix.mat*) *list*
  **assumes** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
      **and** *typing*: $\forall (\sigma, \tau) \in$ *set pairs*.
                $\sigma \in$ *carrier-mat m m* $\wedge\ \tau \in$ *carrier-mat p p*
      **and** *long*: *length pairs* > $2 * (m*p)^2$
  **shows** $\exists\, r$ :: *nat* ⇒ *real*.
          ($\exists\, k <$ *length pairs*. $r\ k \neq 0$) ∧
          ($\forall\, k \geq$ *length pairs*. $r\ k = 0$) ∧
          *cone-list-sum m p*
            (*map* ($\lambda k.$ (*r k*, *fst* (*pairs* ! *k*), *snd* (*pairs* ! *k*)))
                [*0*..<*length pairs*])

$= 0_m\ (m{*}p)\ (m{*}p)$
⟨*proof*⟩

One reduction step: a length-$L$ cone list with $L > (m\ p)^2 + 1$ admits a length-$(L-1)$ list with the same cone sum, weights non-negative, and matrix pairs inherited from the original list.

NOTE on the conclusion: the standard Carathéodory reduction *reweights* rather than *selects*, so we require only that the *matrix pair* $(\sigma, \tau)$ of each new entry occurs in the original list the weight may differ. This is what the downstream closedness argument actually needs (typing + non-negativity inherited from the original cone).

**lemma** *caratheodory-matrix-cone-step*:
**fixes** *m p* :: *nat*
**and** *ts* :: (*real* × *complex Matrix.mat* × *complex Matrix.mat*) *list*
**assumes** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
**and** *ts-typing*: $\forall (w, \sigma, \tau) \in set\ ts.$
$\sigma \in carrier\text{-}mat\ m\ m \wedge \tau \in carrier\text{-}mat\ p\ p$
**and** *ts-nn*: $\forall (w, \text{-}, \text{-}) \in set\ ts.\ 0 \le w$
**and** *ts-long*: $length\ ts > 2 * (m{*}p)^2 + 1$
**shows** $\exists ts'.\ length\ ts' = length\ ts - 1$
$\wedge\ (\forall (w, \sigma, \tau) \in set\ ts'.$
$\sigma \in carrier\text{-}mat\ m\ m \wedge \tau \in carrier\text{-}mat\ p\ p \wedge 0 \le w$
$\wedge\ (\exists w_0.\ (w_0, \sigma, \tau) \in set\ ts))$
$\wedge\ cone\text{-}list\text{-}sum\ m\ p\ ts = cone\text{-}list\text{-}sum\ m\ p\ ts'$
⟨*proof*⟩

The Carathéodory bound follows from the reduction step by strong induction on *length ts*. This part is now closed unconditionally; only the two focused sub-sorries inside *caratheodory-matrix-cone-step* remain.

**lemma** *caratheodory-matrix-cone*:
**fixes** *m p* :: *nat*
**and** *ts* :: (*real* × *complex Matrix.mat* × *complex Matrix.mat*) *list*
**assumes** *m-pos*: $0 < m$ **and** *p-pos*: $0 < p$
**and** *ts-typing*: $\forall (w, \sigma, \tau) \in set\ ts.$
$\sigma \in carrier\text{-}mat\ m\ m \wedge \tau \in carrier\text{-}mat\ p\ p$
**and** *ts-nn*: $\forall (w, \text{-}, \text{-}) \in set\ ts.\ 0 \le w$
**shows** $\exists ts'.\ length\ ts' \le 2 * (m{*}p)^2 + 1$
$\wedge\ (\forall (w, \sigma, \tau) \in set\ ts'.$
$\sigma \in carrier\text{-}mat\ m\ m \wedge \tau \in carrier\text{-}mat\ p\ p \wedge 0 \le w$
$\wedge\ (\exists w_0.\ (w_0, \sigma, \tau) \in set\ ts))$
$\wedge\ cone\text{-}list\text{-}sum\ m\ p\ ts = cone\text{-}list\text{-}sum\ m\ p\ ts'$
⟨*proof*⟩

How the bound is used (Separable_Cone): a sequence in the separable cone has decompositions of length at most $K = 2(mp)^2+1$; padding with zero-weight entries and normalising traces makes the weights bounded, so coordinatewise Bolzano–Weierstrass extracts a convergent subsequence of weights and factors; tensoring and addition are continuous, so the limit lies in the cone. The compactness of the unit-trace positive matrices is obtained from the Euclidean-space instance of Lmat_Euclidean (real dimension $2(mp)^2$) rather than from the sequential-compactness lemma *HS-Cauchy-imp-convergent* proved above.

**end**

**theory** *Lmat-Euclidean*
**imports**
*Vern-Paulsen-QC-3-10.Matrix-Typed*
*HOL−Analysis.Euclidean-Space*
*HOL−Analysis.L2-Norm*

*HOL−Analysis.Elementary-Metric-Spaces*
*HOL−Analysis.Topology-Euclidean-Space*
**begin**

The HOL/Hahn_Banach development imports a locale also called *norm* under the qualifier *Normed-Space*. Hide its short names so that, inside our instantiation blocks, *norm* resolves to the class method from *HOL.Real-Vector-Spaces*.

**hide-const** (**open**) *Normed-Space.norm*
**hide-const** (**open**) *Normed-Space.seminorm*

# 76 Overview

Nothing in this theory is stated in the notes; it is the bridge that lets the HOL-Analysis theorems on convex sets (separating hyperplanes, compactness) be applied to matrices. ('*a*, '*m*, '*n*) *lmat* is given the typeclass tower *real-vector*, *metric-space*, *real-normed-vector*, *real-inner*, *euclidean-space*, *heine-borel*; for '*a* = *complex* the norm is the Hilbert–Schmidt norm, the inner product its real part, and the dimension is $2\,\mathrm{CARD}('m)\,\mathrm{CARD}('n)$.

# 77 Real scalar multiplication and *real-vector* instance

For '*a* :: *real-vector*, lift element-wise scalar multiplication via *map-mat* (*scaleR a*).

**instantiation** *lmat* :: (*real-vector*, *finite*, *finite*) *scaleR*
**begin**

**lift-definition** *scaleR-lmat* ::
  *real* ⇒ ('*a*, '*m* :: *finite*, '*n* :: *finite*) *lmat* ⇒ ('*a*, '*m* :: *finite*, '*n* :: *finite*) *lmat*
  **is** λ*r M*. *map-mat* (λ*x*. *r* ∗$_R$ *x*) *M* ⟨*proof*⟩

**instance** ⟨*proof*⟩

**end**

**lemma** *unwrap-lm-scaleR* [*simp*]:
  *unwrap-lm* ((*r* :: *real*) ∗$_R$ (*M* :: ('*a* :: *real-vector*, '*m* :: *finite*, '*n* :: *finite*) *lmat*))
    = *map-mat* (λ*x*. *r* ∗$_R$ *x*) (*unwrap-lm M*)
  ⟨*proof*⟩

**lemma** *unwrap-lm-scaleR-index* [*simp*]:
  **fixes** *M* :: ('*a* :: *real-vector*, '*m* :: *finite*, '*n* :: *finite*) *lmat*
  **assumes** *i* < *CARD*('*m*) *j* < *CARD*('*n*)
  **shows** *unwrap-lm* (*r* ∗$_R$ *M*) $$ (*i*, *j*) = *r* ∗$_R$ (*unwrap-lm M* $$ (*i*, *j*))
  ⟨*proof*⟩

**instance** *lmat* :: (*real-vector*, *finite*, *finite*) *real-vector*
⟨*proof*⟩

# 78 Auxiliary: pointwise unwrapping (used by later instances)

**lemma** *unwrap-lm-plus-index* [*simp*]:
  **fixes** *M N* :: ('*a* :: *plus*, '*m* :: *finite*, '*n* :: *finite*) *lmat*
  **assumes** *i* < *CARD*('*m*) *j* < *CARD*('*n*)
  **shows** *unwrap-lm* (*M* + *N*) $$ (*i*, *j*) = *unwrap-lm M* $$ (*i*, *j*) + *unwrap-lm N* $$ (*i*, *j*)
  ⟨*proof*⟩

**lemma** *unwrap-lm-minus-index* [*simp*]:
**fixes** $M\ N :: ('a :: ab\text{-}group\text{-}add, 'm :: finite, 'n :: finite)\ lmat$
**assumes** $i < CARD('m)\ j < CARD('n)$
**shows** $unwrap\text{-}lm\ (M - N)\ \$\$\ (i, j) = unwrap\text{-}lm\ M\ \$\$\ (i, j) - unwrap\text{-}lm\ N\ \$\$\ (i, j)$
⟨*proof*⟩

**lemma** *unwrap-lm-zero-index* [*simp*]:
**fixes** $i\ j :: nat$
**assumes** $i < CARD('m :: finite)\ j < CARD('n :: finite)$
**shows** $unwrap\text{-}lm\ (0 :: ('a :: zero, 'm, 'n)\ lmat)\ \$\$\ (i, j) = 0$
⟨*proof*⟩

**lemma** *lmat-eq-iff-entries*:
**fixes** $M\ N :: ('a, 'm :: finite, 'n :: finite)\ lmat$
**shows** $M = N \longleftrightarrow$
$(\forall\, i\ j.\ i < CARD('m) \longrightarrow j < CARD('n)$
$\longrightarrow unwrap\text{-}lm\ M\ \$\$\ (i, j) = unwrap\text{-}lm\ N\ \$\$\ (i, j))$
⟨*proof*⟩

# 79 Distance, uniformity, open: *metric-space* instance

Distance is the L2 sum of entry-wise distances, indexed by $\{0..<CARD('m)\} \times \{0..<CARD('n)\}$. Uniformity and *open* are defined via the standard metric-induced filter so that the *open-uniformity* axiom is discharged by reflexivity.

**instantiation** *lmat* :: (*metric-space*, *finite*, *finite*) *dist*
**begin**

**definition** *dist-lmat* :: $('a, 'm :: finite, 'n :: finite)\ lmat$
$\Rightarrow ('a, 'm :: finite, 'n :: finite)\ lmat \Rightarrow real$ **where**
$dist\text{-}lmat\ M\ N = L2\text{-}set\ (\lambda(i, j).\ dist\ (unwrap\text{-}lm\ M\ \$\$\ (i, j))\ (unwrap\text{-}lm\ N\ \$\$\ (i, j)))$
$(\{0..<CARD('m)\} \times \{0..<CARD('n)\})$

**instance** ⟨*proof*⟩

**end**

**instantiation** *lmat* :: (*metric-space*, *finite*, *finite*) *uniformity-dist*
**begin**

**definition** *uniformity-lmat* ::
$(('a, 'm :: finite, 'n :: finite)\ lmat \times ('a, 'm, 'n)\ lmat)\ filter$ **where**
$(uniformity\text{-}lmat :: (('a, 'm, 'n)\ lmat \times ('a, 'm, 'n)\ lmat)\ filter)$
$= (INF\ e \in \{0<..\}.\ principal\ \{(x, y).\ dist\ x\ y < e\})$

**instance** ⟨*proof*⟩

**end**

**instantiation** *lmat* :: (*metric-space*, *finite*, *finite*) *metric-space*
**begin**

**definition** *open-lmat* :: $('a, 'm :: finite, 'n :: finite)\ lmat\ set \Rightarrow bool$ **where**
$open\text{-}lmat\ U = (\forall\, x \in U.\ \forall_F\ (x', y)\ in\ uniformity.\ x' = x \longrightarrow y \in U)$

**instance** ⟨*proof*⟩

**end**

## 80 Norm and *real-normed-vector* instance

**instantiation** *lmat* :: (*real-normed-vector*, *finite*, *finite*) *real-normed-vector*
**begin**

**definition** *norm-lmat* :: ('*a*, '*m* :: *finite*, '*n* :: *finite*) *lmat* ⇒ *real* **where**
*norm-lmat M* = *L2-set* (λ(*i*, *j*). *norm* (*unwrap-lm M* $$ (*i*, *j*)))
({*0*..<*CARD*('*m*)} × {*0*..<*CARD*('*n*)})

**definition** *sgn-lmat* ::
('*a*, '*m* :: *finite*, '*n* :: *finite*) *lmat* ⇒ ('*a*, '*m*, '*n*) *lmat* **where**
*sgn-lmat* (*M* :: ('*a*, '*m*, '*n*) *lmat*) = *scaleR* (*inverse* (*norm M*)) *M*

**instance** ⟨*proof*⟩

**end**

## 81 Real inner product

For '*a* :: *real-inner*, the lmat inner product is the entry-wise sum of inner products. When '*a* = *complex* with the standard real-inner instance *inner z w* = *Re* (*cnj z* ∗ *w*), this is the real part of the Hilbert-Schmidt inner product.

**instantiation** *lmat* :: (*real-inner*, *finite*, *finite*) *real-inner*
**begin**

**definition** *inner-lmat* :: ('*a*, '*m* :: *finite*, '*n* :: *finite*) *lmat*
⇒ ('*a*, '*m*, '*n*) *lmat* ⇒ *real* **where**
*inner-lmat M N* = (∑ (*i*, *j*) ∈ {*0*..<*CARD*('*m*)} × {*0*..<*CARD*('*n*)}.
*inner* (*unwrap-lm M* $$ (*i*, *j*)) (*unwrap-lm N* $$ (*i*, *j*)))

**instance** ⟨*proof*⟩

**end**

## 82 Basis: matrix units carrying a basis vector

**definition** *lmat-axis* :: *nat* ⇒ *nat* ⇒ '*a* :: *zero*
⇒ ('*a*, '*m* :: *finite*, '*n* :: *finite*) *lmat* **where**
*lmat-axis i j u* = *wrap-lm* (*Matrix.mat* (*CARD*('*m*)) (*CARD*('*n*))
(λ(*r*, *s*). *if r* = *i* ∧ *s* = *j then u else 0*))

**lemma** *unwrap-lmat-axis* [*simp*]:
*unwrap-lm* (*lmat-axis i j u* :: ('*a* :: *zero*, '*m* :: *finite*, '*n* :: *finite*) *lmat*)
= *Matrix.mat* (*CARD*('*m*)) (*CARD*('*n*))
(λ(*r*, *s*). *if r* = *i* ∧ *s* = *j then u else 0*)
⟨*proof*⟩

**lemma** *lmat-axis-index* [*simp*]:
**fixes** *u* :: '*a* :: *zero*
**assumes** *r* < *CARD*('*m* :: *finite*) *s* < *CARD*('*n* :: *finite*)
**shows** *unwrap-lm* (*lmat-axis i j u* :: ('*a*, '*m*, '*n*) *lmat*) $$ (*r*, *s*)
= (*if r* = *i* ∧ *s* = *j then u else 0*)
⟨*proof*⟩

Inner product against a single matrix-unit basis vector picks out one matrix entry.

**lemma** *inner-lmat-axis*:

**fixes** $u :: 'a :: real\text{-}inner$
**and** $M :: ('a, 'm :: finite, 'n :: finite)\ lmat$
**assumes** $ij$: $i < CARD('m)\ j < CARD('n)$
**shows** $inner\ M\ (lmat\text{-}axis\ i\ j\ u :: ('a, 'm, 'n)\ lmat)$
$= inner\ (unwrap\text{-}lm\ M\ \$\$\ (i, j))\ u$
⟨*proof*⟩

Two matrix-unit basis vectors are equal iff their entries and positions match (or both vectors are zero).

**lemma** *lmat-axis-eq-iff*:
**fixes** $u\ v :: 'a :: zero$
**assumes** $i < CARD('m :: finite)\ j < CARD('n :: finite)$
**and** $i' < CARD('m)\ j' < CARD('n)$
**shows** $(lmat\text{-}axis\ i\ j\ u :: ('a, 'm, 'n)\ lmat) = lmat\text{-}axis\ i'\ j'\ v$
$\longleftrightarrow (u = v \wedge i = i' \wedge j = j') \vee (u = 0 \wedge v = 0)$
⟨*proof*⟩

Inner product between two matrix-unit basis vectors: 1 if same position and basis component, 0 otherwise.

**lemma** *inner-lmat-axis-axis*:
**fixes** $u\ v :: 'a :: real\text{-}inner$
**assumes** $ij$: $i < CARD('m :: finite)\ j < CARD('n :: finite)$
**and** $ij'$: $i' < CARD('m)\ j' < CARD('n)$
**shows** $inner\ (lmat\text{-}axis\ i\ j\ u :: ('a, 'm, 'n)\ lmat)\ (lmat\text{-}axis\ i'\ j'\ v)$
$= (if\ i = i' \wedge j = j'\ then\ inner\ u\ v\ else\ 0)$
⟨*proof*⟩

# 83 Basis set

**definition** *Basis-lmat-set* :: $('a :: euclidean\text{-}space, 'm :: finite, 'n :: finite)\ lmat\ set$
**where** $Basis\text{-}lmat\text{-}set = (\bigcup i \in \{0..<CARD('m)\}.$
$\bigcup j \in \{0..<CARD('n)\}.$
$\bigcup u \in Basis.\ \{lmat\text{-}axis\ i\ j\ u :: ('a, 'm, 'n)\ lmat\})$

**lemma** *Basis-lmat-set-finite*:
$finite\ (Basis\text{-}lmat\text{-}set :: ('a :: euclidean\text{-}space, 'm :: finite, 'n :: finite)\ lmat\ set)$
⟨*proof*⟩

**lemma** *Basis-lmat-set-nonempty*:
$Basis\text{-}lmat\text{-}set \neq (\{\} :: ('a :: euclidean\text{-}space, 'm :: finite, 'n :: finite)\ lmat\ set)$
⟨*proof*⟩

**lemma** *Basis-lmat-orthonormal*:
**fixes** $u\ v :: ('a :: euclidean\text{-}space, 'm :: finite, 'n :: finite)\ lmat$
**assumes** $uB$: $u \in Basis\text{-}lmat\text{-}set$ **and** $vB$: $v \in Basis\text{-}lmat\text{-}set$
**shows** $inner\ u\ v = (if\ u = v\ then\ 1\ else\ 0)$
⟨*proof*⟩

**lemma** *axis-mem-Basis-lmat-set*:
**fixes** $u :: 'a :: euclidean\text{-}space$
**assumes** $i\text{-}lt$: $i < CARD('m :: finite)$
**and** $j\text{-}lt$: $j < CARD('n :: finite)$
**and** $uB$: $u \in Basis$
**shows** $(lmat\text{-}axis\ i\ j\ u :: ('a, 'm, 'n)\ lmat) \in Basis\text{-}lmat\text{-}set$
⟨*proof*⟩

**lemma** *Basis-lmat-perp-imp-zero*:

**fixes** $x :: ('a :: \textit{euclidean-space}, 'm :: \textit{finite}, 'n :: \textit{finite})\ \textit{lmat}$
**assumes** $*: \forall u \in \textit{Basis-lmat-set}.\ \textit{inner}\ x\ u = 0$
**shows** $x = 0$
⟨*proof*⟩

**lemma** *zero-inner-Basis-lmat*:
**fixes** $u :: ('a :: \textit{euclidean-space}, 'm :: \textit{finite}, 'n :: \textit{finite})\ \textit{lmat}$
**shows** $\textit{inner}\ (0 :: ('a, 'm, 'n)\ \textit{lmat})\ u = 0$
⟨*proof*⟩

**lemma** *Basis-lmat-perp-zero*:
**fixes** $x :: ('a :: \textit{euclidean-space}, 'm :: \textit{finite}, 'n :: \textit{finite})\ \textit{lmat}$
**shows** $(\forall u \in \textit{Basis-lmat-set}.\ \textit{inner}\ x\ u = 0) \longleftrightarrow x = 0$
⟨*proof*⟩

# 84 Euclidean space instance

**instantiation** $\textit{lmat} :: (\textit{euclidean-space}, \textit{finite}, \textit{finite})\ \textit{euclidean-space}$
**begin**

**definition** $\textit{Basis-lmat} :: ('a, 'm :: \textit{finite}, 'n :: \textit{finite})\ \textit{lmat set}$ **where**
$\textit{Basis-lmat} = \textit{Basis-lmat-set}$

**instance** ⟨*proof*⟩

**end**

# 85 Heine-Borel

**instance** $\textit{lmat} :: (\textit{euclidean-space}, \textit{finite}, \textit{finite})\ \textit{heine-borel}$ ⟨*proof*⟩

# 86 Dimension

Two distinct $(i, j, u)$ with $u \in \textit{Basis-}'a$ give distinct matrix-axis vectors.

**lemma** *lmat-axis-inj-on-Basis-indices*:
**fixes** $u\ v :: 'a :: \textit{euclidean-space}$
**assumes** *ij*: $i < \textit{CARD}('m :: \textit{finite})\ j < \textit{CARD}('n :: \textit{finite})$
**and** *ij'*: $i' < \textit{CARD}('m)\ j' < \textit{CARD}('n)$
**and** *uB*: $u \in \textit{Basis}$ **and** *vB*: $v \in \textit{Basis}$
**shows** $(\textit{lmat-axis}\ i\ j\ u :: ('a, 'm, 'n)\ \textit{lmat}) = \textit{lmat-axis}\ i'\ j'\ v$
$\longleftrightarrow i = i' \wedge j = j' \wedge u = v$
⟨*proof*⟩

**lemma** *DIM-lmat*:
$\textit{DIM}(('a :: \textit{euclidean-space}, 'm :: \textit{finite}, 'n :: \textit{finite})\ \textit{lmat})$
$= \textit{CARD}('m) * \textit{CARD}('n) * \textit{DIM}('a)$
⟨*proof*⟩

# 87 Component-wise bounds via the L2 norm

**lemma** *norm-lmat-index-le*:
**fixes** $M :: ('a :: \textit{real-normed-vector}, 'm :: \textit{finite}, 'n :: \textit{finite})\ \textit{lmat}$
**assumes** $i < \textit{CARD}('m)\ j < \textit{CARD}('n)$
**shows** $\textit{norm}\ (\textit{unwrap-lm}\ M\ \$\$\ (i, j)) \leq \textit{norm}\ M$
⟨*proof*⟩

**lemma** *dist-lmat-index-le*:
**fixes** $M\ N :: ('a :: real\text{-}normed\text{-}vector, 'm :: finite, 'n :: finite)\ lmat$
**assumes** $i < CARD('m)\ j < CARD('n)$
**shows** $dist\ (unwrap\text{-}lm\ M\ \$\$\ (i, j))\ (unwrap\text{-}lm\ N\ \$\$\ (i, j)) \leq dist\ M\ N$
⟨*proof*⟩

## 88 Convergence: lmat sequence convergence implies entry convergence

**lemma** *tendsto-unwrap-lm-index*:
**fixes** $f :: 'b \Rightarrow ('a :: real\text{-}normed\text{-}vector, 'm :: finite, 'n :: finite)\ lmat$
**and** $L :: ('a, 'm, 'n)\ lmat$
**assumes** *lim*: $(f \longrightarrow L)\ F$
**and** *ij*: $i < CARD('m)\ j < CARD('n)$
**shows** $((\lambda x.\ unwrap\text{-}lm\ (f\ x)\ \$\$\ (i, j)) \longrightarrow unwrap\text{-}lm\ L\ \$\$\ (i, j))\ F$
⟨*proof*⟩

**lemma** *LIMSEQ-unwrap-lm-index*:
**fixes** $f :: nat \Rightarrow ('a :: real\text{-}normed\text{-}vector, 'm :: finite, 'n :: finite)\ lmat$
**and** $L :: ('a, 'm, 'n)\ lmat$
**assumes** $f \longrightarrow L\ i < CARD('m)\ j < CARD('n)$
**shows** $(\lambda n.\ unwrap\text{-}lm\ (f\ n)\ \$\$\ (i, j)) \longrightarrow unwrap\text{-}lm\ L\ \$\$\ (i, j)$
⟨*proof*⟩

**end**

# Part V
# Lectures 26–28: separability and entanglement witnesses

**theory** *Separable-Cone*
**imports**
*Lmat-Euclidean*
*Vern-Paulsen-QC-3-10.Quantum-Computing-3-10*
*HOL−Analysis.Convex*
*HOL−Analysis.Convex-Euclidean-Space*
*HOL−Analysis.Polytope*
*HOL−Analysis.Starlike*
**begin**

HOL-Hahn_Banach defines a locale also called *continuous*. Hide its short name so that, in the proofs below, *continuous* resolves to the topological-space predicate from *HOL.Topological-Spaces*.

**hide-const** (**open**) *Function-Norm.continuous*

## 89 Overview (Days 27–28 of the notes)

This theory proves the results restated lecture by lecture in theory Day24_Separability. The notes argue in the real vector space of Hermitian matrices with a closed-cone separation theorem;

formally we work in the typed matrix space (*complex*, $'m \times 'p$, $'m \times 'p$) *lmat*, a real inner-product space (Lmat_Euclidean), and use HOL-Analysis' *separating-hyperplane-closed-point*. Steps: the positive matrices form a closed set; the unit-trace positive matrices form a compact set; the tensor product is continuous, so the product density matrices *Sprime* form a compact set not containing 0; by Carathéodory (Essential_Functional_Analysis) every element of the separable cone is a bounded combination of at most $2(mp)^2 + 1$ generators, and the cone is closed. Separation then yields a real functional, which is converted into a Hermitian witness $W$ with $\Re \operatorname{Tr}(W\rho) < 0 \leq \Re \operatorname{Tr}(W\sigma)$ on the cone.

# 90 Typed PSD predicate

**definition** *psd-lmat* :: (*complex*, $'k$ :: *finite*, $'k$) *lmat* ⇒ *bool* **where**
*psd-lmat M* ⟷ *psd* (*unwrap-lm M*)

**lemma** *psd-lmat-unwrap* [*iff*]:
*psd-lmat* (*M* :: (*complex*, $'k$ :: *finite*, $'k$) *lmat*) ⟷ *psd* (*unwrap-lm M*)
⟨*proof*⟩

# 91 Typed tensor product

Two square matrices with type-level dimensions $'m$, $'p$ tensor to a square matrix with type-level dimension $'m \times 'p$.

**lemma** *tensor-mat-carrier-lmat*:
**fixes** *A B* :: *complex Matrix.mat*
**assumes** *A*: *A* ∈ *carrier-mat* (*CARD*($'m$ :: *finite*)) (*CARD*($'m$))
**and** *B*: *B* ∈ *carrier-mat* (*CARD*($'p$ :: *finite*)) (*CARD*($'p$))
**shows** *A* ⨂ *B* ∈ *carrier-mat* (*CARD*($'m \times 'p$)) (*CARD*($'m \times 'p$))
⟨*proof*⟩

**definition** *tensor-lmat* ::
(*complex*, $'m$ :: *finite*, $'m$) *lmat* ⇒ (*complex*, $'p$ :: *finite*, $'p$) *lmat*
⇒ (*complex*, $'m \times 'p$, $'m \times 'p$) *lmat* **where**
*tensor-lmat* σ τ = *wrap-lm* (*unwrap-lm* σ ⨂ *unwrap-lm* τ)

**lemma** *unwrap-lm-tensor-lmat* [*simp*]:
**fixes** σ :: (*complex*, $'m$ :: *finite*, $'m$) *lmat*
**and** τ :: (*complex*, $'p$ :: *finite*, $'p$) *lmat*
**shows** *unwrap-lm* (*tensor-lmat* σ τ :: (*complex*, $'m \times 'p$, $'m \times 'p$) *lmat*)
= *unwrap-lm* σ ⨂ *unwrap-lm* τ
⟨*proof*⟩

# 92 The separable cone

The set of *separable matrices*: all nonneg combinations of tensor products of typed PSD matrices. Equivalently, the convex conic hull of the product PSD set.

**definition** *separable-lmat-gen* :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*,
$'m \times 'p$) *lmat set* **where**
*separable-lmat-gen* =
{*tensor-lmat* σ τ | σ τ.
(*psd-lmat* σ :: *bool*) ∧ (*psd-lmat* τ :: *bool*)}

**definition** *separable-cone-lmat* :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*,
$'m \times 'p$) *lmat set* **where**

*separable-cone-lmat = convex-cone hull separable-lmat-gen*

Sanity: the cone contains 0, is convex, and is conic.

**lemma** *separable-cone-lmat-convex-cone*:
*convex-cone* (*separable-cone-lmat*
:: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat set*)
⟨*proof*⟩

**lemma** *zero-in-separable-cone-lmat*:
(*0* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*)
∈ *separable-cone-lmat*
⟨*proof*⟩

# 93 Step 1: PSD set is closed

Sub-lemmas for the closedness of PSD: the Hermitian half and the nonneg-quadratic-form half.

The entry-projection $M \mapsto$ *unwrap-lm M $$ (i, j)* is continuous on lmat for valid index $(i, j)$.

**lemma** *isCont-unwrap-lm-index*:
**fixes** *x* :: (*'a* :: *real-normed-vector*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
**assumes** *ij*: *i* < *CARD*(*'m*) *j* < *CARD*(*'n*)
**shows** *isCont* (λ*M*. *unwrap-lm M $$ (i, j)*) *x*
⟨*proof*⟩

**lemma** *continuous-on-unwrap-lm-index*:
**fixes** *S* :: (*'a* :: *real-normed-vector*, *'m* :: *finite*, *'n* :: *finite*) *lmat set*
**assumes** *ij*: *i* < *CARD*(*'m*) *j* < *CARD*(*'n*)
**shows** *continuous-on S* (λ*M*. *unwrap-lm M $$ (i, j)*)
⟨*proof*⟩

**lemma** *closed-entry-hermitian-eq*:
**fixes** *i j* :: *nat*
**assumes** *ij*: *i* < *CARD*(*'k* :: *finite*) *j* < *CARD*(*'k*)
**shows** *closed* {*M* :: (*complex*, *'k*, *'k*) *lmat*.
*unwrap-lm M $$ (i, j)* = *cnj* (*unwrap-lm M $$ (j, i)*)}
⟨*proof*⟩

**lemma** *closed-hermitian-unwrap-lmat*:
*closed* {*M* :: (*complex*, *'k* :: *finite*, *'k*) *lmat*. *hermitian* (*unwrap-lm M*)}
⟨*proof*⟩

**lemma** *closed-nonneg-quad-form-lmat*:
**fixes** *x* :: *complex Matrix.vec*
**assumes** *xdim*: *dim-vec x* = *CARD*(*'k* :: *finite*)
**shows** *closed* {*M* :: (*complex*, *'k*, *'k*) *lmat*.
*Re* (⟨*x* | *unwrap-lm M* ∗$_v$ *x*⟩) ≥ *0*}
⟨*proof*⟩

**lemma** *closed-psd-lmat*:
*closed* {*M* :: (*complex*, *'k* :: *finite*, *'k*) *lmat*. *psd-lmat M*}
⟨*proof*⟩

# 94 Trace on lmat

**definition** *Tr-lmat* :: (*complex*, *'k* :: *finite*, *'k*) *lmat* ⇒ *complex* **where**
*Tr-lmat M* = (∑ *i* = *0*..<*CARD*(*'k*). *unwrap-lm M $$ (i, i)*)

**lemma** *continuous-on-Tr-lmat*:
**fixes** $S :: (complex, 'k :: finite, 'k)\ lmat\ set$
**shows** *continuous-on S Tr-lmat*
⟨*proof*⟩

**lemma** *Tr-lmat-add*:
**fixes** $M\ N :: (complex, 'k :: finite, 'k)\ lmat$
**shows** $Tr\text{-}lmat\ (M + N) = Tr\text{-}lmat\ M + Tr\text{-}lmat\ N$
⟨*proof*⟩

**lemma** *Tr-lmat-scaleR*:
**fixes** $M :: (complex, 'k :: finite, 'k)\ lmat$ **and** $a :: real$
**shows** $Tr\text{-}lmat\ (a *_R M) = a *_R Tr\text{-}lmat\ M$
⟨*proof*⟩

**lemma** *Tr-lmat-zero* [*simp*]:
$Tr\text{-}lmat\ (0 :: (complex, 'k :: finite, 'k)\ lmat) = 0$
⟨*proof*⟩

**lemma** *Tr-lmat-sum-scaleR*:
**fixes** $T :: 'a\ set$ **and** $u :: 'a \Rightarrow real$
**and** $f :: 'a \Rightarrow (complex, 'k :: finite, 'k)\ lmat$
**assumes** *finite T*
**shows** $Tr\text{-}lmat\ (\sum t \in T.\ u\ t *_R f\ t) = (\sum t \in T.\ u\ t *_R Tr\text{-}lmat\ (f\ t))$
⟨*proof*⟩

**lemma** *closed-unit-trace-lmat*:
$closed\ \{M :: (complex, 'k :: finite, 'k)\ lmat.\ Tr\text{-}lmat\ M = 1\}$
⟨*proof*⟩

# 95 Compactness of the unit-trace PSD slice

For PSD $\sigma$ with $Tr\ \sigma = 1$: ${\sigma^2}_{HS} = \sum {\lambda_i}^2 \leq (\sum \lambda_i)^2 = 1$ where $\lambda_i$ are the (nonneg) eigenvalues.

**lemma** *psd-entry-bound-by-diag*:
**fixes** $M :: complex\ Matrix.mat$
**assumes** *Mcar*: $M \in carrier\text{-}mat\ n\ n$
**and** *Mpsd*: *psd M*
**and** *i*: $i < n$ **and** *j*: $j < n$
**shows** $(norm\ (M\ \$\$\ (i, j)))^2 \leq Re\ (M\ \$\$\ (i, i)) * Re\ (M\ \$\$\ (j, j))$
⟨*proof*⟩

**lemma** *norm-lmat-le-of-psd-unit-trace*:
**fixes** $M :: (complex, 'k :: finite, 'k)\ lmat$
**assumes** *psd*: *psd-lmat M* **and** *tr*: $Tr\text{-}lmat\ M = 1$
**shows** $norm\ M \leq 1$
⟨*proof*⟩

**lemma** *compact-unit-trace-psd-lmat*:
$compact\ \{M :: (complex, 'k :: finite, 'k)\ lmat.\ psd\text{-}lmat\ M \wedge Tr\text{-}lmat\ M = 1\}$
⟨*proof*⟩

# 96 Tensor continuity

The typed tensor product is jointly continuous in both factors.

**lemma** *continuous-on-tensor-lmat*:

**fixes** $T :: ((complex, 'm :: finite, 'm)\ lmat \times (complex, 'p :: finite, 'p)\ lmat)\ set$
**shows** *continuous-on* $T$ $(\lambda(\sigma, \tau).\ tensor\text{-}lmat\ \sigma\ \tau :: (complex, 'm \times 'p, 'm \times 'p)\ lmat)$
⟨*proof*⟩

# 97 The unit-trace product slice

**definition** $Sprime :: (complex, 'm :: finite \times 'p :: finite, 'm \times 'p)\ lmat\ set$
**where**
$Sprime =$
$\{\varrho.\ \exists(\sigma :: (complex, 'm, 'm)\ lmat)\ (\tau :: (complex, 'p, 'p)\ lmat).$
$\varrho = tensor\text{-}lmat\ \sigma\ \tau$
$\wedge\ psd\text{-}lmat\ \sigma \wedge Tr\text{-}lmat\ \sigma = 1$
$\wedge\ psd\text{-}lmat\ \tau \wedge Tr\text{-}lmat\ \tau = 1\}$

**lemma** *Sprime-image*:
$(Sprime :: (complex, 'm :: finite \times 'p :: finite, 'm \times 'p)\ lmat\ set)$
$= (\lambda(\sigma, \tau).\ tensor\text{-}lmat\ \sigma\ \tau)\ `$
$(\{\sigma :: (complex, 'm, 'm)\ lmat.\ psd\text{-}lmat\ \sigma \wedge Tr\text{-}lmat\ \sigma = 1\}$
$\times \{\tau :: (complex, 'p, 'p)\ lmat.\ psd\text{-}lmat\ \tau \wedge Tr\text{-}lmat\ \tau = 1\})$
⟨*proof*⟩

**lemma** *compact-Sprime*:
$compact\ (Sprime :: (complex, 'm :: finite \times 'p :: finite, 'm \times 'p)\ lmat\ set)$
⟨*proof*⟩

Reindex $\{0..<n*q\}$ via (div, mod) into $\{0..<n\} \times \{0..<q\}$.

**lemma** *sum-div-mod-split*:
**fixes** $f :: nat \Rightarrow nat \Rightarrow 'a :: comm\text{-}monoid\text{-}add$
**assumes** *q-pos*: $0 < q$
**shows** $(\sum a = 0..<n.\ \sum b = 0..<q.\ f\ a\ b)$
$= (\sum i = 0..<n * q.\ f\ (i\ div\ q)\ (i\ mod\ q))$
⟨*proof*⟩

**lemma** *Tr-lmat-tensor*:
**fixes** $\sigma :: (complex, 'm :: finite, 'm)\ lmat$
**and** $\tau :: (complex, 'p :: finite, 'p)\ lmat$
**shows** $Tr\text{-}lmat\ (tensor\text{-}lmat\ \sigma\ \tau :: (complex, 'm \times 'p, 'm \times 'p)\ lmat)$
$= Tr\text{-}lmat\ \sigma * Tr\text{-}lmat\ \tau$
⟨*proof*⟩

**lemma** *zero-not-in-Sprime*:
$(0 :: (complex, 'm :: finite \times 'p :: finite, 'm \times 'p)\ lmat) \notin Sprime$
⟨*proof*⟩

# 98 Closedness of the separable cone

Proof strategy (uses the typeclass tower from *Lmat-Euclidean*):
Let $S' = \{tensor\text{-}lmat\ \sigma\ \tau \mid \sigma\ \tau.\ psd\text{-}lmat\ \sigma \wedge psd\text{-}lmat\ \tau \wedge Tr\ (unwrap\text{-}lm\ \sigma) = 1 \wedge Tr\ (unwrap\text{-}lm\ \tau) = 1\}$ be the unit-trace product slice.

1. **S' is compact.**
   - $\{\sigma :: lmat.\ psd\text{-}lmat\ \sigma\}$ is closed (entry-wise lift of the Hermitian condition + entry-wise lift of the quadratic-form nonnegativity, both via *tendsto-unwrap-lm-index*).

- $\{\sigma :: lmat.\ Re\ (Tr\ (unwrap\text{-}lm\ \sigma)) = 1\}$ is closed (trace is continuous in entries, via *tendsto-unwrap-lm-index*).
- PSD + unit-trace is bounded: $norm\ \sigma^2 = Re\ Tr\ (\sigma^\dagger\ \sigma) \leq Tr\ \sigma \cdot \lambda_{max} \leq 1$.
- Closed + bounded in finite-dimensional *real-inner* = compact (*bounded-closed-imp-seq-compact* + heine_borel).
- *tensor-lmat* is continuous (entries are bilinear in the entries of the factors, lifts via the typeclass).
- $S'$ is the image of (compact) × (compact) under *tensor-lmat*, hence compact.

2. **convex hull S' is compact** via *compact-convex-hull* (applies because every euclidean_space is a real_inner finite-dimensional space).

3. **$0 \notin$ convex hull S'.** Every element of *convex hull* $S'$ has trace 1 (linear combinations of trace-1 matrices with nonneg coefficients summing to 1), while 0 has trace 0.

4. **separable_lmat_gen = conic hull S'.** Each *tensor-lmat* $\sigma\ \tau$ with $\sigma$, $\tau$ PSD equals $(Tr\ \sigma \cdot Tr\ \tau) \cdot tensor\text{-}lmat\ (\sigma/Tr\ \sigma)\ (\tau/Tr\ \tau)$ when traces are positive, and equals $0$ when one trace is 0 (in which case the corresponding factor is itself 0 because PSD + Tr $= 0 \Rightarrow 0$).

5. **Putting it together.** *separable-cone-lmat* = *convex-cone hull separable-lmat-gen* = *convex-cone hull* (*conic hull* $S'$) = *convex-cone hull* $S'$ = *conic hull* (*convex hull* $S'$) (the last by *convex-cone-hull-separate-nonempty*). Apply *closed-conic-hull* with the (compact, $0 \notin \ldots$) clause.

The infrastructure for all of these steps is now available (*norm*, *inner*, *dist*, *compact*, *convex*, *convex-cone*, etc. all defined for typed lmat via the euclidean_space + heine_borel instances).

The convex cone hulls of *separable-lmat-gen* and *Sprime* coincide: every *tensor-lmat* $\sigma\ \tau$ with $\sigma$, $\tau$ PSD can be rescaled into *Sprime* by dividing by traces (when nonzero) or replaced by 0 when a trace is zero.

The hard direction depends on three facts that each lift a spectral / bilinearity property:
psd_lmat_scaleR PSD preserved under nonneg real scaling psd_lmat_zero_trace_zero PSD with Tr = 0 is the zero matrix tensor_lmat_scaleR_left,right tensor is bilinear over real scaling psd_lmat_Tr_real_nonneg Tr of PSD is a real nonneg complex.
These reduce *tensor* $\sigma\ \tau$ with $\sigma$, $\tau$ PSD to $(Tr\ \sigma * Tr\ \tau) *_R tensor\ (\sigma/Tr\ \sigma)\ (\tau/Tr\ \tau)$ when traces are positive, or to 0 when one is zero both forms in *convex-cone hull Sprime*.

**lemma** *Sprime-subset-separable-lmat-gen*:
(*Sprime* :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat set*)
⊆ *separable-lmat-gen*
⟨*proof*⟩

The four spectral / bilinearity helpers needed for the hard direction. Each is a one-line lift of a JNF fact, but the JNF facts (spectral closure and tensor scaleR) live outside the parent heap so they are stated as targeted sorries here for the moment.

**lemma** *psd-lmat-Tr-real-nonneg*:
**fixes** $M$ :: (*complex*, $'k$ :: *finite*, $'k$) *lmat*
**assumes** *M-psd*: *psd-lmat M*
**shows** $Im\ (Tr\text{-}lmat\ M) = 0 \wedge 0 \leq Re\ (Tr\text{-}lmat\ M)$
⟨*proof*⟩

**lemma** *psd-lmat-zero-trace-zero*:

**fixes** $M :: (complex, 'k :: finite, 'k)\ lmat$
**assumes** *M-psd*: *psd-lmat M* **and** *tr-zero*: *Tr-lmat M = 0*
**shows** *M = 0*
⟨*proof*⟩

**lemma** *psd-smult-nonneg-local*:
**fixes** *A* :: *complex Matrix.mat* **and** *c* :: *real*
**assumes** *c*: *0 ≤ c*
**and** *A*: *psd A* **and** *Ac*: *A ∈ carrier-mat n n*
**shows** *psd* (*complex-of-real c ·$_m$ A*)
⟨*proof*⟩

**lemma** *unwrap-lm-scaleR-eq*:
**fixes** $M :: (complex, 'm :: finite, 'n :: finite)\ lmat$ **and** *a* :: *real*
**shows** *unwrap-lm* (*a *$_R$ M*) = *complex-of-real a ·$_m$ unwrap-lm M*
⟨*proof*⟩

**lemma** *psd-lmat-scaleR-nonneg*:
**fixes** $M :: (complex, 'k :: finite, 'k)\ lmat$ **and** *a* :: *real*
**assumes** *M-psd*: *psd-lmat M* **and** *a-nn*: *0 ≤ a*
**shows** *psd-lmat* (*a *$_R$ M*)
⟨*proof*⟩

**lemma** *tensor-lmat-scaleR-left*:
**fixes** *a* :: *real*
**and** $\sigma :: (complex, 'm :: finite, 'm)\ lmat$
**and** $\tau :: (complex, 'p :: finite, 'p)\ lmat$
**shows** (*tensor-lmat* (*a *$_R$ σ*) *τ*
$:: (complex, 'm \times 'p, 'm \times 'p)\ lmat$)
= *a *$_R$ tensor-lmat σ τ*
⟨*proof*⟩

**lemma** *tensor-lmat-scaleR-right*:
**fixes** *a* :: *real*
**and** $\sigma :: (complex, 'm :: finite, 'm)\ lmat$
**and** $\tau :: (complex, 'p :: finite, 'p)\ lmat$
**shows** (*tensor-lmat σ* (*a *$_R$ τ*)
$:: (complex, 'm \times 'p, 'm \times 'p)\ lmat$)
= *a *$_R$ tensor-lmat σ τ*
⟨*proof*⟩

**lemma** *tensor-lmat-zero-left*:
**fixes** $\tau :: (complex, 'p :: finite, 'p)\ lmat$
**shows** (*tensor-lmat* (*0* :: $(complex, 'm :: finite, 'm)\ lmat$) *τ*
$:: (complex, 'm \times 'p, 'm \times 'p)\ lmat$) = *0*
⟨*proof*⟩

**lemma** *tensor-lmat-zero-right*:
**fixes** $\sigma :: (complex, 'm :: finite, 'm)\ lmat$
**shows** (*tensor-lmat σ* (*0* :: $(complex, 'p :: finite, 'p)\ lmat$)
$:: (complex, 'm \times 'p, 'm \times 'p)\ lmat$) = *0*
⟨*proof*⟩

**lemma** *convex-cone-hull-separable-eq-Sprime*:
(*convex-cone hull*
(*separable-lmat-gen*
$:: (complex, 'm :: finite \times 'p :: finite, 'm \times 'p)\ lmat\ set$))
= *convex-cone hull Sprime*

⟨*proof*⟩

**theorem** *separable-cone-lmat-closed*:
  *closed* (*separable-cone-lmat*
    :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat set*)
⟨*proof*⟩

## 99 The lmat inner product as a real-trace inner product

The real inner product on (*complex*, *'m*, *'n*) *lmat* is the real part of the JNF Hilbert-Schmidt inner product on the underlying matrices. For square matrices it is also *Re* (*Tr* ($M^\dagger$ ∗ *N*)).

**lemma** *inner-lmat-eq-re-hs-inner-prod*:
  **fixes** *M N* :: (*complex*, *'m* :: *finite*, *'n* :: *finite*) *lmat*
  **shows** *inner M N* = *Re* ⟨⟨ *unwrap-lm M* | *unwrap-lm N* ⟩⟩
⟨*proof*⟩

**lemma** *inner-lmat-eq-re-tr-dagger*:
  **fixes** *M N* :: (*complex*, *'k* :: *finite*, *'k*) *lmat*
  **shows** *inner M N* = *Re* (*Tr* ((*unwrap-lm M*)$^\dagger$ ∗ *unwrap-lm N*))
⟨*proof*⟩

## 100 Separating hyperplane for non-separable states

The closedness of the separable cone, combined with HOL-Analysis *separating-hyperplane-closed-point*, yields the geometric Hahn-Banach witness: for every $\varrho$ outside the separable cone there is a linear functional (an HS inner product against some matrix *a*) that is strictly negative on $\varrho$ and nonneg on every separable element. Because the separable set is a CONE, the strict inequality *inner a x* > *b* on the cone forces *b* < *0* and *inner a x* ≥ *0* on the cone.

**theorem** *separable-cone-lmat-separating-hyperplane*:
  **fixes** $\varrho$ :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** *notin*: $\varrho \notin$ *separable-cone-lmat*
  **shows** ∃ *a* :: (*complex*, *'m* × *'p*, *'m* × *'p*) *lmat*.
      *inner a* $\varrho$ < *0*
    ∧ (∀ $\sigma \in$ *separable-cone-lmat*. *0* ≤ *inner a* $\sigma$)
⟨*proof*⟩

## 101 Entanglement witness in real-trace form

Reformulating the separating hyperplane via the *inner* = *Re Tr* identity gives a JNF matrix *W* whose adjoint times $\varrho$ has strictly negative real trace, while $W^\dagger$ ∗ $\sigma$ has nonneg real trace for every $\sigma$ in the separable cone.

**theorem** *entanglement-witness-re-tr*:
  **fixes** $\varrho$ :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** *notin*: $\varrho \notin$ *separable-cone-lmat*
  **shows** ∃ *W* :: *complex Matrix.mat*.
      *W* ∈ *carrier-mat* (*CARD*(*'m* × *'p*)) (*CARD*(*'m* × *'p*))
    ∧ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)) < *0*
    ∧ (∀ $\sigma$ :: (*complex*, *'m* × *'p*, *'m* × *'p*) *lmat*.
        $\sigma \in$ *separable-cone-lmat*
        ⟶ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
⟨*proof*⟩

## 102 Hermitian entanglement witness

For $\varrho$ outside the separable cone with hermitian unwrap, the witness can be chosen hermitian. Direct entrywise construction of $W = (A + A^\dagger)/2$ using *Matrix.mat*; we avoid lemmas about *dagger* distributing over sums or scalar multiplication.

**theorem** *entanglement-witness-hermitian*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** *notin*: $\varrho \notin$ *separable-cone-lmat*
**and** *ϱ-herm*: *hermitian* (*unwrap-lm* $\varrho$)
**shows** ∃ $W$ :: *complex Matrix.mat*.
$W \in$ *carrier-mat* (*CARD*($'m$ × $'p$)) (*CARD*($'m$ × $'p$))
∧ *hermitian* $W$
∧ *Re* (*Tr* ($W$ ∗ *unwrap-lm* $\varrho$)) < 0
∧ (∀ $\sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *separable-cone-lmat*
∧ *hermitian* (*unwrap-lm* $\sigma$)
⟶ 0 ≤ *Re* (*Tr* ($W$ ∗ *unwrap-lm* $\sigma$)))
⟨*proof*⟩

## 103 Hermitian closure of the separable cone

Every element of *separable-cone-lmat* has hermitian unwrap. Generators *tensor-lmat* $\sigma$ $\tau$ with $\sigma$, $\tau$ PSD are hermitian, and the hermitian set is closed under + and nonneg real scaling.

**lemma** *hermitian-zero-mat*:
*hermitian* ($0_m$ $n$ $n$ :: *complex Matrix.mat*)
⟨*proof*⟩

**lemma** *hermitian-add-mat*:
**fixes** $P$ $Q$ :: *complex Matrix.mat*
**assumes** *Pcar*: $P \in$ *carrier-mat* $n$ $n$ **and** *Qcar*: $Q \in$ *carrier-mat* $n$ $n$
**and** *Pherm*: *hermitian* $P$ **and** *Qherm*: *hermitian* $Q$
**shows** *hermitian* ($P$ + $Q$)
⟨*proof*⟩

**lemma** *hermitian-smult-real-mat*:
**fixes** $M$ :: *complex Matrix.mat* **and** $c$ :: *real*
**assumes** *Mcar*: $M \in$ *carrier-mat* $n$ $n$ **and** *Mherm*: *hermitian* $M$
**shows** *hermitian* (*complex-of-real* $c$ ·$_m$ $M$)
⟨*proof*⟩

Tensor product of hermitians is hermitian (entrywise via *index-tensor-mat*).

**lemma** *hermitian-tensor-mat*:
**fixes** $P$ $Q$ :: *complex Matrix.mat*
**assumes** *Pcar*: $P \in$ *carrier-mat* $m$ $m$ **and** *Qcar*: $Q \in$ *carrier-mat* $p$ $p$
**and** *m-pos*: 0 < $m$ **and** *p-pos*: 0 < $p$
**and** *Pherm*: *hermitian* $P$ **and** *Qherm*: *hermitian* $Q$
**shows** *hermitian* ($P$ ⨂ $Q$)
⟨*proof*⟩

Generators of *separable-cone-lmat* have hermitian unwrap.

**lemma** *separable-lmat-gen-hermitian*:
**fixes** $\sigma$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** $\sigma \in$ *separable-lmat-gen*
**shows** *hermitian* (*unwrap-lm* $\sigma$)
⟨*proof*⟩

The hermitian set in lmat is a convex cone, hence contains the hull of the generators.

**lemma** *hermitian-lmat-set-convex-cone*:
  *convex-cone* {*M* :: (*complex*, *'k* :: *finite*, *'k*) *lmat*.
    *hermitian* (*unwrap-lm M*)}
⟨*proof*⟩

**theorem** *separable-cone-lmat-hermitian*:
  **fixes** $\sigma$ :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** $\sigma \in$ *separable-cone-lmat*
  **shows** *hermitian* (*unwrap-lm* $\sigma$)
⟨*proof*⟩

Final form of the entanglement witness theorem: the side condition "$\sigma$ has hermitian unwrap" is automatic for elements of the separable cone.

**theorem** *entanglement-witness-hermitian'*:
  **fixes** $\varrho$ :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** *notin*: $\varrho \notin$ *separable-cone-lmat*
    **and** $\varrho$*-herm*: *hermitian* (*unwrap-lm* $\varrho$)
  **shows** ∃ *W* :: *complex Matrix.mat*.
      *W* ∈ *carrier-mat* (*CARD*(*'m* × *'p*)) (*CARD*(*'m* × *'p*))
    ∧ *hermitian W*
    ∧ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)) < *0*
    ∧ (∀ $\sigma$ :: (*complex*, *'m* × *'p*, *'m* × *'p*) *lmat*.
        $\sigma \in$ *separable-cone-lmat*
        ⟶ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
⟨*proof*⟩

## 104 Main separability characterisation

The Horodecki-style witness characterisation: for hermitian $\varrho$, membership in the separable cone is equivalent to the nonneg-real-trace property holding against every hermitian witness that is itself nonneg-real-trace on the cone.

**theorem** *separable-iff-witness*:
  **fixes** $\varrho$ :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** $\varrho$*-herm*: *hermitian* (*unwrap-lm* $\varrho$)
  **shows** $\varrho \in$ *separable-cone-lmat*
    ⟷
    (∀ *W* :: *complex Matrix.mat*.
      *W* ∈ *carrier-mat* (*CARD*(*'m* × *'p*)) (*CARD*(*'m* × *'p*))
      ∧ *hermitian W*
      ∧ (∀ $\sigma$ :: (*complex*, *'m* × *'p*, *'m* × *'p*) *lmat*.
          $\sigma \in$ *separable-cone-lmat*
          ⟶ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
      ⟶ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)))
⟨*proof*⟩

## 105 Density-matrix specialisation

For an actual quantum density matrix $\varrho$ (PSD with unit trace), the conditions *psd-lmat* $\varrho$ and *Tr-lmat* $\varrho$ = *1* imply *hermitian* (*unwrap-lm* $\varrho$), so the witness theorem and the characterisation specialise cleanly.

**theorem** *entanglement-witness-for-density*:
  **fixes** $\varrho$ :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** $\varrho$*-psd*: *psd-lmat* $\varrho$

**and** *ϱ-tr*: *Tr-lmat ϱ = 1*
**and** *ϱ-not-sep*: *ϱ ∉ separable-cone-lmat*
**shows** ∃ *W* :: *complex Matrix.mat*.
*W ∈ carrier-mat (CARD('m × 'p)) (CARD('m × 'p))*
∧ *hermitian W*
∧ *Re (Tr (W ∗ unwrap-lm ϱ)) < 0*
∧ (∀ *σ* :: (*complex*, *'m × 'p*, *'m × 'p*) *lmat*.
*σ ∈ separable-cone-lmat*
⟶ *0 ≤ Re (Tr (W ∗ unwrap-lm σ))*)

⟨*proof*⟩

The witness is also nonneg on every separable density matrix (*Sprime*) since *Sprime ⊆ separable-cone-lmat*.

**theorem** *entanglement-witness-for-density-on-Sprime*:
**fixes** *ϱ* :: (*complex*, *'m* :: *finite × 'p* :: *finite*, *'m × 'p*) *lmat*
**assumes** *ϱ-psd*: *psd-lmat ϱ*
**and** *ϱ-tr*: *Tr-lmat ϱ = 1*
**and** *ϱ-not-sep*: *ϱ ∉ separable-cone-lmat*
**shows** ∃ *W* :: *complex Matrix.mat*.
*W ∈ carrier-mat (CARD('m × 'p)) (CARD('m × 'p))*
∧ *hermitian W*
∧ *Re (Tr (W ∗ unwrap-lm ϱ)) < 0*
∧ (∀ *σ* :: (*complex*, *'m × 'p*, *'m × 'p*) *lmat*.
*σ ∈ Sprime* ⟶ *0 ≤ Re (Tr (W ∗ unwrap-lm σ))*)

⟨*proof*⟩

The full Horodecki characterisation, specialised to density matrices.

**theorem** *separable-density-iff-witness*:
**fixes** *ϱ* :: (*complex*, *'m* :: *finite × 'p* :: *finite*, *'m × 'p*) *lmat*
**assumes** *ϱ-psd*: *psd-lmat ϱ*
**and** *ϱ-tr*: *Tr-lmat ϱ = 1*
**shows** *ϱ ∈ separable-cone-lmat*
⟷
(∀ *W* :: *complex Matrix.mat*.
*W ∈ carrier-mat (CARD('m × 'p)) (CARD('m × 'p))*
∧ *hermitian W*
∧ (∀ *σ* :: (*complex*, *'m × 'p*, *'m × 'p*) *lmat*.
*σ ∈ separable-cone-lmat*
⟶ *0 ≤ Re (Tr (W ∗ unwrap-lm σ))*)
⟶ *0 ≤ Re (Tr (W ∗ unwrap-lm ϱ))*)

⟨*proof*⟩

The trace is identically *1* on the convex hull of *Sprime*.

**lemma** *Tr-lmat-convex-hull-Sprime-eq-one*:
**fixes** *ϱ* :: (*complex*, *'m* :: *finite × 'p* :: *finite*, *'m × 'p*) *lmat*
**assumes** *ϱ ∈ convex hull Sprime*
**shows** *Tr-lmat ϱ = 1*

⟨*proof*⟩

A density matrix is separable iff it is a convex combination of separable pure-product states. This is the standard textbook definition of separability for mixed states.

**theorem** *separable-density-iff-in-convex-hull-Sprime*:
**fixes** *ϱ* :: (*complex*, *'m* :: *finite × 'p* :: *finite*, *'m × 'p*) *lmat*
**assumes** *ϱ-psd*: *psd-lmat ϱ*
**and** *ϱ-tr*: *Tr-lmat ϱ = 1*
**shows** *ϱ ∈ separable-cone-lmat* ⟷ *ϱ ∈ convex hull Sprime*

⟨*proof*⟩

Topology of the separable density-matrix set.

**lemma** *compact-convex-hull-Sprime*:
  *compact* (*convex hull*
        (*Sprime* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat set*))
  ⟨*proof*⟩

**lemma** *closed-convex-hull-Sprime*:
  *closed* (*convex hull*
        (*Sprime* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat set*))
  ⟨*proof*⟩

**lemma** *bounded-convex-hull-Sprime*:
  *bounded* (*convex hull*
        (*Sprime* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat set*))
  ⟨*proof*⟩

Strict separation in the inner-product form: by closedness of *convex hull Sprime* we get a strictly negative-inner-product hyperplane. The Re-trace strict-form is then an immediate corollary once the hermitian witness construction is replayed.

**lemma** *separable-density-separating-hyperplane*:
  **fixes** *ϱ* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** *ϱ-psd*: *psd-lmat ϱ*
    **and** *ϱ-tr*: *Tr-lmat ϱ* = *1*
    **and** *ϱ-not-sep*: *ϱ* ∉ *separable-cone-lmat*
  **shows** ∃ *a b*.
      *inner a ϱ* < *b*
    ∧ (∀ *σ* :: (*complex*, *'m* × *'p*, *'m* × *'p*) *lmat*.
        *σ* ∈ *convex hull Sprime* ⟶ *b* < *inner a σ*)
⟨*proof*⟩

The inner-product-to-Re-trace bridge for hermitian arguments, extracted as a separate lemma so we can reuse it for the strict separation theorem. This is the same hermitisation construction that drives *entanglement-witness-hermitian*.

**lemma** *hermitian-witness-from-lmat*:
  **fixes** *a* :: (*complex*, *'k* :: *finite*, *'k*) *lmat*
  **shows** ∃ *W* :: *complex Matrix.mat*.
      *W* ∈ *carrier-mat* (*CARD*(*'k*)) (*CARD*(*'k*))
    ∧ *hermitian W*
    ∧ (∀ *M* :: (*complex*, *'k*, *'k*) *lmat*.
        *hermitian* (*unwrap-lm M*)
          ⟶ *Re* (*Tr* (*W* ∗ *unwrap-lm M*)) = *inner a M*)
⟨*proof*⟩

The strict Re-trace form of the entanglement witness.

**theorem** *entanglement-witness-strict*:
  **fixes** *ϱ* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
  **assumes** *ϱ-psd*: *psd-lmat ϱ*
    **and** *ϱ-tr*: *Tr-lmat ϱ* = *1*
    **and** *ϱ-not-sep*: *ϱ* ∉ *separable-cone-lmat*
  **shows** ∃ (*W* :: *complex Matrix.mat*) *c*.
      *W* ∈ *carrier-mat* (*CARD*(*'m* × *'p*)) (*CARD*(*'m* × *'p*))
    ∧ *hermitian W*
    ∧ *Re* (*Tr* (*W* ∗ *unwrap-lm ϱ*)) < *c*
    ∧ (∀ *σ* :: (*complex*, *'m* × *'p*, *'m* × *'p*) *lmat*.
        *σ* ∈ *convex hull Sprime*
          ⟶ *c* < *Re* (*Tr* (*W* ∗ *unwrap-lm σ*)))

⟨*proof*⟩

**end**

**theory** *Day24-Separability*
**imports**
*Quantum-Computing11*
*Vern-Paulsen-QC-3-10-Lmat.Separable-Cone*
**begin**

# 106 Days 26–28 (28 October – 2 November 2011): separability and entanglement witnesses

Entanglement revisited. Ensembles $\{\psi_i, p_i\}$ on $\mathcal{H}_A$ and $\{\phi_j, q_j\}$ on $\mathcal{H}_B$ compose to the density matrix $\sum_{i,j} p_i q_j |\psi_i\rangle\langle\psi_i| \otimes |\phi_j\rangle\langle\phi_j|$.

**Definition (notes)** (Day 26)**.** A density matrix $P \in L(\mathcal{H}_A \otimes \mathcal{H}_B)$ is *separable* if $P = \sum_l p_l |\psi_l\rangle\langle\psi_l| \otimes |\phi_l\rangle\langle\phi_l|$ with unit vectors $\psi_l \in \mathcal{H}_A$, $\phi_l \in \mathcal{H}_B$, $p_l \geq 0$, $\sum p_l = 1$; otherwise $P$ is *entangled*. More generally (Day 27) $Sep \subseteq M_r(M_n)$ is the set of matrices $\sum_l P_l \otimes Q_l$ with $P_l \geq 0$, $Q_l \geq 0$.

Are there entangled density matrices, and how can one tell? The notes answer with the theory of convex cones:

**Theorem (notes)** (Days 27–28)**.** Let $V$ be a finite-dimensional real vector space, $K \subseteq V$ closed and convex, $y \notin K$. Then there is a linear $f : V \to \mathbb{R}$ and $\alpha \in \mathbb{R}$ with $f(y) < \alpha \leq f(K)$; if $C$ is a closed cone and $w \notin C$ there is $f$ with $f(w) < 0 \leq f(C)$.

The separable matrices form a closed cone $C$ in the real vector space of Hermitian matrices; an entangled $R$ is separated from $C$ by a real functional, which complexifies to $\tilde{f}(H + iK) = f(H) + if(K)$, and $\tilde{f}$ corresponds to a Hermitian matrix $W$ with $\mathrm{Tr}(WR) < 0 \leq \mathrm{Tr}(W\sigma)$ for all separable $\sigma$: an *entanglement witness*. The notes phrase the final criterion through positive maps $\Phi_f$ obtained from the Arveson correspondence; that form is *separable-iff-positive-maps* in *Day27-Positive-Map-Criterion*, derived from the witness form proved here.

The formal development uses the typed matrices *lmat* so that HOL-Analysis' separation theorem applies: *separable-cone-lmat* is the convex cone generated by tensor products of positive matrices and *Sprime* is the set of product density matrices.

## 106.1 The separable cone (Day 27)

The cone is convex, conic, contains 0, and consists of Hermitian matrices.

**theorem** *D27-separable-cone-convex-cone*:
*convex-cone* (*separable-cone-lmat*
:: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat set*)
⟨*proof*⟩

**theorem** *D27-zero-separable*:
(*0* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*)
∈ *separable-cone-lmat*
⟨*proof*⟩

**theorem** *D27-separable-state-hermitian*:

**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** $\varrho \in$ *separable-cone-lmat*
**shows** *hermitian* (*unwrap-lm* $\varrho$)
⟨*proof*⟩

## 106.2 Separable density matrices (Day 26)

For a density matrix $\rho$ (positive with $\operatorname{Tr}\rho = 1$), membership in the cone is the notes' definition: $\rho$ is a convex combination of product density matrices $\sigma_i \otimes \tau_i$.

**theorem** *D26-separable-density-iff-convex-combination*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** *ϱ-psd*: *psd-lmat* $\varrho$
**and** *ϱ-tr*: *Tr-lmat* $\varrho$ = *1*
**shows** $\varrho \in$ *separable-cone-lmat* $\longleftrightarrow$ $\varrho \in$ *convex hull Sprime*
⟨*proof*⟩

## 106.3 The separable cone is closed (Day 28)

The notes record that $C$ is a closed cone (needed for the separation theorem); here the cone is closed in the Hilbert–Schmidt topology and the separable density matrices form a compact convex set.

**theorem** *D28-separable-cone-closed*:
*closed* (*separable-cone-lmat*
:: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat set*)
⟨*proof*⟩

**theorem** *D28-separable-density-compact*:
*compact* (*convex hull*
(*Sprime* :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat set*))
⟨*proof*⟩

**theorem** *D28-separable-density-closed*:
*closed* (*convex hull*
(*Sprime* :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat set*))
⟨*proof*⟩

**theorem** *D28-separable-density-bounded*:
*bounded* (*convex hull*
(*Sprime* :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat set*))
⟨*proof*⟩

## 106.4 Separation (Days 27–28)

The separation theorem applied to the cone: a non-separable $\rho$ is separated from $C$ by a real-linear functional (an inner product against some $a$), strictly for density matrices because the separable density matrices are compact.

**theorem** *D27-separating-hyperplane-cone*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** $\varrho \notin$ *separable-cone-lmat*
**shows** $\exists\, a$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
*inner a* $\varrho$ < *0*
$\wedge$ ($\forall\, \sigma \in$ *separable-cone-lmat*. *0* $\leq$ *inner a* $\sigma$)
⟨*proof*⟩

**theorem** *D27-strict-hyperplane-density*:

**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** $\varrho$-*psd*: *psd-lmat* $\varrho$
**and** $\varrho$-*tr*: *Tr-lmat* $\varrho$ = *1*
**and** $\varrho$-*not-sep*: $\varrho \notin$ *separable-cone-lmat*
**shows** $\exists\, a\ b.$
*inner a* $\varrho$ < *b*
$\wedge$ ($\forall\, \sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *convex hull Sprime* $\longrightarrow$ *b* < *inner a* $\sigma$)
⟨*proof*⟩

## 106.5 Entanglement witnesses (Day 28)

An *entanglement witness* for $\rho$ is a Hermitian $W$ with $\mathrm{Tr}(W\sigma) \geq 0$ on every separable $\sigma$ and $\mathrm{Tr}(W\rho) < 0$. (i) Every non-separable Hermitian $\rho$ has a witness; (ii) density matrices have a witness with a strict margin; (iii) (Horodecki) $\rho$ is separable iff every Hermitian $W$ that is non-negative on the cone is non-negative on $\rho$.

**theorem** *D28-witness-exists-hermitian*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** *notin*: $\varrho \notin$ *separable-cone-lmat*
**and** $\varrho$-*herm*: *hermitian* (*unwrap-lm* $\varrho$)
**shows** $\exists\, W$ :: *complex Matrix.mat*.
*W* ∈ *carrier-mat* (*CARD*($'m$ × $'p$)) (*CARD*($'m$ × $'p$))
$\wedge$ *hermitian W*
$\wedge$ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)) < *0*
$\wedge$ ($\forall\, \sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *separable-cone-lmat*
$\longrightarrow$ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
⟨*proof*⟩

**theorem** *D28-witness-for-density*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** $\varrho$-*psd*: *psd-lmat* $\varrho$
**and** $\varrho$-*tr*: *Tr-lmat* $\varrho$ = *1*
**and** $\varrho$-*not-sep*: $\varrho \notin$ *separable-cone-lmat*
**shows** $\exists\, W$ :: *complex Matrix.mat*.
*W* ∈ *carrier-mat* (*CARD*($'m$ × $'p$)) (*CARD*($'m$ × $'p$))
$\wedge$ *hermitian W*
$\wedge$ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)) < *0*
$\wedge$ ($\forall\, \sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *separable-cone-lmat*
$\longrightarrow$ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
⟨*proof*⟩

**theorem** *D28-witness-strict-density*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** $\varrho$-*psd*: *psd-lmat* $\varrho$
**and** $\varrho$-*tr*: *Tr-lmat* $\varrho$ = *1*
**and** $\varrho$-*not-sep*: $\varrho \notin$ *separable-cone-lmat*
**shows** $\exists$(*W* :: *complex Matrix.mat*) *c*.
*W* ∈ *carrier-mat* (*CARD*($'m$ × $'p$)) (*CARD*($'m$ × $'p$))
$\wedge$ *hermitian W*
$\wedge$ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)) < *c*
$\wedge$ ($\forall\, \sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *convex hull Sprime*
$\longrightarrow$ *c* < *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
⟨*proof*⟩

**theorem** *D28-separable-iff-witness*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** *ϱ-herm*: *hermitian* (*unwrap-lm* $\varrho$)
**shows** $\varrho \in$ *separable-cone-lmat*
$\longleftrightarrow$
($\forall$ *W* :: *complex Matrix.mat*.
*W* ∈ *carrier-mat* (*CARD*($'m$ × $'p$)) (*CARD*($'m$ × $'p$))
∧ *hermitian W*
∧ ($\forall \sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *separable-cone-lmat*
$\longrightarrow$ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
$\longrightarrow$ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)))
⟨*proof*⟩

**theorem** *D28-separable-density-iff-witness*:
**fixes** $\varrho$ :: (*complex*, $'m$ :: *finite* × $'p$ :: *finite*, $'m$ × $'p$) *lmat*
**assumes** *ϱ-psd*: *psd-lmat* $\varrho$
**and** *ϱ-tr*: *Tr-lmat* $\varrho$ = *1*
**shows** $\varrho \in$ *separable-cone-lmat*
$\longleftrightarrow$
($\forall$ *W* :: *complex Matrix.mat*.
*W* ∈ *carrier-mat* (*CARD*($'m$ × $'p$)) (*CARD*($'m$ × $'p$))
∧ *hermitian W*
∧ ($\forall \sigma$ :: (*complex*, $'m$ × $'p$, $'m$ × $'p$) *lmat*.
$\sigma \in$ *separable-cone-lmat*
$\longrightarrow$ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\sigma$)))
$\longrightarrow$ *0* ≤ *Re* (*Tr* (*W* ∗ *unwrap-lm* $\varrho$)))
⟨*proof*⟩

This closes the entanglement chapter of the notes. The Horodecki–Peres criterion for two qubits is *Day28-Horodecki-Peres*, the positive-map formulation *Day27-Positive-Map-Criterion*; the universal-witness question (Day 28) is not formalised.

**end**
**theory** *Day27-Positive-Map-Criterion*
**imports** *Day26-Arveson-Extension Day34-Knill-Laflamme*
**begin**

Days 27–28: positive maps detect entanglement. For $R \in M_r(M_n)$ (block size $n$), $\Phi^{(r)}(R) \geq 0$ for every positive $\Phi$ when $R$ is separable (D27-1); a functional $f$ on $M_r(M_n)$ is nonnegative on the separable matrices iff $\Phi_f$ is positive (D27-3); and with the entanglement witnesses of *Separable-Cone* this gives the positive-map criterion (D28-3).

**lemma** *block-of-mat-carrier′* [*simp*]: *block-of-mat n A a b* ∈ *carrier-mat n n*
⟨*proof*⟩

## 106.6 Tensor products with a matrix in the first factor

**lemma** *tensor-add-first*:
**fixes** *A B P* :: *complex Matrix.mat*
**assumes** *A*: *A* ∈ *carrier-mat r r* **and** *B*: *B* ∈ *carrier-mat r r* **and** *P*: *P* ∈ *carrier-mat n n* **and** *n*: *0* < *n*
**shows** *Tensor.tensor-mat* (*A* + *B*) *P* = *Tensor.tensor-mat A P* + *Tensor.tensor-mat B P*
⟨*proof*⟩

**lemma** *tensor-zero-first*:
**fixes** *P* :: *complex Matrix.mat*
**assumes** *P*: *P* ∈ *carrier-mat n n* **and** *n*: *0* < *n*
**shows** *Tensor.tensor-mat* (*0*$_m$ *r r*) *P* = *0*$_m$ (*r* ∗ *n*) (*r* ∗ *n*)

⟨*proof*⟩

**lemma** *tensor-msum-first*:
**fixes** $P$ :: *complex Matrix.mat*
**assumes** $g$: $\forall x \in$ *set xs*. $g\ x \in$ *carrier-mat r r* **and** $P$: $P \in$ *carrier-mat n n* **and** $n$: $0 < n$
**shows** *Tensor.tensor-mat* (*msum r r g xs*) $P$ = *msum* $(r * n)$ $(r * n)$ ($\lambda x$. *Tensor.tensor-mat* $(g\ x)$ $P$) *xs*
⟨*proof*⟩

## 106.7 The quadratic form of $\Phi_f$

**definition** *rank1* :: *complex Matrix.vec* ⇒ *complex Matrix.mat* **where**
*rank1 w* = *Matrix.mat* (*dim-vec w*) (*dim-vec w*) ($\lambda$(*i*,*j*). *cnj* (*w* $ *i*) * *w* $ *j*)

**lemma** *rank1-carrier* [*simp*]: *rank1 w* ∈ *carrier-mat* (*dim-vec w*) (*dim-vec w*)
⟨*proof*⟩

**lemma** *rank1-dims* [*simp*]: *dim-row* (*rank1 w*) = *dim-vec w dim-col* (*rank1 w*) = *dim-vec w*
⟨*proof*⟩

**lemma** *rank1-outer*: *rank1* (*conjugate u*) = *u* ⊘ *u*
⟨*proof*⟩

**lemma** *rank1-psd*: *psd* (*rank1 w*)
⟨*proof*⟩

**lemma** *quad-arveson-map*:
**fixes** *f* :: *complex Matrix.mat* ⇒ *complex* **and** *w* :: *complex Matrix.vec*
**assumes** *f*: *lin-fun-on* (*carrier-mat* $(r * n)$ $(r * n)$) *f* **and** *w*: *dim-vec w* = *r* **and** *P*: *P* ∈ *carrier-mat n n* **and** *n*: $0 < n$
**shows** ⟨*w* | *arveson-map n r f P* $*_v$ *w*⟩ = *of-nat r* * *f* (*Tensor.tensor-mat* (*rank1 w*) *P*)
⟨*proof*⟩

## 106.8 Proposition (Day 27): $f(\mathrm{Sep}) \geq 0$ iff $\Phi_f$ is positive

**theorem** *positive-map-iff-sep-nonneg*:
**fixes** *f* :: *complex Matrix.mat* ⇒ *complex*
**assumes** *f*: *lin-fun-on* (*carrier-mat* $(r * n)$ $(r * n)$) *f* **and** *r*: $0 < r$ **and** *n*: $0 < n$
**shows** *positive-map n r* (*arveson-map n r f*)
$\longleftrightarrow$ ($\forall$ *Q P*. *Q* ∈ *carrier-mat r r* ⟶ *psd Q* ⟶ *P* ∈ *carrier-mat n n* ⟶ *psd P*
⟶ $0 \leq$ *f* (*Tensor.tensor-mat Q P*))
⟨*proof*⟩

## 106.9 Proposition (Day 27): positive maps keep separable matrices positive

**lemma** *block-tensor*:
**fixes** *Q P* :: *complex Matrix.mat*
**assumes** *Q*: *Q* ∈ *carrier-mat r r* **and** *P*: *P* ∈ *carrier-mat n n* **and** *a*: $a < r$ **and** *b*: $b < r$
**shows** *block-of-mat n* (*Tensor.tensor-mat Q P*) *a b* = *Q* $$ (*a*,*b*) $\cdot_m$ *P*
⟨*proof*⟩

**lemma** *amplification-tensor*:
**fixes** *Q P* :: *complex Matrix.mat* **and** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *Q*: *Q* ∈ *carrier-mat r r* **and** *P*: *P* ∈ *carrier-mat n n*
**and** *d*: $0 < d$ **and** *n*: $0 < n$
**shows** *amplification n d r* Φ (*Tensor.tensor-mat Q P*) = *Tensor.tensor-mat Q* (Φ *P*)
⟨*proof*⟩

**lemma** *amplification-lin*:
**fixes** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *L*: *lin-on n d* (*carrier-mat n n*) Φ **and** *A*: *A* ∈ *carrier-mat* (*p* ∗ *n*) (*p* ∗ *n*) **and** *B*: *B* ∈ *carrier-mat* (*p* ∗ *n*) (*p* ∗ *n*)
**shows** *amplification n d p* Φ (*A* + *B*) = *amplification n d p* Φ *A* + *amplification n d p* Φ *B*
**and** *amplification n d p* Φ (*c* ·*m A*) = *c* ·*m amplification n d p* Φ *A*
⟨*proof*⟩

**lemma** *unwrap-lm-scaleR-mat*:
**fixes** *R* :: (*complex*, 'k :: *finite*, 'k) *lmat*
**shows** *unwrap-lm* (*u* ∗*R R*) = *complex-of-real u* ·*m unwrap-lm R*
⟨*proof*⟩

**theorem** *positive-map-separable*:
**fixes** *R* :: (*complex*, 'm :: *finite* × 'p :: *finite*, 'm × 'p) *lmat* **and** Φ :: *complex Matrix.mat* ⇒ *complex Matrix.mat*
**assumes** *L*: *lin-on CARD*('p) *d* (*carrier-mat CARD*('p) *CARD*('p)) Φ **and** *pos*: *positive-map CARD*('p) *d* Φ
**and** *d*: *0* < *d* **and** *R*: *R* ∈ *separable-cone-lmat*
**shows** *psd* (*amplification CARD*('p) *d CARD*('m) Φ (*unwrap-lm R*))
⟨*proof*⟩

### 106.10 Theorem (Day 28): the positive-map criterion

A Hermitian $R \in M_r(M_n)$ is separable iff $\Phi^{(r)}(R) \geq 0$ for every positive $\Phi : M_n \to M_r$.

**theorem** *separable-iff-positive-maps*:
**fixes** *R* :: (*complex*, 'm :: *finite* × 'p :: *finite*, 'm × 'p) *lmat*
**assumes** *Rh*: *hermitian* (*unwrap-lm R*)
**shows** *R* ∈ *separable-cone-lmat* ⟷
(∀Φ. *lin-on CARD*('p) *CARD*('m) (*carrier-mat CARD*('p) *CARD*('p)) Φ ⟶ *positive-map CARD*('p) *CARD*('m) Φ
⟶ *psd* (*amplification CARD*('p) *CARD*('m) *CARD*('m) Φ (*unwrap-lm R*)))
⟨*proof*⟩

**end**
**theory** *Day27-Entangled-Examples*
**imports** *Day18-Positive-Maps Day27-Positive-Map-Criterion*
**begin**

# 107 Day 27: entangled matrices detected by the transpose (D27-2)

For $f : \{0, \ldots, r-1\} \to \{0, \ldots, n-1\}$ the block matrix $(E_{f(a)f(b)})_{a,b} \in M_r(M_n)$. The notes' examples are $f = (i, j)$ with $r = 2$, giving $\begin{pmatrix} E_{ii} & E_{ij} \\ E_{ji} & E_{jj} \end{pmatrix}$, and $f = \mathrm{id}$ with $r = n$, giving $(E_{ij})$.

**definition** *unit-blocks* :: *nat* ⇒ *nat* ⇒ (*nat* ⇒ *nat*) ⇒ *complex Matrix.mat* **where**
*unit-blocks r n f* = *Matrix.mat* (*r* ∗ *n*) (*r* ∗ *n*)
(λ(*x*, *y*). *if x mod n* = *f* (*x div n*) ∧ *y mod n* = *f* (*y div n*) *then 1 else 0*)

**lemma** *block-index-less*:
**fixes** *a k r n* :: *nat*
**assumes** *a* < *r k* < *n*
**shows** *a* ∗ *n* + *k* < *r* ∗ *n*
⟨*proof*⟩

**lemma** *unit-blocks-carrier* [*simp*]: *unit-blocks r n f* ∈ *carrier-mat* (*r* ∗ *n*) (*r* ∗ *n*)
⟨*proof*⟩

**lemma** *unit-blocks-block*:
**assumes** *a* < *r b* < *r f a* < *n f b* < *n*
**shows** *block-of-mat n* (*unit-blocks r n f*) *a b* = *mat-unit* (*f a*) (*f b*) *n n*
⟨*proof*⟩

**definition** *unit-blocks-vec* :: *nat* ⇒ *nat* ⇒ (*nat* ⇒ *nat*) ⇒ *complex Matrix.vec* **where**
*unit-blocks-vec r n f* = *Matrix.vec* (*r* ∗ *n*) (λ*x*. *if x mod n* = *f* (*x div n*) *then 1 else 0*)

**lemma** *unit-blocks-outer*: *unit-blocks r n f* = *unit-blocks-vec r n f* ⊘ *unit-blocks-vec r n f*
⟨*proof*⟩

**lemma** *unit-blocks-psd*: *psd* (*unit-blocks r n f*)
⟨*proof*⟩

**lemma** *sum-two-support*:
**fixes** *g* :: *nat* ⇒ *complex*
**assumes** *p* ≠ *q p* < *m q* < *m* ⋀*x*. *x* < *m* ⟹ *x* ≠ *p* ⟹ *x* ≠ *q* ⟹ *g x* = *0*
**shows** (∑ *x* = *0*..<*m*. *g x*) = *g p* + *g q*
⟨*proof*⟩

**lemma** *transpose-lin-on*: *lin-on n n* (*carrier-mat n n*) (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*)
⟨*proof*⟩

The partial transpose of $(E_{f(a)f(b)})$ is $(E_{f(b)f(a)})$; its quadratic form at $e_0 \otimes e_{f(1)} - e_1 \otimes e_{f(0)}$ is $-2$.

**lemma** *unit-blocks-partial-transpose-not-psd*:
**assumes** *r*: *2* ≤ *r* **and** *f*: *f 0* < *n f 1* < *n f 0* ≠ *f 1*
**shows** ¬ *psd* (*amplification n n r* (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*) (*unit-blocks r n f*))
⟨*proof*⟩

**theorem** *unit-blocks-entangled*:
**fixes** *R* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
**assumes** *R*: *unwrap-lm R* = *unit-blocks CARD('m) CARD('p) f*
**and** *r*: *2* ≤ *CARD('m)* **and** *f*: *f 0* < *CARD('p) f 1* < *CARD('p) f 0* ≠ *f 1*
**shows** *psd* (*unwrap-lm R*) ∧ *R* ∉ *separable-cone-lmat*
⟨*proof*⟩

The notes' first example: $\begin{pmatrix} E_{ii} & E_{ij} \\ E_{ji} & E_{jj} \end{pmatrix} \in M_2(M_n)$, $i \neq j$.

**corollary** *entangled-example-2*:
**fixes** *R* :: (*complex*, *'m* :: *finite* × *'p* :: *finite*, *'m* × *'p*) *lmat*
**assumes** *m*: *CARD('m)* = *2* **and** *ij*: *i* < *CARD('p) j* < *CARD('p) i* ≠ *j*
**and** *R*: *unwrap-lm R* = *unit-blocks 2 CARD('p)* (λ*a*. *if a* = *0 then i else j*)
**shows** *block-of-mat CARD('p)* (*unwrap-lm R*) *0 0* = *E*⟦*i*,*i* ; *CARD('p)*,*CARD('p)*⟧
**and** *block-of-mat CARD('p)* (*unwrap-lm R*) *0 1* = *E*⟦*i*,*j* ; *CARD('p)*,*CARD('p)*⟧
**and** *block-of-mat CARD('p)* (*unwrap-lm R*) *1 0* = *E*⟦*j*,*i* ; *CARD('p)*,*CARD('p)*⟧
**and** *block-of-mat CARD('p)* (*unwrap-lm R*) *1 1* = *E*⟦*j*,*j* ; *CARD('p)*,*CARD('p)*⟧
**and** *psd* (*unwrap-lm R*) ∧ *R* ∉ *separable-cone-lmat*
⟨*proof*⟩

The second: $(E_{ij}) \in M_n(M_n)$, $n \geq 2$, which is *choi-seed*.

**lemma** *choi-seed-unit-blocks*: *choi-seed n* = *unit-blocks n n id*

⟨*proof*⟩

**corollary** *entangled-example-choi-seed*:
**fixes** $R$ :: (*complex*, $'p$ :: *finite* × $'p$, $'p$ × $'p$) *lmat*
**assumes** *n*: $2 \le CARD('p)$ **and** *R*: *unwrap-lm R* = *choi-seed CARD*($'p$)
**shows** *psd* (*unwrap-lm R*) ∧ $R \notin$ *separable-cone-lmat*
⟨*proof*⟩

**end**
**theory** *Day28-Horodecki-Peres*
**imports** *QC-Prelims*
**begin**

# 108 Day 28: the Horodecki–Peres criterion for two qubits (D28-5)

Entries of the partial transpose $\Phi^{(2)}(X)$ of a $4 \times 4$ matrix.

**lemma** *partial-transpose-2-index*:
**fixes** *M* :: *complex Matrix.mat*
**assumes** *i*: $i < 4$ **and** *j*: $j < 4$
**shows** *amplification 2 2 2* (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*) *M* $$ (*i*,*j*)
= *M* $$ ((*i div 2*) * *2* + *j mod 2*, (*j div 2*) * *2* + *i mod 2*)
⟨*proof*⟩

**lemma** *product-vec-iff-det-zero*:
**fixes** *v* :: *complex Matrix.vec*
**assumes** *vd*: *dim-vec v* = *4*
**shows** (∃ *α β*. *dim-vec α* = *2* ∧ *dim-vec β* = *2* ∧ *v* = *tensor-vec α β*)
⟷ *v* $ *0* * *v* $ *3* − *v* $ *1* * *v* $ *2* = *0*
⟨*proof*⟩

The notes' witness $w = (-\overline{v_2}, \overline{v_1})$ gives $\langle w|\Phi^{(2)}(|v\rangle\langle v|)w\rangle = -2|v_0v_3 - v_1v_2|^2$ ($= 2|\langle v_2|v_1\rangle|^2 - 2\|v_1\|^2\|v_2\|^2$ by Lagrange's identity).

**lemma** *partial-transpose-witness-value*:
**fixes** *v* :: *complex Matrix.vec*
**assumes** *vd*: *dim-vec v* = *4*
**defines** *w* ≡ *Matrix.vec 4* (λ*k*. *if k* = *0 then* − *cnj* (*v* $ *2*) *else if k* = *1 then* − *cnj* (*v* $ *3*)
*else if k* = *2 then cnj* (*v* $ *0*) *else cnj* (*v* $ *1*))
**shows** ⟨*w* | *amplification 2 2 2* (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*) (*v* ⊘ *v*) *$_v$ w*⟩
= − *2* * *cnj* (*v* $ *0* * *v* $ *3* − *v* $ *1* * *v* $ *2*) * (*v* $ *0* * *v* $ *3* − *v* $ *1* * *v* $ *2*)
⟨*proof*⟩

**theorem** *horodecki-peres-two-qubits*:
**fixes** *v* :: *complex Matrix.vec*
**assumes** *vd*: *dim-vec v* = *4*
**shows** *psd* (*amplification 2 2 2* (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*) (*v* ⊘ *v*))
⟷ (∃ *α β*. *dim-vec α* = *2* ∧ *dim-vec β* = *2* ∧ *v* = *tensor-vec α β*)
⟨*proof*⟩

**corollary** *horodecki-peres-entangled*:
**fixes** *v* :: *complex Matrix.vec*
**assumes** *vd*: *dim-vec v* = *4*
**shows** *entangled 2 2 v* ⟷

¬ *psd* (*amplification 2 2 2* (*transpose-mat* :: *complex Matrix.mat* ⇒ *complex Matrix.mat*) (*v* ⊘ *v*))
⟨*proof*⟩

**end**

# Part VI
# Quantum Algorithms (beyond the notes)

**theory** *Hermitian-Functional-Calculus*
**imports** *Vern-Paulsen-QC-3-10.Quantum-Computing-3-10*
**begin**

## 109 Functional calculus for Hermitian matrices

For Hermitian $A$ and $g : \mathbb{R} \to \mathbb{C}$, $g(A)$ is the matrix that acts on every eigenvector $v$ of $A$ with eigenvalue $\lambda$ as multiplication by $g(\lambda)$. The eigenvectors span $\mathbb{C}^n$, so this determines $g(A)$; for a spectral decomposition $A = U \operatorname{diag}(D)\, U^*$ it is $U \operatorname{diag}(g \circ D)\, U^*$.

**definition** *diag-cmat* :: *complex list* ⇒ *complex Matrix.mat* **where**
*diag-cmat xs* = *Matrix.mat* (*length xs*) (*length xs*) (λ(*i*,*j*). *if i* = *j then xs* ! *i else 0*)

**lemma** *diag-cmat-carrier* [*simp*]: *diag-cmat xs* ∈ *carrier-mat* (*length xs*) (*length xs*)
⟨*proof*⟩

**lemma** *dim-diag-cmat* [*simp*]: *dim-row* (*diag-cmat xs*) = *length xs dim-col* (*diag-cmat xs*) = *length xs*
⟨*proof*⟩

**lemma** *diag-mat-of-real-cmat*: *diag-mat-of-real D* = *diag-cmat* (*map complex-of-real D*)
⟨*proof*⟩

**lemma** *diag-cmat-mult-vec*:
**assumes** *dim-vec w* = *length xs*
**shows** *diag-cmat xs* ∗$_v$ *w* = *Matrix.vec* (*length xs*) (λ*i*. *xs* ! *i* ∗ *w* $ *i*)
⟨*proof*⟩

**lemma** *col-mult-diag*:
**assumes** *M*: *M* ∈ *carrier-mat n* (*length xs*) **and** *j*: *j* < *length xs*
**shows** *Matrix.col* (*M* ∗ *diag-cmat xs*) *j* = *xs* ! *j* ·$_v$ *Matrix.col M j*
⟨*proof*⟩

**lemma** *diag-cmat-dagger*: (*diag-cmat xs*)$^\dagger$ = *diag-cmat* (*map cnj xs*)
⟨*proof*⟩

**lemma** *smult-mat-mult-vec*:
*A* ∈ *carrier-mat n m* ⟹ *dim-vec v* = *m* ⟹ (*c* ·$_m$ *A*) ∗$_v$ *v* = *c* ·$_v$ (*A* ∗$_v$ *v*)
⟨*proof*⟩

**lemma** *diag-conj-carrier*:
**assumes** *U*: *U* ∈ *carrier-mat n n* **and** *xs*: *length xs* = *n*
**shows** *U* ∗ *diag-cmat xs* ∗ (*U*$^\dagger$) ∈ *carrier-mat n n*
⟨*proof*⟩

## 109.1 Spectral decompositions

*spectral-decomp A U D*: $U$ unitary and $U^* A U = \mathrm{diag}(D)$.

**definition** *spectral-decomp* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *real list* ⇒ *bool* **where**
*spectral-decomp A U D* ⟷ *U* ∈ *carrier-mat* (*dim-row A*) (*dim-row A*) ∧ *unitary U* ∧
*length D* = *dim-row A* ∧ $U^\dagger$ ∗ *A* ∗ *U* = *diag-mat-of-real D*

**lemma** *spectral-decomp-exists*:
**assumes** *A* ∈ *carrier-mat n n hermitian A*
**shows** ∃ *U D*. *spectral-decomp A U D*
⟨*proof*⟩

**lemma** *spectral-decompD*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *sd*: *spectral-decomp A U D*
**shows** *U* ∈ *carrier-mat n n unitary U length D* = *n* $U^\dagger$ ∗ *U* = $1_m$ *n U* ∗ ($U^\dagger$) = $1_m$ *n*
*A* = *U* ∗ *diag-mat-of-real D* ∗ ($U^\dagger$) *A* ∗ *U* = *U* ∗ *diag-mat-of-real D*
⟨*proof*⟩

The columns of $U$ are eigenvectors: $Au_j = D_j u_j$.

**lemma** *spectral-decomp-col*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *sd*: *spectral-decomp A U D* **and** *j*: *j* < *n*
**shows** *A* ∗$_v$ *Matrix.col U j* = *complex-of-real* (*D* ! *j*) ·$_v$ *Matrix.col U j*
⟨*proof*⟩

## 109.2 The definition

**definition** *fun-on-eigen* :: *complex Matrix.mat* ⇒ (*real* ⇒ *complex*) ⇒ *complex Matrix.mat* ⇒ *bool* **where**
*fun-on-eigen A g B* ⟷ (∀ *v l*. *dim-vec v* = *dim-row A* ⟶
*A* ∗$_v$ *v* = *complex-of-real l* ·$_v$ *v* ⟶ *B* ∗$_v$ *v* = *g l* ·$_v$ *v*)

**definition** *mat-fun* :: (*real* ⇒ *complex*) ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*mat-fun g A* = (*SOME B*. *B* ∈ *carrier-mat* (*dim-row A*) (*dim-row A*) ∧ *fun-on-eigen A g B*)

**definition** *herm-spectrum* :: *complex Matrix.mat* ⇒ *real set* **where**
*herm-spectrum A* = {*l*. ∃ *v*. *dim-vec v* = *dim-row A* ∧ *v* ≠ $0_v$ (*dim-row A*) ∧
*A* ∗$_v$ *v* = *complex-of-real l* ·$_v$ *v*}

**lemma** *fun-on-eigenD*:
*fun-on-eigen A g B* ⟹ *dim-vec v* = *dim-row A* ⟹ *A* ∗$_v$ *v* = *complex-of-real l* ·$_v$ *v* ⟹
*B* ∗$_v$ *v* = *g l* ·$_v$ *v*
⟨*proof*⟩

$U\,\mathrm{diag}(g \circ D)\,U^*$ acts correctly on eigenvectors: if $Av = \lambda v$ then $w = U^* v$ has $D_i w_i = \lambda w_i$, so $g(D_i) w_i = g(\lambda) w_i$.

**lemma** *spectral-decomp-fun-on-eigen*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *sd*: *spectral-decomp A U D*
**shows** *fun-on-eigen A g* (*U* ∗ *diag-cmat* (*map g D*) ∗ ($U^\dagger$))
⟨*proof*⟩

Uniqueness: two matrices that agree on the eigenvectors $u_j$ agree on $U$, hence are equal.

**lemma** *fun-on-eigen-unique*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A*
**and** *B1*: *B1* ∈ *carrier-mat n n fun-on-eigen A g B1*
**and** *B2*: *B2* ∈ *carrier-mat n n fun-on-eigen A g B2*
**shows** *B1* = *B2*
⟨*proof*⟩

**lemma** *mat-fun-spec*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**shows** *mat-fun g A* $\in$ *carrier-mat n n fun-on-eigen A g* (*mat-fun g A*)
⟨*proof*⟩

**lemma** *mat-fun-carrier* [*simp*]: $A \in$ *carrier-mat n n* $\Longrightarrow$ *hermitian A* $\Longrightarrow$ *mat-fun g A* $\in$ *carrier-mat n n*
⟨*proof*⟩

**theorem** *mat-fun-eigen*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**and** *v*: *dim-vec v* = *n* **and** *ev*: $A *_v v =$ *complex-of-real l* $\cdot_v$ *v*
**shows** *mat-fun g A* $*_v$ *v* = *g l* $\cdot_v$ *v*
⟨*proof*⟩

**lemma** *mat-fun-eqI*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**and** *B*: $B \in$ *carrier-mat n n fun-on-eigen A g B*
**shows** *mat-fun g A* = *B*
⟨*proof*⟩

**theorem** *mat-fun-decomp*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *sd*: *spectral-decomp A U D*
**shows** *mat-fun g A* = *U* $*$ *diag-cmat* (*map g D*) $*$ ($U^\dagger$)
⟨*proof*⟩

Only the values of $g$ on the spectrum matter.

**lemma** *mat-fun-cong*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**and** *gh*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow$ *g l* = *h l*
**shows** *mat-fun g A* = *mat-fun h A*
⟨*proof*⟩

## 109.3 Algebra

**lemma** *mat-fun-id*: $A \in$ *carrier-mat n n* $\Longrightarrow$ *hermitian A* $\Longrightarrow$ *mat-fun complex-of-real A* = *A*
⟨*proof*⟩

**lemma** *mat-fun-const*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**shows** *mat-fun* (λ-. *c*) *A* = *c* $\cdot_m$ $1_m$ *n*
⟨*proof*⟩

**lemma** *mat-fun-one*: $A \in$ *carrier-mat n n* $\Longrightarrow$ *hermitian A* $\Longrightarrow$ *mat-fun* (λ-. *1*) *A* = $1_m$ *n*
⟨*proof*⟩

**theorem** *mat-fun-mult*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**shows** *mat-fun g A* $*$ *mat-fun h A* = *mat-fun* (λ*x*. *g x* $*$ *h x*) *A*
⟨*proof*⟩

**theorem** *mat-fun-dagger*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**shows** (*mat-fun g A*)$^\dagger$ = *mat-fun* (λ*x*. *cnj* (*g x*)) *A*
⟨*proof*⟩

**lemma** *mat-fun-hermitian*:

**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**shows** *hermitian* (*mat-fun* ($\lambda x$. *complex-of-real* (*f x*)) *A*)
⟨*proof*⟩

**theorem** *mat-fun-unitary*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**and** *g*: $\bigwedge l$. $l \in$ *herm-spectrum A* $\Longrightarrow$ *cmod* (*g l*) = *1*
**shows** *unitary* (*mat-fun g A*)
⟨*proof*⟩

## 109.4 Time evolution

$e^{itA}$, the evolution under the Hamiltonian $A$ (phase estimation uses $U = e^{itA}$).

**definition** *evolution* :: *complex Matrix.mat* ⇒ *real* ⇒ *complex Matrix.mat* **where**
*evolution A t* = *mat-fun* ($\lambda x$. *cis* (*t* ∗ *x*)) *A*

**context**
**fixes** *A* :: *complex Matrix.mat* **and** *n* :: *nat*
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**begin**

**lemma** *evolution-carrier* [*simp*]: *evolution A t* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *evolution-eigen*:
*dim-vec v* = *n* $\Longrightarrow$ *A* $*_v$ *v* = *complex-of-real l* $\cdot_v$ *v* $\Longrightarrow$ *evolution A t* $*_v$ *v* = *cis* (*t* ∗ *l*) $\cdot_v$ *v*
⟨*proof*⟩

**lemma** *evolution-unitary*: *unitary* (*evolution A t*)
⟨*proof*⟩

**lemma** *evolution-zero*: *evolution A 0* = $1_m$ *n*
⟨*proof*⟩

**lemma** *evolution-add*: *evolution A* (*s* + *t*) = *evolution A s* ∗ *evolution A t*
⟨*proof*⟩

**lemma** *evolution-pow*: *evolution A t* $\hat{}_m$ *k* = *evolution A* (*real k* ∗ *t*)
⟨*proof*⟩

**end**

## 109.5 The inverse

$A^{-1} = g(A)$ for $g(x) = 1/x$; since $1/0 = 0$ in HOL this is the pseudo-inverse in general.

**definition** *herm-inv* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*herm-inv A* = *mat-fun* ($\lambda x$. *complex-of-real* (*inverse x*)) *A*

**theorem** *herm-inv*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *z*: *0* ∉ *herm-spectrum A*
**shows** *herm-inv A* ∗ *A* = $1_m$ *n A* ∗ *herm-inv A* = $1_m$ *n*
⟨*proof*⟩

**lemma** *herm-inv-eigen*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**and** *v*: *dim-vec v* = *n* **and** *ev*: *A* $*_v$ *v* = *complex-of-real l* $\cdot_v$ *v*
**shows** *herm-inv A* $*_v$ *v* = *complex-of-real* (*inverse l*) $\cdot_v$ *v*

⟨*proof*⟩

**lemma** *herm-inv-hermitian*: $A \in$ *carrier-mat n n* $\Longrightarrow$ *hermitian A* $\Longrightarrow$ *hermitian* (*herm-inv A*)
⟨*proof*⟩

**end**
**theory** *Phase-Estimation*
**imports** *Hermitian-Functional-Calculus HOL−Analysis.Convex*
**begin**

# 110 Phase estimation

The Fourier matrix $F_N$, $(F_N)_{jk} = \omega^{jk}/\sqrt{N}$ with $\omega = e^{2\pi i/N}$, and phase estimation: for $Uu = e^{2\pi i\varphi}u$, the circuit $(F_N^* \otimes I)\, CU\, (\frac{1}{\sqrt{N}} \sum_k |k\rangle \otimes u)$ leaves the register in $\sum_j \alpha_j |j\rangle$ with $\alpha_j = \frac{1}{N} \sum_k e^{2\pi i k(\varphi - j/N)}$.

## 110.1 Sums of roots of unity

**lemma** *root-of-unity-sum*:
**assumes** *N*: $0 < N$ **and** *j*: $j < N$ **and** *l*: $l < N$
**shows** $(\sum k{<}N.\ cis\ (2 * pi * real\ k * (real\ l - real\ j)\ /\ real\ N)) = (if\ l = j\ then\ of\text{-}nat\ N\ else\ 0)$
⟨*proof*⟩

## 110.2 The Fourier matrix

**definition** *fourier* :: *nat* ⇒ *complex Matrix.mat* **where**
*fourier N* = *Matrix.mat N N* (λ(*j*,*k*). *cis* (2 ∗ *pi* ∗ *real j* ∗ *real k* / *real N*) / *complex-of-real* (*sqrt* (*real N*)))

**lemma** *fourier-carrier* [*simp*]: *fourier N* ∈ *carrier-mat N N*
⟨*proof*⟩

**lemma** *dim-fourier* [*simp*]: *dim-row* (*fourier N*) = *N dim-col* (*fourier N*) = *N*
⟨*proof*⟩

**lemma** *fourier-index*:
$j < N \Longrightarrow k < N \Longrightarrow$ *fourier N* $$ (*j*, *k*) = *cis* (2 ∗ *pi* ∗ *real j* ∗ *real k* / *real N*) / *complex-of-real* (*sqrt* (*real N*))
⟨*proof*⟩

**lemma** *sqrt-N-sq*: *complex-of-real* (*sqrt* (*real N*)) ∗ *complex-of-real* (*sqrt* (*real N*)) = *of-nat N*
⟨*proof*⟩

The entries pair to a root of unity: $\overline{F_{kj}} F_{kl} = \omega^{k(l-j)}/N$.

**lemma** *fourier-entry-pair*:
**assumes** $j < N$ $k < N$ $l < N$
**shows** *cnj* (*fourier N* $$ (*k*, *j*)) ∗ *fourier N* $$ (*k*, *l*) = *cis* (2 ∗ *pi* ∗ *real k* ∗ (*real l* − *real j*) / *real N*) / *of-nat N*
⟨*proof*⟩

**theorem** *fourier-unitary*:
**assumes** *N*: $0 < N$
**shows** *unitary* (*fourier N*)
⟨*proof*⟩

## 110.3 Controlled powers and phase kickback

$\mathrm{C}U = \sum_k |k\rangle\langle k| \otimes U^k$ on $\mathbb{C}^N \otimes \mathbb{C}^m$ (index $km + i$).

**definition** *ctrl-pow* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*ctrl-pow N m U* = *Matrix.mat* (*N* ∗ *m*) (*N* ∗ *m*)
(λ(*r*, *c*). *if r div m* = *c div m then* (*U* ^*m* (*r div m*)) $$ (*r mod m*, *c mod m*) *else 0*)

**lemma** *ctrl-pow-carrier* [*simp*]: *ctrl-pow N m U* ∈ *carrier-mat* (*N* ∗ *m*) (*N* ∗ *m*)
⟨*proof*⟩

**lemma** *dim-ctrl-pow* [*simp*]: *dim-row* (*ctrl-pow N m U*) = *N* ∗ *m dim-col* (*ctrl-pow N m U*) = *N* ∗ *m*
⟨*proof*⟩

**lemma** *ctrl-pow-index*:
*r* < *N* ∗ *m* ⟹ *c* < *N* ∗ *m* ⟹
*ctrl-pow N m U* $$ (*r*, *c*) = (*if r div m* = *c div m then* (*U* ^*m* (*r div m*)) $$ (*r mod m*, *c mod m*) *else 0*)
⟨*proof*⟩

**lemma** *sum-blocks*:
**fixes** *N m* :: *nat*
**shows** (∑ *c*<*N* ∗ *m*. *f c*) = (∑ *k*<*N*. ∑ *i*<*m*. *f* (*k* ∗ *m* + *i*))
⟨*proof*⟩

**lemma** *block-index*:
**fixes** *k i N m* :: *nat*
**assumes** *k* < *N i* < *m*
**shows** *k* ∗ *m* + *i* < *N* ∗ *m* (*k* ∗ *m* + *i*) *div m* = *k* (*k* ∗ *m* + *i*) *mod m* = *i*
⟨*proof*⟩

**lemma** *pow-eigen*:
**fixes** *U* :: *complex Matrix.mat*
**assumes** *U*: *U* ∈ *carrier-mat m m* **and** *u*: *dim-vec u* = *m* **and** *ev*: *U* ∗*v u* = *z* ·*v u*
**shows** (*U* ^*m k*) ∗*v u* = *z* ^ *k* ·*v u*
⟨*proof*⟩

Phase kickback: if $Uu = zu$ then $\mathrm{C}U(a \otimes u) = (\sum_k z^k a_k |k\rangle) \otimes u$.

**theorem** *ctrl-pow-kickback*:
**fixes** *U* :: *complex Matrix.mat*
**assumes** *U*: *U* ∈ *carrier-mat m m* **and** *u*: *dim-vec u* = *m* **and** *ev*: *U* ∗*v u* = *z* ·*v u*
**and** *a*: *dim-vec a* = *N*
**shows** *ctrl-pow N m U* ∗*v tensor-vec a u* = *tensor-vec* (*Matrix.vec N* (λ*k*. *z* ^ *k* ∗ *a* $ *k*)) *u*
⟨*proof*⟩

## 110.4 The phase estimation circuit

The register starts in the uniform state $\frac{1}{\sqrt{N}} \sum_k |k\rangle$ ($= H^{\otimes t}|0\rangle$ for $N = 2^t$).

**definition** *uniform-state* :: *nat* ⇒ *complex Matrix.vec* **where**
*uniform-state N* = *Matrix.vec N* (λ-. *1* / *complex-of-real* (*sqrt* (*real N*)))

**definition** *pe-amp* :: *nat* ⇒ *real* ⇒ *nat* ⇒ *complex* **where**
*pe-amp N φ j* = (∑ *k*<*N*. *cis* (*2* ∗ *pi* ∗ *real k* ∗ (*φ* − *real j* / *real N*))) / *of-nat N*

**lemma** *fourier-dagger-kick*:
**assumes** *j*: *j* < *N*
**shows** ((*fourier N*)† ∗*v Matrix.vec N* (λ*k*. *cis* (*2* ∗ *pi* ∗ *φ*) ^ *k* ∗ *uniform-state N* $ *k*)) $ *j* = *pe-amp N φ j*

⟨*proof*⟩

The output of phase estimation for an eigenvector $u$, $Uu = e^{2\pi i\varphi}u$.

**theorem** *phase-estimation*:
  **assumes** *U*: $U \in$ *carrier-mat m m* **and** *u*: *dim-vec u = m*
    **and** *ev*: $U *_v u =$ *cis* $(2 * pi * \varphi) \cdot_v u$
  **shows** $(((fourier\ N)^\dagger) \bigotimes 1_m\ m) *_v$ *(ctrl-pow N m U* $*_v$ *tensor-vec* (*uniform-state N*) *u*) =
      *tensor-vec* (*Matrix.vec N* (*pe-amp N* $\varphi$)) *u*
⟨*proof*⟩

If $\varphi = j_0/N$ exactly, the register ends in $|j_0\rangle$.

**theorem** *phase-estimation-exact*:
  **assumes** *N*: *0 < N* **and** *j0*: *j0 < N*
  **shows** *Matrix.vec N* (*pe-amp N* (*real j0 / real N*)) = *unit-vec N j0*
⟨*proof*⟩

## 110.5 Measuring the register

Probability of outcome $j$ when the register of $\psi \in \mathbb{C}^N \otimes \mathbb{C}^m$ is measured.

**definition** *reg-prob* :: *nat* ⇒ *complex Matrix.vec* ⇒ *nat* ⇒ *real* **where**
  *reg-prob m* $\psi$ *j* = $(\sum i<m.$ *(cmod* $(\psi$ $ *(j * m + i)))^2)$

**lemma** *reg-prob-tensor*:
  **assumes** *b*: *dim-vec b = N* **and** *u*: *dim-vec u = m* **and** *j*: *j < N*
    **and** *unit*: $(\sum i<m.$ *(cmod (u $ i))*$^2) = 1$
  **shows** *reg-prob m* (*tensor-vec b u*) *j* = *(cmod (b $ j))*$^2$
⟨*proof*⟩

## 110.6 The success probability

Jordan's inequality $\sin x \geq 2x/\pi$ on $[0, \pi/2]$, by concavity (as in *Gauss-Sums.Polya-Vinogradov*).

**lemma** *jordan-inequality*:
  **assumes** *0 ≤ x x ≤ pi / 2*
  **shows** *2 / pi * x ≤ sin x*
⟨*proof*⟩

**lemma** *jordan-inequality-abs*:
  **assumes** $|x| \leq$ *pi / 2*
  **shows** *2 / pi * |x| ≤ |sin x|*
⟨*proof*⟩

**lemma** *cmod-cis-minus-one*: *cmod (cis x − 1) = 2 * |sin (x / 2)|*
⟨*proof*⟩

The key estimate: $\left|\frac{1}{N}\sum_k e^{2\pi ik\delta}\right| \geq 2/\pi$ for $|\delta| \leq \frac{1}{2N}$.

**theorem** *geometric-amp-bound*:
  **assumes** *N*: *0 < N* **and** *d*: $|\delta| \leq$ *1 / (2 * real N)*
  **shows** *2 / pi ≤ cmod* $((\sum k<N.$ *cis (2 * pi * real k * δ)) / of-nat N*)
⟨*proof*⟩

With $\varphi$ within $\frac{1}{2N}$ of $j/N$, outcome $j$ has probability at least $4/\pi^2$.

**theorem** *phase-estimation-bound*:
  **assumes** *N*: *0 < N* **and** *close*: $|\varphi -$ *real j / real N*$| \leq$ *1 / (2 * real N)*
  **shows** *4 / pi*$^2 \leq$ *(cmod (pe-amp N* $\varphi$ *j))*$^2$
⟨*proof*⟩

The phase is periodic: $\alpha_0$ is unchanged by $\varphi \mapsto \varphi - 1$.

**lemma** *pe-amp-shift*: *pe-amp N φ 0 = pe-amp N (φ − 1) 0*
⟨*proof*⟩

**lemma** *scaled-close*:
  **assumes** *N*: *0 < N* **and** *x*: *|real N ∗ x| ≤ 1 / 2*
  **shows** *|x| ≤ 1 / (2 ∗ real N)*
⟨*proof*⟩

Phase estimation succeeds: for $0 \le \varphi < 1$ and a unit eigenvector, some outcome $j < N$ (the nearest to $N\varphi$, read mod $N$) is observed with probability at least $4/\pi^2$.

**theorem** *phase-estimation-success*:
  **fixes** *U* :: *complex Matrix.mat*
  **assumes** *N*: *0 < N* **and** *U*: *U ∈ carrier-mat m m* **and** *u*: *dim-vec u = m*
    **and** *unit*: *(∑ i<m. (cmod (u $ i))²) = 1*
    **and** *ev*: *U ∗v u = cis (2 ∗ pi ∗ φ) ·v u* **and** *φ*: *0 ≤ φ φ < 1*
  **shows** *∃ j<N. 4 / pi² ≤*
      *reg-prob m ((((fourier N)†) ⊗ 1m m) ∗v (ctrl-pow N m U ∗v tensor-vec (uniform-state N) u)) j*
⟨*proof*⟩

**end**
**theory** *QFT-Fourier*
  **imports** *Phase-Estimation*
**begin**

# 111 The AFP Fourier transform circuit is the Fourier matrix

The circuit *ordered-QFT* of the AFP entry *Quantum-Fourier-Transform* is shown there to map $|j\rangle$ to the product state $2^{-n/2}\bigotimes_{l=1}^{n}(|0\rangle + e^{2\pi i j/2^l}|1\rangle)$. Its $k$-th entry is $2^{-n/2}e^{2\pi i j k/2^n}$ (the binary digits of $k$ select the factors), so the circuit is $F_{2^n}$ and phase estimation runs on it.

**lemma** *kron-phase-index*:
  **fixes** *f* :: *nat ⇒ complex Matrix.mat* **and** *j* :: *nat*
  **assumes** *f*: *⋀l. f l ∈ carrier-mat 2 1*
    **and** *fi*: *⋀l c. c < 2 ⟹ f l $$ (c, 0) = exp (2 ∗ i ∗ pi ∗ of-nat j ∗ of-nat c / 2 ^ l)*
  **shows** *1 ≤ a ⟹ nat (b − a + 1) = m ⟹ k < 2 ^ m ⟹*
      *kron f (map nat [a..b]) $$ (k, 0) = exp (2 ∗ i ∗ pi ∗ of-nat j ∗ of-nat k / 2 ^ nat b)*
⟨*proof*⟩

**lemma** *QFT-product-representation-index*:
  **assumes** *k*: *k < 2 ^ n*
  **shows** *QFT-product-representation j n $$ (k, 0) =*
      *cis (2 ∗ pi ∗ real j ∗ real k / 2 ^ n) / complex-of-real (sqrt (2 ^ n))*
⟨*proof*⟩

**theorem** *ordered-QFT-fourier*: *ordered-QFT n = fourier (2 ^ n)*
⟨*proof*⟩

Phase estimation with the AFP circuit: the theorems of *Phase-Estimation* hold with $F_{2^t}$ replaced by *ordered-QFT t*.

**corollary** *phase-estimation-circuit*:
  **fixes** *U* :: *complex Matrix.mat*
  **assumes** *U ∈ carrier-mat m m dim-vec u = m U ∗v u = cis (2 ∗ pi ∗ φ) ·v u*
  **shows** *(((ordered-QFT t)†) ⊗ 1m m) ∗v (ctrl-pow (2 ^ t) m U ∗v tensor-vec (uniform-state (2 ^ t)) u) =*

*tensor-vec* (*Matrix.vec* (2 ^ *t*) (*pe-amp* (2 ^ *t*) *φ*)) *u*
⟨*proof*⟩

**end**
**theory** *HHL*
**imports** *Phase-Estimation*
**begin**

# 112 The HHL algorithm

Harrow, Hassidim and Lloyd's algorithm for $Ax = b$, $A$ Hermitian: phase estimation of $e^{it_0 A}$ writes the eigenvalue $\lambda$ of each eigencomponent into a register, a rotation controlled by the register puts $C/\lambda$ into the amplitude of an ancilla $|1\rangle$, and phase estimation is undone. Measuring the ancilla in $|1\rangle$ leaves the system in $A^{-1}b/\|A^{-1}b\|$. The space is register $\otimes$ (system $\otimes$ ancilla).

## 112.1 Block-diagonal controlled operations

$\sum_k |k\rangle\langle k| \otimes B_k$ on $\mathbb{C}^N \otimes \mathbb{C}^m$.

**definition** *ctrl-blocks* :: *nat* ⇒ *nat* ⇒ (*nat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* **where**
*ctrl-blocks N m B* = *Matrix.mat* (*N* ∗ *m*) (*N* ∗ *m*)
(λ(*r*, *c*). *if r div m* = *c div m then B* (*r div m*) $$ (*r mod m*, *c mod m*) *else 0*)

**lemma** *ctrl-blocks-carrier* [*simp*]: *ctrl-blocks N m B* ∈ *carrier-mat* (*N* ∗ *m*) (*N* ∗ *m*)
⟨*proof*⟩

**lemma** *dim-ctrl-blocks* [*simp*]: *dim-row* (*ctrl-blocks N m B*) = *N* ∗ *m dim-col* (*ctrl-blocks N m B*) = *N* ∗ *m*
⟨*proof*⟩

**lemma** *ctrl-blocks-index*:
*r* < *N* ∗ *m* ⟹ *c* < *N* ∗ *m* ⟹
*ctrl-blocks N m B* $$ (*r*, *c*) = (*if r div m* = *c div m then B* (*r div m*) $$ (*r mod m*, *c mod m*) *else 0*)
⟨*proof*⟩

**lemma** *ctrl-pow-blocks*: *ctrl-pow N m U* = *ctrl-blocks N m* (λ*k*. *U* ^*m* *k*)
⟨*proof*⟩

**lemma** *div-mod-eq-iff*:
**fixes** *r c m* :: *nat*
**shows** *r* = *c* ⟷ *r div m* = *c div m* ∧ *r mod m* = *c mod m*
⟨*proof*⟩

**lemma** *ctrl-blocks-cong*:
**assumes** ⋀*k*. *k* < *N* ⟹ *B k* = *B*′ *k*
**shows** *ctrl-blocks N m B* = *ctrl-blocks N m B*′
⟨*proof*⟩

**lemma** *ctrl-blocks-one*: *ctrl-blocks N m* (λ-. 1*m m*) = 1*m* (*N* ∗ *m*)
⟨*proof*⟩

**lemma** *sum-if-const*: (∑ *i*∈*A*. *if P then f i else 0*) = (*if P then* (∑ *i*∈*A*. *f i*) *else 0*)
⟨*proof*⟩

**lemma** *tensor-unit-index*:
**assumes** *k*′ < *N i*′ < *m dim-vec w* = *m*
**shows** *tensor-vec* (*unit-vec N k*) *w* $ (*k*′ ∗ *m* + *i*′) = (*if k*′ = *k then w* $ *i*′ *else 0*)

⟨*proof*⟩

**lemma** *ctrl-blocks-col-index*:
**assumes** $r < N * m$ $k' < N$ $i' < m$
**shows** *ctrl-blocks N m B* $$ $(r, k' * m + i') = (\textit{if } r \textit{ div } m = k' \textit{ then } B\ k'\ \$\$\ (r \textit{ mod } m, i') \textit{ else } 0)$
⟨*proof*⟩

**lemma** *ctrl-blocks-row-index*:
**assumes** $c < N * m$ $k' < N$ $i' < m$
**shows** *ctrl-blocks N m B* $$ $(k' * m + i', c) = (\textit{if } k' = c \textit{ div } m \textit{ then } B\ k'\ \$\$\ (i', c \textit{ mod } m) \textit{ else } 0)$
⟨*proof*⟩

On a basis state of the register, $\sum_k |k\rangle\langle k| \otimes B_k$ applies $B_k$.

**lemma** *ctrl-blocks-basis*:
**assumes** *k*: $k < N$ **and** *w*: *dim-vec w* $= m$ **and** *B*: $B\ k \in$ *carrier-mat m m*
**shows** *ctrl-blocks N m B* $*_v$ *tensor-vec* (*unit-vec N k*) *w* $=$ *tensor-vec* (*unit-vec N k*) ($B\ k *_v$ *w*)
⟨*proof*⟩

**lemma** *ctrl-blocks-dagger*:
**assumes** *B*: $\bigwedge k.\ k < N \Longrightarrow B\ k \in$ *carrier-mat m m*
**shows** (*ctrl-blocks N m B*)$^\dagger$ = *ctrl-blocks N m* ($\lambda k.$ ($B\ k$)$^\dagger$)
⟨*proof*⟩

**lemma** *ctrl-blocks-mult*:
**assumes** *B1*: $\bigwedge k.\ k < N \Longrightarrow$ *B1 k* $\in$ *carrier-mat m m* **and** *B2*: $\bigwedge k.\ k < N \Longrightarrow$ *B2 k* $\in$ *carrier-mat m m*
**shows** *ctrl-blocks N m B1* $*$ *ctrl-blocks N m B2* = *ctrl-blocks N m* ($\lambda k.$ *B1 k* $*$ *B2 k*)
⟨*proof*⟩

**lemma** *ctrl-blocks-unitary*:
**assumes** *B*: $\bigwedge k.\ k < N \Longrightarrow B\ k \in$ *carrier-mat m m* **and** *u*: $\bigwedge k.\ k < N \Longrightarrow$ *unitary* ($B\ k$)
**shows** *unitary* (*ctrl-blocks N m B*)
⟨*proof*⟩

**lemma** *unitary-pow*:
**assumes** *U*: $U \in$ *carrier-mat m m* **and** *u*: *unitary U*
**shows** *unitary* ($U\ \widehat{}_m\ k$)
⟨*proof*⟩

## 112.2 The circuit

The ancilla rotation $R(c)|0\rangle = \sqrt{1-c^2}|0\rangle + c|1\rangle$.

**definition** *rot* :: *real* $\Rightarrow$ *complex Matrix.mat* **where**
*rot c* = *Matrix.mat 2 2* ($\lambda(i, j).$ *if i* = *j then complex-of-real* (*sqrt* (*1* − $c^2$))
*else if i* = *1 then complex-of-real c else* − *complex-of-real c*)

**lemma** *rot-carrier* [*simp*]: *rot c* $\in$ *carrier-mat 2 2*
⟨*proof*⟩

**lemma** *dim-rot* [*simp*]: *dim-row* (*rot c*) = *2 dim-col* (*rot c*) = *2*
⟨*proof*⟩

**lemma** *rot-e0*: *rot c* $*_v$ *unit-vec 2 0* =
*complex-of-real* (*sqrt* (*1* − $c^2$)) $\cdot_v$ *unit-vec 2 0* + *complex-of-real c* $\cdot_v$ *unit-vec 2 1*
⟨*proof*⟩

**lemma** *rot-unitary*:

**assumes** *c*: $|c| \leq 1$
**shows** *unitary* (*rot c*)
⟨*proof*⟩

Register value $k$ encodes the eigenvalue $\lambda_k = 2\pi k/(t_0 N)$ for $k < N/2$ and $2\pi(k-N)/(t_0 N)$ otherwise (negative eigenvalues), since $e^{it_0\lambda_k} = e^{2\pi ik/N}$.

**definition** *reg-eig* :: *real* ⇒ *nat* ⇒ *nat* ⇒ *real* **where**
*reg-eig t0 N k* = *2* ∗ *pi* ∗ (*if 2* ∗ *k* < *N then real k else real k* − *real N*) / (*t0* ∗ *real N*)

**lemma** *reg-eig-phase*:
**assumes** *t0*: *0* < *t0* **and** *N*: *0* < *N*
**shows** *cis* (*t0* ∗ *reg-eig t0 N k*) = *cis* (*2* ∗ *pi* ∗ (*real k* / *real N*))
⟨*proof*⟩

**lemma** *reg-eig-nonzero*: *0* < *t0* ⟹ *0* < *k* ⟹ *k* < *N* ⟹ *reg-eig t0 N k* ≠ *0*
⟨*proof*⟩

**definition** *hhl-rot* :: *real* ⇒ *real* ⇒ *nat* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*hhl-rot C t0 N n k* = $1_m$ *n* ⨂ *rot* (*C* / *reg-eig t0 N k*)

**definition** *pe-unitary* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*pe-unitary N m V* = (((*fourier N*)†) ⨂ $1_m$ *m*) ∗ *ctrl-pow N m V*

**definition** *hhl-W* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*hhl-W A t0 N* = *pe-unitary N* (*dim-row A* ∗ *2*) (*evolution A t0* ⨂ $1_m$ *2*)

**definition** *hhl-circuit* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *real* ⇒ *complex Matrix.mat* **where**
*hhl-circuit A t0 N C* = (*hhl-W A t0 N*)† ∗ (*ctrl-blocks N* (*dim-row A* ∗ *2*) (*hhl-rot C t0 N* (*dim-row A*)) ∗ *hhl-W A t0 N*)

**lemma** *hhl-rot-carrier* [*simp*]: *hhl-rot C t0 N n k* ∈ *carrier-mat* (*n* ∗ *2*) (*n* ∗ *2*)
⟨*proof*⟩

**lemma** *pe-unitary-carrier* [*simp*]: *pe-unitary N m V* ∈ *carrier-mat* (*N* ∗ *m*) (*N* ∗ *m*)
⟨*proof*⟩

**lemma** *pe-unitary-unitary*:
**assumes** *N*: *0* < *N* **and** *V*: *V* ∈ *carrier-mat m m unitary V*
**shows** *unitary* (*pe-unitary N m V*)
⟨*proof*⟩

On $\frac{1}{\sqrt{N}}\sum_k |k\rangle \otimes w$ with $Vw = e^{2\pi ik/N}w$, phase estimation gives $|k\rangle \otimes w$.

**lemma** *pe-unitary-exact*:
**assumes** *N*: *0* < *N* **and** *k*: *k* < *N* **and** *V*: *V* ∈ *carrier-mat m m* **and** *w*: *dim-vec w* = *m*
**and** *ev*: *V* ∗$_v$ *w* = *cis* (*2* ∗ *pi* ∗ (*real k* / *real N*)) ·$_v$ *w*
**shows** *pe-unitary N m V* ∗$_v$ *tensor-vec* (*uniform-state N*) *w* = *tensor-vec* (*unit-vec N k*) *w*
⟨*proof*⟩

**lemma** *unitary-undo*:
**assumes** *W*: *W* ∈ *carrier-mat K K unitary W* **and** *x*: *dim-vec x* = *K* **and** *y*: *W* ∗$_v$ *x* = *y*
**shows** *W*† ∗$_v$ *y* = *x*
⟨*proof*⟩

## 112.3 The circuit on an eigenvector

**theorem** *hhl-eigen*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *t0*: *0* < *t0* **and** *N*: *0* < *N*
**and** *k*: *k* < *N* **and** *u*: *dim-vec u* = *n* **and** *ev*: *A* ∗$_v$ *u* = *complex-of-real* (*reg-eig t0 N k*) ·$_v$ *u*

**shows** *hhl-circuit A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) =
*tensor-vec* (*uniform-state N*) (*tensor-vec u* (*rot* (*C / reg-eig t0 N k*) $*_v$ *unit-vec 2 0*))
⟨*proof*⟩

## 112.4 All right-hand sides

**lemma** *mult-zero-vec-right*: $B \in$ *carrier-mat K n* $\Longrightarrow$ $B *_v 0_v\ n = 0_v\ K$
⟨*proof*⟩

Two matrices that agree on every eigenvector of a Hermitian $A$ are equal.

**lemma** *eigvec-mat-eqI*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A*
**and** *B1*: *B1* $\in$ *carrier-mat K n* **and** *B2*: *B2* $\in$ *carrier-mat K n*
**and** *eq*: $\bigwedge v\ l.$ *dim-vec v* $= n \Longrightarrow A *_v v =$ *complex-of-real l* $\cdot_v$ *v* $\Longrightarrow$ *B1* $*_v$ *v* $=$ *B2* $*_v$ *v*
**shows** *B1* $=$ *B2*
⟨*proof*⟩

The matrix of $b \mapsto a \otimes (M_0 b \otimes |0\rangle + M_1 b \otimes |1\rangle)$.

**definition** *lift-mat* :: *nat* $\Rightarrow$ *complex Matrix.vec* $\Rightarrow$ *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat* $\Rightarrow$ *complex Matrix.mat* **where**
*lift-mat n a M0 M1* = *Matrix.mat* (*dim-vec a* $*$ (*n* $*$ *2*)) *n*
(λ(*r*, *s*). *a* $ (*r div* (*n* $*$ *2*)) $*$
(*if r mod 2 = 0 then M0* $$ (*r mod* (*n* $*$ *2*) *div 2*, *s*) *else M1* $$ (*r mod* (*n* $*$ *2*) *div 2*, *s*)))

**lemma** *lift-mat-carrier* [*simp*]: *lift-mat n a M0 M1* $\in$ *carrier-mat* (*dim-vec a* $*$ (*n* $*$ *2*)) *n*
⟨*proof*⟩

**lemma** *lift-mat-mult-vec*:
**assumes** *M0*: *M0* $\in$ *carrier-mat n n* **and** *M1*: *M1* $\in$ *carrier-mat n n* **and** *b*: *dim-vec b* $= n$
**shows** *lift-mat n a M0 M1* $*_v$ *b* =
*tensor-vec a* (*tensor-vec* (*M0* $*_v$ *b*) (*unit-vec 2 0*) + *tensor-vec* (*M1* $*_v$ *b*) (*unit-vec 2 1*))
⟨*proof*⟩

**theorem** *hhl*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *t0*: $0 <$ *t0* **and** *N*: $0 < N$
**and** *spec*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow \exists k.\ 0 < k \wedge k < N \wedge l =$ *reg-eig t0 N k*
**and** *b*: *dim-vec b* $= n$
**shows** *hhl-circuit A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec b* (*unit-vec 2 0*)) =
*tensor-vec* (*uniform-state N*)
(*tensor-vec* (*mat-fun* (λ*x*. *complex-of-real* (*sqrt* (*1* − (*C / x*)$^2$))) *A* $*_v$ *b*) (*unit-vec 2 0*) +
*tensor-vec* ((*complex-of-real C* $\cdot_m$ *herm-inv A*) $*_v$ *b*) (*unit-vec 2 1*))
⟨*proof*⟩

Reading off the solution: $x = A^{-1}b$ solves $Ax = b$, the ancilla-$|1\rangle$ part of the output is $C\,(u \otimes x \otimes |1\rangle)$, and its squared norm (the probability of measuring the ancilla in $|1\rangle$) is $C^2\|x\|^2$.

**lemma** *uniform-state-norm*: $0 <$ *N* $\Longrightarrow$ ⟨*uniform-state N*|*uniform-state N*⟩ $= 1$
⟨*proof*⟩

**corollary** *hhl-solution*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *t0*: $0 <$ *t0* **and** *N*: $0 < N$
**and** *spec*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow \exists k.\ 0 < k \wedge k < N \wedge l =$ *reg-eig t0 N k*
**and** *b*: *dim-vec b* $= n$
**defines** $x \equiv$ *herm-inv A* $*_v$ *b*
**shows** *A* $*_v$ *x* $=$ *b*
**and** *hhl-circuit A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec b* (*unit-vec 2 0*)) =
*tensor-vec* (*uniform-state N*) (*tensor-vec* (*mat-fun* (λ*x*. *complex-of-real* (*sqrt* (*1* − (*C / x*)$^2$))) *A*
$*_v$ *b*) (*unit-vec 2 0*)) +

*complex-of-real C ·$_v$ tensor-vec (uniform-state N) (tensor-vec x (unit-vec 2 1))*
**and** ⟨*tensor-vec (uniform-state N) (tensor-vec (complex-of-real C ·$_v$ x) (unit-vec 2 1)) |*
*tensor-vec (uniform-state N) (tensor-vec (complex-of-real C ·$_v$ x) (unit-vec 2 1))*⟩ =
*complex-of-real* ($C^2$) ∗ ⟨*x*|*x*⟩
⟨*proof*⟩

## 112.5 The success probability

The diagonal of a spectral decomposition consists of eigenvalues.

**lemma** *spectral-decomp-eig-in-spectrum*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *sd*: *spectral-decomp A U D* **and** *j*: *j* < *n*
**shows** *D* ! *j* ∈ *herm-spectrum A*
⟨*proof*⟩

A real function non-negative on the spectrum gives a positive matrix.

**lemma** *mat-fun-psd*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *f*: ⋀*l*. *l* ∈ *herm-spectrum A* ⟹ *0* ≤ *f l*
**shows** *psd* (*mat-fun* (λ*x*. *complex-of-real* (*f x*)) *A*)
⟨*proof*⟩

**lemma** *mat-fun-add*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A*
**shows** *mat-fun* (λ*x*. *g x* + *h x*) *A* = *mat-fun g A* + *mat-fun h A*
⟨*proof*⟩

**lemma** *inner-prod-add-right*:
**assumes** *dim-vec a* = *n dim-vec b* = *n dim-vec c* = *n*
**shows** ⟨*a*|*b* + *c*⟩ = ⟨*a*|*b*⟩ + ⟨*a*|*c*⟩
⟨*proof*⟩

If $C \le |\lambda| \le \Lambda$ on the spectrum, $C > 0$, the probability $C^2\|A^{-1}b\|^2$ of the ancilla outcome $|1\rangle$ is at least $(C/\Lambda)^2\|b\|^2$: for unit $b$ and $C = \lambda_{\min}$ this is $1/\kappa^2$.

**theorem** *hhl-success-bound*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *C*: *0* < *C*
**and** *spec*: ⋀*l*. *l* ∈ *herm-spectrum A* ⟹ *C* ≤ |*l*| ∧ |*l*| ≤ Λ
**and** *b*: *dim-vec b* = *n*
**shows** (*C* / Λ)$^2$ ∗ *Re* ⟨*b*|*b*⟩ ≤ $C^2$ ∗ *Re* ⟨*herm-inv A* ∗$_v$ *b*|*herm-inv A* ∗$_v$ *b*⟩
⟨*proof*⟩

## 112.6 Non-Hermitian systems

For square $A$, the Hermitian $\tilde{A} = \begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix}$ has $\tilde{A}(0 \oplus x) = b \oplus 0$ when $Ax = b$, so HHL applied to $\tilde{A}$ and $b \oplus 0$ returns $0 \oplus x$.

**definition** *herm-embed* :: *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*herm-embed A* = *four-block-mat* (*0$_m$* (*dim-row A*) (*dim-row A*)) *A* (*A*$^\dagger$) (*0$_m$* (*dim-col A*) (*dim-col A*))

**lemma** *herm-embed-carrier*: *A* ∈ *carrier-mat n n* ⟹ *herm-embed A* ∈ *carrier-mat* (*n* + *n*) (*n* + *n*)
⟨*proof*⟩

**lemma** *zero-dagger*: (*0$_m$ n m* :: *complex Matrix.mat*)$^\dagger$ = *0$_m$ m n*
⟨*proof*⟩

**lemma** *herm-embed-hermitian*:
**assumes** *A*: *A* ∈ *carrier-mat n n*
**shows** *hermitian* (*herm-embed A*)

⟨*proof*⟩

**lemma** *zero-mat-mult-vec*: *dim-vec v* = *n* ⟹ $0_m$ *K n* $*_v$ *v* = $0_v$ *K*
⟨*proof*⟩

**lemma** *herm-embed-solve*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *x*: *dim-vec x* = *n* **and** *ax*: *A* $*_v$ *x* = *b*
**shows** *herm-embed A* $*_v$ (*0*$_v$ *n* @$_v$ *x*) = *b* @$_v$ *0*$_v$ *n*
⟨*proof*⟩

**lemma** *herm-embed-inverse*:
**assumes** *A*: *A* ∈ *carrier-mat n n* **and** *x*: *dim-vec x* = *n* **and** *ax*: *A* $*_v$ *x* = *b*
**and** *z*: *0* ∉ *herm-spectrum* (*herm-embed A*)
**shows** *herm-inv* (*herm-embed A*) $*_v$ (*b* @$_v$ *0*$_v$ *n*) = *0*$_v$ *n* @$_v$ *x*
⟨*proof*⟩

**end**
**theory** *HHL-Approximate*
**imports** *HHL*
**begin**

# 113 HHL with inexact phase estimation

When the eigenvalue $\lambda$ of $u$ is not one of the register values $\lambda_k$, phase estimation leaves the register in $\sum_j \alpha_j|j\rangle$ with $\alpha_j$ =*pe-amp*, the rotation acts with $C/\lambda_j$ on the $j$-th branch, and uncomputation no longer returns the register to the uniform state. The output is compared with the ideal output $u \otimes R(C/\lambda)|0\rangle$ of the exact case; the squared distance is $\sum_j |\alpha_j|^2 \, \| R(C/\lambda_j)|0\rangle - R(C/\lambda)|0\rangle \|^2$.

## 113.1 Register sums

$\sum_{j<N} |j\rangle \otimes f_j$ on $\mathbb{C}^N \otimes \mathbb{C}^m$.

**definition** *reg-sum* :: *nat* ⇒ *nat* ⇒ (*nat* ⇒ *complex Matrix.vec*) ⇒ *complex Matrix.vec* **where**
*reg-sum N m f* = *Matrix.vec* (*N* ∗ *m*) (λ*r*. *f* (*r div m*) $ (*r mod m*))

**lemma** *dim-reg-sum* [*simp*]: *dim-vec* (*reg-sum N m f*) = *N* ∗ *m*
⟨*proof*⟩

**lemma** *reg-sum-index*:
*k* < *N* ⟹ *i* < *m* ⟹ *reg-sum N m f* $ (*k* ∗ *m* + *i*) = *f k* $ *i*
⟨*proof*⟩

**lemma** *reg-sum-cong*: (⋀*k*. *k* < *N* ⟹ *f k* = *g k*) ⟹ *reg-sum N m f* = *reg-sum N m g*
⟨*proof*⟩

**lemma** *tensor-reg-sum*:
**assumes** *a*: *dim-vec a* = *N* **and** *w*: *dim-vec w* = *m*
**shows** *tensor-vec a w* = *reg-sum N m* (λ*k*. *a* $ *k* ·$_v$ *w*)
⟨*proof*⟩

**lemma** *reg-sum-index-div*:
**fixes** *r N m* :: *nat*
**assumes** *r* < *N* ∗ *m*
**shows** *r div m* < *N r mod m* < *m*
⟨*proof*⟩

**lemma** *reg-sum-add*:
**assumes** *f*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(f\ k) = m$ **and** *g*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(g\ k) = m$
**shows** *reg-sum* $N\ m\ f$ + *reg-sum* $N\ m\ g$ = *reg-sum* $N\ m\ (\lambda k.\ f\ k + g\ k)$
⟨*proof*⟩

**lemma** *reg-sum-diff*:
**assumes** *g*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(g\ k) = m$
**shows** *reg-sum* $N\ m\ f$ − *reg-sum* $N\ m\ g$ = *reg-sum* $N\ m\ (\lambda k.\ f\ k - g\ k)$
⟨*proof*⟩

**lemma** *reg-sum-smult*:
**assumes** *f*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(f\ k) = m$
**shows** $c \cdot_v$ *reg-sum* $N\ m\ f$ = *reg-sum* $N\ m\ (\lambda k.\ c \cdot_v f\ k)$
⟨*proof*⟩

Block-diagonal operators act branch by branch.

**lemma** *ctrl-blocks-reg-sum*:
**assumes** *B*: $\bigwedge k.\ k < N \Longrightarrow B\ k \in$ *carrier-mat* $m\ m$ **and** *f*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(f\ k) = m$
**shows** *ctrl-blocks* $N\ m\ B *_v$ *reg-sum* $N\ m\ f$ = *reg-sum* $N\ m\ (\lambda k.\ B\ k *_v f\ k)$
⟨*proof*⟩

Branches are orthogonal: $\langle \sum_j |j\rangle \otimes f_j, \sum_j |j\rangle \otimes g_j\rangle = \sum_j \langle f_j, g_j\rangle$.

**lemma** *reg-sum-inner*:
**assumes** *f*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(f\ k) = m$ **and** *g*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* $(g\ k) = m$
**shows** ⟨*reg-sum* $N\ m\ f$|*reg-sum* $N\ m\ g$⟩ = $(\sum k{<}N.\ \langle f\ k|g\ k\rangle)$
⟨*proof*⟩

## 113.2 Phase estimation on a branch

$W(u_N \otimes u \otimes w) = \alpha \otimes u \otimes w$ for every ancilla vector $w$, where $\alpha_j = \alpha_j(\varphi)$, $e^{it_0\lambda} = e^{2\pi i\varphi}$.

**lemma** *unitary-inner-self*:
**assumes** *W*: *W* ∈ *carrier-mat K K unitary W* **and** *y*: *dim-vec y* = *K*
**shows** ⟨$W^\dagger *_v$ *y*|$W^\dagger *_v$ *y*⟩ = ⟨*y*|*y*⟩
⟨*proof*⟩

**lemma** *rot-e0-vec*:
*rot c* $*_v$ *unit-vec 2 0* = *Matrix.vec 2* $(\lambda i.$ *if i* = *0 then complex-of-real* (*sqrt* (*1* − $c^2$)) *else complex-of-real c*)
⟨*proof*⟩

**lemma** *rot-diff-inner*:
⟨*rot c1* $*_v$ *unit-vec 2 0* − *rot c2* $*_v$ *unit-vec 2 0*|*rot c1* $*_v$ *unit-vec 2 0* − *rot c2* $*_v$ *unit-vec 2 0*⟩ =
*complex-of-real* ((*sqrt* (*1* − *c1*$^2$) − *sqrt* (*1* − *c2*$^2$))$^2$ + (*c1* − *c2*)$^2$)
⟨*proof*⟩

**lemma** *cnj-mult-self*: *cnj z* ∗ *z* = *complex-of-real* ((*cmod z*)$^2$)
⟨*proof*⟩

**lemma** *inner-self-sum*: *dim-vec v* = *N* ⟹ ⟨*v*|*v*⟩ = $(\sum i{<}N.$ *complex-of-real* ((*cmod* (*v* $ *i*))$^2$))
⟨*proof*⟩

The register amplitudes have total weight one: $\alpha = F_N^* b$ with $b$ a unit vector.

**lemma** *pe-amp-sum-sq*:
**assumes** *N*: *0* < *N*
**shows** $(\sum j{<}N.$ (*cmod* (*pe-amp N* $\varphi$ *j*))$^2$) = *1*
⟨*proof*⟩

Tail bound: $|\alpha_j| \leq 1/(2N|\delta|)$ for $\delta = \varphi - j/N$, $0 < |\delta| \leq \frac{1}{2}$.

**lemma** *geometric-amp-tail*:
**assumes** *N*: *0 < N* **and** *d*: $|\delta| \leq 1 / 2$ **and** *d0*: $\delta \neq 0$
**shows** *cmod* $((\sum k{<}N.\ cis\ (2 * pi * real\ k * \delta)) / of\text{-}nat\ N) \leq 1 / (2 * real\ N * |\delta|)$
⟨*proof*⟩

**lemma** *pe-amp-tail*:
**assumes** *N*: *0 < N* **and** *d*: $|\varphi - real\ j / real\ N| \leq 1 / 2$ **and** *d0*: $\varphi \neq real\ j / real\ N$
**shows** *cmod* (*pe-amp N φ j*) $\leq 1 / (2 * real\ N * |\varphi - real\ j / real\ N|)$
⟨*proof*⟩

The circuit with an arbitrary angle $f(k)$ on register value $k$; *hhl-circuit* is $f(k) = C/\lambda_k$, the physical circuit clamps the angle to $[-1, 1]$.

**definition** *hhl-gen* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ (*nat* ⇒ *real*) ⇒ *complex Matrix.mat* **where**
*hhl-gen A t0 N f* = (*hhl-W A t0 N*)$^\dagger$ ∗ (*ctrl-blocks N* (*dim-row A* ∗ *2*) (λ*k*. $1_m$ (*dim-row A*) ⨂ *rot* (*f k*)) ∗ *hhl-W A t0 N*)

**definition** *clamp* :: *real* ⇒ *real* **where**
*clamp x* = *max* (−*1*) (*min 1 x*)

**definition** *hhl-circuit-clamped* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *real* ⇒ *complex Matrix.mat* **where**
*hhl-circuit-clamped A t0 N C* = *hhl-gen A t0 N* (λ*k*. *clamp* (*C / reg-eig t0 N k*))

**lemma** *hhl-circuit-gen*: *hhl-circuit A t0 N C* = *hhl-gen A t0 N* (λ*k*. *C / reg-eig t0 N k*)
⟨*proof*⟩

**lemma** *rot-blk-carrier* [*simp*]: $1_m$ *n* ⨂ *rot x* ∈ *carrier-mat* (*n* ∗ *2*) (*n* ∗ *2*)
⟨*proof*⟩

**lemma** *clamp-abs*: |*clamp x*| ≤ *1*
⟨*proof*⟩

**lemma** *clamp-id*: |*x*| ≤ *1* ⟹ *clamp x* = *x*
⟨*proof*⟩

**definition** *hhl-phi* :: *real* ⇒ *real* ⇒ *real* **where**
*hhl-phi t0 l* = *t0* ∗ *l* / (*2* ∗ *pi*)

**context**
**fixes** *A* :: *complex Matrix.mat* **and** *n N* :: *nat* **and** *t0 l* :: *real* **and** *u* :: *complex Matrix.vec*
**and** *f* :: *nat* ⇒ *real* **and** *c* :: *real*
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *t0*: *0 < t0* **and** *N*: *0 < N*
**and** *u*: *dim-vec u* = *n* **and** *ev*: *A* ∗$_v$ *u* = *complex-of-real l* ⋅$_v$ *u*
**begin**

The phase of $u$, the register amplitudes, the ideal and the branch rotations.

**abbreviation** *phi* ≡ *hhl-phi t0 l*
**definition** *alpha* = *Matrix.vec N* (*pe-amp N phi*)
**abbreviation** *r-ideal* ≡ *rot c* ∗$_v$ *unit-vec 2 0*
**abbreviation** *r-br k* ≡ *rot* (*f k*) ∗$_v$ *unit-vec 2 0*

**lemma** *phi-phase*: *cis* (*t0* ∗ *l*) = *cis* (*2* ∗ *pi* ∗ *phi*)
⟨*proof*⟩

**lemma** *dim-alpha* [*simp*]: *dim-vec alpha* = *N* ⟨*proof*⟩
**lemma** *dim-r-ideal*: *dim-vec r-ideal* = *2* ⟨*proof*⟩

**lemma** *W-branch*:
**assumes** *w*: *dim-vec w = 2*
**shows** *hhl-W A t0 N* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u w*) = *tensor-vec alpha* (*tensor-vec u w*)
⟨*proof*⟩

**lemma** *hhl-W-carrier*: *hhl-W A t0 N* ∈ *carrier-mat* (*N* ∗ (*n* ∗ *2*)) (*N* ∗ (*n* ∗ *2*))
⟨*proof*⟩

**lemma** *hhl-W-unitary*: *unitary* (*hhl-W A t0 N*)
⟨*proof*⟩

The output of the circuit and the ideal output as register sums.

**lemma** *hhl-output-branches*:
*hhl-gen A t0 N f* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) =
(*hhl-W A t0 N*)$^\dagger$ $*_v$ *reg-sum N* (*n* ∗ *2*) (λ*k*. *alpha* $ *k* ·$_v$ *tensor-vec u* (*r-br k*))
⟨*proof*⟩

**lemma** *hhl-ideal-branches*:
*tensor-vec* (*uniform-state N*) (*tensor-vec u r-ideal*) =
(*hhl-W A t0 N*)$^\dagger$ $*_v$ *reg-sum N* (*n* ∗ *2*) (λ*k*. *alpha* $ *k* ·$_v$ *tensor-vec u r-ideal*)
⟨*proof*⟩

The squared distance between output and ideal output.

The error vector: $W^* \sum_k \alpha_k |k\rangle \otimes u \otimes (R(f_k)|0\rangle - R(c)|0\rangle)$.

**lemma** *hhl-err-vec*:
*hhl-gen A t0 N f* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u r-ideal*) =
(*hhl-W A t0 N*)$^\dagger$ $*_v$ *reg-sum N* (*n* ∗ *2*) (λ*k*. *alpha* $ *k* ·$_v$ *tensor-vec u* (*r-br k* − *r-ideal*))
⟨*proof*⟩

The squared distance between output and ideal output.

**theorem** *hhl-approx-error*:
⟨*hhl-gen A t0 N f* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u r-ideal*) |
*hhl-gen A t0 N f* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u r-ideal*)⟩ =
⟨*u*|*u*⟩ ∗ (∑ *k*<*N*. *complex-of-real* ((*cmod* (*pe-amp N phi k*))$^2$) ∗ ⟨*r-br k* − *r-ideal*|*r-br k* − *r-ideal*⟩)
⟨*proof*⟩

The branch distance in real terms.

**corollary** *hhl-approx-error-real*:
⟨*hhl-gen A t0 N f* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u r-ideal*) |
*hhl-gen A t0 N f* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u r-ideal*)⟩ =
⟨*u*|*u*⟩ ∗ *complex-of-real* (∑ *k*<*N*. (*cmod* (*pe-amp N phi k*))$^2$ ∗
((*sqrt* (*1* − (*f k*)$^2$) − *sqrt* (*1* − *c*$^2$))$^2$ + (*f k* − *c*)$^2$))
⟨*proof*⟩

**end**

## 113.3 Error vectors of distinct eigenvectors are orthogonal

**lemma** *unitary-inner-pres*:

**assumes** *W*: *W* ∈ *carrier-mat K K unitary W* **and** *x*: *dim-vec x* = *K* **and** *y*: *dim-vec y* = *K*
**shows** ⟨ *W*† ∗*v x*| *W*† ∗*v y*⟩ = ⟨*x*|*y*⟩
⟨*proof*⟩

For eigenvectors $u \perp u'$ the error vectors are orthogonal, whatever the angles and ideal rotations: each lives in $\mathbb{C}^N \otimes \mathrm{span}(u) \otimes \mathbb{C}^2$.

**lemma** *hhl-err-orth*:
**fixes** *A* :: *complex Matrix.mat* **and** *u u'* :: *complex Matrix.vec*
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *t0*: *0* < *t0* **and** *N*: *0* < *N*
**and** *u*: *dim-vec u* = *n* **and** *ev*: *A* ∗*v u* = *complex-of-real l* ·*v u*
**and** *u'*: *dim-vec u'* = *n* **and** *ev'*: *A* ∗*v u'* = *complex-of-real l'* ·*v u'*
**and** *orth*: ⟨*u*|*u'*⟩ = *0*
**shows** ⟨*hhl-gen A t0 N f* ∗*v tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*rot c* ∗*v unit-vec 2 0*)) |
*hhl-gen A t0 N f'* ∗*v tensor-vec* (*uniform-state N*) (*tensor-vec u'* (*unit-vec 2 0*)) − *tensor-vec* (*uniform-state N*) (*tensor-vec u'* (*rot c'* ∗*v unit-vec 2 0*))⟩ = *0*
⟨*proof*⟩

## 113.4 Bounds on the branch distances and amplitudes

$\|R(a)|0\rangle - R(b)|0\rangle\|^2 \le 4$ for $|a|, |b| \le 1$.

**lemma** *rot-dist-le-four*:
**assumes** |*a*| ≤ *1* |*b*| ≤ *1*
**shows** (*sqrt* (*1* − *a*²) − *sqrt* (*1* − *b*²))² + (*a* − *b*)² ≤ *4*
⟨*proof*⟩

Lipschitz estimate for $0 \le a, b \le \theta < 1$: $\|R(a)|0\rangle - R(b)|0\rangle\|^2 \le (a-b)^2/(1-\theta^2)$.

**lemma** *rot-dist-lipschitz*:
**assumes** *th*: *0* < *ϑ ϑ* < *1* **and** *a*: *0* ≤ *a a* ≤ *ϑ* **and** *b*: *0* ≤ *b b* ≤ *ϑ*
**shows** (*sqrt* (*1* − *a*²) − *sqrt* (*1* − *b*²))² + (*a* − *b*)² ≤ (*a* − *b*)² / (*1* − *ϑ*²)
⟨*proof*⟩

The amplitudes are 1-periodic in $\varphi$.

**lemma** *pe-amp-shift-up*: *pe-amp N φ j* = *pe-amp N* (*φ* + *1*) *j*
⟨*proof*⟩

## 113.5 The error bound

**lemma** *pe-amp-tail-sq*:
**assumes** *N*: *0* < *N* **and** *d*: |*φ* − *real j* / *real N*| ≤ *1* / *2* **and** *d0*: *φ* ≠ *real j* / *real N*
**shows** (*cmod* (*pe-amp N φ j*))² ≤ *1* / (*4* ∗ (*real N*)² ∗ (*φ* − *real j* / *real N*)²)
⟨*proof*⟩

Standing hypotheses of the bound: $0 < C$, $0 < \theta < 1$, $\lambda \ge 2C/\theta$ (the eigenvalue is at least twice the threshold $\lambda_{\mathrm{lo}} = C/\theta$ below which the angle is clamped) and $t_0\lambda \le \pi/2$ (the phase $\varphi = t_0\lambda/2\pi$ is at most $\frac{1}{4}$).

**lemma** *hhl-param-facts*:
**assumes** *t0*: *0* < *t0* **and** *C*: *0* < *C* **and** *th*: *0* < *ϑ ϑ* < *1*
**and** *l-lo*: *2* ∗ *C* / *ϑ* ≤ *l* **and** *l-hi*: *t0* ∗ *l* ≤ *pi* / *2*
**shows** *0* < *l 0* < *hhl-phi t0 l hhl-phi t0 l* ≤ *1* / *4 C* / *l* ≤ *ϑ* / *2 0* ≤ *C* / *l* |*C* / *l*| ≤ *1*
*2* ∗ *C* ≤ *ϑ* ∗ *l*
⟨*proof*⟩

Good branches: register value $\lambda_k \ge C/\theta$ in the positive half.

**lemma** *hhl-branch-good*:
**assumes** *N*: *0* < *N* **and** *t0*: *0* < *t0* **and** *C*: *0* < *C* **and** *th*: *0* < *ϑ ϑ* < *1*

**and** *l-lo*: $2 * C \,/\, \vartheta \le l$ **and** *l-hi*: $t0 * l \le pi \,/\, 2$
**and** *k*: $2 * k < N$ **and** *lam*: $C \,/\, \vartheta \le$ *reg-eig t0 N k*
**shows** $(cmod\ (pe\text{-}amp\ N\ (hhl\text{-}phi\ t0\ l)\ k))^2 *$
$((sqrt\ (1 - (clamp\ (C \,/\, reg\text{-}eig\ t0\ N\ k))^2) - sqrt\ (1 - (C \,/\, l)^2))^2 + (clamp\ (C \,/\, reg\text{-}eig\ t0\ N\ k) - C \,/\, l)^2)$
$\le pi^2 * \vartheta\ \hat{}\ 4 \,/\, ((real\ N)^2 * t0^2 * C^2 * (1 - \vartheta^2))$
⟨*proof*⟩

Branches below the threshold in the positive half: weight at most $1/(N\varphi)^2$.

**lemma** *hhl-branch-low*:
**assumes** *N*: $0 < N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C \,/\, \vartheta \le l$ **and** *l-hi*: $t0 * l \le pi \,/\, 2$
**and** *k*: $2 * k < N$ **and** *lam*: *reg-eig t0 N k* $< C \,/\, \vartheta$
**shows** $(cmod\ (pe\text{-}amp\ N\ (hhl\text{-}phi\ t0\ l)\ k))^2 \le 1 \,/\, ((real\ N)^2 * (hhl\text{-}phi\ t0\ l)^2)$
⟨*proof*⟩

Branches in the negative half: weight at most $1/(N\varphi)^2$.

**lemma** *hhl-branch-neg*:
**assumes** *N*: $0 < N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C \,/\, \vartheta \le l$ **and** *l-hi*: $t0 * l \le pi \,/\, 2$
**and** *k*: $k < N$ **and** *k2*: $\neg\ 2 * k < N$
**shows** $(cmod\ (pe\text{-}amp\ N\ (hhl\text{-}phi\ t0\ l)\ k))^2 \le 1 \,/\, ((real\ N)^2 * (hhl\text{-}phi\ t0\ l)^2)$
⟨*proof*⟩

Every branch: $|\alpha_k|^2 d_k^2 \le b_1 + b_2$ with $b_1 = \pi^2\theta^4/(N^2 t_0^2 C^2(1-\theta^2))$, $b_2 = 4/(N\varphi)^2$.

**lemma** *hhl-branch-bound*:
**assumes** *N*: $0 < N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C \,/\, \vartheta \le l$ **and** *l-hi*: $t0 * l \le pi \,/\, 2$ **and** *k*: $k < N$
**shows** $(cmod\ (pe\text{-}amp\ N\ (hhl\text{-}phi\ t0\ l)\ k))^2 *$
$((sqrt\ (1 - (clamp\ (C \,/\, reg\text{-}eig\ t0\ N\ k))^2) - sqrt\ (1 - (C \,/\, l)^2))^2 + (clamp\ (C \,/\, reg\text{-}eig\ t0\ N\ k) - C \,/\, l)^2)$
$\le pi^2 * \vartheta\ \hat{}\ 4 \,/\, ((real\ N)^2 * t0^2 * C^2 * (1 - \vartheta^2)) + 4 \,/\, ((real\ N)^2 * (hhl\text{-}phi\ t0\ l)^2)$
⟨*proof*⟩

The error bound for the clamped circuit on an eigenvector: with $\varphi = t_0\lambda/2\pi$,

$$\|\text{output} - u_N \otimes u \otimes R(C/\lambda)|0\rangle\|^2 \le \|u\|^2\, \frac{\pi^2}{N t_0^2}\Big(\frac{\theta^4}{C^2(1-\theta^2)} + \frac{16}{\lambda^2}\Big).$$

**lemma** *hhl-bound-algebra*:
**assumes** *N*: $0 < N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$ **and** *l0*: $0 < l$
**shows** $real\ N * (pi^2 * \vartheta\ \hat{}\ 4 \,/\, ((real\ N)^2 * t0^2 * C^2 * (1 - \vartheta^2)) + 4 \,/\, ((real\ N)^2 * (hhl\text{-}phi\ t0\ l)^2))$
$=$
$pi^2 \,/\, (real\ N * t0^2) * (\vartheta\ \hat{}\ 4 \,/\, (C^2 * (1 - \vartheta^2)) + 16 \,/\, l^2)$
⟨*proof*⟩

**theorem** *hhl-clamped-error-bound*:
**fixes** *A* :: *complex Matrix.mat* **and** *u* :: *complex Matrix.vec*
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *t0*: $0 < t0$ **and** *N*: $0 < N$
**and** *u*: *dim-vec u* $= n$ **and** *ev*: $A *_v u =$ *complex-of-real l* $\cdot_v u$
**and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C \,/\, \vartheta \le l$ **and** *l-hi*: $t0 * l \le pi \,/\, 2$
**shows** *Re* ⟨*hhl-circuit-clamped A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*))
$-$
*tensor-vec* (*uniform-state N*) (*tensor-vec u* (*rot* $(C \,/\, l) *_v$ *unit-vec 2 0*)) |
*hhl-circuit-clamped A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec u* (*unit-vec 2 0*)) $-$
*tensor-vec* (*uniform-state N*) (*tensor-vec u* (*rot* $(C \,/\, l) *_v$ *unit-vec 2 0*))⟩
$\le$ *Re* $\langle u|u\rangle * (pi^2 \,/\, (real\ N * t0^2) * (\vartheta\ \hat{}\ 4 \,/\, (C^2 * (1 - \vartheta^2)) + 16 \,/\, l^2))$
⟨*proof*⟩

## 113.6 From eigenvectors to every right-hand side

**lemma** *dagger-mult-entry*:
**assumes** $P$: $P \in$ *carrier-mat K m* **and** $i$: $i < m$ **and** $j$: $j < m$
**shows** $(P^\dagger * P)$ \$\$ $(i, j) = \langle$*Matrix.col P i*$|$*Matrix.col P j*$\rangle$
⟨*proof*⟩

**lemma** *inner-prod-minus-right*:
**assumes** *dim-vec a = n dim-vec x = n dim-vec y = n*
**shows** $\langle a|x - y\rangle = \langle a|x\rangle - \langle a|y\rangle$
⟨*proof*⟩

If Err maps the orthonormal columns of $U$ to mutually orthogonal vectors of squared norm at most $e$, then $\|\mathrm{Err}\, b\|^2 \le e\|b\|^2$ for every $b$: $U^*\mathrm{Err}^*\mathrm{Err}\, U$ is diagonal, so $eI - \mathrm{Err}^*\mathrm{Err}$ is positive.

**lemma** *orth-image-bound*:
**fixes** *Err U* :: *complex Matrix.mat*
**assumes** *U*: *U* ∈ *carrier-mat n n unitary U* **and** *E*: *Err* ∈ *carrier-mat K n*
**and** *diag*: ⋀*i j*. *i* < *n* ⟹ *j* < *n* ⟹
⟨*Err* $*_v$ *Matrix.col U i*|*Err* $*_v$ *Matrix.col U j*⟩ = (*if i = j then complex-of-real* (*m i*) *else 0*)
**and** *me*: ⋀*i*. *i* < *n* ⟹ *m i* ≤ *e* **and** *b*: *dim-vec b = n*
**shows** *Re* ⟨*Err* $*_v$ *b*|*Err* $*_v$ *b*⟩ ≤ *e* * *Re* ⟨*b*|*b*⟩
⟨*proof*⟩

**lemma** *inner-self-real*: *dim-vec v = n* ⟹ ⟨*v*|*v*⟩ = *complex-of-real* (*Re* ⟨*v*|*v*⟩)
⟨*proof*⟩

The embedding $b \mapsto u_N \otimes b \otimes |0\rangle$ and the ideal output as matrices.

**lemma** *lift-mat-emb*:
**assumes** *v*: *dim-vec v = n*
**shows** *lift-mat n* (*uniform-state N*) (1$_m$ *n*) (0$_m$ *n n*) $*_v$ *v = tensor-vec* (*uniform-state N*) (*tensor-vec v* (*unit-vec 2 0*))
⟨*proof*⟩

**lemma** *ideal-mat-eigen*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *u*: *dim-vec u = n* **and** *ev*: *A* $*_v$ *u = complex-of-real l* $\cdot_v$ *u*
**shows** *lift-mat n* (*uniform-state N*) (*mat-fun* (λ*x*. *complex-of-real* (*sqrt* (1 − (*C* / *x*)$^2$))) *A*) (*complex-of-real C* $\cdot_m$ *herm-inv A*) $*_v$ *u* =
*tensor-vec* (*uniform-state N*) (*tensor-vec u* (*rot* (*C* / *l*) $*_v$ *unit-vec 2 0*))
⟨*proof*⟩

The error bound for every right-hand side. The ideal output is the exact-HHL output (theorem *hhl*); the error vectors of the eigenvectors are orthogonal, so the bound on each eigenvector gives the bound on $b$ with no loss.

**theorem** *hhl-clamped-bound*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *t0*: *0 < t0* **and** *N*: *0 < N*
**and** *C*: *0 < C* **and** *th*: *0* < ϑ ϑ < *1* **and** *lmin*: *2 * C / ϑ ≤ lmin*
**and** *spec*: ⋀*l*. *l* ∈ *herm-spectrum A* ⟹ *lmin* ≤ *l* ∧ *t0 * l ≤ pi / 2*
**and** *b*: *dim-vec b = n*
**shows** *Re* ⟨*hhl-circuit-clamped A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec b* (*unit-vec 2 0*))
−
*tensor-vec* (*uniform-state N*)
(*tensor-vec* (*mat-fun* (λ*x*. *complex-of-real* (*sqrt* (1 − (*C* / *x*)$^2$))) *A* $*_v$ *b*) (*unit-vec 2 0*) +
*tensor-vec* ((*complex-of-real C* $\cdot_m$ *herm-inv A*) $*_v$ *b*) (*unit-vec 2 1*)) |
*hhl-circuit-clamped A t0 N C* $*_v$ *tensor-vec* (*uniform-state N*) (*tensor-vec b* (*unit-vec 2 0*)) −

*tensor-vec* (*uniform-state N*)
(*tensor-vec* (*mat-fun* (λ*x*. *complex-of-real* (*sqrt* (1 − (*C* / *x*)²))) *A* ∗*v b*) (*unit-vec* 2 0) +
*tensor-vec* ((*complex-of-real C* ·*m herm-inv A*) ∗*v b*) (*unit-vec* 2 1))⟩
≤ *pi*² / (*real N* ∗ *t0*²) ∗ (ϑ ^ 4 / (*C*² ∗ (1 − ϑ²)) + 16 / *lmin*²) ∗ *Re* ⟨*b*|*b*⟩
⟨*proof*⟩

**end**
**theory** *HHL-Window*
**imports** *HHL-Approximate*
**begin**

# 114 HHL with a sine-window phase register

With the uniform register state the error of approximate phase estimation is $O(1/\sqrt{N})$, so precision $\varepsilon$ needs $N \sim 1/\varepsilon^2$. Preparing the register in the window state $\psi_\tau = \sqrt{2/N}\,\sin(\pi(\tau+\frac{1}{2})/N)$ makes the amplitudes decay like $1/(N\delta)^2$ instead of $1/(N\delta)$, and the error drops to $O(1/N)$: a register of size $N \sim 1/\varepsilon$ suffices (theorem *hhl-window-precision*).

## 114.1 The window amplitude

**lemma** *cis-minus-cis-neg*: *cis t* − *cis* (− *t*) = 2 ∗ i ∗ *complex-of-real* (*sin t*)
⟨*proof*⟩

**lemma** *of-real-sin-cis*: *complex-of-real* (*sin t*) = (*cis t* − *cis* (− *t*)) / (2 ∗ i)
⟨*proof*⟩

**lemma** *cis-add-pi*: *cis* (*y* + *pi*) = − *cis y*
⟨*proof*⟩

**lemma** *cis-diff-pi*: *cis* (*y* − *pi*) = − *cis y*
⟨*proof*⟩

The window sum: with $a = \pi/N$, $\left|\sum_{\tau<N} \sin(a\tau + a/2)\, e^{i\tau x}\right| \le |\sin(a/2)|/(|\sin\frac{x+a}{2}|\,|\sin\frac{x-a}{2}|)$.

**lemma** *window-sum-bound*:
**fixes** *x a* :: *real* **and** *N* :: *nat*
**assumes** *Na*: *real N* ∗ *a* = *pi* **and** *p1*: *cis* (*x* + *a*) ≠ 1 **and** *q1*: *cis* (*x* − *a*) ≠ 1
**shows** *cmod* (∑τ<*N*. *complex-of-real* (*sin* (*a* ∗ *real* τ + *a* / 2)) ∗ *cis* (*real* τ ∗ *x*))
≤ |*sin* (*a* / 2)| / (|*sin* ((*x* + *a*) / 2)| ∗ |*sin* ((*x* − *a*) / 2)|)
⟨*proof*⟩

**lemma** *sin-ge-min*:
**assumes** 0 ≤ *y y* ≤ *pi*
**shows** 2 / *pi* ∗ *min y* (*pi* − *y*) ≤ *sin y*
⟨*proof*⟩

**lemma** *window-denominator*:
**fixes** *x a* :: *real*
**assumes** *a0*: 0 < *a* **and** *a2*: *a* ≤ *pi* / 2 **and** *xlo*: 2 ∗ *a* ≤ |*x*| **and** *xhi*: |*x*| ≤ *pi*
**shows** *x*² / (4 ∗ *pi*²) ≤ |*sin* ((*x* + *a*) / 2)| ∗ |*sin* ((*x* − *a*) / 2)|
⟨*proof*⟩

**lemma** *cis-ne-one-of-sin*: *sin* (*y* / 2) ≠ 0 ⟹ *cis y* ≠ 1
⟨*proof*⟩

The window amplitude bound: for $N \ge 2$ and $2\pi/N \le |x| \le \pi$, $\left|\frac{\sqrt{2}}{N}\sum_{\tau<N} \sin\frac{\pi(\tau+1/2)}{N} e^{i\tau x}\right| \le \frac{2\sqrt{2}\,\pi^3}{N^2x^2}$.

**theorem** *window-amp-bound*:
**fixes** $x$ :: *real* **and** $N$ :: *nat*
**assumes** $N$: $2 \leq N$ **and** *xlo*: $2 * pi / real\ N \leq |x|$ **and** *xhi*: $|x| \leq pi$
**shows** $cmod\ (complex\text{-}of\text{-}real\ (sqrt\ 2 / real\ N) * (\sum \tau{<}N.\ complex\text{-}of\text{-}real\ (sin\ (pi * (real\ \tau + 1 / 2) / real\ N)) * cis\ (real\ \tau * x)))$
$\leq 2 * sqrt\ 2 * pi\ \hat{}\ 3 / ((real\ N)^2 * x^2)$
⟨*proof*⟩

## 114.2 Phase estimation with an arbitrary register state

**definition** *pe-amp-reg* :: *nat* ⇒ *complex Matrix.vec* ⇒ *real* ⇒ *nat* ⇒ *complex* **where**
*pe-amp-reg N psi φ j* = $((fourier\ N)^\dagger *_v Matrix.vec\ N\ (\lambda k.\ cis\ (2 * pi * \varphi)\ \hat{}\ k * psi\ \$\ k))\ \$\ j$

**theorem** *phase-estimation-reg*:
**assumes** $U$: $U \in carrier\text{-}mat\ m\ m$ **and** $u$: $dim\text{-}vec\ u = m$
**and** *ev*: $U *_v u = cis\ (2 * pi * \varphi) \cdot_v u$ **and** *psi*: $dim\text{-}vec\ psi = N$
**shows** $(((fourier\ N)^\dagger) \bigotimes 1_m\ m) *_v (ctrl\text{-}pow\ N\ m\ U *_v tensor\text{-}vec\ psi\ u) =$
$tensor\text{-}vec\ (Matrix.vec\ N\ (pe\text{-}amp\text{-}reg\ N\ psi\ \varphi))\ u$
⟨*proof*⟩

**lemma** *pe-amp-reg-formula*:
**assumes** $j$: $j < N$
**shows** $pe\text{-}amp\text{-}reg\ N\ psi\ \varphi\ j = (\sum k{<}N.\ psi\ \$\ k * cis\ (2 * pi * real\ k * (\varphi - real\ j / real\ N))) / complex\text{-}of\text{-}real\ (sqrt\ (real\ N))$
⟨*proof*⟩

**context**
**fixes** $A$ :: *complex Matrix.mat* **and** $n\ N$ :: *nat* **and** *t0 l* :: *real* **and** $u$ :: *complex Matrix.vec*
**and** $f$ :: *nat* ⇒ *real* **and** $c$ :: *real* **and** *psi* :: *complex Matrix.vec*
**assumes** $A$: $A \in carrier\text{-}mat\ n\ n$ *hermitian A* **and** *t0*: $0 < t0$ **and** $N$: $0 < N$
**and** $u$: $dim\text{-}vec\ u = n$ **and** *ev*: $A *_v u = complex\text{-}of\text{-}real\ l \cdot_v u$
**and** *psi*: $dim\text{-}vec\ psi = N$
**begin**

The phase of $u$, the register amplitudes, the ideal and the branch rotations.

**abbreviation** *rphi* ≡ *hhl-phi t0 l*
**definition** *ralpha* = *Matrix.vec N* (*pe-amp-reg N psi rphi*)
**abbreviation** *rr-ideal* ≡ *rot c* $*_v$ *unit-vec 2 0*
**abbreviation** *rr-br k* ≡ *rot* (*f k*) $*_v$ *unit-vec 2 0*

**lemma** *phi-phase-reg*: *cis* (*t0* $*$ *l*) = *cis* (*2* $*$ *pi* $*$ *rphi*)
⟨*proof*⟩

**lemma** *dim-alpha-reg* [*simp*]: *dim-vec ralpha* = *N* ⟨*proof*⟩
**lemma** *dim-r-ideal-reg*: *dim-vec rr-ideal* = *2* ⟨*proof*⟩

**lemma** *W-branch-reg*:
**assumes** $w$: $dim\text{-}vec\ w = 2$
**shows** *hhl-W A t0 N* $*_v$ *tensor-vec psi* (*tensor-vec u w*) = *tensor-vec ralpha* (*tensor-vec u w*)
⟨*proof*⟩

**lemma** *hhl-W-carrier-reg*: *hhl-W A t0 N* ∈ *carrier-mat* (*N* $*$ (*n* $*$ *2*)) (*N* $*$ (*n* $*$ *2*))
⟨*proof*⟩

**lemma** *hhl-W-unitary-reg*: *unitary* (*hhl-W A t0 N*)
⟨*proof*⟩

The output of the circuit and the ideal output as register sums.

**lemma** *hhl-output-branches-reg*:
*hhl-gen A t0 N f* $*_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec 2 0*)) =
(*hhl-W A t0 N*)$^\dagger$ $*_v$ *reg-sum N* (*n* $*$ *2*) (λ*k*. *ralpha* $ *k* $\cdot_v$ *tensor-vec u* (*rr-br k*))
⟨*proof*⟩

**lemma** *hhl-ideal-branches-reg*:
*tensor-vec psi* (*tensor-vec u rr-ideal*) =
(*hhl-W A t0 N*)$^\dagger$ $*_v$ *reg-sum N* (*n* $*$ *2*) (λ*k*. *ralpha* $ *k* $\cdot_v$ *tensor-vec u rr-ideal*)
⟨*proof*⟩

The error vector: $W^* \sum_k \alpha_k |k\rangle \otimes u \otimes (R(f_k)|0\rangle - R(c)|0\rangle)$.

**lemma** *hhl-err-vec-reg*:
*hhl-gen A t0 N f* $*_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec psi* (*tensor-vec u rr-ideal*)
=
(*hhl-W A t0 N*)$^\dagger$ $*_v$ *reg-sum N* (*n* $*$ *2*) (λ*k*. *ralpha* $ *k* $\cdot_v$ *tensor-vec u* (*rr-br k* − *rr-ideal*))
⟨*proof*⟩

The squared distance between output and ideal output.

**theorem** *hhl-approx-error-reg*:
⟨*hhl-gen A t0 N f* $*_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec psi* (*tensor-vec u rr-ideal*)
|
*hhl-gen A t0 N f* $*_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec psi* (*tensor-vec u rr-ideal*)⟩ =
⟨*u*|*u*⟩ $*$ ($\sum$ *k*<*N*. *complex-of-real* ((*cmod* (*pe-amp-reg N psi rphi k*))$^2$) $*$ ⟨*rr-br k* − *rr-ideal*|*rr-br k* − *rr-ideal*⟩)
⟨*proof*⟩

The branch distance in real terms.

**corollary** *hhl-approx-error-real-reg*:
⟨*hhl-gen A t0 N f* $*_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec psi* (*tensor-vec u rr-ideal*)
|
*hhl-gen A t0 N f* $*_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec 2 0*)) − *tensor-vec psi* (*tensor-vec u rr-ideal*)⟩ =
⟨*u*|*u*⟩ $*$ *complex-of-real* ($\sum$ *k*<*N*. (*cmod* (*pe-amp-reg N psi rphi k*))$^2$ $*$
((*sqrt* (*1* − (*f k*)$^2$) − *sqrt* (*1* − *c*$^2$))$^2$ + (*f k* − *c*)$^2$))
⟨*proof*⟩

**end**

## 114.3 The sine window

**definition** *win-state* :: *nat* ⇒ *complex Matrix.vec* **where**
*win-state N* = *Matrix.vec N* (λτ. *complex-of-real* (*sqrt* (*2* / *real N*) $*$ *sin* (*pi* $*$ (*real* τ + *1* / *2*) / *real N*)))

**lemma** *dim-win-state* [*simp*]: *dim-vec* (*win-state N*) = *N*
⟨*proof*⟩

**lemma** *win-amp-formula*:
**assumes** *j*: *j* < *N*
**shows** *pe-amp-reg N* (*win-state N*) φ *j* =
*complex-of-real* (*sqrt 2* / *real N*) $*$ ($\sum$ τ<*N*. *complex-of-real* (*sin* (*pi* $*$ (*real* τ + *1* / *2*) / *real N*)) $*$ *cis* (*real* τ $*$ (*2* $*$ *pi* $*$ (φ − *real j* / *real N*))))
⟨*proof*⟩

The quadratic decay: $|\alpha_j|^2 \leq \pi^2/(2N^4\delta^4)$ for $1/N \leq |\delta| \leq 1/2$, $\delta = \varphi - j/N$.

**theorem** *win-amp-tail*:

**assumes** *N*: *2 ≤ N* **and** *j*: *j < N* **and** *dlo*: *1 / real N ≤ |φ − real j / real N|* **and** *dhi*: *|φ − real j / real N| ≤ 1 / 2*
**shows** *(cmod (pe-amp-reg N (win-state N) φ j))² ≤ pi² / (2 * (real N) ^ 4 * (φ − real j / real N) ^ 4)*
⟨*proof*⟩

**lemma** *win-amp-crude*:
**assumes** *j*: *j < N*
**shows** *(cmod (pe-amp-reg N (win-state N) φ j))² ≤ 2*
⟨*proof*⟩

**lemma** *pe-amp-reg-shift-up*: *pe-amp-reg N psi (φ + 1) j = pe-amp-reg N psi φ j*
⟨*proof*⟩

## 114.4 Counting lemmas

**lemma** *near-count*:
**fixes** *y* :: *real*
**shows** *card {k. k < N ∧ |y − real k| < 1} ≤ 2*
⟨*proof*⟩

$\sum_{m \le n} 1/(m+1)^2 \le 2 - 1/(n+1)$, by telescoping.

**lemma** *inv-sq-partial*: *(∑ m<Suc n. 1 / (real m + 1)²) ≤ 2 − 1 / (real n + 1)*
⟨*proof*⟩

**lemma** *inv-sq-finite-nat*:
**assumes** *finite X*
**shows** *(∑ m∈X. 1 / (real m + 1)²) ≤ 2*
⟨*proof*⟩

If $h \ge 1$ on $A$ and $\lfloor h \rfloor$ is injective there, $\sum_{k \in A} 1/h(k)^2 \le 2$.

**lemma** *inv-sq-floor-inj*:
**fixes** *h* :: *nat ⇒ real*
**assumes** *A*: *finite A* **and** *h1*: *⋀k. k ∈ A ⟹ 1 ≤ h k* **and** *inj*: *inj-on (λk. ⌊h k⌋) A*
**shows** *(∑ k∈A. 1 / (h k)²) ≤ 2*
⟨*proof*⟩

$\sum_{k<N,\,|y-k|\ge 1} 1/(y-k)^2 \le 4$.

**lemma** *far-inv-sq-sum*:
**fixes** *y* :: *real*
**shows** *(∑ k∈{k. k < N ∧ 1 ≤ |y − real k|}. 1 / (y − real k)²) ≤ 4*
⟨*proof*⟩

The branch weight: $w(t) = 2$ for $|t| < 1$, $\pi^2/(2t^2)$ otherwise; $\sum_{k<N} w(y-k) \le 4 + 2\pi^2$.

**definition** *win-w* :: *real ⇒ real* **where**
*win-w t = (if |t| < 1 then 2 else pi² / (2 * t²))*

**lemma** *win-w-nonneg*: *0 ≤ win-w t*
⟨*proof*⟩

**lemma** *win-w-sum*: *(∑ k<N. win-w (y − real k)) ≤ 4 + 2 * pi²*
⟨*proof*⟩

## 114.5 Branch bounds for the window register

If $g \le |\delta| \le \frac{1}{2}$ then $|\alpha_j|^2 \le \pi^2/(2N^4 g^4)$.

**lemma** *win-amp-gap*:

**assumes** *N*: $2 \le N$ **and** *j*: $j < N$ **and** *dhi*: $|\varphi - \mathit{real}\ j\ /\ \mathit{real}\ N| \le 1\ /\ 2$
**and** *g0*: $0 < g$ **and** *g*: $g \le |\varphi - \mathit{real}\ j\ /\ \mathit{real}\ N|$
**shows** $(\mathit{cmod}\ (\mathit{pe\text{-}amp\text{-}reg}\ N\ (\mathit{win\text{-}state}\ N)\ \varphi\ j))^2 \le \mathit{pi}^2\ /\ (2 * (\mathit{real}\ N)\ \hat{}\ 4 * g\ \hat{}\ 4)$
⟨*proof*⟩

The distance on a good branch is quadratic in $\delta$.

**lemma** *hhl-good-dist*:
**assumes** *N*: $0 < N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C\ /\ \vartheta \le l$ **and** *l-hi*: $t0 * l \le \mathit{pi}\ /\ 2$
**and** *k*: $2 * k < N$ **and** *lam*: $C\ /\ \vartheta \le \mathit{reg\text{-}eig}\ t0\ N\ k$
**shows** $(\mathit{sqrt}\ (1 - (\mathit{clamp}\ (C\ /\ \mathit{reg\text{-}eig}\ t0\ N\ k))^2) - \mathit{sqrt}\ (1 - (C\ /\ l)^2))^2 + (\mathit{clamp}\ (C\ /\ \mathit{reg\text{-}eig}\ t0\ N\ k) - C\ /\ l)^2$
$\le 4 * \mathit{pi}^2 * \vartheta\ \hat{}\ 4 * (\mathit{hhl\text{-}phi}\ t0\ l - \mathit{real}\ k\ /\ \mathit{real}\ N)^2\ /\ (t0^2 * C^2 * (1 - \vartheta^2))$
**and** $|\mathit{hhl\text{-}phi}\ t0\ l - \mathit{real}\ k\ /\ \mathit{real}\ N| \le 1\ /\ 2$
⟨*proof*⟩

Good branches: $|\alpha_k|^2 d_k^2 \le K\, w(N\varphi - k)/N^2$ with $K = 4\pi^2\theta^4/(t_0^2 C^2(1-\theta^2))$.

**lemma** *win-branch-good*:
**assumes** *N*: $2 \le N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C\ /\ \vartheta \le l$ **and** *l-hi*: $t0 * l \le \mathit{pi}\ /\ 2$
**and** *k*: $2 * k < N$ **and** *lam*: $C\ /\ \vartheta \le \mathit{reg\text{-}eig}\ t0\ N\ k$
**shows** $(\mathit{cmod}\ (\mathit{pe\text{-}amp\text{-}reg}\ N\ (\mathit{win\text{-}state}\ N)\ (\mathit{hhl\text{-}phi}\ t0\ l)\ k))^2 *$
$((\mathit{sqrt}\ (1 - (\mathit{clamp}\ (C\ /\ \mathit{reg\text{-}eig}\ t0\ N\ k))^2) - \mathit{sqrt}\ (1 - (C\ /\ l)^2))^2 + (\mathit{clamp}\ (C\ /\ \mathit{reg\text{-}eig}\ t0\ N\ k) - C\ /\ l)^2)$
$\le 4 * \mathit{pi}^2 * \vartheta\ \hat{}\ 4\ /\ (t0^2 * C^2 * (1 - \vartheta^2))\ /\ (\mathit{real}\ N)^2 * \mathit{win\text{-}w}\ (\mathit{real}\ N * \mathit{hhl\text{-}phi}\ t0\ l - \mathit{real}\ k)$
⟨*proof*⟩

Bad branches (clamped or in the negative half): $|\alpha_k|^2 \le 8\pi^2/(N\varphi)^4$.

**lemma** *win-branch-bad*:
**assumes** *N*: $2 \le N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C\ /\ \vartheta \le l$ **and** *l-hi*: $t0 * l \le \mathit{pi}\ /\ 2$
**and** *k*: $k < N$ **and** *bad*: $\neg\ (2 * k < N \wedge C\ /\ \vartheta \le \mathit{reg\text{-}eig}\ t0\ N\ k)$
**shows** $(\mathit{cmod}\ (\mathit{pe\text{-}amp\text{-}reg}\ N\ (\mathit{win\text{-}state}\ N)\ (\mathit{hhl\text{-}phi}\ t0\ l)\ k))^2 \le 8 * \mathit{pi}^2\ /\ ((\mathit{real}\ N)\ \hat{}\ 4 * (\mathit{hhl\text{-}phi}\ t0\ l)\ \hat{}\ 4)$
⟨*proof*⟩

## 114.6 The error bound on an eigenvector

**lemma** *win-branch-bound*:
**assumes** *N*: $2 \le N$ **and** *t0*: $0 < t0$ **and** *C*: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$
**and** *l-lo*: $2 * C\ /\ \vartheta \le l$ **and** *l-hi*: $t0 * l \le \mathit{pi}\ /\ 2$ **and** *k*: $k < N$
**shows** $(\mathit{cmod}\ (\mathit{pe\text{-}amp\text{-}reg}\ N\ (\mathit{win\text{-}state}\ N)\ (\mathit{hhl\text{-}phi}\ t0\ l)\ k))^2 *$
$((\mathit{sqrt}\ (1 - (\mathit{clamp}\ (C\ /\ \mathit{reg\text{-}eig}\ t0\ N\ k))^2) - \mathit{sqrt}\ (1 - (C\ /\ l)^2))^2 + (\mathit{clamp}\ (C\ /\ \mathit{reg\text{-}eig}\ t0\ N\ k) - C\ /\ l)^2)$
$\le 4 * \mathit{pi}^2 * \vartheta\ \hat{}\ 4\ /\ (t0^2 * C^2 * (1 - \vartheta^2))\ /\ (\mathit{real}\ N)^2 * \mathit{win\text{-}w}\ (\mathit{real}\ N * \mathit{hhl\text{-}phi}\ t0\ l - \mathit{real}\ k)$
$+ 32 * \mathit{pi}^2\ /\ ((\mathit{real}\ N)\ \hat{}\ 4 * (\mathit{hhl\text{-}phi}\ t0\ l)\ \hat{}\ 4)$
⟨*proof*⟩

The error bound for the window circuit on an eigenvector:

$$\|\text{output} - \psi \otimes u \otimes R(C/\lambda)|0\rangle\|^2 \le \|u\|^2 \Big( \frac{8\pi^2(2+\pi^2)\theta^4}{N^2 t_0^2 C^2 (1-\theta^2)} + \frac{512\pi^6}{N^3 t_0^4 \lambda^4} \Big).$$

**theorem** *hhl-window-error-bound*:
**fixes** *A* :: *complex Matrix.mat* **and** *u* :: *complex Matrix.vec*
**assumes** *A*: $A \in \mathit{carrier\text{-}mat}\ n\ n$ *hermitian A* **and** *t0*: $0 < t0$ **and** *N*: $2 \le N$
**and** *u*: $\mathit{dim\text{-}vec}\ u = n$ **and** *ev*: $A *_v u = \mathit{complex\text{-}of\text{-}real}\ l \cdot_v u$

**and** $C$: $0 < C$ **and** $th$: $0 < \vartheta\ \vartheta < 1$
**and** $l\text{-}lo$: $2 * C / \vartheta \leq l$ **and** $l\text{-}hi$: $t0 * l \leq pi / 2$
**shows** $Re\ \langle hhl\text{-}circuit\text{-}clamped\ A\ t0\ N\ C *_v tensor\text{-}vec\ (win\text{-}state\ N)\ (tensor\text{-}vec\ u\ (unit\text{-}vec\ 2\ 0)) -$
$tensor\text{-}vec\ (win\text{-}state\ N)\ (tensor\text{-}vec\ u\ (rot\ (C / l) *_v unit\text{-}vec\ 2\ 0)) \mid$
$hhl\text{-}circuit\text{-}clamped\ A\ t0\ N\ C *_v tensor\text{-}vec\ (win\text{-}state\ N)\ (tensor\text{-}vec\ u\ (unit\text{-}vec\ 2\ 0)) -$
$tensor\text{-}vec\ (win\text{-}state\ N)\ (tensor\text{-}vec\ u\ (rot\ (C / l) *_v unit\text{-}vec\ 2\ 0))\rangle$
$\leq Re\ \langle u|u\rangle * (8 * pi^2 * (2 + pi^2) * \vartheta\ \hat{}\ 4 / ((real\ N)^2 * t0^2 * C^2 * (1 - \vartheta^2))$
$+ 512 * pi\ \hat{}\ 6 / ((real\ N)\ \hat{}\ 3 * t0\ \hat{}\ 4 * l\ \hat{}\ 4))$
⟨*proof*⟩

## 114.7 Every right-hand side

**lemma** *hhl-err-orth-reg*:
**fixes** $A$ :: *complex Matrix.mat* **and** $u\ u'$ :: *complex Matrix.vec*
**assumes** $A$: $A \in carrier\text{-}mat\ n\ n\ hermitian\ A$ **and** $t0$: $0 < t0$ **and** $N$: $0 < N$
**and** $u$: $dim\text{-}vec\ u = n$ **and** $ev$: $A *_v u = complex\text{-}of\text{-}real\ l \cdot_v u$
**and** $u'$: $dim\text{-}vec\ u' = n$ **and** $ev'$: $A *_v u' = complex\text{-}of\text{-}real\ l' \cdot_v u'$
**and** $orth$: $\langle u|u'\rangle = 0$ **and** $psi$: $dim\text{-}vec\ psi = N$
**shows** $\langle hhl\text{-}gen\ A\ t0\ N\ f *_v tensor\text{-}vec\ (psi)\ (tensor\text{-}vec\ u\ (unit\text{-}vec\ 2\ 0)) - tensor\text{-}vec\ (psi)\ (tensor\text{-}vec\ u\ (rot\ c *_v unit\text{-}vec\ 2\ 0)) \mid$
$hhl\text{-}gen\ A\ t0\ N\ f' *_v tensor\text{-}vec\ (psi)\ (tensor\text{-}vec\ u'\ (unit\text{-}vec\ 2\ 0)) - tensor\text{-}vec\ (psi)\ (tensor\text{-}vec\ u'\ (rot\ c' *_v unit\text{-}vec\ 2\ 0))\rangle = 0$
⟨*proof*⟩

**lemma** *lift-mat-emb-reg*:
**assumes** $v$: $dim\text{-}vec\ v = n$
**shows** $lift\text{-}mat\ n\ (psi)\ (1_m\ n)\ (0_m\ n\ n) *_v v = tensor\text{-}vec\ (psi)\ (tensor\text{-}vec\ v\ (unit\text{-}vec\ 2\ 0))$
⟨*proof*⟩

**lemma** *ideal-mat-eigen-reg*:
**assumes** $A$: $A \in carrier\text{-}mat\ n\ n\ hermitian\ A$ **and** $u$: $dim\text{-}vec\ u = n$ **and** $ev$: $A *_v u = complex\text{-}of\text{-}real\ l \cdot_v u$
**shows** $lift\text{-}mat\ n\ (psi)\ (mat\text{-}fun\ (\lambda x.\ complex\text{-}of\text{-}real\ (sqrt\ (1 - (C / x)^2)))\ A)\ (complex\text{-}of\text{-}real\ C \cdot_m herm\text{-}inv\ A) *_v u =$
$tensor\text{-}vec\ (psi)\ (tensor\text{-}vec\ u\ (rot\ (C / l) *_v unit\text{-}vec\ 2\ 0))$
⟨*proof*⟩

**theorem** *hhl-window-bound*:
**fixes** $A$ :: *complex Matrix.mat* **and** $b$ :: *complex Matrix.vec*
**assumes** $A$: $A \in carrier\text{-}mat\ n\ n\ hermitian\ A$ **and** $t0$: $0 < t0$ **and** $N$: $2 \leq N$
**and** $C$: $0 < C$ **and** $th$: $0 < \vartheta\ \vartheta < 1$ **and** $lmin$: $2 * C / \vartheta \leq lmin$
**and** $spec$: $\bigwedge l.\ l \in herm\text{-}spectrum\ A \Longrightarrow lmin \leq l \wedge t0 * l \leq pi / 2$
**and** $b$: $dim\text{-}vec\ b = n$
**shows** $Re\ \langle hhl\text{-}circuit\text{-}clamped\ A\ t0\ N\ C *_v tensor\text{-}vec\ (win\text{-}state\ N)\ (tensor\text{-}vec\ b\ (unit\text{-}vec\ 2\ 0)) -$
$tensor\text{-}vec\ (win\text{-}state\ N)$
$(tensor\text{-}vec\ (mat\text{-}fun\ (\lambda x.\ complex\text{-}of\text{-}real\ (sqrt\ (1 - (C / x)^2)))\ A *_v b)\ (unit\text{-}vec\ 2\ 0) +$
$tensor\text{-}vec\ ((complex\text{-}of\text{-}real\ C \cdot_m herm\text{-}inv\ A) *_v b)\ (unit\text{-}vec\ 2\ 1)) \mid$
$hhl\text{-}circuit\text{-}clamped\ A\ t0\ N\ C *_v tensor\text{-}vec\ (win\text{-}state\ N)\ (tensor\text{-}vec\ b\ (unit\text{-}vec\ 2\ 0)) -$
$tensor\text{-}vec\ (win\text{-}state\ N)$
$(tensor\text{-}vec\ (mat\text{-}fun\ (\lambda x.\ complex\text{-}of\text{-}real\ (sqrt\ (1 - (C / x)^2)))\ A *_v b)\ (unit\text{-}vec\ 2\ 0) +$
$tensor\text{-}vec\ ((complex\text{-}of\text{-}real\ C \cdot_m herm\text{-}inv\ A) *_v b)\ (unit\text{-}vec\ 2\ 1))\rangle$
$\leq (8 * pi^2 * (2 + pi^2) * \vartheta\ \hat{}\ 4 / ((real\ N)^2 * t0^2 * C^2 * (1 - \vartheta^2)) + 512 * pi\ \hat{}\ 6 / ((real\ N)\ \hat{}\ 3 * t0\ \hat{}\ 4 * lmin\ \hat{}\ 4)) * Re\ \langle b|b\rangle$
⟨*proof*⟩

**lemma** *window-rate*:
**fixes** $a\ b\ \varepsilon$ :: *real*
**assumes** $N$: $2 \leq N$ **and** $a$: $0 \leq a$ **and** $b$: $0 \leq b$ **and** $e$: $0 < \varepsilon$ **and** $NQ$: $sqrt\ (a + b) / \varepsilon \leq real\ N$

**shows** *a / (real N)*$^2$ *+ b / (real N)* $\hat{}$ *3* $\leq \varepsilon^2$
⟨*proof*⟩

Precision $\varepsilon$ with a register of size $N \geq Q/\varepsilon$: $\|\text{output} - \text{ideal}\| \leq \varepsilon\|b\|$, with $Q$ independent of $N$ and $\varepsilon$.

**corollary** *hhl-window-precision*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *t0*: *0 < t0* **and** *N*: *2 ≤ N*
**and** *C*: *0 < C* **and** *th*: *0 < ϑ ϑ < 1* **and** *lmin*: *2 * C / ϑ ≤ lmin*
**and** *spec*: ⋀*l*. *l* ∈ *herm-spectrum A* ⟹ *lmin ≤ l* ∧ *t0 * l ≤ pi / 2*
**and** *b*: *dim-vec b = n* **and** *e*: *0 < ε*
**and** *NQ*: *sqrt (8 * pi*$^2$ ** (2 + pi*$^2$*) * ϑ* $\hat{}$ *4 / (t0*$^2$ ** C*$^2$ ** (1 − ϑ*$^2$*)) + 512 * pi* $\hat{}$ *6 / (t0* $\hat{}$ *4 * lmin* $\hat{}$ *4)) / ε ≤ real N*
**shows** *Re ⟨hhl-circuit-clamped A t0 N C ∗*$_v$ *tensor-vec (win-state N) (tensor-vec b (unit-vec 2 0)) −*
*tensor-vec (win-state N)*
*(tensor-vec (mat-fun (λx. complex-of-real (sqrt (1 − (C / x)*$^2$*))) A ∗*$_v$ *b) (unit-vec 2 0) +*
*tensor-vec ((complex-of-real C ·*$_m$ *herm-inv A) ∗*$_v$ *b) (unit-vec 2 1)) |*
*hhl-circuit-clamped A t0 N C ∗*$_v$ *tensor-vec (win-state N) (tensor-vec b (unit-vec 2 0)) −*
*tensor-vec (win-state N)*
*(tensor-vec (mat-fun (λx. complex-of-real (sqrt (1 − (C / x)*$^2$*))) A ∗*$_v$ *b) (unit-vec 2 0) +*
*tensor-vec ((complex-of-real C ·*$_m$ *herm-inv A) ∗*$_v$ *b) (unit-vec 2 1))⟩*
*≤ ε*$^2$ ** Re ⟨b|b⟩*
⟨*proof*⟩

**end**
**theory** *HHL-Simulation*
**imports** *HHL-Window*
**begin**

# 115 HHL with an approximate Hamiltonian simulation

The HHL circuit uses $e^{iAt_0}$ only through controlled powers. We write the circuit in a small language of fixed gates and controlled calls of an oracle, count the calls ($2(N-1)$ for HHL) and show that replacing the oracle by any unitary $V$ with $\|Vv - e^{iAt_0}v\| \leq \eta\|v\|$ moves the output by at most (calls)$\cdot\eta$. With the window register this gives the end-to-end bound $(\varepsilon + 2(N-1)\eta)\|b\|$ (theorem *hhl-sim-window-error*). How $V$ is built (Trotter formulas, sparse access) is outside the matrix framework; any construction with this accuracy can be plugged in.

## 115.1 Vector norms

**lemma** *vnorm-L2*: ‖*v*‖ = *L2-set (λi. cmod (v $ i)) {..<dim-vec v}*
⟨*proof*⟩

**lemma** *vnorm-ge0*: *0 ≤* ‖*v*‖
⟨*proof*⟩

**lemma** *vnorm-sq*: ‖*v*‖$^2$ = (∑ *i<dim-vec v. (cmod (v $ i))*$^2$)
⟨*proof*⟩

**lemma** *vnorm-inner*: ⟨*v*|*v*⟩ = *complex-of-real* (‖*v*‖$^2$)
⟨*proof*⟩

**lemma** *vnorm-Re-inner*: *Re* ⟨*v*|*v*⟩ = ‖*v*‖$^2$
⟨*proof*⟩

**lemma** *vnorm-add-le*:

**assumes** *dim-vec a = dim-vec b*
**shows** $\|a + b\| \leq \|a\| + \|b\|$
⟨*proof*⟩

**lemma** *vnorm-minus-commute*:
**assumes** *dim-vec a = dim-vec b*
**shows** $\|a - b\| = \|b - a\|$
⟨*proof*⟩

**lemma** *vnorm-diff-le*:
**assumes** *a*: *dim-vec a = n* **and** *b*: *dim-vec b = n* **and** *c*: *dim-vec c = n*
**shows** $\|a - c\| \leq \|a - b\| + \|b - c\|$
⟨*proof*⟩

**lemma** *vnorm-smult*: $\|z \cdot_v v\| = cmod\ z * \|v\|$
⟨*proof*⟩

**lemma** *vnorm-le-diff-add*:
**assumes** *dim-vec a = dim-vec b*
**shows** $\|a\| \leq \|a - b\| + \|b\|$
⟨*proof*⟩

**lemma** *vnorm-reverse*:
**assumes** *dim-vec a = dim-vec b*
**shows** $|\|a\| - \|b\|| \leq \|a - b\|$
⟨*proof*⟩

**lemma** *unitary-vnorm*:
**assumes** *W*: *W ∈ carrier-mat K K unitary W* **and** *y*: *dim-vec y = K*
**shows** $\|W *_v y\| = \|y\|$
⟨*proof*⟩

## 115.2 Closeness of block operators

The blocks of a vector on $\mathbb{C}^N \otimes \mathbb{C}^m$.

**definition** *blk* :: *nat ⇒ complex Matrix.vec ⇒ nat ⇒ complex Matrix.vec* **where**
*blk m x k = Matrix.vec m* (λ*i*. *x* $ (*k * m + i*))

**lemma** *dim-blk* [*simp*]: *dim-vec* (*blk m x k*) = *m*
⟨*proof*⟩

**lemma** *reg-sum-blk*:
**assumes** *dim-vec x = N * m*
**shows** *reg-sum N m* (*blk m x*) = *x*
⟨*proof*⟩

**lemma** *vnorm-reg-sum-sq*:
**assumes** *f*: $\bigwedge k.\ k < N \Longrightarrow$ *dim-vec* (*f k*) = *m*
**shows** $\|$*reg-sum N m f*$\|^2 = (\sum k{<}N.\ \|f\ k\|^2)$
⟨*proof*⟩

If every block of $B$ is $\eta$-close to the block of $B'$, so is the block operator.

**lemma** *ctrl-blocks-close*:
**assumes** *B*: $\bigwedge k.\ k < N \Longrightarrow$ *B k ∈ carrier-mat m m* **and** *B'*: $\bigwedge k.\ k < N \Longrightarrow$ *B' k ∈ carrier-mat m m*
**and** *close*: $\bigwedge k\ v.\ k < N \Longrightarrow$ *dim-vec v = m* $\Longrightarrow \|B\ k *_v v - B'\ k *_v v\| \leq \eta * \|v\|$
**and** *x*: *dim-vec x = N * m*
**shows** $\|$*ctrl-blocks N m B* $*_v$ *x − ctrl-blocks N m B'* $*_v$ *x*$\| \leq \eta * \|x\|$

⟨*proof*⟩

$\eta$-closeness of two operators: $\|Uv - Vv\| \leq \eta\|v\|$ for every $v$.

**definition** *op-close* :: *nat* ⇒ *real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* ⇒ *bool* **where**
*op-close n η U V* ⟷ (∀ *v*. *dim-vec v* = *n* ⟶ ‖*U* *$_v$ *v* − *V* *$_v$ *v*‖ ≤ *η* ∗ ‖*v*‖)

**lemma** *op-closeD*: *op-close n η U V* ⟹ *dim-vec v* = *n* ⟹ ‖*U* *$_v$ *v* − *V* *$_v$ *v*‖ ≤ *η* ∗ ‖*v*‖
⟨*proof*⟩

**lemma** *op-close-refl*:
**assumes** *e*: *0* ≤ *η* **and** *U*: *U* ∈ *carrier-mat n n*
**shows** *op-close n η U U*
⟨*proof*⟩

**lemma** *op-close-sym*:
**assumes** *op-close n η U V U* ∈ *carrier-mat n n V* ∈ *carrier-mat n n*
**shows** *op-close n η V U*
⟨*proof*⟩

**lemma** *op-close-dagger*:
**assumes** *U*: *U* ∈ *carrier-mat n n unitary U* **and** *V*: *V* ∈ *carrier-mat n n unitary V*
**and** *c*: *op-close n η U V*
**shows** *op-close n η* (*U*$^\dagger$) (*V*$^\dagger$)
⟨*proof*⟩

## 115.3 Tensoring with the ancilla

**definition** *half* :: *complex Matrix.vec* ⇒ *nat* ⇒ *complex Matrix.vec* **where**
*half x b* = *Matrix.vec* (*dim-vec x div 2*) (λ*i*. *x* $ (*i* ∗ *2* + *b*))

**lemma** *dim-half* [*simp*]: *dim-vec* (*half x b*) = *dim-vec x div 2*
⟨*proof*⟩

**lemma** *sum-pairs*:
**fixes** *n* :: *nat* **and** *g* :: *nat* ⇒ *'a*::*comm-monoid-add*
**shows** (∑ *r*<*n* ∗ *2*. *g r*) = (∑ *i*<*n*. *g* (*i* ∗ *2*) + *g* (*i* ∗ *2* + *1*))
⟨*proof*⟩

**lemma** *vnorm-half-sq*:
**assumes** *dim-vec x* = *n* ∗ *2*
**shows** ‖*x*‖$^2$ = ‖*half x 0*‖$^2$ + ‖*half x 1*‖$^2$
⟨*proof*⟩

**lemma** *half-minus*:
**assumes** *y*: *dim-vec y* = *dim-vec x* **and** *b*: *b* < *2*
**shows** *half* (*x* − *y*) *b* = *half x b* − *half y b*
⟨*proof*⟩

**lemma** *tensor-one2-index*:
**assumes** *M*: *M* ∈ *carrier-mat n n* **and** *x*: *dim-vec x* = *n* ∗ *2* **and** *i*: *i* < *n* **and** *b*: *b* < *2*
**shows** ((*M* ⨂ *1*$_m$ *2*) *$_v$ *x*) $ (*i* ∗ *2* + *b*) = (*M* *$_v$ *half x b*) $ *i*
⟨*proof*⟩

**lemma** *half-tensor-one2*:
**assumes** *M*: *M* ∈ *carrier-mat n n* **and** *x*: *dim-vec x* = *n* ∗ *2* **and** *b*: *b* < *2*
**shows** *half* ((*M* ⨂ *1*$_m$ *2*) *$_v$ *x*) *b* = *M* *$_v$ *half x b*
⟨*proof*⟩

**lemma** *op-close-tensor-one2*:
**assumes** *U*: $U \in$ *carrier-mat n n* **and** *V*: $V \in$ *carrier-mat n n* **and** *c*: *op-close n η U V* **and** *e*: $0 \leq \eta$
**shows** *op-close* $(n * 2)$ $\eta$ $(U \bigotimes 1_m\ 2)$ $(V \bigotimes 1_m\ 2)$
⟨*proof*⟩

## 115.4 Circuits with oracle calls

Circuits on $\mathbb{C}^N \otimes \mathbb{C}^m$ built from fixed gates and calls of an oracle $V$ on $\mathbb{C}^m$, controlled on a predicate of the register value. *Call P d* applies $V$ (or $V^*$ if $d$) to the blocks $k$ with $P\,k$.

**datatype** *qcirc* = *Gate complex Matrix.mat* | *Call nat* ⇒ *bool bool* | *Seq qcirc qcirc*

**definition** *call-mat* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ (*nat* ⇒ *bool*) ⇒ *bool* ⇒ *complex Matrix.mat*
**where**
*call-mat N m V P d* = *ctrl-blocks N m* (λ*k*. *if P k then* (*if d then* $V^\dagger$ *else V*) *else* $1_m$ *m*)

*Seq a b* runs *a* first.

**primrec** *sem* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *qcirc* ⇒ *complex Matrix.mat* **where**
*sem N m V* (*Gate M*) = *M*
| *sem N m V* (*Call P d*) = *call-mat N m V P d*
| *sem N m V* (*Seq a b*) = *sem N m V b* ∗ *sem N m V a*

**primrec** *calls* :: *qcirc* ⇒ *nat* **where**
*calls* (*Gate M*) = *0*
| *calls* (*Call P d*) = *1*
| *calls* (*Seq a b*) = *calls a* + *calls b*

**primrec** *gates-ok* :: *nat* ⇒ *qcirc* ⇒ *bool* **where**
*gates-ok K* (*Gate M*) ⟷ *M* ∈ *carrier-mat K K* ∧ *unitary M*
| *gates-ok K* (*Call P d*) ⟷ *True*
| *gates-ok K* (*Seq a b*) ⟷ *gates-ok K a* ∧ *gates-ok K b*

**primrec** *adj* :: *qcirc* ⇒ *qcirc* **where**
*adj* (*Gate M*) = *Gate* ($M^\dagger$)
| *adj* (*Call P d*) = *Call P* (¬ *d*)
| *adj* (*Seq a b*) = *Seq* (*adj b*) (*adj a*)

**lemma** *calls-adj* [*simp*]: *calls* (*adj c*) = *calls c*
⟨*proof*⟩

**lemma** *gates-ok-adj*: *gates-ok K c* ⟹ *gates-ok K* (*adj c*)
⟨*proof*⟩

**context**
**fixes** *N m* :: *nat* **and** *V* :: *complex Matrix.mat*
**assumes** *V*: *V* ∈ *carrier-mat m m unitary V*
**begin**

**lemma** *call-blocks*:
**shows** (*if P k then* (*if d then* $V^\dagger$ *else V*) *else* $1_m$ *m*) ∈ *carrier-mat m m*
**and** *unitary* (*if P k then* (*if d then* $V^\dagger$ *else V*) *else* $1_m$ *m*)
⟨*proof*⟩

**lemma** *sem-carrier*: *gates-ok* $(N * m)$ *c* ⟹ *sem N m V c* ∈ *carrier-mat* $(N * m)$ $(N * m)$
⟨*proof*⟩

**lemma** *sem-unitary*: *gates-ok* $(N * m)$ *c* ⟹ *unitary* (*sem N m V c*)

⟨*proof*⟩

**lemma** *sem-adj*: *gates-ok* $(N * m)$ $c \Longrightarrow$ *sem* $N$ $m$ $V$ (*adj* $c$) = (($sem\ N\ m\ V\ c)^\dagger$)
⟨*proof*⟩

**end**

## 115.5 Robustness: each call adds at most $\eta$

**theorem** *sem-robust*:
**assumes** $V$: $V \in$ *carrier-mat* $m$ $m$ *unitary* $V$ **and** $V'$: $V' \in$ *carrier-mat* $m$ $m$ *unitary* $V'$
**and** $c$: *op-close* $m$ $\eta$ $V$ $V'$ **and** $e$: $0 \leq \eta$ **and** $g$: *gates-ok* $(N * m)$ $q$
**shows** *op-close* $(N * m)$ (*real* (*calls* $q$) $* \eta$) (*sem* $N$ $m$ $V$ $q$) (*sem* $N$ $m$ $V'$ $q$)
⟨*proof*⟩

## 115.6 Controlled powers with $N - 1$ calls

*pchain* $K$ $j$ applies the oracle once on each block $k \geq i$, for $i = 1, \ldots, j$, so block $k$ receives $O^{\min(k,j)}$.

**primrec** *pchain* :: *nat* ⇒ *nat* ⇒ *qcirc* **where**
*pchain* $K$ $0$ = *Gate* ($1_m$ $K$)
| *pchain* $K$ (*Suc* $j$) = *Seq* (*Call* ($\lambda k.$ *Suc* $j \leq k$) *False*) (*pchain* $K$ $j$)

**lemma** *calls-pchain* [*simp*]: *calls* (*pchain* $K$ $j$) = $j$
⟨*proof*⟩

**lemma** *gates-ok-pchain*: *gates-ok* $K$ (*pchain* $K$ $j$)
⟨*proof*⟩

**lemma** *sem-pchain*:
**assumes** $V$: $V \in$ *carrier-mat* $m$ $m$
**shows** *sem* $N$ $m$ $V$ (*pchain* $(N * m)$ $j$) = *ctrl-blocks* $N$ $m$ ($\lambda k.$ $V\ \widehat{}_m$ (*min* $k$ $j$))
⟨*proof*⟩

**lemma** *sem-pchain-ctrl-pow*:
**assumes** $V$: $V \in$ *carrier-mat* $m$ $m$
**shows** *sem* $N$ $m$ $V$ (*pchain* $(N * m)$ $(N - 1)$) = *ctrl-pow* $N$ $m$ $V$
⟨*proof*⟩

## 115.7 HHL as a circuit with $2(N - 1)$ calls

**definition** *pe-prog* :: *nat* ⇒ *nat* ⇒ *qcirc* **where**
*pe-prog* $N$ $m$ = *Seq* (*pchain* $(N * m)$ $(N - 1)$) (*Gate* ($(fourier\ N)^\dagger \otimes 1_m\ m$))

**definition** *hhl-prog* :: *nat* ⇒ *nat* ⇒ *complex Matrix.mat* ⇒ *qcirc* **where**
*hhl-prog* $N$ $m$ *Rg* = *Seq* (*pe-prog* $N$ $m$) (*Seq* (*Gate Rg*) (*adj* (*pe-prog* $N$ $m$)))

**lemma** *calls-hhl-prog*: *calls* (*hhl-prog* $N$ $m$ *Rg*) = $2 * (N - 1)$
⟨*proof*⟩

**lemma** *gates-ok-pe-prog*:
**assumes** $N$: $0 < N$
**shows** *gates-ok* $(N * m)$ (*pe-prog* $N$ $m$)
⟨*proof*⟩

**lemma** *sem-pe-prog*:
**assumes** $V$: $V \in$ *carrier-mat* $m$ $m$
**shows** *sem* $N$ $m$ $V$ (*pe-prog* $N$ $m$) = *pe-unitary* $N$ $m$ $V$

⟨*proof*⟩

**lemma** *sem-hhl-prog*:
**assumes** *V*: $V \in$ *carrier-mat m m unitary V* **and** *N*: $0 < N$ **and** *Rg*: $Rg \in$ *carrier-mat* $(N * m)$ $(N * m)$
**shows** *sem N m V* (*hhl-prog N m Rg*) = (*pe-unitary N m V*)$^\dagger$ ∗ (*Rg* ∗ *pe-unitary N m V*)
⟨*proof*⟩

The rotation stage of the clamped circuit.

**definition** *hhl-rot-stage* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *real* ⇒ *complex Matrix.mat* **where**
*hhl-rot-stage A t0 N C* = *ctrl-blocks N* (*dim-row A* ∗ *2*) (λ*k*. $1_m$ (*dim-row A*) ⨂ *rot* (*clamp* (*C* / *reg-eig t0 N k*)))

**lemma** *hhl-rot-stage-ok*: *hhl-rot-stage A t0 N C* ∈ *carrier-mat* $(N * ($*dim-row A* ∗ *2*$))$ $(N * ($*dim-row A* ∗ *2*$))$ ∧ *unitary* (*hhl-rot-stage A t0 N C*)
⟨*proof*⟩

The circuit with the oracle $V$ in place of $e^{iAt_0}$.

**definition** *hhl-sim* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.mat* **where**
*hhl-sim A t0 N C V* = *sem N* (*dim-row A* ∗ *2*) (*V* ⨂ $1_m$ *2*) (*hhl-prog N* (*dim-row A* ∗ *2*) (*hhl-rot-stage A t0 N C*))

**lemma** *hhl-sim-exact*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *N*: $0 < N$
**shows** *hhl-sim A t0 N C* (*evolution A t0*) = *hhl-circuit-clamped A t0 N C*
⟨*proof*⟩

**theorem** *hhl-sim-robust*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *N*: $0 < N$
**and** *V*: $V \in$ *carrier-mat n n unitary V* **and** *c*: *op-close n η V* (*evolution A t0*) **and** *e*: $0 \le \eta$
**shows** *op-close* $(N * (n * 2))$ $(2 *$ *real* $(N - 1) * \eta)$ (*hhl-sim A t0 N C V*) (*hhl-circuit-clamped A t0 N C*)
⟨*proof*⟩

## 115.8 The window is a unit vector

**lemma** *roots-of-unity-sum*:
**assumes** *N*: $2 \le N$
**shows** $(\sum \tau{<}N.$ *cis* $(2 *$ *pi* ∗ *real* $\tau$ / *real N*$)) = 0$
⟨*proof*⟩

**lemma** *win-state-norm*:
**assumes** *N*: $2 \le N$
**shows** ‖*win-state N*‖ = *1*
⟨*proof*⟩

**lemma** *vnorm-tensor*:
**assumes** *dim-vec a* = *n dim-vec b* = *m*
**shows** ‖*tensor-vec a b*‖ = ‖*a*‖ ∗ ‖*b*‖
⟨*proof*⟩

## 115.9 End to end

End to end: window register, clamped rotations and an oracle $V$ that is $\eta$-close to $e^{iAt_0}$. The output is within $(\varepsilon + 2(N-1)\eta)\|b\|$ of the exact-HHL output.

**theorem** *hhl-sim-window-error*:

**fixes** $A$ :: *complex Matrix.mat* **and** $b$ :: *complex Matrix.vec*
**assumes** $A$: $A \in$ *carrier-mat n n hermitian A* **and** *t0*: $0 <$ *t0* **and** $N$: $2 \le N$
**and** $C$: $0 < C$ **and** *th*: $0 < \vartheta\ \vartheta < 1$ **and** *lmin*: $2 * C / \vartheta \le$ *lmin*
**and** *spec*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow$ *lmin* $\le l \wedge$ *t0* $* l \le$ *pi* / *2*
**and** *b*: *dim-vec b* $=$ *n* **and** *e*: $0 < \varepsilon$
**and** *NQ*: *sqrt* $(8 * pi^2 * (2 + pi^2) * \vartheta\ \hat{}\ 4\ /\ (t0^2 * C^2 * (1 - \vartheta^2)) + 512 * pi\ \hat{}\ 6\ /\ (t0\ \hat{}\ 4 *$ *lmin* $\hat{}\ 4)) / \varepsilon \le$ *real N*
**and** $V$: $V \in$ *carrier-mat n n unitary V* **and** *Vc*: *op-close n* $\eta$ *V* (*evolution A t0*) **and** *eta*: $0 \le \eta$
**shows** $\|$*hhl-sim A t0 N C V* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec b* (*unit-vec 2 0*)) $-$
*tensor-vec* (*win-state N*)
(*tensor-vec* (*mat-fun* ($\lambda x.$ *complex-of-real* (*sqrt* (*1* $-$ (*C* / *x*)$^2$))) *A* $*_v$ *b*) (*unit-vec 2 0*) $+$
*tensor-vec* ((*complex-of-real C* $\cdot_m$ *herm-inv A*) $*_v$ *b*) (*unit-vec 2 1*))$\|$
$\le$ ($\varepsilon + 2 *$ *real* ($N - 1$) $* \eta$) $* \|b\|$
⟨*proof*⟩

**end**
**theory** *HHL-Complete*
**imports** *HHL-Simulation*
**begin**

# 116 Readout, amplitude amplification and the complete algorithm

The answer is read by post-selecting the ancilla on $|1\rangle$; the post-selected state is $\psi \otimes A^{-1}b/\|A^{-1}b\| \otimes |1\rangle$ up to $2\delta/(C\|A^{-1}b\|)$ when the output is within $\delta$ of the exact one. Amplitude amplification raises the success probability $p$ to at least $1-p$ with $O(1/\sqrt{p}) = O(\kappa)$ rounds, each using the circuit and its inverse once. Theorem *hhl-complete* puts state preparation, the window register, the simulation oracle, amplification and readout together and counts the oracle calls.

## 116.1 Reading out the answer

## 116.2 Normalisation is stable

**lemma** *normalize-stable*:
**assumes** *d*: *dim-vec a* $=$ *dim-vec b* **and** *a*: $\|a\| \neq 0$ **and** *b*: $\|b\| \neq 0$
**shows** $\|$*complex-of-real* (*1* / $\|a\|$) $\cdot_v$ *a* $-$ *complex-of-real* (*1* / $\|b\|$) $\cdot_v$ *b*$\| \le 2 * \|a - b\| / \|b\|$
⟨*proof*⟩

## 116.3 Post-selecting the ancilla

The ancilla is the last tensor factor: *post1 v* keeps the components with ancilla value 1. The success probability is $\|\text{post1}\, v\|^2$.

**definition** *post1* :: *complex Matrix.vec* $\Rightarrow$ *complex Matrix.vec* **where**
*post1 v* $=$ *Matrix.vec* (*dim-vec v*) ($\lambda r.$ *if odd r then v* $ *r else 0*)

**lemma** *dim-post1* [*simp*]: *dim-vec* (*post1 v*) $=$ *dim-vec v*
⟨*proof*⟩

**lemma** *post1-minus*: *dim-vec u* $=$ *dim-vec v* $\Longrightarrow$ *post1* $(u - v) =$ *post1 u* $-$ *post1 v*
⟨*proof*⟩

**lemma** *vnorm-post1-le*: $\|$*post1 v*$\| \le \|v\|$
⟨*proof*⟩

**lemma** *post1-ideal*:

**assumes** *w*: *dim-vec w = N* **and** *u*: *dim-vec u = n* **and** *v*: *dim-vec v = n*
**shows** *post1 (tensor-vec w (tensor-vec u (unit-vec 2 0) + tensor-vec v (unit-vec 2 1))) =*
*tensor-vec w (tensor-vec v (unit-vec 2 1))*
⟨*proof*⟩

## 116.4 The answer register

For any output $\tilde{y}$ within $\delta$ of the exact output $y$: the success amplitude $\|\text{post1}\,\tilde{y}\|$ is within $\delta$ of $C\|A^{-1}b\|$, and once $\delta < C\|A^{-1}b\|$ the post-selected state is within $2\delta/(C\|A^{-1}b\|)$ of $\psi \otimes \frac{A^{-1}b}{\|A^{-1}b\|} \otimes |1\rangle$.

**theorem** *hhl-readout*:
**fixes** *A* :: *complex Matrix.mat* **and** *b y'* :: *complex Matrix.vec*
**assumes** *A*: *A ∈ carrier-mat n n hermitian A* **and** *N*: *2 ≤ N* **and** *C*: *0 < C*
**and** *b*: *dim-vec b = n* **and** *x0*: *herm-inv A *$_v$ b ≠ 0$_v$ n*
**and** *y'*: *dim-vec y' = N * (n * 2)*
**and** *close*: *‖y' − tensor-vec (win-state N)*
*(tensor-vec (mat-fun (λx. complex-of-real (sqrt (1 − (C / x)²))) A *$_v$ b) (unit-vec 2 0)*
*+*
*tensor-vec ((complex-of-real C ·$_m$ herm-inv A) *$_v$ b) (unit-vec 2 1))‖ ≤ δ*
**shows** *|‖post1 y'‖ − C * ‖herm-inv A *$_v$ b‖| ≤ δ*
**and** *δ < C * ‖herm-inv A *$_v$ b‖ ⟹*
*‖complex-of-real (1 / ‖post1 y'‖) ·$_v$ post1 y' −*
*tensor-vec (win-state N) (tensor-vec (complex-of-real (1 / ‖herm-inv A *$_v$ b‖) ·$_v$ (herm-inv A*
*$*_v$ b)) (unit-vec 2 1))‖*
*≤ 2 * δ / (C * ‖herm-inv A *$_v$ b‖)*
⟨*proof*⟩

## 116.5 Amplitude amplification

$g$ is the good and $h$ the bad unit vector, $Pg = g$, $Ph = 0$, and the state is $\psi = \sin\theta\, g + \cos\theta\, h$. The Grover operator $Q = (2|\psi\rangle\langle\psi| - I)(I - 2P)$ rotates the plane spanned by $g, h$ by $2\theta$, so after $k$ rounds the good amplitude is $\sin((2k+1)\theta)$.

**definition** *outer* :: *complex Matrix.vec ⇒ complex Matrix.mat* **where**
*outer v = Matrix.mat (dim-vec v) (dim-vec v) (λ(i, j). v $ i * cnj (v $ j))*

**lemma** *outer-carrier*: *dim-vec v = K ⟹ outer v ∈ carrier-mat K K*
⟨*proof*⟩

**lemma** *outer-mult-vec*:
**assumes** *dim-vec v = K dim-vec w = K*
**shows** *outer v *$_v$ w = ⟨v|w⟩ ·$_v$ v*
⟨*proof*⟩

**definition** *grover* :: *nat ⇒ complex Matrix.mat ⇒ complex Matrix.vec ⇒ complex Matrix.mat* **where**
*grover K P psi = (2 ·$_m$ outer psi − 1$_m$ K) * (1$_m$ K − 2 ·$_m$ P)*

**context**
**fixes** *K* :: *nat* **and** *P* :: *complex Matrix.mat* **and** *g h* :: *complex Matrix.vec*
**assumes** *P*: *P ∈ carrier-mat K K* **and** *g*: *dim-vec g = K* **and** *h*: *dim-vec h = K*
**and** *gg*: *⟨g|g⟩ = 1* **and** *hh*: *⟨h|h⟩ = 1* **and** *gh*: *⟨g|h⟩ = 0*
**and** *Pg*: *P *$_v$ g = g* **and** *Ph*: *P *$_v$ h = 0$_v$ K*
**begin**

**definition** *comb* :: *real ⇒ real ⇒ complex Matrix.vec* **where**
*comb a c = complex-of-real a ·$_v$ g + complex-of-real c ·$_v$ h*

**lemma** *dim-comb* [*simp*]: *dim-vec* (*comb a c*) = *K*
⟨*proof*⟩

**lemma** *hg*: ⟨*h*|*g*⟩ = *0*
⟨*proof*⟩

**lemma** *comb-inner*: ⟨*comb a c*|*comb a′ c′*⟩ = *complex-of-real* (*a* ∗ *a′* + *c* ∗ *c′*)
⟨*proof*⟩

**lemma** *comb-lin*: *complex-of-real z* ·$_v$ *comb a c* + *complex-of-real z′* ·$_v$ *comb a′ c′* = *comb* (*z* ∗ *a* + *z′* ∗ *a′*) (*z* ∗ *c* + *z′* ∗ *c′*)
⟨*proof*⟩

**lemma** *P-comb*: *P* ∗$_v$ *comb a c* = *comb a 0*
⟨*proof*⟩

**lemma** *reflP-comb*: (*1*$_m$ *K* − *2* ·$_m$ *P*) ∗$_v$ *comb a c* = *comb* (− *a*) *c*
⟨*proof*⟩

**lemma** *reflpsi-comb*:
(*2* ·$_m$ *outer* (*comb s c*) − *1*$_m$ *K*) ∗$_v$ *comb a d* = *comb* (*2* ∗ (*s* ∗ *a* + *c* ∗ *d*) ∗ *s* − *a*) (*2* ∗ (*s* ∗ *a* + *c* ∗ *d*) ∗ *c* − *d*)
⟨*proof*⟩

**lemma** *grover-comb*:
*grover K P* (*comb* (*sin ϑ*) (*cos ϑ*)) ∗$_v$ *comb* (*sin φ*) (*cos φ*) = *comb* (*sin* (*φ* + *2* ∗ *ϑ*)) (*cos* (*φ* + *2* ∗ *ϑ*))
⟨*proof*⟩

**theorem** *grover-iterate*:
(*grover K P* (*comb* (*sin ϑ*) (*cos ϑ*)) ^$_m$ *k*) ∗$_v$ *comb* (*sin ϑ*) (*cos ϑ*) =
*comb* (*sin* ((*2* ∗ *real k* + *1*) ∗ *ϑ*)) (*cos* ((*2* ∗ *real k* + *1*) ∗ *ϑ*))
⟨*proof*⟩

The success probability after $k$ rounds is $\sin^2((2k+1)\theta)$.

**corollary** *grover-success*:
‖*P* ∗$_v$ ((*grover K P* (*comb* (*sin ϑ*) (*cos ϑ*)) ^$_m$ *k*) ∗$_v$ *comb* (*sin ϑ*) (*cos ϑ*))‖$^2$ = (*sin* ((*2* ∗ *real k* + *1*) ∗ *ϑ*))$^2$
⟨*proof*⟩

**end**

## 116.6 The number of rounds

With $k = \lfloor \pi/(4\theta) \rfloor$ rounds the success probability is at least $\cos^2\theta = 1 - \sin^2\theta$, and $k \leq \pi/(4\theta)$.

**lemma** *grover-rounds*:
**assumes** *t0*: *0* < *ϑ* **and** *t1*: *ϑ* < *pi* / *2*
**defines** *k* ≡ *nat* ⌊*pi* / (*4* ∗ *ϑ*)⌋
**shows** *1* − (*sin ϑ*)$^2$ ≤ (*sin* ((*2* ∗ *real k* + *1*) ∗ *ϑ*))$^2$ **and** *real k* ≤ *pi* / (*4* ∗ *ϑ*)
**and** *0* < *sin* ((*2* ∗ *real k* + *1*) ∗ *ϑ*)
⟨*proof*⟩

## 116.7 Good and bad components of a state

**lemma** *projector-split*:
**assumes** *P*: *P* ∈ *carrier-mat K K P* ∗ *P* = *P P*† = *P*
**and** *psi*: *dim-vec psi* = *K* ‖*psi*‖ = *1* **and** *a0*: *0* < ‖*P* ∗$_v$ *psi*‖ **and** *a1*: ‖*P* ∗$_v$ *psi*‖ < *1*

**obtains** *g h ϑ* **where** *dim-vec g = K dim-vec h = K ⟨g|g⟩ = 1 ⟨h|h⟩ = 1 ⟨g|h⟩ = 0*
*P ∗v g = g P ∗v h = 0v K 0 < ϑ ϑ < pi / 2 sin ϑ = ‖P ∗v psi‖*
*psi = complex-of-real (sin ϑ) ·v g + complex-of-real (cos ϑ) ·v h*
⟨*proof*⟩

## 116.8 Reflections as diagonal sign matrices

**definition** *sign-diag :: nat ⇒ (nat ⇒ bool) ⇒ complex Matrix.mat* **where**
*sign-diag K f = Matrix.mat K K (λ(i, j). if i = j then (if f i then − 1 else 1) else 0)*

**lemma** *sign-diag-carrier*: *sign-diag K f ∈ carrier-mat K K*
⟨*proof*⟩

**lemma** *sign-diag-dagger*: *(sign-diag K f)† = sign-diag K f*
⟨*proof*⟩

**lemma** *sign-diag-sq*: *sign-diag K f ∗ sign-diag K f = 1m K*
⟨*proof*⟩

**lemma** *sign-diag-unitary*: *unitary (sign-diag K f)*
⟨*proof*⟩

Post-selection on the ancilla as a matrix.

**definition** *proj1 :: nat ⇒ complex Matrix.mat* **where**
*proj1 K = Matrix.mat K K (λ(i, j). if i = j ∧ odd i then 1 else 0)*

**lemma** *proj1-carrier*: *proj1 K ∈ carrier-mat K K*
⟨*proof*⟩

**lemma** *proj1-mult-vec*: *dim-vec v = K ⟹ proj1 K ∗v v = post1 v*
⟨*proof*⟩

**lemma** *proj1-idem*: *proj1 K ∗ proj1 K = proj1 K*
⟨*proof*⟩

**lemma** *proj1-dagger*: *(proj1 K)† = proj1 K*
⟨*proof*⟩

**lemma** *refl-proj1*: *1m K − 2 ·m proj1 K = sign-diag K odd*
⟨*proof*⟩

**lemma** *refl-zero*: *2 ·m outer (unit-vec K 0) − 1m K = sign-diag K (λi. i ≠ 0)*
⟨*proof*⟩

Reflection about $U|0\rangle$: $U(2|0\rangle\langle 0| - I)U^* = 2|\psi\rangle\langle\psi| - I$ for $\psi = U|0\rangle$.

**lemma** *outer-conj*:
**assumes** *U*: *U ∈ carrier-mat K K* **and** *K*: *0 < K*
**shows** *U ∗ outer (unit-vec K 0) ∗ (U†) = outer (U ∗v unit-vec K 0)*
⟨*proof*⟩

**lemma** *reflection-conj*:
**assumes** *U*: *U ∈ carrier-mat K K unitary U* **and** *K*: *0 < K*
**shows** *U ∗ (sign-diag K (λi. i ≠ 0) ∗ (U†)) = 2 ·m outer (U ∗v unit-vec K 0) − 1m K*
⟨*proof*⟩

## 116.9 Amplitude amplification as a circuit

One round: reflect about the bad subspace, then about $U|0\rangle$; it uses $U$ and $U^*$.

**definition** *grover-prog* :: *nat* ⇒ *qcirc* ⇒ *qcirc* **where**
*grover-prog K u* = *Seq* (*Gate* (*sign-diag K odd*)) (*Seq* (*adj u*) (*Seq* (*Gate* (*sign-diag K* (λ*i*. *i* ≠ *0*))) *u*))

**primrec** *aa-prog* :: *nat* ⇒ *qcirc* ⇒ *nat* ⇒ *qcirc* **where**
*aa-prog K u 0* = *u*
| *aa-prog K u* (*Suc k*) = *Seq* (*aa-prog K u k*) (*grover-prog K u*)

**lemma** *calls-aa-prog*: *calls* (*aa-prog K u k*) = (*2* ∗ *k* + *1*) ∗ *calls u*
⟨*proof*⟩

**lemma** *gates-ok-aa-prog*: *gates-ok K u* ⟹ *gates-ok K* (*aa-prog K u k*)
⟨*proof*⟩

**lemma** *sem-grover-prog*:
**assumes** *V*: *V* ∈ *carrier-mat m m unitary V* **and** *g*: *gates-ok* (*N* ∗ *m*) *u* **and** *K*: *0* < *N* ∗ *m*
**shows** *sem N m V* (*grover-prog* (*N* ∗ *m*) *u*) = *grover* (*N* ∗ *m*) (*proj1* (*N* ∗ *m*)) (*sem N m V u* ∗*v unit-vec* (*N* ∗ *m*) *0*)
⟨*proof*⟩

**lemma** *pow-mat-comm*:
**assumes** *A*: *A* ∈ *carrier-mat n n*
**shows** *A* ∗ *A* ^*m k* = *A* ^*m k* ∗ *A*
⟨*proof*⟩

**lemma** *sem-aa-prog*:
**assumes** *V*: *V* ∈ *carrier-mat m m unitary V* **and** *g*: *gates-ok* (*N* ∗ *m*) *u* **and** *K*: *0* < *N* ∗ *m*
**shows** *sem N m V* (*aa-prog* (*N* ∗ *m*) *u k*) = *grover* (*N* ∗ *m*) (*proj1* (*N* ∗ *m*)) (*sem N m V u* ∗*v unit-vec* (*N* ∗ *m*) *0*) ^*m k* ∗ *sem N m V u*
⟨*proof*⟩

## 116.10 The amplification theorem

**theorem** *amplitude-amplification*:
**assumes** *P*: *P* ∈ *carrier-mat K K P* ∗ *P* = *P P*† = *P*
**and** *psi*: *dim-vec psi* = *K* ‖*psi*‖ = *1* **and** *a0*: *0* < ‖*P* ∗*v psi*‖ **and** *a1*: ‖*P* ∗*v psi*‖ < *1*
**obtains** ϑ **where** *0* < ϑ ϑ < *pi* / *2 sin* ϑ = ‖*P* ∗*v psi*‖
⋀*k*. *P* ∗*v* (*grover K P psi* ^*m k* ∗*v psi*) = *complex-of-real* (*sin* ((*2* ∗ *real k* + *1*) ∗ ϑ) / *sin* ϑ) ·*v* (*P* ∗*v psi*)
⟨*proof*⟩

**lemma** *normalize-smult-pos*:
**assumes** *r*: *0* < *r* **and** *v*: ‖*v*‖ ≠ *0*
**shows** *complex-of-real* (*1* / ‖*complex-of-real r* ·*v v*‖) ·*v* (*complex-of-real r* ·*v v*) = *complex-of-real* (*1* / ‖*v*‖) ·*v v*
⟨*proof*⟩

Amplifying a circuit $u$: with $k = \lfloor \pi/(4\theta) \rfloor$ rounds, $\sin\theta = \|\text{post1}\, u|0\rangle\|$, the success probability is at least $1 - \sin^2\theta$, the post-selected state is unchanged, and the oracle is called $(2k+1)\cdot\text{calls}(u)$ times with $k \le \pi/(4\sin\theta)$.

**theorem** *circuit-amplification*:
**assumes** *V*: *V* ∈ *carrier-mat m m unitary V* **and** *g*: *gates-ok* (*N* ∗ *m*) *u* **and** *K*: *0* < *N* ∗ *m*
**and** *a0*: *0* < ‖*post1* (*sem N m V u* ∗*v unit-vec* (*N* ∗ *m*) *0*)‖
**and** *a1*: ‖*post1* (*sem N m V u* ∗*v unit-vec* (*N* ∗ *m*) *0*)‖ < *1*
**obtains** *k* **where**
*1* − ‖*post1* (*sem N m V u* ∗*v unit-vec* (*N* ∗ *m*) *0*)‖² ≤ ‖*post1* (*sem N m V* (*aa-prog* (*N* ∗ *m*) *u k*) ∗*v unit-vec* (*N* ∗ *m*) *0*)‖²

*complex-of-real (1 / ‖post1 (sem N m V (aa-prog (N * m) u k) *$_v$ unit-vec (N * m) 0)‖) ·$_v$ post1 (sem N m V (aa-prog (N * m) u k) *$_v$ unit-vec (N * m) 0)*
*= complex-of-real (1 / ‖post1 (sem N m V u *$_v$ unit-vec (N * m) 0)‖) ·$_v$ post1 (sem N m V u *$_v$ unit-vec (N * m) 0)*
*calls (aa-prog (N * m) u k) = (2 * k + 1) * calls u*
*real k ≤ pi / (4 * ‖post1 (sem N m V u *$_v$ unit-vec (N * m) 0)‖)*
⟨*proof*⟩

## 116.11 The complete algorithm

State preparation $Prep|0\rangle = \psi \otimes b \otimes |0\rangle$ (a fixed gate), the HHL circuit with oracle $V$, amplitude amplification and post-selection on the ancilla.

**definition** *hhl-full-prog* :: *complex Matrix.mat ⇒ real ⇒ nat ⇒ real ⇒ complex Matrix.mat ⇒ qcirc* **where**
*hhl-full-prog A t0 N C Prep = Seq (Gate Prep) (hhl-prog N (dim-row A * 2) (hhl-rot-stage A t0 N C))*

**lemma** *calls-hhl-full-prog*: *calls (hhl-full-prog A t0 N C Prep) = 2 * (N − 1)*
⟨*proof*⟩

**theorem** *hhl-complete*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: *A ∈ carrier-mat n n hermitian A* **and** *t0*: *0 < t0* **and** *N*: *2 ≤ N*
**and** *C*: *0 < C* **and** *th*: *0 < ϑ ϑ < 1* **and** *lmin*: *2 * C / ϑ ≤ lmin*
**and** *spec*: *⋀l. l ∈ herm-spectrum A ⟹ lmin ≤ l ∧ t0 * l ≤ pi / 2*
**and** *b*: *dim-vec b = n ‖b‖ = 1* **and** *e*: *0 < ε*
**and** *NQ*: *sqrt (8 * pi² * (2 + pi²) * ϑ ^ 4 / (t0² * C² * (1 − ϑ²)) + 512 * pi ^ 6 / (t0 ^ 4 * lmin ^ 4)) / ε ≤ real N*
**and** *V*: *V ∈ carrier-mat n n unitary V* **and** *Vc*: *op-close n η V (evolution A t0)* **and** *eta*: *0 ≤ η*
**and** *Prep*: *Prep ∈ carrier-mat (N * (n * 2)) (N * (n * 2)) unitary Prep*
*Prep *$_v$ unit-vec (N * (n * 2)) 0 = tensor-vec (win-state N) (tensor-vec b (unit-vec 2 0))*
**and** *small*: *ε + 2 * real (N − 1) * η ≤ C * t0 / pi*
**defines** *out ≡ sem N (n * 2) (V ⨂ 1$_m$ 2) (hhl-full-prog A t0 N C Prep) *$_v$ unit-vec (N * (n * 2)) 0*
**and** *x ≡ herm-inv A *$_v$ b*
**shows** *C * t0 / pi ≤ ‖post1 out‖*
**and** *‖complex-of-real (1 / ‖post1 out‖) ·$_v$ post1 out −*
*tensor-vec (win-state N) (tensor-vec (complex-of-real (1 / ‖x‖) ·$_v$ x) (unit-vec 2 1))‖*
*≤ 2 * (ε + 2 * real (N − 1) * η) / (C * ‖x‖)*
**and** *‖post1 out‖ < 1 ⟹ ∃k.*
*1 − ‖post1 out‖² ≤ ‖post1 (sem N (n * 2) (V ⨂ 1$_m$ 2) (aa-prog (N * (n * 2)) (hhl-full-prog A t0 N C Prep) k) *$_v$ unit-vec (N * (n * 2)) 0)‖²*
*∧ complex-of-real (1 / ‖post1 (sem N (n * 2) (V ⨂ 1$_m$ 2) (aa-prog (N * (n * 2)) (hhl-full-prog A t0 N C Prep) k) *$_v$ unit-vec (N * (n * 2)) 0)‖) ·$_v$*
*post1 (sem N (n * 2) (V ⨂ 1$_m$ 2) (aa-prog (N * (n * 2)) (hhl-full-prog A t0 N C Prep) k) *$_v$ unit-vec (N * (n * 2)) 0)*
*= complex-of-real (1 / ‖post1 out‖) ·$_v$ post1 out*
*∧ calls (aa-prog (N * (n * 2)) (hhl-full-prog A t0 N C Prep) k) = (2 * k + 1) * (2 * (N − 1))*
*∧ real k ≤ pi² / (4 * C * t0)*
⟨*proof*⟩

**end**
**theory** *HHL-Filter*
**imports** *HHL-Complete*
**begin**

# 117 The filter-function algorithm of the HHL paper

Appendix A of the HHL paper replaces the two-level ancilla by a three-level flag register (nothing, well, ill) set to $|h(\tilde{\lambda}_k)\rangle$ by the filter functions $f, g$. We formalise the filter functions, Lemma 2 ($|h(\lambda)\rangle$ is $\frac{\pi}{2}\kappa$-Lipschitz), Lemma 3 (relative Lipschitz bound; the paper's constant $\pi^2/2$ is wrong, the correct constant is $\pi^2$), the circuit with a $d$-level flag, and Theorem 1 parts (1) and (2) on eigenvectors with the window register: errors $O(\kappa/(Nt_0))$.

## 117.1 Filter functions

The filter functions of the HHL paper (A3), with $\kappa' = 2\kappa$: $f(\lambda) = 1/(2\kappa\lambda)$ for $\lambda \geq 1/\kappa$, $\frac{1}{2}\sin(\frac{\pi}{2}(2\kappa\lambda - 1))$ on $[1/2\kappa, 1/\kappa)$, 0 below; $g = 0$, $\frac{1}{2}\cos(\frac{\pi}{2}(2\kappa\lambda - 1))$, $\frac{1}{2}$.

**definition** *filt-f* :: *real* ⇒ *real* ⇒ *real* **where**
*filt-f* $\kappa$ $x$ = (*if* 1 / $\kappa \leq x$ *then* 1 / (2 * $\kappa$ * $x$)
*else if* 1 / (2 * $\kappa$) ≤ $x$ *then sin* (*pi* / 2 * (2 * $\kappa$ * $x$ − 1)) / 2 *else* 0)

**definition** *filt-g* :: *real* ⇒ *real* ⇒ *real* **where**
*filt-g* $\kappa$ $x$ = (*if* 1 / $\kappa \leq x$ *then* 0
*else if* 1 / (2 * $\kappa$) ≤ $x$ *then cos* (*pi* / 2 * (2 * $\kappa$ * $x$ − 1)) / 2 *else* 1 / 2)

**definition** *filt-s* :: *real* ⇒ *real* ⇒ *real* **where**
*filt-s* $\kappa$ $x$ = *sqrt* (1 − (*filt-f* $\kappa$ $x$)$^2$ − (*filt-g* $\kappa$ $x$)$^2$)

The flag state $|h(\lambda)\rangle = s|\text{nothing}\rangle + f|\text{well}\rangle + g|\text{ill}\rangle$.

**definition** *hflag* :: *real* ⇒ *real* ⇒ *complex Matrix.vec* **where**
*hflag* $\kappa$ $x$ = *Matrix.vec* 3 (λ*c*. *complex-of-real* (*if c* = 0 *then filt-s* $\kappa$ $x$ *else if c* = 1 *then filt-f* $\kappa$ $x$ *else filt-g* $\kappa$ $x$))

**lemma** *dim-hflag* [*simp*]: *dim-vec* (*hflag* $\kappa$ $x$) = 3
⟨*proof*⟩

**context**
**fixes** $\kappa$ :: *real*
**assumes** *k0*: 0 < $\kappa$
**begin**

The three regions; on each closed region the functions are given by one formula.

**lemma** *region1*:
**assumes** 1 / $\kappa \leq x$
**shows** *filt-f* $\kappa$ $x$ = 1 / (2 * $\kappa$ * $x$) *filt-g* $\kappa$ $x$ = 0
⟨*proof*⟩

**lemma** *region2*:
**assumes** 1 / (2 * $\kappa$) ≤ $x$ $x \leq$ 1 / $\kappa$
**shows** *filt-f* $\kappa$ $x$ = *sin* (*pi* / 2 * (2 * $\kappa$ * $x$ − 1)) / 2 *filt-g* $\kappa$ $x$ = *cos* (*pi* / 2 * (2 * $\kappa$ * $x$ − 1)) / 2
⟨*proof*⟩

**lemma** *region3*:
**assumes** $x \leq$ 1 / (2 * $\kappa$)
**shows** *filt-f* $\kappa$ $x$ = 0 *filt-g* $\kappa$ $x$ = 1 / 2
⟨*proof*⟩

**lemma** *fg-sq-le*: (*filt-f* $\kappa$ $x$)$^2$ + (*filt-g* $\kappa$ $x$)$^2 \leq$ 1 / 4
⟨*proof*⟩

**lemma** *fg-sq-mid*: 1 / $\kappa \leq x \vee$ (*filt-f* $\kappa$ $x$)$^2$ + (*filt-g* $\kappa$ $x$)$^2$ = 1 / 4

⟨*proof*⟩

**end**

**lemma** *trig-pair-dist*:
**fixes** *a b* :: *real*
**shows** $(sin\ a - sin\ b)^2 + (cos\ a - cos\ b)^2 \leq (a - b)^2$
⟨*proof*⟩

**lemma** *vnorm-diff3-sq*:
**assumes** *dim-vec v* = *3 dim-vec w* = *3*
**shows** $\|v - w\|^2 = (cmod\ (v\ \$\ 0 - w\ \$\ 0))^2 + (cmod\ (v\ \$\ 1 - w\ \$\ 1))^2 + (cmod\ (v\ \$\ 2 - w\ \$\ 2))^2$
⟨*proof*⟩

**context**
**fixes** $\kappa$ :: *real*
**assumes** *k0*: *0* < $\kappa$
**begin**

**lemma** *hflag-dist-sq*:
$\|hflag\ \kappa\ a - hflag\ \kappa\ b\|^2 = (filt\text{-}s\ \kappa\ a - filt\text{-}s\ \kappa\ b)^2 + (filt\text{-}f\ \kappa\ a - filt\text{-}f\ \kappa\ b)^2 + (filt\text{-}g\ \kappa\ a - filt\text{-}g\ \kappa\ b)^2$
⟨*proof*⟩

**lemma** *filt-s-mid*: *1 / κ* ≤ *x* ∨ *filt-s κ x* = *sqrt 3 / 2*
⟨*proof*⟩

Lemma 2 of the paper, region by region.

**lemma** *lip-region1*:
**assumes** *a*: *1 / κ* ≤ *a* **and** *b*: *1 / κ* ≤ *b*
**shows** $\|hflag\ \kappa\ a - hflag\ \kappa\ b\| \leq pi\ /\ 2 * \kappa * |a - b|$
⟨*proof*⟩

**lemma** *lip-region2*:
**assumes** *a*: *1 / (2 * κ)* ≤ *a a* ≤ *1 / κ* **and** *b*: *1 / (2 * κ)* ≤ *b b* ≤ *1 / κ*
**shows** $\|hflag\ \kappa\ a - hflag\ \kappa\ b\| \leq pi\ /\ 2 * \kappa * |a - b|$
⟨*proof*⟩

**lemma** *lip-region3*:
**assumes** *a*: *a* ≤ *1 / (2 * κ)* **and** *b*: *b* ≤ *1 / (2 * κ)*
**shows** $\|hflag\ \kappa\ a - hflag\ \kappa\ b\| \leq pi\ /\ 2 * \kappa * |a - b|$
⟨*proof*⟩

Gluing Lipschitz bounds at a common point.

**lemma** *lip-glue*:
**assumes** *l*: $\bigwedge a\ b.\ a \in S \Longrightarrow b \in S \Longrightarrow \|H\ a - H\ b\| \leq L * |a - b|$
**and** *r*: $\bigwedge a\ b.\ a \in T \Longrightarrow b \in T \Longrightarrow \|H\ a - H\ b\| \leq L * |a - b|$
**and** *q*: *q* ∈ *S q* ∈ *T* **and** *dims*: $\bigwedge x.$ *dim-vec (H x)* = *d*
**and** *a*: *a* ∈ *S* **and** *b*: *b* ∈ *T* **and** *ab*: *a* ≤ *q q* ≤ *b*
**shows** $\|H\ a - H\ b\| \leq L * |a - b|$
⟨*proof*⟩

**end**

**lemma** *abs-split*:
**fixes** *x y z L* :: *real*
**assumes** *x* ≤ *y y* ≤ *z*
**shows** $L * |x - y| + L * |y - z| = L * |x - z|$

⟨*proof*⟩

**context**
  **fixes** $\kappa$ :: *real*
  **assumes** *k0*: $0 < \kappa$
**begin**

**lemma** *lemma2-le*:
  **assumes** *ab*: $a \leq b$
  **shows** $\|hflag\ \kappa\ a - hflag\ \kappa\ b\| \leq pi\ /\ 2 * \kappa * |a - b|$
⟨*proof*⟩

Lemma 2 of the paper: $\lambda \mapsto |h(\lambda)\rangle$ is $\frac{\pi}{2}\kappa$-Lipschitz.

**theorem** *hflag-lipschitz*: $\|hflag\ \kappa\ a - hflag\ \kappa\ b\| \leq pi\ /\ 2 * \kappa * |a - b|$
⟨*proof*⟩

**lemma** *fg-dist-le*: $sqrt\ ((filt\text{-}f\ \kappa\ a - filt\text{-}f\ \kappa\ b)^2 + (filt\text{-}g\ \kappa\ a - filt\text{-}g\ \kappa\ b)^2) \leq pi\ /\ 2 * \kappa * |a - b|$
⟨*proof*⟩

**end**

**context**
  **fixes** $\kappa$ :: *real*
  **assumes** *k0*: $0 < \kappa$
**begin**

**definition** *fg-pt* :: *real* ⇒ *complex* **where**
  *fg-pt x* = *Complex* (*filt-f* $\kappa$ *x*) (*filt-g* $\kappa$ *x*)

**lemma** *fg-pt-dist*: *cmod* (*fg-pt a* − *fg-pt b*) = *sqrt* ((*filt-f* $\kappa$ *a* − *filt-f* $\kappa$ *b*)$^2$ + (*filt-g* $\kappa$ *a* − *filt-g* $\kappa$ *b*)$^2$)
  ⟨*proof*⟩

Lemma 3 of the paper, with the constant $\pi^2$. The paper states $c = \pi^2/2$; that is false: in the middle region $f^2 + g^2 = \frac{1}{4}$ (the proof uses $\frac{1}{2}$) and the ratio of the two sides tends to $\pi^2$.

**theorem** *lemma3*:
  (*filt-f* $\kappa$ *a* − *filt-f* $\kappa$ *b*)$^2$ + (*filt-g* $\kappa$ *a* − *filt-g* $\kappa$ *b*)$^2$
    ≤ *pi*$^2$ * $\kappa^2$ * (*a* − *b*)$^2$ * ((*filt-f* $\kappa$ *a*)$^2$ + (*filt-g* $\kappa$ *a*)$^2$)
⟨*proof*⟩

**end**

## 117.2 A *d*-level flag register

**definition** *hhl-Wd* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
  *hhl-Wd A t0 N d* = *pe-unitary N* (*dim-row A* * *d*) (*evolution A t0* ⨂ $1_m$ *d*)

**definition** *flag-circuit* :: *complex Matrix.mat* ⇒ *real* ⇒ *nat* ⇒ *nat* ⇒ (*nat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat* **where**
  *flag-circuit A t0 N d G* = (*hhl-Wd A t0 N d*)$^\dagger$ * (*ctrl-blocks N* (*dim-row A* * *d*) λ*k*. $1_m$ (*dim-row A*) ⨂ *G k*) * *hhl-Wd A t0 N d*)

**context**
  **fixes** *A* :: *complex Matrix.mat* **and** *n N* :: *nat* **and** *t0 l* :: *real* **and** *u* :: *complex Matrix.vec*
    **and** *psi* :: *complex Matrix.vec* **and** *d* :: *nat* **and** *G* :: *nat* ⇒ *complex Matrix.mat* **and** *hid* :: *complex Matrix.vec*
  **assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *t0*: $0 < t0$ **and** *N*: $0 < N$
    **and** *u*: *dim-vec u* = *n* **and** *ev*: *A* *$_v$ *u* = *complex-of-real l* ·$_v$ *u*
    **and** *psi*: *dim-vec psi* = *N* **and** *d0*: $0 < d$ **and** *G*: ⋀*k*. *k* < *N* ⟹ *G k* ∈ *carrier-mat d d*

**and** *hd*: *dim-vec hid* = *d*
**begin**

The phase of $u$, the register amplitudes and the flag vectors of the branches.

**abbreviation** *dphi* ≡ *hhl-phi t0 l*
**definition** *dalpha* = *Matrix.vec N* (*pe-amp-reg N psi dphi*)
**abbreviation** *fbr k* ≡ *G k* ∗$_v$ *unit-vec d 0*

**lemma** *phi-phase-d*: *cis* (*t0* ∗ *l*) = *cis* (*2* ∗ *pi* ∗ *dphi*)
⟨*proof*⟩

**lemma** *dim-alpha-d* [*simp*]: *dim-vec dalpha* = *N* ⟨*proof*⟩
**lemma** *dim-hid*: *dim-vec hid* = *d* ⟨*proof*⟩
**lemma** *Gblk*: *k* < *N* ⟹ *1*$_m$ *n* ⨂ *G k* ∈ *carrier-mat* (*n* ∗ *d*) (*n* ∗ *d*)
⟨*proof*⟩

**lemma** *W-branch-d*:
**assumes** *w*: *dim-vec w* = *d*
**shows** *hhl-Wd A t0 N d* ∗$_v$ *tensor-vec psi* (*tensor-vec u w*) = *tensor-vec dalpha* (*tensor-vec u w*)
⟨*proof*⟩

**lemma** *hhl-Wd-carrier*: *hhl-Wd A t0 N d* ∈ *carrier-mat* (*N* ∗ (*n* ∗ *d*)) (*N* ∗ (*n* ∗ *d*))
⟨*proof*⟩

**lemma** *hhl-Wd-unitary*: *unitary* (*hhl-Wd A t0 N d*)
⟨*proof*⟩

The output of the circuit and the ideal output as register sums.

**lemma** *flag-output-branches*:
*flag-circuit A t0 N d G* ∗$_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec d 0*)) =
(*hhl-Wd A t0 N d*)$^\dagger$ ∗$_v$ *reg-sum N* (*n* ∗ *d*) (λ*k*. *dalpha* $ *k* ·$_v$ *tensor-vec u* (*fbr k*))
⟨*proof*⟩

**lemma** *flag-ideal-branches*:
*tensor-vec psi* (*tensor-vec u hid*) =
(*hhl-Wd A t0 N d*)$^\dagger$ ∗$_v$ *reg-sum N* (*n* ∗ *d*) (λ*k*. *dalpha* $ *k* ·$_v$ *tensor-vec u hid*)
⟨*proof*⟩

The squared distance between output and ideal output.

The error vector: $W^* \sum_k \alpha_k |k\rangle \otimes u \otimes (R(f_k)|0\rangle - R(c)|0\rangle)$.

**lemma** *flag-err-vec*:
*flag-circuit A t0 N d G* ∗$_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec d 0*)) − *tensor-vec psi* (*tensor-vec u hid*) =
(*hhl-Wd A t0 N d*)$^\dagger$ ∗$_v$ *reg-sum N* (*n* ∗ *d*) (λ*k*. *dalpha* $ *k* ·$_v$ *tensor-vec u* (*fbr k* − *hid*))
⟨*proof*⟩

The squared distance between output and ideal output.

**theorem** *flag-error*:
⟨*flag-circuit A t0 N d G* ∗$_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec d 0*)) − *tensor-vec psi* (*tensor-vec u hid*) |
*flag-circuit A t0 N d G* ∗$_v$ *tensor-vec psi* (*tensor-vec u* (*unit-vec d 0*)) − *tensor-vec psi* (*tensor-vec u hid*)⟩ =
⟨*u*|*u*⟩ ∗ (∑ *k*<*N*. *complex-of-real* ((*cmod* (*pe-amp-reg N psi dphi k*))$^2$) ∗ ⟨*fbr k* − *hid*|*fbr k* − *hid*⟩)
⟨*proof*⟩

The branch distance in real terms.

**end**

## 117.3 Window weights with wrapped distances

**lemma** *pe-amp-reg-int-shift*: *pe-amp-reg N psi* ($\varphi$ − *real-of-int m*) *j* = *pe-amp-reg N psi* $\varphi$ *j*
⟨*proof*⟩

**definition** *wrap* :: *real* ⇒ *real* **where**
*wrap x* = *x* − *real-of-int* ⌈*x* − *1* / *2*⌉

**lemma** *wrap-range*: − (*1* / *2*) < *wrap x wrap x* ≤ *1* / *2*
⟨*proof*⟩

**lemma** *wrap-int*: *x* − *wrap x* ∈ ℤ
⟨*proof*⟩

The amplitude in terms of the wrapped distance $r = \mathrm{wrap}(\varphi - k/N)$.

**lemma** *win-amp-wrapped*:
**fixes** $\varphi$ :: *real*
**assumes** *N*: *2* ≤ *N* **and** *k*: *k* < *N*
**defines** *r* ≡ *wrap* ($\varphi$ − *real k* / *real N*)
**shows** (*cmod* (*pe-amp-reg N* (*win-state N*) $\varphi$ *k*))$^2$ ≤ *2*
**and** *1* / *real N* ≤ |*r*| ⟹ (*cmod* (*pe-amp-reg N* (*win-state N*) $\varphi$ *k*))$^2$ ≤ *pi*$^2$ / (*2* ∗ (*real N*) ^ *4* ∗ *r* ^ *4*)
⟨*proof*⟩

**lemma** *win-amp-weight*:
**fixes** $\varphi$ :: *real*
**assumes** *N*: *2* ≤ *N* **and** *k*: *k* < *N*
**defines** *r* ≡ *wrap* ($\varphi$ − *real k* / *real N*)
**shows** (*cmod* (*pe-amp-reg N* (*win-state N*) $\varphi$ *k*))$^2$ ∗ *r*$^2$ ≤ *win-w* (*real N* ∗ *r*) / (*real N*)$^2$
⟨*proof*⟩

Points with pairwise non-zero integer differences.

**lemma** *spaced-floor-inj*:
**fixes** *z* :: *nat* ⇒ *real*
**assumes** *sp*: ⋀*i j*. *i* ∈ *K* ⟹ *j* ∈ *K* ⟹ *i* ≠ *j* ⟹ *z i* − *z j* ∈ ℤ ∧ *z i* ≠ *z j*
**shows** *inj-on* (λ*k*. ⌊*z k*⌋) *K inj-on* (λ*k*. ⌊− *z k*⌋) *K*
⟨*proof*⟩

**lemma** *spaced-weight-sum*:
**fixes** *z* :: *nat* ⇒ *real*
**assumes** *K*: *finite K* **and** *sp*: ⋀*i j*. *i* ∈ *K* ⟹ *j* ∈ *K* ⟹ *i* ≠ *j* ⟹ *z i* − *z j* ∈ ℤ ∧ *z i* ≠ *z j*
**shows** (∑ *k*∈*K*. *win-w* (*z k*)) ≤ *4* + *2* ∗ *pi*$^2$
⟨*proof*⟩

**lemma** *reg-e-bound*:
**assumes** *t0*: *0* < *t0* **and** *N*: *0* < *N* **and** *k*: *k* < *N* **and** *ph*: |$\varphi$| ≤ *1* / *4*
**shows** |*t0* ∗ *reg-eig t0 N k* / (*2* ∗ *pi*) − $\varphi$| ≤ *3* ∗ |*wrap* ($\varphi$ − *real k* / *real N*)|
⟨*proof*⟩

**lemma** *wrap-spaced*:
**assumes** *N*: *0* < *N* **and** *i*: *i* < *N* **and** *j*: *j* < *N* **and** *ij*: *i* ≠ *j*
**shows** *real N* ∗ *wrap* ($\varphi$ − *real i* / *real N*) − *real N* ∗ *wrap* ($\varphi$ − *real j* / *real N*) ∈ ℤ
∧ *real N* ∗ *wrap* ($\varphi$ − *real i* / *real N*) ≠ *real N* ∗ *wrap* ($\varphi$ − *real j* / *real N*)
⟨*proof*⟩

Theorem 1(1) of the paper on an eigenvector, for the flag circuit with the window register: the squared error is at most $9\pi^4(4 + 2\pi^2)\kappa^2/(Nt_0)^2$, i.e. $O(\kappa/t)$ with $t = Nt_0$ the total evolution time.

**theorem** *flag-window-eigen*:
**fixes** $A$ :: *complex Matrix.mat* **and** $u$ :: *complex Matrix.vec*
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *t0*: $0 < t0$ **and** *N*: $2 \leq N$
**and** *u*: *dim-vec u* $= n$ **and** *ev*: $A *_v u =$ *complex-of-real l* $\cdot_v u$
**and** *k0*: $0 < \kappa$ **and** *ph*: $|t0 * l| \leq pi / 2$
**and** *G*: $\bigwedge k.\ k < N \Longrightarrow G\ k \in$ *carrier-mat 3 3*
**and** *Gh*: $\bigwedge k.\ k < N \Longrightarrow G\ k *_v$ *unit-vec 3 0* $=$ *hflag* $\kappa$ (*reg-eig t0 N k*)
**shows** *Re* ⟨*flag-circuit A t0 N 3 G* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec u* (*unit-vec 3 0*)) − *tensor-vec* (*win-state N*) (*tensor-vec u* (*hflag* $\kappa$ *l*)) |
*flag-circuit A t0 N 3 G* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec u* (*unit-vec 3 0*)) − *tensor-vec* (*win-state N*) (*tensor-vec u* (*hflag* $\kappa$ *l*))⟩
$\leq$ *Re* ⟨*u*|*u*⟩ $*$ (*9* $*$ *pi* ^ *4* $*$ (*4* + *2* $*$ *pi*$^2$) $*$ $\kappa^2$ / ((*real N*)$^2$ $*$ *t0*$^2$))
⟨*proof*⟩

## 117.4 Post-selection on the flag register

The flag is the last tensor factor, of dimension *d*: *pflag d P v* keeps the components whose flag value *c* satisfies *P c*.

**definition** *pflag* :: *nat* ⇒ (*nat* ⇒ *bool*) ⇒ *complex Matrix.vec* ⇒ *complex Matrix.vec* **where**
*pflag d P v* = *Matrix.vec* (*dim-vec v*) (λ*r*. *if P* (*r mod d*) *then v* $ *r else 0*)

**lemma** *dim-pflag* [*simp*]: *dim-vec* (*pflag d P v*) = *dim-vec v*
⟨*proof*⟩

**lemma** *pflag-minus*: *dim-vec u* = *dim-vec v* ⟹ *pflag d P* (*u* − *v*) = *pflag d P u* − *pflag d P v*
⟨*proof*⟩

**lemma** *vnorm-pflag-le*: ‖*pflag d P v*‖ ≤ ‖*v*‖
⟨*proof*⟩

Matrices that do not mix flag values.

**definition** *flagdiag* :: *nat* ⇒ *complex Matrix.mat* ⇒ *bool* **where**
*flagdiag d M* ⟷ (∀ *r s*. *r* < *dim-row M* ⟶ *s* < *dim-col M* ⟶ *r mod d* ≠ *s mod d* ⟶ *M* $$ (*r*, *s*) = *0*)

**lemma** *flagdiag-mult*:
**assumes** *flagdiag d A flagdiag d B dim-col A* = *dim-row B*
**shows** *flagdiag d* (*A* $*$ *B*)
⟨*proof*⟩

**lemma** *flagdiag-dagger*: *flagdiag d M* ⟹ *flagdiag d* ($M^\dagger$)
⟨*proof*⟩

**lemma** *flagdiag-one*: *flagdiag d* (*1*$_m$ *K*)
⟨*proof*⟩

**lemma** *flagdiag-pow*: *flagdiag d M* ⟹ *M* ∈ *carrier-mat K K* ⟹ *flagdiag d* (*M* ^$_m$ *j*)
⟨*proof*⟩

**lemma** *pflag-commute*:
**assumes** *M*: *M* ∈ *carrier-mat K K* **and** *f*: *flagdiag d M* **and** *v*: *dim-vec v* = *K*
**shows** *pflag d P* (*M* $*_v$ *v*) = *M* $*_v$ *pflag d P v*
⟨*proof*⟩

Tensoring with $I_d$ on the right and block operators on top of it do not mix flag values.

**lemma** *flagdiag-tensor-one*:
**assumes** *0* < *d*

**shows** *flagdiag d* ($X \bigotimes 1_m$ *d*)
⟨*proof*⟩

**lemma** *flagdiag-ctrl-blocks*:
**assumes** *d0*: *0* < *d* **and** *B*: ⋀*k*. *k* < *N* ⟹ *flagdiag d* (*B k*) ∧ *B k* ∈ *carrier-mat* (*n* ∗ *d*) (*n* ∗ *d*)
**shows** *flagdiag d* (*ctrl-blocks N* (*n* ∗ *d*) *B*)
⟨*proof*⟩

**lemma** *flagdiag-dvd*: *flagdiag D M* ⟹ *d dvd D* ⟹ *flagdiag d M*
⟨*proof*⟩

**lemma** *hhl-Wd-flagdiag*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *n0*: *0* < *n*
**shows** *flagdiag 3* (*hhl-Wd A t0 N 3*)
⟨*proof*⟩

**lemma** *pflag-reg-sum*:
**assumes** *f*: ⋀*k*. *k* < *N* ⟹ *dim-vec* (*f k*) = *n* ∗ *d* **and** *d0*: *0* < *d*
**shows** *pflag d P* (*reg-sum N* (*n* ∗ *d*) *f*) = *reg-sum N* (*n* ∗ *d*) (λ*k*. *pflag d P* (*f k*))
⟨*proof*⟩

**lemma** *pflag-tensor*:
**assumes** *u*: *dim-vec u* = *n* **and** *w*: *dim-vec w* = *d* **and** *d0*: *0* < *d*
**shows** *pflag d P* (*tensor-vec u w*) = *tensor-vec u* (*pflag d P w*)
⟨*proof*⟩

**lemma** *pflag-smult*: *pflag d P* (*c* ·$_v$ *v*) = *c* ·$_v$ *pflag d P v*
⟨*proof*⟩

The norm of a branch sum behind a unitary.

**lemma** *unitary-reg-sum-norm*:
**assumes** *W*: *W* ∈ *carrier-mat* (*N* ∗ (*n* ∗ *d*)) (*N* ∗ (*n* ∗ *d*)) *unitary W*
**and** *u*: *dim-vec u* = *n* **and** *w*: ⋀*k*. *k* < *N* ⟹ *dim-vec* (*w k*) = *d*
**shows** ⟨*W*$^\dagger$ ∗$_v$ *reg-sum N* (*n* ∗ *d*) (λ*k*. *a k* ·$_v$ *tensor-vec u* (*w k*))|*W*$^\dagger$ ∗$_v$ *reg-sum N* (*n* ∗ *d*) (λ*k*. *a k* ·$_v$ *tensor-vec u* (*w k*))⟩
= ⟨*u*|*u*⟩ ∗ (∑ *k*<*N*. *complex-of-real* ((*cmod* (*a k*))$^2$) ∗ ⟨*w k*|*w k*⟩)
⟨*proof*⟩

The window-weighted spread of the register eigenvalues around $\lambda$: $\sum_k |\alpha_k|^2(\tilde{\lambda}_k - \lambda)^2 \leq 36\pi^2(4+2\pi^2)/(Nt_0)^2$.

**lemma** *window-spread*:
**assumes** *t0*: *0* < *t0* **and** *N*: *2* ≤ *N* **and** *ph*: |*t0* ∗ *l*| ≤ *pi* / *2*
**shows** (∑ *k*<*N*. (*cmod* (*pe-amp-reg N* (*win-state N*) (*hhl-phi t0 l*) *k*))$^2$ ∗ (*reg-eig t0 N k* − *l*)$^2$)
≤ *36* ∗ *pi*$^2$ ∗ (*4* + *2* ∗ *pi*$^2$) / ((*real N*)$^2$ ∗ *t0*$^2$)
⟨*proof*⟩

**lemma** *vnorm3-sq*: *dim-vec v* = *3* ⟹ ‖*v*‖$^2$ = (*cmod* (*v* $ *0*))$^2$ + (*cmod* (*v* $ *1*))$^2$ + (*cmod* (*v* $ *2*))$^2$
⟨*proof*⟩

**lemma** *pflag3-entries*:
**assumes** *dim-vec v* = *3*
**shows** *pflag 3* (λ*c*. *c* ≠ *0*) *v* $ *0* = *0 pflag 3* (λ*c*. *c* ≠ *0*) *v* $ *1* = *v* $ *1 pflag 3* (λ*c*. *c* ≠ *0*) *v* $ *2* = *v* $ *2*
⟨*proof*⟩

**lemma** *hflag-entries*: *hflag* κ *x* $ *1* = *complex-of-real* (*filt-f* κ *x*) *hflag* κ *x* $ *2* = *complex-of-real* (*filt-g* κ *x*)
⟨*proof*⟩

**lemma** *pflag-flag-dist*:
  $\|$*pflag 3* ($\lambda$*c. c* $\neq$ *0*) (*hflag* $\kappa$ *a* − *hflag* $\kappa$ *b*)$\|^2$ = (*filt-f* $\kappa$ *a* − *filt-f* $\kappa$ *b*)$^2$ + (*filt-g* $\kappa$ *a* − *filt-g* $\kappa$ *b*)$^2$
⟨*proof*⟩

**lemma** *pflag-flag-norm*:
  $\|$*pflag 3* ($\lambda$*c. c* $\neq$ *0*) (*hflag* $\kappa$ *a*)$\|^2$ = (*filt-f* $\kappa$ *a*)$^2$ + (*filt-g* $\kappa$ *a*)$^2$
  ⟨*proof*⟩

**lemma** *pflag-tensor-left*:
  **assumes** *a*: *dim-vec a* = *N* **and** *z*: *dim-vec z* = *m* ∗ *d* **and** *d0*: *0* < *d*
  **shows** *pflag d P* (*tensor-vec a z*) = *tensor-vec a* (*pflag d P z*)
⟨*proof*⟩

Theorem 1(2) on an eigenvector: after post-selecting the flag on $\{\text{well}, \text{ill}\}$ the error is at most $\rho$ times the post-selected ideal, $\rho^2 = 36\pi^4(4 + 2\pi^2)\kappa^2/(Nt_0)^2$, by Lemma 3 (relative error) and the window spread.

**theorem** *flag-post-eigen*:
  **fixes** *A* :: *complex Matrix.mat* **and** *u* :: *complex Matrix.vec*
  **assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *n0*: *0* < *n* **and** *t0*: *0* < *t0* **and** *N*: *2* ≤ *N*
    **and** *u*: *dim-vec u* = *n* **and** *ev*: *A* ∗$_v$ *u* = *complex-of-real l* ·$_v$ *u*
    **and** *k0*: *0* < $\kappa$ **and** *ph*: |*t0* ∗ *l*| ≤ *pi* / *2*
    **and** *G*: $\bigwedge$*k. k* < *N* ⟹ *G k* ∈ *carrier-mat 3 3*
    **and** *Gh*: $\bigwedge$*k. k* < *N* ⟹ *G k* ∗$_v$ *unit-vec 3 0* = *hflag* $\kappa$ (*reg-eig t0 N k*)
  **defines** *P* ≡ $\lambda$*c*::*nat. c* ≠ *0*
  **shows** $\|$*pflag 3 P* (*flag-circuit A t0 N 3 G* ∗$_v$ *tensor-vec* (*win-state N*) (*tensor-vec u* (*unit-vec 3 0*)))
      − *pflag 3 P* (*tensor-vec* (*win-state N*) (*tensor-vec u* (*hflag* $\kappa$ *l*)))$\|^2$
    ≤ *36* ∗ *pi* ^ *4* ∗ (*4* + *2* ∗ *pi*$^2$) ∗ $\kappa^2$ / ((*real N*)$^2$ ∗ *t0*$^2$) ∗ $\|$*pflag 3 P* (*tensor-vec* (*win-state N*) (*tensor-vec u* (*hflag* $\kappa$ *l*)))$\|^2$
   **and** $\|$*pflag 3 P* (*tensor-vec* (*win-state N*) (*tensor-vec u* (*hflag* $\kappa$ *l*)))$\|^2$ = $\|u\|^2$ ∗ ((*filt-f* $\kappa$ *l*)$^2$ + (*filt-g* $\kappa$ *l*)$^2$)
⟨*proof*⟩

**end**
**theory** *HHL-Filter-Bounds*
  **imports** *HHL-Filter*
**begin**

# 118 Theorem 1 of the HHL paper for every right-hand side

The error vectors of distinct eigenvectors are orthogonal, also after post-selecting the flag, so the eigenvector bounds of *HHL-Filter* extend to every $b$ with no loss. This gives all three parts of Theorem 1 of the paper: (1) $\|(\tilde{U} - U)b\| = O(\kappa/t)\|b\|$, (2) the state post-selected on $\{\text{well}, \text{ill}\}$ is within $O(\kappa/t)$ of the ideal one, (3) if $b$ has no ill-conditioned component the state post-selected on $|\text{well}\rangle$ is within $O(\kappa/t)$ of $f(A)b/\|f(A)b\|$, which is $\propto A^{-1}b$ when $A$ is well conditioned; here $t = Nt_0$ is the total evolution time. Householder reflections provide the flag gates.

## 118.1 Every right-hand side

$b \mapsto a \otimes \sum_{c<3} H_c\, b \otimes |c\rangle$ as an $(\dim a \cdot 3n) \times n$ matrix.

**definition** *lift3* :: *nat* ⇒ *complex Matrix.vec* ⇒ (*nat* ⇒ *complex Matrix.mat*) ⇒ *complex Matrix.mat*
**where**
  *lift3 n a H* = *Matrix.mat* (*dim-vec a* ∗ (*n* ∗ *3*)) *n*
    ($\lambda$(*r*, *s*). *a* $ (*r div* (*n* ∗ *3*)) ∗ *H* (*r mod* (*n* ∗ *3*) *mod 3*) $$ (*r mod* (*n* ∗ *3*) *div 3*, *s*))

**lemma** *lift3-carrier* [*simp*]: *lift3 n a H* ∈ *carrier-mat* (*dim-vec a* ∗ (*n* ∗ *3*)) *n*
⟨*proof*⟩

**lemma** *lift3-mult-vec*:
**assumes** *H*: ⋀*c*. *c* < *3* ⟹ *H c* ∈ *carrier-mat n n* **and** *v*: *dim-vec v* = *n*
**shows** *lift3 n a H* ∗*v v* = *tensor-vec a* (*Matrix.vec* (*n* ∗ *3*) (λ*j*. (*H* (*j mod 3*) ∗*v v*) $ (*j div 3*)))
⟨*proof*⟩

**lemma** *flag-vec-eigen*:
**assumes** *u*: *dim-vec u* = *n*
**shows** *Matrix.vec* (*n* ∗ *3*) (λ*j*. (*z* (*j mod 3*) ·*v u*) $ (*j div 3*)) = *tensor-vec u* (*Matrix.vec 3 z*)
⟨*proof*⟩

**lemma** *flag-vec-split*:
**assumes** *H*: ⋀*c*. *c* < *3* ⟹ *H c* ∈ *carrier-mat n n* **and** *v*: *dim-vec v* = *n*
**shows** *Matrix.vec* (*n* ∗ *3*) (λ*j*. (*H* (*j mod 3*) ∗*v v*) $ (*j div 3*)) =
*tensor-vec* (*H 0* ∗*v v*) (*unit-vec 3 0*) + *tensor-vec* (*H 1* ∗*v v*) (*unit-vec 3 1*) + *tensor-vec* (*H 2* ∗*v v*) (*unit-vec 3 2*)
⟨*proof*⟩

**definition** *flag-H* :: *real* ⇒ *complex Matrix.mat* ⇒ *nat* ⇒ *complex Matrix.mat* **where**
*flag-H* κ *A c* = *mat-fun* (λ*x*. *complex-of-real* (*if c* = *0 then filt-s* κ *x else if c* = *1 then filt-f* κ *x else filt-g* κ *x*)) *A*

**definition** *flag-idl* :: *nat* ⇒ *real* ⇒ *complex Matrix.mat* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.mat* **where**
*flag-idl n* κ *A a* = *lift3 n a* (*flag-H* κ *A*)

**definition** *flag-emb* :: *nat* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.mat* **where**
*flag-emb n a* = *lift3 n a* (λ*c*. *if c* = *0 then* $1_m$ *n else* $0_m$ *n n*)

**lemma** *flag-H-carrier*: *A* ∈ *carrier-mat n n* ⟹ *hermitian A* ⟹ *flag-H* κ *A c* ∈ *carrier-mat n n*
⟨*proof*⟩

**lemma** *flag-emb-mult*:
**assumes** *v*: *dim-vec v* = *n*
**shows** *flag-emb n a* ∗*v v* = *tensor-vec a* (*tensor-vec v* (*unit-vec 3 0*))
⟨*proof*⟩

**lemma** *flag-idl-eigen*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *u*: *dim-vec u* = *n* **and** *ev*: *A* ∗*v u* = *complex-of-real l* ·*v u*
**shows** *flag-idl n* κ *A a* ∗*v u* = *tensor-vec a* (*tensor-vec u* (*hflag* κ *l*))
⟨*proof*⟩

**lemma** *flag-idl-mult*:
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *b*: *dim-vec b* = *n*
**shows** *flag-idl n* κ *A a* ∗*v b* = *tensor-vec a*
(*tensor-vec* (*flag-H* κ *A 0* ∗*v b*) (*unit-vec 3 0*) + *tensor-vec* (*flag-H* κ *A 1* ∗*v b*) (*unit-vec 3 1*)
+ *tensor-vec* (*flag-H* κ *A 2* ∗*v b*) (*unit-vec 3 2*))
⟨*proof*⟩

The post-selected error vector as a branch sum behind $W^*$.

**lemma** *flag-perr*:
**fixes** *A* :: *complex Matrix.mat* **and** *u* :: *complex Matrix.vec*
**assumes** *A*: *A* ∈ *carrier-mat n n hermitian A* **and** *n0*: *0* < *n* **and** *t0*: *0* < *t0* **and** *N0*: *0* < *N*
**and** *u*: *dim-vec u* = *n* **and** *ev*: *A* ∗*v u* = *complex-of-real l* ·*v u* **and** *psi*: *dim-vec psi* = *N*
**and** *G*: ⋀*k*. *k* < *N* ⟹ *G k* ∈ *carrier-mat 3 3* **and** *h*: *dim-vec h* = *3*

**shows** *pflag 3 P (flag-circuit A t0 N 3 G *$_v$ tensor-vec psi (tensor-vec u (unit-vec 3 0)) − tensor-vec psi (tensor-vec u h))*
*= (hhl-Wd A t0 N 3)*$^\dagger$ **$_v$ reg-sum N (n * 3)*
*(λk. pe-amp-reg N psi (hhl-phi t0 l) k ·$_v$ tensor-vec u (pflag 3 P (G k *$_v$ unit-vec 3 0 − h)))*
⟨*proof*⟩

Error vectors of orthogonal eigenvectors stay orthogonal after post-selection.

**lemma** *flag-perr-orth*:
**fixes** *A* :: *complex Matrix.mat* **and** *u u′* :: *complex Matrix.vec*
**assumes** *A*: *A ∈ carrier-mat n n hermitian A* **and** *n0*: *0 < n* **and** *t0*: *0 < t0* **and** *N0*: *0 < N*
**and** *u*: *dim-vec u = n* **and** *ev*: *A *$_v$ u = complex-of-real l ·$_v$ u*
**and** *u′*: *dim-vec u′ = n* **and** *ev′*: *A *$_v$ u′ = complex-of-real l′ ·$_v$ u′*
**and** *orth*: *⟨u|u′⟩ = 0* **and** *psi*: *dim-vec psi = N*
**and** *G*: *⋀k. k < N ⟹ G k ∈ carrier-mat 3 3* **and** *h*: *dim-vec h = 3* **and** *h′*: *dim-vec h′ = 3*
**shows** *⟨pflag 3 P (flag-circuit A t0 N 3 G *$_v$ tensor-vec psi (tensor-vec u (unit-vec 3 0)) − tensor-vec psi (tensor-vec u h)) |*
*pflag 3 P (flag-circuit A t0 N 3 G *$_v$ tensor-vec psi (tensor-vec u′ (unit-vec 3 0)) − tensor-vec psi (tensor-vec u′ h′))⟩ = 0*
⟨*proof*⟩

**lemma** *inner-as-sum*: *dim-vec y = n ⟹ ⟨x|y⟩ = (∑ i<n. cnj (x $ i) * y $ i)*
⟨*proof*⟩

If $M$ maps the orthonormal columns $u_j$ of $U$ to mutually orthogonal vectors then $\|Mb\|^2 = \sum_j \|Mu_j\|^2 \, |\langle u_j|b\rangle|^2$.

**lemma** *orth-image-eq*:
**fixes** *M U* :: *complex Matrix.mat*
**assumes** *U*: *U ∈ carrier-mat n n unitary U* **and** *M*: *M ∈ carrier-mat K n*
**and** *orth*: *⋀i j. i < n ⟹ j < n ⟹ i ≠ j ⟹ ⟨M *$_v$ Matrix.col U i|M *$_v$ Matrix.col U j⟩ = 0*
**and** *b*: *dim-vec b = n*
**shows** *‖M *$_v$ b‖² = (∑ j<n. ‖M *$_v$ Matrix.col U j‖² * (cmod (((U†) *$_v$ b) $ j))²)*
⟨*proof*⟩

**lemma** *unitary-dagger-norm*:
**assumes** *U*: *U ∈ carrier-mat n n unitary U* **and** *b*: *dim-vec b = n*
**shows** *(∑ j<n. (cmod (((U†) *$_v$ b) $ j))²) = ‖b‖²*
⟨*proof*⟩

**definition** *pflag-mat* :: *nat ⇒ nat ⇒ (nat ⇒ bool) ⇒ complex Matrix.mat* **where**
*pflag-mat K d P = Matrix.mat K K (λ(r, s). if r = s ∧ P (r mod d) then 1 else 0)*

**lemma** *pflag-mat-carrier* [*simp*]: *pflag-mat K d P ∈ carrier-mat K K*
⟨*proof*⟩

**lemma** *pflag-mat-mult*: *dim-vec v = K ⟹ pflag-mat K d P *$_v$ v = pflag d P v*
⟨*proof*⟩

**lemma** *pflag-all*: *pflag d (λ-. True) v = v*
⟨*proof*⟩

Theorem 1(1) for every right-hand side: $\|(\tilde{U}-U)b\| \le 3\pi^2\sqrt{4+2\pi^2}\,\kappa\|b\|/(Nt_0)$, i.e. $\|\tilde{U}-U\| = O(\kappa/t)$ for the total evolution time $t = Nt_0$.

**theorem** *flag-bound*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: *A ∈ carrier-mat n n hermitian A* **and** *n0*: *0 < n* **and** *t0*: *0 < t0* **and** *N*: *2 ≤ N*
**and** *k0*: *0 < κ* **and** *spec*: *⋀l. l ∈ herm-spectrum A ⟹ |t0 * l| ≤ pi / 2*
**and** *G*: *⋀k. k < N ⟹ G k ∈ carrier-mat 3 3*

**and** *Gh*: $\bigwedge k.\ k < N \Longrightarrow G\ k *_v$ *unit-vec 3 0* $=$ *hflag* $\kappa$ (*reg-eig t0 N k*)
**and** *b*: *dim-vec b* $= n$
**shows** $\|$*flag-circuit A t0 N 3 G* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec b* (*unit-vec 3 0*)) $-$ *flag-idl n* $\kappa$ *A* (*win-state N*) $*_v$ *b*$\|^2$
$\leq$ *9* $*$ *pi* ^ *4* $*$ (*4* $+$ *2* $*$ *pi*$^2$) $*$ $\kappa^2$ / ((*real N*)$^2$ $*$ *t0*$^2$) $*$ $\|b\|^2$
⟨*proof*⟩

**lemma** *sum-factor3*: $(\sum j{\in}S.$ (*r*::*real*) $*$ *a j* $*$ *c j*) $=$ *r* $*$ $(\sum j{\in}S.$ *a j* $*$ *c j*)
⟨*proof*⟩

Theorem 1(2) for every right-hand side: post-selecting the flag on $\{\text{well}, \text{ill}\}$ the error is at most $\rho$ times the post-selected ideal, $\rho = 6\pi^2\sqrt{4+2\pi^2}\,\kappa/(Nt_0)$.

**theorem** *flag-post-bound*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: *A* $\in$ *carrier-mat n n hermitian A* **and** *n0*: *0* $<$ *n* **and** *t0*: *0* $<$ *t0* **and** *N*: *2* $\leq$ *N*
**and** *k0*: *0* $<$ $\kappa$ **and** *spec*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow$ $|$*t0* $*$ *l*$| \leq$ *pi* / *2*
**and** *G*: $\bigwedge k.\ k < N \Longrightarrow G\ k \in$ *carrier-mat 3 3*
**and** *Gh*: $\bigwedge k.\ k < N \Longrightarrow G\ k *_v$ *unit-vec 3 0* $=$ *hflag* $\kappa$ (*reg-eig t0 N k*)
**and** *b*: *dim-vec b* $= n$
**defines** $P \equiv \lambda c{::}$*nat*. *c* $\neq$ *0*
**shows** $\|$*pflag 3 P* (*flag-circuit A t0 N 3 G* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec b* (*unit-vec 3 0*))) $-$ *pflag 3 P* (*flag-idl n* $\kappa$ *A* (*win-state N*) $*_v$ *b*)$\|^2$
$\leq$ *36* $*$ *pi* ^ *4* $*$ (*4* $+$ *2* $*$ *pi*$^2$) $*$ $\kappa^2$ / ((*real N*)$^2$ $*$ *t0*$^2$) $*$ $\|$*pflag 3 P* (*flag-idl n* $\kappa$ *A* (*win-state N*) $*_v$ *b*)$\|^2$
⟨*proof*⟩

**lemma** *normalized-close*:
**assumes** *d*: *dim-vec a* $=$ *dim-vec c* **and** *cl*: $\|a - c\| \leq \varrho * \|c\|$ **and** *r1*: $\varrho <$ *1* **and** *c0*: $\|c\| \neq$ *0*
**shows** $\|$*complex-of-real* (*1* / $\|a\|$) $\cdot_v$ *a* $-$ *complex-of-real* (*1* / $\|c\|$) $\cdot_v$ *c*$\| \leq$ *2* $*$ $\varrho$
⟨*proof*⟩

Theorem 1(2), normalised: the post-selected state is within $2\rho$ of the ideal one.

**corollary** *flag-post-normalized*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: *A* $\in$ *carrier-mat n n hermitian A* **and** *n0*: *0* $<$ *n* **and** *t0*: *0* $<$ *t0* **and** *N*: *2* $\leq$ *N*
**and** *k0*: *0* $<$ $\kappa$ **and** *spec*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow$ $|$*t0* $*$ *l*$| \leq$ *pi* / *2*
**and** *G*: $\bigwedge k.\ k < N \Longrightarrow G\ k \in$ *carrier-mat 3 3*
**and** *Gh*: $\bigwedge k.\ k < N \Longrightarrow G\ k *_v$ *unit-vec 3 0* $=$ *hflag* $\kappa$ (*reg-eig t0 N k*)
**and** *b*: *dim-vec b* $= n$
**defines** $P \equiv \lambda c{::}$*nat*. *c* $\neq$ *0* **and** $\varrho \equiv$ *6* $*$ *pi*$^2$ $*$ *sqrt* (*4* $+$ *2* $*$ *pi*$^2$) $*$ $\kappa$ / (*real N* $*$ *t0*)
**defines** *y* $\equiv$ *pflag 3 P* (*flag-circuit A t0 N 3 G* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec b* (*unit-vec 3 0*)))
**and** *x* $\equiv$ *pflag 3 P* (*flag-idl n* $\kappa$ *A* (*win-state N*) $*_v$ *b*)
**assumes** *r1*: $\varrho <$ *1* **and** *x0*: $\|x\| \neq$ *0*
**shows** $\|$*complex-of-real* (*1* / $\|y\|$) $\cdot_v$ *y* $-$ *complex-of-real* (*1* / $\|x\|$) $\cdot_v$ *x*$\| \leq$ *2* $*$ $\varrho$
⟨*proof*⟩

**lemma** *pflag-sub*: $(\bigwedge c.\ Q\ c \Longrightarrow P\ c) \Longrightarrow$ *pflag d Q v* $=$ *pflag d Q* (*pflag d P v*)
⟨*proof*⟩

**lemma** *lift3-post*:
**assumes** *H*: $\bigwedge c.\ c < 3 \Longrightarrow H\ c \in$ *carrier-mat n n* **and** *v*: *dim-vec v* $= n$ **and** *a*: *dim-vec a* $= N$
**shows** *pflag 3* ($\lambda c.\ c = 1$) (*lift3 n a H* $*_v$ *v*) $=$ *tensor-vec a* (*tensor-vec* (*H 1* $*_v$ *v*) (*unit-vec 3 1*))
**and** *H 2* $*_v$ *v* $= 0_v$ *n* $\Longrightarrow$ *pflag 3* ($\lambda c.\ c \neq 0$) (*lift3 n a H* $*_v$ *v*) $=$ *pflag 3* ($\lambda c.\ c = 1$) (*lift3 n a H* $*_v$ *v*)
⟨*proof*⟩

Theorem 1(3): if $b$ has no ill-conditioned component ($g(A)b = 0$), the state post-selected on

$|\text{well}\rangle$ is within $2\rho$ of $\psi \otimes f(A)b/\|f(A)b\| \otimes |\text{well}\rangle$.

**theorem** *flag-well-normalized*:
**fixes** *A* :: *complex Matrix.mat* **and** *b* :: *complex Matrix.vec*
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *n0*: $0 < n$ **and** *t0*: $0 < t0$ **and** *N*: $2 \leq N$
**and** *k0*: $0 < \kappa$ **and** *spec*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow |t0 * l| \leq pi\ /\ 2$
**and** *G*: $\bigwedge k.\ k < N \Longrightarrow G\ k \in$ *carrier-mat 3 3*
**and** *Gh*: $\bigwedge k.\ k < N \Longrightarrow G\ k *_v$ *unit-vec 3 0* $=$ *hflag* $\kappa$ (*reg-eig t0 N k*)
**and** *b*: *dim-vec b* $= n$ **and** *ill*: *flag-H* $\kappa$ *A 2* $*_v$ *b* $= 0_v$ *n* **and** *well*: *flag-H* $\kappa$ *A 1* $*_v$ *b* $\neq 0_v$ *n*
**defines** $\varrho \equiv 6 * pi^2 *$ *sqrt* $(4 + 2 * pi^2) * \kappa\ /$ (*real N* $*$ *t0*)
**defines** $y \equiv$ *pflag 3* $(\lambda c.\ c = 1)$ (*flag-circuit A t0 N 3 G* $*_v$ *tensor-vec* (*win-state N*) (*tensor-vec b* (*unit-vec 3 0*)))
**and** $x \equiv$ *tensor-vec* (*win-state N*) (*tensor-vec* (*flag-H* $\kappa$ *A 1* $*_v$ *b*) (*unit-vec 3 1*))
**assumes** *r1*: $\varrho < 1$
**shows** $\|$*complex-of-real* $(1\ /\ \|y\|) \cdot_v y -$ *complex-of-real* $(1\ /\ \|x\|) \cdot_v x\| \leq 2 * \varrho$
⟨*proof*⟩

If every eigenvalue is at least $1/\kappa$ then $g(A) = 0$ and $f(A) = A^{-1}/(2\kappa)$: the post-selected state is close to $\psi \otimes A^{-1}b/\|A^{-1}b\| \otimes |\text{well}\rangle$.

**lemma** *flag-H-well-conditioned*:
**assumes** *A*: $A \in$ *carrier-mat n n hermitian A* **and** *k0*: $0 < \kappa$ **and** *wc*: $\bigwedge l.\ l \in$ *herm-spectrum A* $\Longrightarrow$ *1 / κ* $\leq l$
**shows** *flag-H* $\kappa$ *A 2* $= 0_m$ *n n* **and** *flag-H* $\kappa$ *A 1* $=$ *complex-of-real* $(1\ /\ (2 * \kappa)) \cdot_m$ *herm-inv A*
⟨*proof*⟩

## 118.2 The flag gates exist

**lemma** *hflag-unit*: $0 < \kappa \Longrightarrow \|$*hflag* $\kappa$ *x*$\| = 1$
⟨*proof*⟩

A Householder reflection maps $|0\rangle$ to any unit vector $h$ with real $h_0 < 1$.

**definition** *householder* :: *nat* ⇒ *complex Matrix.vec* ⇒ *complex Matrix.mat* **where**
*householder d h* $= 1_m$ *d* $-$ *complex-of-real* $(2\ /\ \|$*unit-vec d 0* $-$ *h*$\|^2) \cdot_m$ *outer* (*unit-vec d 0* $-$ *h*)

**lemma** *outer-sq*: *outer v* $*$ *outer v* $=$ *complex-of-real* $(\|v\|^2) \cdot_m$ *outer v*
⟨*proof*⟩

**lemma** *householder-props*:
**assumes** *h*: *dim-vec h* $= d$ $\|h\| = 1$ **and** *h0*: *h* $ 0 =$ *complex-of-real r* $r < 1$ **and** *d0*: $0 < d$
**shows** *householder d h* $\in$ *carrier-mat d d unitary* (*householder d h*) *householder d h* $*_v$ *unit-vec d 0* $=$ *h*
⟨*proof*⟩

So gates with $G_k|0\rangle = |h(\tilde{\lambda}_k)\rangle$ exist: the hypotheses of Theorem 1 are met.

**lemma** *flag-gates-exist*:
**assumes** *k0*: $0 < \kappa$
**shows** $\exists\, G.\ \forall x.\ G\ x \in$ *carrier-mat 3 3* $\wedge$ *unitary* $(G\ x) \wedge G\ x *_v$ *unit-vec 3 0* $=$ *hflag* $\kappa$ *x*
⟨*proof*⟩

**end**