

Acute stress modulates the use of hippocampal subsystems during learning








Brynn E. Sherman¹, Isabella Huang¹, Elaine G. Wijaya¹, Nicholas B. Turk-Browne^{1,2}, Elizabeth V. Goldfarb^{1,2,3} ¹Department of Psychology; ²Wu Tsai Institute; ³Department of Psychiatry, Yale University


Introduction Stress tends to negatively affect the hippocampus & episodic memory 1,2 However, stress may have diverging effects across hipp. subregions^{3,4} Different kinds of memory may rely on Pathway (TSP) Pathway (MSP) distinct hippocampal pathways (episodic memory: TSP, statistical learning: MSP)⁵ — stress Stress response is temporally dynamic⁶; delay between stress and encoding can Entorhinal Cortex affect how stress influences memory Episodic Memory Statistical Learning Hypothesis: How (and when) does acute stress Acute stress will enhance differentially affect episodic memory statistical learning and & statistical learning?

impair episodic memory









Discussion

Preliminary analyses suggest that acute stress (particularly when encountered ~15 min before learning) may promote the adaptive use of learned statistical regularities

- greater RT facilitation
- enhanced retention of regularities 24h later
- enhanced memory (consolidation?) of predictable items

Strongest effects with ~15 minute stress-learning delay; may suggest role of glucocorticoids in mediating effects

Suggests a broader role for stress in promoting integrative representations, in line with the role of stress in biasing behavior towards stimulus-response learning⁹

How does acute stress modulate connectivity among hippocampal subregions (i.e., the pathways supporting these processes)?