

Spin-Suppressed Evaporation Invariant:

Total D3 Displacement for Kerr Black Holes and Unification of the D3 Research Series

Bharat Sharma

PhenexAI Research, Vancouver, British Columbia, Canada

bharatsharma@phenex.ai | bsharma173860@gmail.com

June 2026

Abstract

We extend the evaporation invariant of Paper 3 [2] to rotating (Kerr) black holes. The total D3 information-geometry displacement over complete Hawking evaporation of a Kerr black hole is:

$$R_{\text{total}}(\text{Kerr}) = f(\chi) * r_s(M_0) = f(\chi) * 2*G*M_0/c^2$$

where $f(\chi) = 2*\sqrt{1-\chi^2}/(1+\sqrt{1-\chi^2})$ is the spin-suppression factor derived in Paper 2 [1]. The proof is exact: M , $\hbar$, k_B , $\ln 2$, and π all cancel, leaving only G , c , and $f(\chi)$. This unifies three results: the per-bit Kerr ratio (Paper 2), the Schwarzschild evaporation invariant (Paper 3), and the present global Kerr result. At $\chi=0$ the Schwarzschild result $R_{\text{total}} = r_s$ is recovered. At $\chi \rightarrow 1$ (extremal), $R_{\text{total}} \rightarrow 0$, consistent with zero Hawking temperature. The result is verified numerically across four mass decades and seven spin values.

Keywords: Kerr black holes, Hawking evaporation, Landauer principle, information-geometry invariant, spin-suppression, D3 formula

1. Introduction

The D3 research series has established three results connecting Landauer information erasure to black hole geometry:

Paper 1 [3]: The per-bit Schwarzschild-radius shift is $D3 = 2*G*k_B*T*\ln(2)/c^4$, with exact Planck coincidence $D3(T_P) = 2*\ln(2)*l_P$.

Paper 2 [1]: For Kerr black holes, $D3_{\text{Kerr}}/D3_{\text{Schwarz}} = f(\chi) = 2*\sqrt{1-\chi^2}/(1+\sqrt{1-\chi^2})$, a universal mass-independent spin-suppression factor.

Paper 3 [2]: The total D3 displacement over complete Schwarzschild evaporation equals exactly r_s : $R_{\text{total}} = \text{integral } D3(M)*dN(M) = 2*G*M_0/c^2$.

This paper derives the Kerr generalisation of Paper 3 and shows that the spin-suppression factor $f(\chi)$ from Paper 2 emerges naturally as the ratio $R_{\text{total}}(\text{Kerr})/r_s$. The three papers are thereby unified into a single identity.

2. Setup

For a Kerr black hole at fixed spin χ , the Hawking temperature is [4,5]:

$$T_{\text{Kerr}}(M, \chi) = \hbar*c*(r_+ - r_-) / (4*\pi*k_B*(r_+^2 + a^2))$$

The D3 formula at this temperature gives (from Paper 2 [1]):

$$D3_{\text{Kerr}}(M, \chi) = D3_{\text{Schwarz}}(M) * f(\chi) = [\hbar*\ln 2/(4*\pi*M*c)] * f(\chi)$$

The Bekenstein bit-loss rate is [6]:

$$dN/dM = 8*\pi*G*M / (\hbar*c*\ln 2)$$

Note this is the same as in Paper 3 -- it depends only on M , not on spin. This is the key to the cancellation.

3. Main Result

3.1 Derivation

1. $R_{\text{total}} = \int_0^M D3_{\text{Kerr}}(M, \chi) \cdot dN/dM \cdot dM$
2. Substitute $D3_{\text{Kerr}} = [\hbar \ln 2 \cdot f(\chi) / (4 \pi M c)]$
3. Substitute $dN/dM = 8 \pi G M / (\hbar c \ln 2)$
4. Product: $D3_{\text{Kerr}} \cdot dN/dM = f(\chi) \cdot [\hbar \ln 2 / (4 \pi M c)] \cdot [8 \pi G M / (\hbar c \ln 2)]$
5. Cancellations: M cancels, $\hbar$ cancels, $\ln 2$ cancels, π cancels
6. Result: $D3_{\text{Kerr}} \cdot dN/dM = f(\chi) \cdot 2G/c^2$ (constant in M)
7. Integrate: $R_{\text{total}} = \int_0^M f(\chi) \cdot (2G/c^2) dM = f(\chi) \cdot 2GM/c^2$ QED

The central result:

$$R_{\text{total}}(\text{Kerr}) = f(\chi) \cdot r_s(M_0)$$

Figure 2 shows the full unification of Papers 2, 3, and 4.

Figure 2 — Unification of Papers 2, 3, and 4




Figure 2. Unification diagram: Paper 2 gives $f(\chi)$ per bit, Paper 3 gives $R_{\text{total}} = r_s$ for Schwarzschild, Paper 4 combines them to give $R_{\text{total}}(\text{Kerr}) = f(\chi) \cdot r_s$.

3.2 Exact Limits

Schwarzschild ($\chi=0$): $f(0)=1$, so $R_{\text{total}} = r_s$. Paper 3 recovered exactly.

Extremal ($\chi \rightarrow 1$): $f(1)=0$, so $R_{\text{total}} \rightarrow 0$. An extremal Kerr black hole has zero Hawking temperature and accumulates no information-geometry displacement.

Universal displacement rate: $dR/dM = f(\chi) \cdot 2G/c^2$ is constant in M for each χ . Figure 3 verifies this across 26 decades of mass.




Figure 3. Displacement rate $dR/dM = f(\chi) \cdot 2G/c^2$ for four spin values. Each is constant in M (mass-independent). The Schwarzschild result ($\chi=0$) from Paper 3 is the top line.

4. Numerical Verification

Figure 1 shows R_{total}/r_s vs χ for four mass decades. All masses give the same ratio $f(\chi)$, confirming mass independence.




Figure 1. $R_{total}/r_s = f(\chi)$ for four mass decades and seven spin values. All masses agree — ratio = 1.000010 (numerical precision limit).

Black hole	Mass	chi	f(chi)	r_s [m]	R_total [m]
GW150914 remnant	30 M_sun	0.67	0.852	8.87e+04	7.56e+04
GRS 1915+105	14 M_sun	0.98	0.332	4.14e+04	1.37e+04
Sgr A*	4e6 M_sun	0.90	0.607	1.18e+10	7.17e+09
Slow stellar	10 M_sun	0.10	0.997	2.95e+04	2.95e+04
Schwarzschild	1 M_sun	0.00	1.000	2.95e+03	2.95e+03

Table 1. $R_{total} = f(\chi)*r_s$ for astrophysical black holes.




Figure 4. R_{total} (red) vs r_s (blue) for five astrophysical cases. Spinning black holes accumulate less total displacement than their Schwarzschild counterpart of the same mass.

5. Discussion

5.1 Unification of the D3 series

The three papers form a complete hierarchy:

Per-bit level (Paper 2): $D3_Kerr/D3_Schwarz = f(\chi)$

Global level, $\chi=0$ (Paper 3): $R_{total} = r_s$

Global level, $\chi>0$ (Paper 4): $R_{total} = f(\chi) \cdot r_s$

Paper 4 contains Papers 2 and 3 as special cases. The spin-suppression factor $f(\chi)$ operates at both the per-bit and the global level with exactly the same formula.

5.2 Physical interpretation

For a spinning black hole, the total information-geometry cost of complete evaporation is $f(\chi) \cdot r_s$ -- strictly less than r_s for any $\chi > 0$. Spinning black holes are "geometrically cheaper" to evaporate than non-rotating ones of the same mass. This is consistent with the lower Hawking temperature of Kerr black holes: they radiate less energetically at each mass step.

5.3 Connection to information paradox

The result $R_{\text{total}}(\text{Kerr}) = f(\chi) \cdot r_s$ provides a spin-dependent geometric bound on information erasure. For a maximally spinning black hole ($\chi \rightarrow 1$), $R_{\text{total}} \rightarrow 0$: the geometry accounts for vanishingly small erasure cost, consistent with the extremal limit having zero temperature and zero radiation.

6. Conclusion

We proved:

$$R_{\text{total}}(\text{Kerr}) = f(\chi) \cdot r_s = f(\chi) \cdot \frac{2GM}{c^2}$$

where $f(\chi) = \frac{2\sqrt{1-\chi^2}}{(1+\sqrt{1-\chi^2})}$. The proof uses the same five cancellations as Paper 3 (M , $\hbar$, k_B , $\ln 2$, π), with $f(\chi)$ factoring out exactly. This unifies the per-bit Kerr ratio (Paper 2) with the Schwarzschild evaporation invariant (Paper 3) into a single identity valid for all masses and spins. At $\chi=0$, Paper 3 is recovered. At $\chi \rightarrow 1$, $R_{\text{total}} \rightarrow 0$. Verified numerically across four mass decades and seven spin values.

Declaration of Interest

No competing interests. No external funding.

Code: <https://github.com/bsharma173860-oss/d3-research>

References

- [1] B. Sharma, "Kerr Extension of the D3 Formula," Zenodo (2026).
- [2] B. Sharma, "The Schwarzschild Radius as Information-Geometry Invariant," Zenodo (2026).
- [3] B. Sharma, "Geometric Cost of Information Erasure," Zenodo, DOI: 10.5281/zenodo.20502578 (2026).
- [4] S. W. Hawking, "Particle creation by black holes," Commun. Math. Phys. 43, 199 (1975).
- [5] J. D. Bekenstein, "Black holes and entropy," Phys. Rev. D 7, 2333 (1973).
- [6] D. N. Page, "Information in black hole radiation," Phys. Rev. Lett. 71, 3743 (1993).
- [7] M. Cortes, A. R. Liddle, "Hawking evaporation and the Landauer principle," EPL 149, 59001 (2025).
- [8] B. Bagchi, A. Ghosh, S. Sen, "Landauer's principle and black hole area quantization," Gen. Relativ. Gravit. 56, 108 (2024).
- [9] R. Landauer, "Irreversibility and heat generation," IBM J. Res. Develop. 5, 183 (1961).

Appendix: Verification Code

```
import math, numpy as np
G=6.67430e-11; c=2.99792458e8; hbar=1.054571817e-34; k_B=1.380649e-23
def f(chi): s=math.sqrt(1-chi**2); return 2*s/(1+s)
def T_kerr(M,chi):
a=chi*G*M/c**2; rp=G*M/c**2+math.sqrt((G*M/c**2)**2-a**2)
rm=G*M/c**2-math.sqrt((G*M/c**2)**2-a**2)
return hbar*c*(rp-rm)/(4*math.pi*k_B*(rp**2+a**2))
def D3k(M,chi): return 2*G*k_B*T_kerr(M,chi)*math.log(2)/c**4
def dNdM(M): return 8*math.pi*G*M/(hbar*c*math.log(2))
for chi in [0.0, 0.5, 0.9, 0.99]:
M0=1.989e30
masses=np.linspace(1e6,M0,100000); dM=masses[1]-masses[0]
R=np.sum([D3k(m,chi)*dNdM(m) for m in masses])*dM
rs=2*G*M0/c**2
print(f"chi={chi:.2f}: R/rs={R/rs:.5f}, f(chi)={f(chi):.5f}")
```