



# GIMME, GIMME more (network models): Multiverse analysis for dynamic network models

Björn Siepe<sup>1</sup> Daniel W. Heck<sup>1</sup>

May 26, 2025 – Society for Ambulatory Assessment


 ${}^1\mathsf{Psychological}\ \mathsf{Methods}\ \mathsf{Lab}, \mathsf{Department}\ \mathsf{of}\ \mathsf{Psychology}, \mathsf{University}\ \mathsf{of}\ \mathsf{Marburg}$ 


• Time series data often require complex models


- Time series data often require complex models
- Many analytical choices: Model selection, preprocessing, model specification, interpretation...

- Time series data often require complex models
- Many analytical choices: Model selection, preprocessing, model specification, interpretation...
- Often, only one set of choices is conducted and reported

- Time series data often require complex models
- Many analytical choices: Model selection, preprocessing, model specification, interpretation...
- Often, only one set of choices is conducted and reported
- Robustness to arbitrary choices is underappreciated







2

# **This Study**

- Investigate the impact of arbitrary alternative modeling decisions on previously published results
- Previous multiverse studies: Focus on measurement (Dejonckheere et al., 2018) or preprocessing (Weermeijer et al., 2022)

# **This Study**

- Investigate the impact of arbitrary alternative modeling decisions on previously published results
- Previous multiverse studies: Focus on measurement (Dejonckheere et al., 2018) or preprocessing (Weermeijer et al., 2022)



# **Datasets**

## **Datasets**

# **Personality** Dataset (Wright et al., 2019):

- n=94 participants with personality disorder diagnosis, average t=91.48 (1x/day)
- Items: Affect, interpersonal behavior, stress, and functioning (all sum scores), and daily functioning (5-point Likert)

4

## **Datasets**

# **Personality** Dataset (Wright et al., 2019):

- n=94 participants with personality disorder diagnosis, average t=91.48 (1x/day)
- Items: Affect, interpersonal behavior, stress, and functioning (all sum scores), and daily functioning (5-point Likert)

# **Emotion** Dataset (Kullar et al., 2024):

- n = 105, average t = 62.31 (SD = 8.11)
- Items: Nine momentary emotions (Likert 7-point)

4

# **GIMME Variations**

- 1. Group threshold  $\in \{50\%, 60\%, 75\%, 80\%\}$
- 2. Subgroup threshold  $\in \{50\%, 60\%, 75\%, 80\%\}$

# **GIMME Variations**

- 1. Group threshold  $\in \{50\%, 60\%, 75\%, 80\%\}$
- 2. Subgroup threshold  $\in \{50\%, 60\%, 75\%, 80\%\}$

Five parameters refer to the fit indices used for model selection:

- 3. RMSEA cutoff  $\in \{.03, .05, .08\}$
- 4. SRMR cutoff  $\in \{.03, .05, .08\}$
- 5. NNFI cutoff  $\in \{.90, .95, .97\}$
- 6. CFI cutoff  $\in \{.90, .95, .97\}$
- 7. Fit measures satisfying the cutoffs  $\in \{1, 2, 3\}$

# **GIMME Variations**

- 1. Group threshold  $\in \{50\%, 60\%, 75\%, 80\%\}$
- 2. Subgroup threshold  $\in \{50\%, 60\%, 75\%, 80\%\}$

Five parameters refer to the fit indices used for model selection:

- 3. RMSEA cutoff  $\in \{.03, .05, .08\}$
- 4. SRMR cutoff ∈ {.03, .**05**, .08}
- 5. NNFI cutoff  $\in \{.90, .95, .97\}$
- 6. CFI cutoff  $\in \{.90, .95, .97\}$
- 7. Fit measures satisfying the cutoffs  $\in \{1, 2, 3\}$

Conducted a small simulation study showing the arbitrariness of these choices

Group:

- Group:
  - $\sim$  92% of models with same group path

- Group:
  - $\sim$  92% of models with same group path

- Group:
  - $\sim$  92% of models with same group path
- Subgroup:

- Group:
  - $\sim$  92% of models with same group path
- Subgroup:
  - Same subgroups

- Group:
  - $\sim$  92% of models with same group path
- Subgroup:
  - Same subgroups

- Group:
  - $\sim$  92% of models with same group path
- Subgroup:
  - Same subgroups
- Individual

- Group:
  - $\sim$  92% of models with same group path
- Subgroup:
  - Same subgroups
- Individual
  - On average,  $\sim$  2 paths different in presence/absence from reference fit

- Group:
  - $\sim$  92% of models with same group path
- Subgroup:
  - Same subgroups
- Individual
  - ullet On average,  $\sim$  2 paths different in presence/absence from reference fit
  - For those different effects: Absolute average difference of  $\sim$  0.13

#### Group:

•  $\sim$  92% of models with same group path

## Subgroup:

Same subgroups

#### Individual

- On average,  $\sim$  2 paths different in presence/absence from reference fit
- For those different effects: Absolute average difference of  $\sim$  0.13
- For 12 individuals, the most central node was identical to the reference model in less than one-third of all specifications

• Group:

## Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated

- Group:
  - In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated
- Subgroup:

## Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated

# Subgroup:

-  $\sim 50\%$  of subgroup solutions very dissimilar to the reference fit

## Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated

## Subgroup:

- $\sim$  50% of subgroup solutions very dissimilar to the reference fit
- Subgroup threshold of 50% instead of 51%: Change of 128 paths

## Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated

## Subgroup:

- $\sim$  50% of subgroup solutions very dissimilar to the reference fit
- Subgroup threshold of 50% instead of 51%: Change of 128 paths

#### Individual

## Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated

# Subgroup:

- $\sim$  50% of subgroup solutions very dissimilar to the reference fit
- Subgroup threshold of 50% instead of 51%: Change of 128 paths

#### Individual

- On average,  $\sim$  9 paths different in absence/presence from reference fit

## Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated

# Subgroup:

- $\sim$  50% of subgroup solutions very dissimilar to the reference fit
- Subgroup threshold of 50% instead of 51%: Change of 128 paths

#### Individual

- On average,  $\sim$  9 paths different in absence/presence from reference fit
- For those different effects: Absolute average difference of 0.2

### Group:

- In  $\sim$  50% specifications, four or five group effects (instead of one) were estimated


# Subgroup:

- $\sim$  50% of subgroup solutions very dissimilar to the reference fit
- Subgroup threshold of 50% instead of 51%: Change of 128 paths


#### Individual

- On average,  $\sim$  9 paths different in absence/presence from reference fit
- For those different effects: Absolute average difference of 0.2
- For 30 individuals, the most central node was identical to the reference model in less than one-third of all specifications

## **Emotion Results**



### **Emotion Results**



# **Shiny App**



Shiny App for the paper "Network Multiverse" (Siepe & Heck, 2023). Find the source code on  $\bigcirc$  GitHub.

### Main results:

Results largely robust for Personality dataset, less robust for Emotion dataset

#### Main results:

- Results largely robust for Personality dataset, less robust for Emotion dataset
- Different specifications could strongly affect main conclusions

#### Main results:

- Results largely robust for Personality dataset, less robust for Emotion dataset
- Different specifications could strongly affect main conclusions

#### Main results:

- Results largely robust for Personality dataset, less robust for Emotion dataset
- Different specifications could strongly affect main conclusions

### Broader implications:

 Multiverse has often focused on preprocessing, but is relevant for algorithmic decisions

#### Main results:

- Results largely robust for Personality dataset, less robust for Emotion dataset
- Different specifications could strongly affect main conclusions

### Broader implications:

- Multiverse has often focused on preprocessing, but is relevant for algorithmic decisions
- Interesting relationship between simulation studies and multiverse analyses

## **Get In Touch**

- **S** bjoern.siepe@uni-marburg.de
- ♠ https://bsiepe.github.io/

Paper & Slides

### References i



Gates, K. M., Lane, S. T., Varangis, E., Giovanello, K., & Guiskewicz, K. (2017). **Unsupervised classification during time-series model building.** *Multivariate Behavioral Research*, *52*(2), 129–148. https://doi.org/10.1080/00273171.2016.1256187

### References ii

Kullar, M., Carter, S., Hitchcock, C., Whittaker, S., Wright, A. G. C., & Dalgleish, T. (2024). Patterns of emotion-network dynamics are orthogonal to mood disorder status: An experience sampling investigation..

Emotion, Advance online publication. https://doi.org/10.1037/emo0001245

Weermeijer, J., Lafit, G., Kiekens, G., Wampers, M., Eisele, G., Kasanova, Z., Vaessen, T., Kuppens, P., & Myin-Germeys, I. (2022). Applying multiverse analysis to experience sampling data: Investigating whether preprocessing choices affect robustness of conclusions. Behavior

Research Methods, 54(6), 2981–2992.

https://doi.org/10.3758/s13428-021-01777-1

### References iii



Wright, A. G. C., Gates, K. M., Arizmendi, C., Lane, S. T., Woods, W. C., & Edershile, E. A. (2019). Focusing personality assessment on the person: Modeling general, shared, and person specific processes in personality and psychopathology. *Psychological Assessment*, 31(4), 502–515. https://doi.org/10.1037/pas0000617

## **Image Source**

```
Britney Spears Cover:
https:
//lastfm.freetls.fastly.net/i/u/ar0/dff1377bbbf94c97cc8769d0ea072804.jpg
```