

# Lab1\_151

September 1, 2022

## 0.1 MATH 151 Lab 1

Section Number: 568

Members:

- Brighton Sikarskie
- Colton Hesser
- Gabriel Gonzalez
- Gabriel Cuevas

```
[1]: from sympy import *  
from sympy.plotting import (plot, plot_parametric)
```

### 0.1.1 Question 1

1a

```
[2]: print((79 * (E ** 1.29 + 11.1 ** 2)) / (2026 - 5.1 ** 3))
```

5.29251613828432

1b

```
[3]: def to_rad(deg): return deg * pi / 180  
  
equation = (cos((11 * pi) / 12) * sec(to_rad(75)) + tan((7 * pi) / 12))  
print(f"approximate: {equation.simplify()}")  
print(f"exact: {equation.evalf()}")
```

approximate:  $-4 - 2\sqrt{3}$

exact: -7.46410161513775

### 0.1.2 Question 2

2a

```
[4]: x = Symbol('x')  
equations = (sqrt(x ** 2 - 4) / (x - 2))  
def f(o): return equations.subs(x, o).evalf()  
  
print(f(-10), f(-100), f(-1000000))
```

-0.816496580927726 -0.980196058819607 -0.999998000002000

**2b**

```
[5]: print(f(2.01), f(2.0001), f(2.000001))
```

20.0249843945009 200.002499984149 2000.00024988243

**2c** Based on part (a), what happens to the y-values of  $f$  when  $x$  gets REALLY large in the negative direction?

When  $x$  gets REALLY large in the negative direction the y-values of  $f$  become close to -1.

**2d** Based on part (b), what happens to the y-values of  $f$  when  $x$  gets close to 2 from the right?

When  $x$  gets close to 2 from the right the y-values of  $f$  becomes increasingly larger. It seems for every 0 we put behind the decimal the outputted value gets increased by a power of 10.

### 0.1.3 Question 3

**3a**

```
[6]: a, v, h = symbols('a v h')

equation = (-16 / (v ** 2 * (cos(a) ** 2))) * x ** 2 + tan(a) * x + h
ball_height = equation.subs([a, to_rad(26)], (v, 130), (h, 3), (x, 409)).
    ↪evalf()
print(f"ball height: {ball_height}")
```

ball height: 6.43593602529508

Since the wall is 10ft high and the ball is only 6.4ft high it will hit the wall and not clear it.

**3b**

```
[7]: equation = (-16 / (v ** 2 * (cos(a) ** 2))) * x ** 2 + tan(a) * x + h - 10
h = solve(equation.subs([a, to_rad(54.2)], (v, 24), (x, 15)]), h)[0]
print(f"He released it from {h} feet in the air")
```

He released it from 7.46746878686940 feet in the air