

Search for Contact Interactions Using Inclusive Jet p_T Spectrum @13 TeV

Suman Beri¹, Suneel Dutt², Pampa Ghose³, Bipen
Kotwal², and Harrison B. Prosper³

¹Panjab University, ²Shoolini University,
³Florida State University

Exotica Jets+X Working Group Meeting

3 October 2016

Outline

1. Overview
2. Analysis
3. Results
4. Plans for 2016 Data

Overview

Goal

Search for new QCD-like interactions by looking for deviations from QCD in the inclusive jet p_T spectrum.

Inclusive: all jets that fall within a given phase space.

Basic Assumption

At LHC energies, the “true” Lagrangian L can be written as

$$L = L_{SM}^{(0)} + \frac{1}{\Lambda} L^{(1)} + \frac{1}{\Lambda^2} L^{(2)} + \dots$$

in which the $L^{(2)}$ term contains dim-6 operators that describe 4-quark interactions, which can be approximated as contact interactions (CI).

Overview

Experimental Input

- Inclusive jet p_T spectrum in $|y| < 0.5$, $638 \leq p_T \leq 2000$ GeV¹, anti- k_T R = 0.7 jets.
- Jet response function (JRF)
- Jet energy scale (JES) uncertainty:
Summer15_50nsV5 for AK8PF jets
- Jet energy resolution (JER) uncertainty: 10%

¹Eur. Phys. J. C **76** (2016) no.8, 451, SMP-15-007

Overview

Theoretical Input

- PDFs (LHAPDF-6.1.6)
CT14nlo, MMHT2014nlo68cl, NNPDF30_nlo_as_0118_1000
- Program to calculate QCD@NLO
fastnlo_toolkit-2.3.1pre-1871
InclusiveNJets_fnl5332g_v23_fix.tab
- Program to calculate CI@NLO
CIJET-1.1
(J. Gao, Comput.Phys.Commun. 184 (2013) 2362)
- Non-perturbative corrections (NP)
- Electroweak corrections (EWK)

Overview

Many thanks to the Inclusive Jet Group!

Sourav Dev

Giannis Flouris

Hannes Jung

Panos Kokkas

Ksenia Shchelina

and especially to [Paolo Gunnellini](#) who provided the details of the measured spectrum and the NP and EWK correction functions.

Overview: Models

We consider the following discrete set of values for the κ parameters:

Model	η_{LL}	η_{RL}	η_{RR}
LL	± 1	0	0
RR	0	0	± 1
VV	± 1	± 1	± 1
AA	± 1	∓ 1	± 1
V-A	0	± 1	0

where $\kappa_1 = \eta_{LL}$, $\kappa_3 = 2\eta_{RL}$, $\kappa_5 = \eta_{RR}$, and $\kappa_2 = \kappa_4 = \kappa_6 = 0$.

Overview: Models

The NLO QCD+CI cross section per jet p_T bin can be written as

$$\sigma = \sigma_{\text{QCD}} +$$

$$+ \lambda \sum_{i=1}^6 \kappa_i [b_i + a_i g + a_i f]$$

$$+ \lambda^2 \sum_{i=1}^6 \kappa_i^2 [b_{ii} + a_{ii} g + a_{ii} f]$$

$$+ \lambda^2 \sum_{i=1,3,5} \kappa_i \kappa_{i+1} [b_{ii+1} + a_{ii+1} g + a_{ii+1} f]$$

$$+ \lambda^2 \sum_{i=1,2,5,6} \kappa_i \kappa_4 [b_{i4} + a_{i4} g + a_{i4} f]$$

where $g = -\ln(\mu_0 \sqrt{k})$

and $\lambda = 1/\Lambda^2$

$f = \ln(\sqrt{k / \lambda})$

The CI term comprises 57

coefficients


Analysis

Big Picture

1. Integrate multinomial likelihood $p(D|\lambda, \mathbf{v})$ over nuisance parameters \mathbf{v} associated with PDFs, renormalization and factorization scales, jet energy scale, and jet energy resolution. ($\lambda = 1/\Lambda^2$ and D denotes the observed counts.)
2. Compute expected limits by setting
“observed” counts = expected counts
3. Compute observed limits using observed counts.



See backup for details.

Data: 71.52 pb⁻¹ @ 13TeV





Data vs. QCD

PDF + JES + JER (Summer15 V5) uncertainties



Data vs. LL Model ($\Lambda=20\text{TeV}$)



PDF + JES + JER (Summer15 V5) uncertainties



Constructive

Data vs. LL Model ($\Lambda=20\text{TeV}$)



PDF + JES + JER (Summer15 V5) uncertainties





Destructive

LIMITS

LL Limits @ 95% CL: Expected



LL Limits @ 95% CL: Observed



Limits @ 95% CL

$L = 0.072 \text{ fb}^{-1}$ $L = 2.6 \text{ fb}^{-1}$
 AN-16-338 AN-15-245*

Model	Observed (TeV)	Expected (TeV)	Expected (TeV)
$\Lambda^+(\text{LL/RR})$	10.5	9.9	12.1
$\Lambda^-(\text{LL/RR})$	15.1	13.7	17.3
$\Lambda^+(\text{VV})$	12.0	11.5	13.9
$\Lambda^-(\text{VV})$	19.5	17.5	22.2
$\Lambda^+(\text{AA})$	12.1	11.5	13.9
$\Lambda^-(\text{AA})$	19.3	17.5	22.1
$\Lambda^+(\text{V-A})$	8.7	8.0	9.5
$\Lambda^-(\text{V-A})$	8.7	8.0	9.5

*Search for quark contact interactions and extra spatial dimensions in the dijet angular distributions at 13 TeV, AN-15-245, L. Apanasevich *et al.*

Plans for 2016 Data



We would like to work on a publishable result using the 2016 data set.

However, this only makes sense if the analysis is not severely limited by systematic uncertainties.

We've made a rough estimate for the LL model (assuming JEC uncertainties become \ll PDF uncertainties) using CT14.

Plans for 2016 Data: Λ^{95} vs Lumi

These plots suggest that we can use up to $\sim 2.0/\text{fb}$ for destructive



interference models and up to $\sim 20/\text{fb}$ for constructive interference models with 2014 PDFs.

Plans for 2016 Data

1. Work with inclusive jet group on ongoing measurement of inclusive jet p_T spectrum using 2016 data.
2. Repeat analysis using 2016 measurement. Extend cross section predictions beyond 2TeV.
3. Write AN/paper.

Timescale $\sim 6 - 8$ months.

Immediate plans: complete current AN, then write a PAS.

BACKUP

Analysis

Steps

1. For each PDF generate 200 *randomly* sampled PDF sets using `hessian2replicas` in LHAPDF6.1.6, except for NNPDF30_nlo for which the sample already exists.
2. For each sampled PDF set and 7 combinations of the renormalization and factorization scales, compute the QCD inclusive jet p_T spectrum (yielding 4200 spectra).
3. Repeat 2., but instead use CIJET to compute the 57 differential coefficients needed to compute the CI spectra for *arbitrary* values of κ and Λ (57×4200 spectra).

CI Differential Coefficients (CT14, 0)



Analysis

Steps

4. Convolve each differential spectrum, $f(p_T) = d^2F/dp_T d|y|$ (either QCD or the 57 CI coefficients),

$$f_{obs}(p_T) = \int_0^\infty R(p_T | z) f(z) dz$$

with the jet response function R , while accounting for the uncertainty in the jet energy scale (JES) and jet energy resolution (JER), taking care to maintain the correlations across all bins and all spectra.

Analysis

Steps

5. Compute the marginal likelihood

$$p(D \mid \lambda, \kappa) = \int \text{multinomial}(D \mid \lambda, \kappa, v) \pi(v) dv$$
$$\simeq \frac{1}{K} \sum_{i=1}^{K=4200} \text{multinomial}(D \mid \lambda, \kappa, v_i)$$



where D denotes the counts per bin, and v denotes the JES, JER, and PDF nuisance parameters.

6. Compute upper limit on λ by solving

$$\int_0^{\lambda^{UP}} p(\lambda \mid D, \kappa) d\lambda = 0.95,$$

where $p(\lambda \mid D, \kappa) = p(D \mid \lambda, \kappa) p(\lambda) / p(D, \kappa)$

Contact Interaction Search @ 7 TeV



PHYSICAL REVIEW D **87**, 052017 (2013)

Search for contact interactions using the inclusive jet p_T spectrum in pp collisions at $\sqrt{s} = 7 \text{ TeV}$

S. Chatrchyan *et al.**

(CMS Collaboration)

(Received 21 January 2013; published 26 March 2013)