

# Brief Introduction to Bayesian Inference


Byron J. Smith

BMI 206

2020-11-19

Slides: <https://bit.ly/36GJrRE>

Code: <https://bit.ly/36M3YV1>



With help  
from XKCD

Byron J. Smith  
BMI 206  
2020-11-19

Slides: <https://bit.ly/36GJRrE>

Code: <https://bit.ly/36M3YV1>

What is science?

# What is science?

- Learn discrete facts about the world.
  - “COVID-19 is caused by SARS-CoV-2.”
  - Karl Popper
  - “Strong Inference”: Rule out all other plausibilities

# What is science?

- Learn discrete facts about the world.
  - “COVID-19 is caused by SARS-CoV-2.”
  - Karl Popper
  - “Strong Inference”: Rule out all other plausibilities
- Measure (continuous) parameters
  - “Wearing a mask decreases your risk of becoming infected with SARS-CoV-2 by 55.2%.”

# What is science?

- Learn discrete facts about the world.
  - "COVID-19 is caused by SARS-CoV-2."
  - Karl Popper
  - "Strong Inference": Rule out all other plausibilities
- Measure (continuous) parameters
  - "Wearing a mask decreases your risk of becoming infected with SARS-CoV-2 by 55.2%."
- ...But what about questions somewhere in between these two extremes?
  - "Does wearing a mask affect your risk of becoming infected?"

# “Traditional”: Null hypothesis statistical testing

*“Does wearing a mask affect your risk of becoming infected?”*

- Run an (e.g.) t-test
  - Implies a null hypothesis:  $H_0 : \mu_{\text{mask}} = \mu_{\text{not}}$
- Calculate p-value
- If  $p < 0.05$  : we “reject the null hypothesis”
  - Conclude that wearing a mask *does* affect your infection risk
- Else  $p > 0.05$  : we cannot reject

# “Traditional”: Null hypothesis statistical testing

*“Does wearing a mask affect your risk of becoming infected?”*

- Run an (e.g.) t-test
  - Implies a null hypothesis:  $H_0 : \mu_{\text{mask}} = \mu_{\text{not}}$
- Calculate p-value
- If  $p < 0.05$  : we “reject the null hypothesis”
  - Conclude that wearing a mask *does* affect your infection risk
- Else  $p > 0.05$  : we cannot reject

**What does the p-value represent?**



# What does the p-value represent?

- Probability of seeing a test statistic as extreme as the one we observed in a world where the null hypothesis is true.
  - NOT: The probability of the null hypothesis being true.

# What does the p-value represent?

- Probability of seeing a test statistic as extreme as the one we observed in a world where the null hypothesis is true.
  - NOT: The probability of the null hypothesis being true.

<u>P-VALUE</u>	<u>INTERPRETATION</u>
0.001	HIGHLY SIGNIFICANT
0.01	
0.02	
0.03	
0.04	SIGNIFICANT
0.049	
0.050	OH CRAP. REDO CALCULATIONS.
0.051	ON THE EDGE OF SIGNIFICANCE
0.06	
0.07	HIGHLY SUGGESTIVE, SIGNIFICANT AT THE $P < 0.10$ LEVEL
0.08	
0.09	
0.099	HEY, LOOK AT
$\geq 0.1$	THIS INTERESTING SUBGROUP ANALYSIS

# “Science by p-value”


## Shortcomings:

- Misleading when:
  - Intuition/reality does not match test assumptions (E.g. small sample size)
  - When our null hypothesis is trivially wrong
- Ignores prior information
- Ignores effect size

# “Science by p-value”

## Shortcomings:

- Misleading when:
  - Intuition/reality does not match test assumptions (E.g. small sample size)
  - When our null hypothesis is trivially wrong
- Ignores prior information
- Ignores effect size



# “Science by p-value”

## Shortcomings:

- Misleading when:
  - Intuition/reality does not match test assumptions (E.g. small sample size)
  - When our null hypothesis is trivially wrong
- Ignores prior information
- Ignores effect size

Odds Are,  
**It's Wrong**

### STATISTICS

## *Measurement error and the replication crisis*

The assumption that measurement error always reduces effect sizes is false

By **Eric Loken<sup>1</sup>** and **Andrew Gelman<sup>2</sup>**

reliable measurement. In epidemiology, it is the only way to know that a treatment is effective.

# “Science by p-value”

## Shortcomings:

- Misleading when:
  - Intuition/reality does not match test assumptions (E.g. small sample size)
  - When our null hypothesis is trivially wrong
- Ignores prior information
- Ignores effect size

Odds Are,  
**It's Wrong**

STATISTICS

*Measurement error and  
the replication crisis*

**STATISTICAL ERRORS**

*P values, the ‘gold standard’ of statistical validity, are not as reliable as many scientists assume.*

BY REGINA NUZZO

# “Science by p-value”

## Shortcomings:

- Misleading when:
  - Intuition/reality does not match test assumptions (E.g. small sample size)
  - When our null hypothesis is trivially wrong
- Ignores prior information
- Ignores effect size

## What we would prefer:

*"How does wearing a mask affect the risk of becoming infected?"*



*There is a 95% probability that wearing a mask decreases your risk of becoming infected by 50% or more*


# Bayes's Rule

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$



# Bayes's Rule

Parameters


$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Bayes's Rule

Data



$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Bayes's Rule

“Posterior”

$$\boxed{P(\theta|X)} = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Bayes's Rule

Have you seen this before?

$$P(\theta|X) = \frac{P(X|\theta)P(\theta)}{P(X)}$$

# Bayes's Rule

“Likelihood”

$$P(\theta|X) = \frac{\mathcal{L}(\theta|X)P(\theta)}{P(X)}$$

# Bayes's Rule

“Prior”

$$P(\theta|X) = \frac{P(X|\theta)P(\theta)}{P(X)}$$

# Bayes's Rule

“Normalizing Constant”

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{\boxed{P(X)}}$$


# Bayes's Rule

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$



# Bayes's Rule

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$



# Does wearing a mask affect COVID-19 risk?

Rephrase: What is the relative risk to wearers vs. non-wearers?

$\theta$  Odds ratio of COVID-19 among wearers and non-wearers

$X$

$P(X|\theta)$

$P(\theta)$

$P(X)$

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Does wearing a mask affect COVID-19 risk?

Rephrase: What is the relative risk to wearers vs. non-wearers?

$\theta$  Odds ratio of COVID-19 among wearers and non-wearers

$X$  Observed individuals and cases among both groups

$$P(X|\theta)$$

$$P(\theta)$$

$$P(X)$$

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Does wearing a mask affect COVID-19 risk?

Rephrase: What is the relative risk to wearers vs. non-wearers?

$\theta$  Odds ratio of COVID-19 among wearers and non-wearers

$X$  Observed individuals and cases among both groups

$P(X|\theta)$  Probability of the observed numbers, given the odds ratio

$P(\theta)$

$P(X)$

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Does wearing a mask affect COVID-19 risk?

Rephrase: What is the relative risk to wearers vs. non-wearers?

$\theta$  Odds ratio of COVID-19 among wearers and non-wearers

$X$  Observed individuals and cases among both groups

$P(X|\theta)$  Probability of the observed numbers, given the odds ratio

$P(\theta)$  Prior probability of the given odds ratio

$P(X)$

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$

# Does wearing a mask affect COVID-19 risk?

Rephrase: What is the relative risk to wearers vs. non-wearers?

$\theta$  Odds ratio of COVID-19 among wearers and non-wearers

$X$  Observed individuals and cases among both groups

$P(X|\theta)$  Probability of the observed numbers, given the odds ratio


$P(\theta)$  Prior probability of the given odds ratio

$P(X)$  Marginal probability of this observation

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$


# Does wearing a mask affect COVID-19 risk?

- As we collect more data, the posterior distribution converges on the observed odds ratio



# Does wearing a mask affect COVID-19 risk?


- As we collect more data, the posterior distribution converges on the observed odds ratio






# Does wearing a mask affect COVID-19 risk?

- As we collect more data, the posterior distribution converges on the observed odds ratio




# Does wearing a mask affect COVID-19 risk?

- As we collect more data, the posterior distribution converges on the observed odds ratio




# Does wearing a mask affect COVID-19 risk?

- As we collect more data, the posterior distribution converges on the observed odds ratio



# Does wearing a mask affect COVID-19 risk?

- As we collect more data, the posterior distribution converges on the observed odds ratio



How to calculate the posterior probability?

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$


# How to calculate the posterior probability?

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{P(X)}$$



# How to calculate the posterior probability?

- Hard

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{\int_{\Theta} P(X|\theta') P(\theta') d\theta'}$$


# How to calculate the posterior probability?

- Hard
- Easier: *Sampling* from the posterior

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{\int_{\Theta} P(X|\theta') P(\theta') d\theta'} \propto P(X|\theta) P(\theta)$$



# How to calculate the posterior probability?


- Hard
- Easier: *Sampling* from the posterior
- Markov-Chain Monte Carlo (Metropolis-Hastings algorithm)

$$P(\theta|X) = \frac{P(X|\theta) P(\theta)}{\int_{\Theta} P(X|\theta') P(\theta') d\theta'} \propto P(X|\theta) P(\theta)$$

# Sampling $\theta$ from the posterior



- Markov-Chain Monte Carlo (Metropolis-Hastings algorithm)

$$\theta_1, \theta_2, \theta_3, \dots, \theta_n$$



# Sampling $\theta$ from the posterior

- Markov-Chain Monte Carlo (Metropolis-Hastings algorithm)
- Use samples from the posterior to calculate
  - Expectations
  - Credible intervals



# Why doesn't everyone do it this way?

- Computation
  - Scales unfavorably with number of parameters, size of data
  - May require many samples due to “poor mixing”
- The Prior
  - “Introduces bias”
  - Subject to criticism
- Alternatives
  - Maximum likelihood

# Why doesn't everyone do it this way?

- Computation

- Scales unfavorably with number of parameters, size of data
- May require many samples due to “poor mixing”

- The Prior

- “Introduces bias”
- Subject to criticism

- Alternatives

- Maximum likelihood

- Computation

- Huge progress in last 20 years.
- HMC, Variational Inference

- The Prior

- Useful for incorporating expert knowledge, constraints, intuition.
- Makes assumptions explicit

- Alternatives

- Challenging to assess uncertainty from point estimates

# Logistic Regression

$$y_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_K x_{iK}$$

$$\beta_k \sim \text{Normal}(0, 1)$$

Try it out!

<https://bit.ly/36M3YVl>

# Logistic Regression

$$y_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_K x_{iK}$$

$$\beta_k \sim \text{Normal}(0, 10)$$



# Logistic Regression

$$y_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_K x_{iK}$$

$$\beta_k \sim \text{Laplace}(1)$$

# Democratized statistical modeling

- Emphasizes parameter estimates and uncertainty over p-values
- “Inference Button”: flexible, well built software to sample from and interpret models
  - STAN, PyMC3, Pyro, Turing.jl
- Only scratched the surface in this session
- Every Statistics XKCD (44 and counting): <https://bit.ly/32YH4Jc>

