

Machine Learning I

Review of Probability and Statistics

Souhaib Ben Taieb

University of Mons



Overview

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

References

- ▶ **Introduction to Probability for Data Science**, Stanley H. Chan.
[Link] (Book, slides and videos)
- ▶ Probability Theory Review for Machine Learning, Samuel leong.
[Link]
- ▶ *All of Statistics*, Larry Wasserman. [Link]

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Sample space and events

- ▶ When we speak about probability, we often refer to the probability of **an event of uncertain nature** taking place.
- ▶ We first need to clarify what the **possible events** are to which we want to attach probability.
- ▶ We often conduct an experiment, i.e. take some measurements of a **random (stochastic) process**.
- ▶ Our measurements take values in some set Ω , the **sample space** (or the outcome space)., which defines *all possible outcomes* of our measurements.

Sample space and events

- ▶ We toss one coin heads (H) or tails (T)
 - ▶ $\Omega = \{H, T\}$
- ▶ We toss two coins
 - ▶ $\Omega = \{HH, HT, TH, TT\}$
- ▶ We measure the reaction time to some stimulus
 - ▶ $\Omega = (0, \infty)$

Sample space and events

An **event** A is a subset of Ω ($A \subseteq \Omega$), i.e., it is a subset of possible outcomes of our experiment. We say that an event A **occurs** if the outcome of our experiment belongs to the set A .

- ▶ Let $\Omega = \{HH, HT, TH, TT\}$, and consider the following events:
 - ▶ $A_1 = \{HH, TH, TT\}$ and $A_2 = \{TH, TT\}$.

We observe $\omega = HT$. Which events did occur?

- ▶ Let $\Omega = (0, \infty)$, and consider the following events:
 - ▶ $A_1 = (3, 6)$, $A_2 = (1, 2)$ and $A_3 = (2, 8)$.

We observe $\omega = 4$. Which events did occur?

Sample space and events

An **event** A is a subset of Ω ($A \subseteq \Omega$), i.e., it is a subset of possible outcomes of our experiment. We say that an event A **occurs** if the outcome of our experiment belongs to the set A .

- ▶ Let $\Omega = \{HH, HT, TH, TT\}$, and consider the following events:
 - ▶ $A_1 = \{HH, TH, TT\}$ and $A_2 = \{TH, TT\}$.

We observe $\omega = HT$. Which events did occur?

- ▶ Let $\Omega = (0, \infty)$, and consider the following events:
 - ▶ $A_1 = (3, 6)$, $A_2 = (1, 2)$ and $A_3 = (2, 8)$.

We observe $\omega = 4$. Which events did occur?

Sample space and events

An **event** A is a subset of Ω ($A \subseteq \Omega$), i.e., it is a subset of possible outcomes of our experiment. We say that an event A **occurs** if the outcome of our experiment belongs to the set A .

- ▶ Let $\Omega = \{HH, HT, TH, TT\}$, and consider the following events:
 - ▶ $A_1 = \{HH, TH, TT\}$ and $A_2 = \{TH, TT\}$.

We observe $\omega = HT$. Which events did occur?

- ▶ Let $\Omega = (0, \infty)$, and consider the following events:
 - ▶ $A_1 = (3, 6)$, $A_2 = (1, 2)$ and $A_3 = (2, 8)$.

We observe $\omega = 4$. Which events did occur?

Probability space

A **probability space** is defined by the triple $(\Omega, \mathcal{F}, \mathbb{P})$ where

- ▶ Ω is the **sample space**
- ▶ $\mathcal{F} = 2^\Omega$ is the **space of events** (or event space)¹
- ▶ \mathbb{P} is the **probability measure/distribution** that maps an event $A \in \mathcal{F}$ to a real value between zero and one

¹ 2^Ω is the set of all subsets of S including S and the empty set \emptyset . Note that $\mathcal{F} = 2^\Omega$ is not fully general, but it is often sufficient for practical purposes.

Probability axioms

A **probability distribution** is a mapping from events to real numbers that satisfy certain **axioms**:

1. *Non-negativity*:

$$\mathbb{P}(A) \geq 0, \quad \forall A \subseteq \Omega$$

2. *Unity of Ω* :

$$\mathbb{P}(\Omega) = 1$$

3. *Additivity*. For all disjoint events $A, B \in \mathcal{F}$ (i.e. $A \cap B = \emptyset$), we have that,

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$$

Probability axioms

A **probability distribution** is a mapping from events to real numbers that satisfy certain **axioms**:

1. *Non-negativity*:

$$\mathbb{P}(A) \geq 0, \quad \forall A \subseteq \Omega$$

2. *Unity of Ω* :

$$\mathbb{P}(\Omega) = 1$$

3. *Additivity*. For all disjoint events $A, B \in \mathcal{F}$ (i.e. $A \cap B = \emptyset$), we have that,

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$$

Probability axioms

A **probability distribution** is a mapping from events to real numbers that satisfy certain **axioms**:

1. *Non-negativity*:

$$\mathbb{P}(A) \geq 0, \quad \forall A \subseteq \Omega$$

2. *Unity of Ω* :

$$\mathbb{P}(\Omega) = 1$$

3. *Additivity*. For all disjoint events $A, B \in \mathcal{F}$ (i.e. $A \cap B = \emptyset$), we have that,

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$$

Probability properties

Using set theory and the probability axioms, we can show several useful and intuitive properties of probability distributions.

- ▶ $\mathbb{P}(\emptyset) = 0$
- ▶ $A \subset B \implies \mathbb{P}(A) \leq \mathbb{P}(B)$
- ▶ $0 \leq \mathbb{P}(A) \leq 1$
- ▶ $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- ▶ $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

All of these properties can be understood via a **Venn diagram**.

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\text{Axiom 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\text{Axiom 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\text{Axiom 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\text{Axiom 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\text{Axiom 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\text{Axiom 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\text{Axiom 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\text{Axiom 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\textbf{Axiom 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\textbf{Axiom 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\textbf{Axiom 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\textbf{Axiom 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\textbf{Axiom 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\textbf{Axiom 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\textbf{Axiom 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\textbf{Axiom 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\textbf{Axiom 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\textbf{Axiom 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\textbf{Axiom 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\textbf{Axiom 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\mathbf{Axiom\ 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\mathbf{Axiom\ 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\mathbf{Axiom\ 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\mathbf{Axiom\ 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\mathbf{Axiom\ 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\mathbf{Axiom\ 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\mathbf{Axiom\ 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\mathbf{Axiom\ 1})$$

Probability properties

► $\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$

$$\mathbb{P}(\Omega) = 1 \quad (\mathbf{Axiom\ 2})$$

$$\iff \mathbb{P}(A \cup A^c) = 1, \quad \forall A \subseteq \Omega$$

$$\iff \mathbb{P}(A^c) + \mathbb{P}(A) = 1 \quad (\mathbf{Axiom\ 3})$$

$$\iff \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

► $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B).$

$$A \subseteq B$$

$$\implies B = A \cup (B \setminus A) \quad (A \cap (B \setminus A) = \emptyset)$$

$$\implies \mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \quad (\mathbf{Axiom\ 3})$$

$$\implies \mathbb{P}(B) \geq \mathbb{P}(A) \quad (\mathbf{Axiom\ 1})$$

Probability of an event (discrete case)

- The probability of any event $A = \{\omega_1, \omega_2, \dots, \omega_k\}$ ($\omega \in \Omega$) is the sum of the probabilities of its elements:

$$\mathbb{P}(A) = \mathbb{P}(\{\omega_1, \omega_2, \dots, \omega_k\}) = \sum_{i=1}^k \mathbb{P}(\{\omega_i\})$$

- If Ω consists of n equally likely outcomes (i.e. a uniform distribution), then the probability of any event A is

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|} = \frac{|A|}{n}$$

- Suppose we toss a **fair** dice twice. The sample space is $\Omega = \{(t_1, t_2) : t_1, t_2 = 1, 2, \dots, 6\}$. Let A be the event that the sum of two tosses being less than five. What is $\mathbb{P}(A)$?

Probability of an event (discrete case)

- ▶ The probability of any event $A = \{\omega_1, \omega_2, \dots, \omega_k\}$ ($\omega \in \Omega$) is the sum of the probabilities of its elements:

$$\mathbb{P}(A) = \mathbb{P}(\{\omega_1, \omega_2, \dots, \omega_k\}) = \sum_{i=1}^k \mathbb{P}(\{\omega_i\})$$

- ▶ If Ω consists of n equally likely outcomes (i.e. a uniform distribution), then the probability of any event A is

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|} = \frac{|A|}{n}$$

- ▶ Suppose we toss a **fair** dice twice. The sample space is $\Omega = \{(t_1, t_2) : t_1, t_2 = 1, 2, \dots, 6\}$. Let A be the event that the sum of two tosses being less than five. What is $\mathbb{P}(A)$?

Probability of an event (discrete case)

- The probability of any event $A = \{\omega_1, \omega_2, \dots, \omega_k\}$ ($\omega \in \Omega$) is the sum of the probabilities of its elements:

$$\mathbb{P}(A) = \mathbb{P}(\{\omega_1, \omega_2, \dots, \omega_k\}) = \sum_{i=1}^k \mathbb{P}(\{\omega_i\})$$

- If Ω consists of n equally likely outcomes (i.e. a uniform distribution), then the probability of any event A is

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|} = \frac{|A|}{n}$$

- Suppose we toss a **fair** dice twice. The sample space is $\Omega = \{(t_1, t_2) : t_1, t_2 = 1, 2, \dots, 6\}$. Let A be the event that the sum of two tosses being less than five. What is $\mathbb{P}(A)$?

Conditional probability

If $\mathbb{P}(B) > 0$, the **conditional probability** of A *given* B is

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

Note: $\mathbb{P}(A|B) \neq \mathbb{P}(B|A)$ (in general)

The **chain rule** can be obtained by rewriting the above expression as follows:

$$\mathbb{P}(A \cap B) = \mathbb{P}(B)\mathbb{P}(A|B) = \mathbb{P}(A)\mathbb{P}(B|A)$$

More generally, we have

$$\mathbb{P}(A_1 \cap A_2 \cap A_3 \dots) = \mathbb{P}(A_1)\mathbb{P}(A_2|A_1)\mathbb{P}(A_3|A_1A_2)\dots$$

Independence of events

Two events A and B are called **independent** if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B).$$

A set of events $A_j (j \in J)$ are called **mutually independent** if

$$\mathbb{P}\left(\bigcap_{j \in J} A_j\right) = \prod_{j \in J} \mathbb{P}(A_j).$$

Conditional probability gives another interpretation of independence: A and B are independent if the *unconditional probability* is the **same** as the *conditional probability*.

When combined with other properties of probability, independence can sometimes **simplify the calculation** of the probability of certain events.

Example

Consider a **fair** coin. What is the probability of *at least one head in the first 10 (**independent**) tosses*?

Let A be the event “at least one head in 10 tosses”. Then, A^c is the event “No heads in 10 tosses” (all 10 tosses being tails).

We have

$$\begin{aligned}\mathbb{P}(A) &= 1 - \mathbb{P}(A^c) \\ &= 1 - \mathbb{P}(\underbrace{T \cap T \cap T \cap \dots \cap T}_{10 \text{ times}}) \\ &= 1 - \prod_{i=1}^{10} \mathbb{P}(T) \quad (\text{independent tosses}) \\ &= 1 - (1/2)^{10} \quad (\text{fair coin})\end{aligned}$$

Example

Consider a **fair** coin. What is the probability of *at least one head in the first 10 (**independent**) tosses*?

Let A be the event “at least one head in 10 tosses”. Then, A^c is the event “No heads in 10 tosses” (all 10 tosses being tails).

We have

$$\begin{aligned}\mathbb{P}(A) &= 1 - \mathbb{P}(A^c) \\ &= 1 - \mathbb{P}(\underbrace{T \cap T \cap T \cap \dots \cap T}_{10 \text{ times}}) \\ &= 1 - \prod_{i=1}^{10} \mathbb{P}(T) \quad (\text{independent tosses}) \\ &= 1 - (1/2)^{10} \quad (\text{fair coin})\end{aligned}$$

Example

Consider a **fair** coin. What is the probability of *at least one head in the first 10 (**independent**) tosses*?

Let A be the event “at least one head in 10 tosses”. Then, A^c is the event “No heads in 10 tosses” (all 10 tosses being tails).

We have

$$\begin{aligned}\mathbb{P}(A) &= 1 - \mathbb{P}(A^c) \\ &= 1 - \mathbb{P}(\underbrace{T \cap T \cap T \cap \dots \cap T}_{10 \text{ times}}) \\ &= 1 - \prod_{i=1}^{10} \mathbb{P}(T) \quad (\text{independent tosses}) \\ &= 1 - (1/2)^{10} \quad (\text{fair coin})\end{aligned}$$

Example

Consider a **fair** coin. What is the probability of *at least one head in the first 10 (**independent**) tosses*?

Let A be the event “at least one head in 10 tosses”. Then, A^c is the event “No heads in 10 tosses” (all 10 tosses being tails).

We have

$$\begin{aligned}\mathbb{P}(A) &= 1 - \mathbb{P}(A^c) \\ &= 1 - \mathbb{P}(\underbrace{T \cap T \cap T \cap \dots \cap T}_{10 \text{ times}}) \\ &= 1 - \prod_{i=1}^{10} \mathbb{P}(T) \quad (\text{independent tosses}) \\ &= 1 - (1/2)^{10} \quad (\text{fair coin})\end{aligned}$$

Example

Consider a **fair** coin. What is the probability of *at least one head in the first 10 (**independent**) tosses*?

Let A be the event “at least one head in 10 tosses”. Then, A^c is the event “No heads in 10 tosses” (all 10 tosses being tails).

We have

$$\begin{aligned}\mathbb{P}(A) &= 1 - \mathbb{P}(A^c) \\ &= 1 - \mathbb{P}(\underbrace{T \cap T \cap T \cap \dots \cap T}_{10 \text{ times}}) \\ &= 1 - \prod_{i=1}^{10} \mathbb{P}(T) \quad (\text{independent tosses}) \\ &= 1 - (1/2)^{10} \quad (\text{fair coin})\end{aligned}$$

Example

Consider a **fair** coin. What is the probability of *at least one head in the first 10 (**independent**) tosses*?

Let A be the event “at least one head in 10 tosses”. Then, A^c is the event “No heads in 10 tosses” (all 10 tosses being tails).

We have

$$\begin{aligned}\mathbb{P}(A) &= 1 - \mathbb{P}(A^c) \\ &= 1 - \mathbb{P}(\underbrace{T \cap T \cap T \cap \dots \cap T}_{10 \text{ times}}) \\ &= 1 - \prod_{i=1}^{10} \mathbb{P}(T) \quad (\text{independent tosses}) \\ &= 1 - (1/2)^{10} \quad (\text{fair coin})\end{aligned}$$

Exercise

Consider tossing a **fair** dice. Let A be the event that the result is an odd number, and $B = \{1, 2, 3\}$.

- ▶ Compute $\mathbb{P}(A|B)$
- ▶ Compute $\mathbb{P}(A)$
- ▶ Are A and B independent?

Law of total probability

Let A_1, A_2, \dots, A_n be a partition of Ω . Then, for any $B \subseteq \Omega$, we have that

$$\mathbb{P}(B) = \sum_{i=1}^n \mathbb{P}(B \cap A_i) = \sum_{i=1}^n \mathbb{P}(B|A_i)\mathbb{P}(A_i)$$




Image source: https://mathwiki.cs.ut.ee/probability/04_total_probability

Law of total probability

The **law of total probability** is a combination of **additivity** and **conditional probability**. In fact, we have

$$\begin{aligned}\mathbb{P}(B) &= \mathbb{P}((B \cap A_1) \cup (B \cap A_2) \cup \dots \cup (B \cap A_k)) \\ &= \sum_{i=1}^n \mathbb{P}(B \cap A_i) \\ &= \sum_{i=1}^n \mathbb{P}(B|A_i)\mathbb{P}(A_i)\end{aligned}$$

Bayes' Rule

Let A_1, A_2, \dots, A_n be a partition of Ω . **Bayes' Rule** states that

$$\mathbb{P}(A_i|B) = \frac{\mathbb{P}(B|A_i)\mathbb{P}(A_i)}{\sum_{i=1}^n \mathbb{P}(B|A_i)\mathbb{P}(A_i)}.$$

Roughly, Bayes' rule allows us to calculate $\mathbb{P}(A_i|B)$ from $\mathbb{P}(B|A_i)$. This is useful when $\mathbb{P}(A_i|B)$ is *not obvious to calculate* but $\mathbb{P}(B|A_i)$ and $\mathbb{P}(A_i)$ are easy to obtain.

Bayes' Rule is a combination of **conditional probability** and the **law of total probability**:

$$\mathbb{P}(A_i|B) = \frac{\mathbb{P}(A_i \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A_i)\mathbb{P}(A_i)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A_i)\mathbb{P}(A_i)}{\sum_{i=1}^n \mathbb{P}(B|A_i)\mathbb{P}(A_i)}.$$

Bayes' Rule

Let A_1, A_2, \dots, A_n be a partition of Ω . **Bayes' Rule** states that

$$\mathbb{P}(A_i|B) = \frac{\mathbb{P}(B|A_i)\mathbb{P}(A_i)}{\sum_{i=1}^n \mathbb{P}(B|A_i)\mathbb{P}(A_i)}.$$

Roughly, Bayes' rule allows us to calculate $\mathbb{P}(A_i|B)$ from $\mathbb{P}(B|A_i)$. This is useful when $\mathbb{P}(A_i|B)$ is *not obvious to calculate* but $\mathbb{P}(B|A_i)$ and $\mathbb{P}(A_i)$ are easy to obtain.

Bayes' Rule is a combination of **conditional probability** and the **law of total probability**:

$$\mathbb{P}(A_i|B) = \frac{\mathbb{P}(A_i \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A_i)\mathbb{P}(A_i)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A_i)\mathbb{P}(A_i)}{\sum_{i=1}^n \mathbb{P}(B|A_i)\mathbb{P}(A_i)}.$$

Example

Suppose there are three types of emails:

- ▶ $A_1 = \text{SPAM}$
- ▶ $A_2 = \text{Low Priority}$
- ▶ $A_3 = \text{High Priority.}$

Based on previous experience, we have

$$\mathbb{P}(A_1) = 0.85, \mathbb{P}(A_2) = 0.1, \mathbb{P}(A_3) = 0.05.$$

Let B the event that an email contains the word “free”, then

$$\mathbb{P}(B|A_1) = 0.9, \mathbb{P}(B|A_2) = 0.1, \mathbb{P}(B|A_3) = 0.1.$$

When we receive an email containing the word “free”, **what is the probability that it is a spam?**

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Random variables

Often we are interested in dealing with *summaries of experiments* rather than the actual *outcome*.

For instance, suppose we toss a coin three times. But we may only be interested in a summary such as the number of heads. We have

$$\begin{array}{cccccccc} \Omega = \{ & \underbrace{HHH} & , & \underbrace{HHT} & , & \underbrace{HTH} & , & \underbrace{THH} & , & \underbrace{TTH} & , & \underbrace{THT} & , & \underbrace{HTT} & , & \underbrace{TTT} & \} \\ \downarrow & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathbb{R} & 3 & & 2 & & 2 & & 2 & & 1 & & 1 & & 1 & & 0 \end{array}$$

These summary statistics are called **random variables**. Specifically, a random variable X is a *function* from the sample space Ω to the reals:

$$X : \Omega \rightarrow \mathbb{R}$$

Random variables

A random variable (r. v.) can be seen as a **mapping** between

- ▶ a distribution on Ω ,

to

- ▶ a distribution on the reals (or the *range* of the r. v., $\mathcal{X} \subseteq \mathbb{R}$)

Formally, we have that for some subset $S \subseteq \mathcal{X}$,

$$\mathbb{P}_X(X \in S) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \in S\}).$$

Random variables

$$\Omega = \{\underbrace{HHH}_{\downarrow 3}, \underbrace{HHT}_{\downarrow 2}, \underbrace{HTH}_{\downarrow 2}, \underbrace{THH}_{\downarrow 2}, \underbrace{TTH}_{\downarrow 1}, \underbrace{THT}_{\downarrow 1}, \underbrace{HTT}_{\downarrow 1}, \underbrace{TTT}_{\downarrow 0}\}$$

For the previous example, we have

$$\mathbb{P}_X(X = 0) = 1/8, \quad \mathbb{P}_X(X = 1) = 3/8,$$

$$\mathbb{P}_X(X = 2) = 3/8, \quad \mathbb{P}_X(X = 3) = 1/8.$$

In the following, we will use \mathbb{P} to denote probability.

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables


Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Probability mass function



The **probability mass function** (PMF) of a random variable X is a function which specifies the *probability* of obtaining a number x . We denote the PMF as

$$p_X(x) = \mathbb{P}(X = x).$$

A function p_X is a PMF if and only if

1. $p_X(x) \geq 0, \forall x \in \mathcal{X}$
2. $\sum_{x \in \mathcal{X}} p_X(x) = 1$

What is the PMF of the previous coin flip example?

Some important discrete distributions

- ▶ Discrete **uniform** distribution on K categories. The PMF of $X \in \{C_1, C_2, \dots, C_K\}$ is given by

$$p_X(x) = \frac{1}{K}, \quad \forall x \in \{C_1, C_2, \dots, C_K\}$$

- ▶ The **Bernoulli** distribution with parameter $p \in [0, 1]$. The PMF of $X \in \{0, 1\}$ is given by

$$p_X(x) = \begin{cases} p, & \text{if } x = 1 \\ 1 - p, & \text{if } x = 0 \end{cases} = p^x(1 - p)^{1-x}$$

It can represent a coin toss when the coin has bias p where 1 denotes heads and 0 denotes tails.

- ▶ Other important distributions: Binomial, Geometric, Poisson, etc.
- ▶ The symbol “ \sim ” denotes “distributed as”, i.e. $X \sim \text{Ber}(p)$ means that X has a Bernoulli distribution with parameter p .

Some important discrete distributions

- ▶ Discrete **uniform** distribution on K categories. The PMF of $X \in \{C_1, C_2, \dots, C_K\}$ is given by

$$p_X(x) = \frac{1}{K}, \quad \forall x \in \{C_1, C_2, \dots, C_K\}$$

- ▶ The **Bernoulli** distribution with parameter $p \in [0, 1]$. The PMF of $X \in \{0, 1\}$ is given by

$$p_X(x) = \begin{cases} p, & \text{if } x = 1 \\ 1 - p, & \text{if } x = 0 \end{cases} = p^x(1 - p)^{1-x}$$

It can represent a coin toss when the coin has bias p where 1 denotes heads and 0 denotes tails.

- ▶ Other important distributions: Binomial, Geometric, Poisson, etc.
- ▶ The symbol “ \sim ” denotes “distributed as”, i.e. $X \sim \text{Ber}(p)$ means that X has a Bernoulli distribution with parameter p .

Some important discrete distributions

- ▶ Discrete **uniform** distribution on K categories. The PMF of $X \in \{C_1, C_2, \dots, C_K\}$ is given by

$$p_X(x) = \frac{1}{K}, \quad \forall x \in \{C_1, C_2, \dots, C_K\}$$

- ▶ The **Bernoulli** distribution with parameter $p \in [0, 1]$. The PMF of $X \in \{0, 1\}$ is given by

$$p_X(x) = \begin{cases} p, & \text{if } x = 1 \\ 1 - p, & \text{if } x = 0 \end{cases} = p^x(1 - p)^{1-x}$$

It can represent a coin toss when the coin has bias p where 1 denotes heads and 0 denotes tails.

- ▶ Other important distributions: Binomial, Geometric, Poisson, etc.
- ▶ The symbol “ \sim ” denotes “distributed as”, i.e. $X \sim \text{Ber}(p)$ means that X has a Bernoulli distribution with parameter p .

Some important discrete distributions

- ▶ Discrete **uniform** distribution on K categories. The PMF of $X \in \{C_1, C_2, \dots, C_K\}$ is given by

$$p_X(x) = \frac{1}{K}, \quad \forall x \in \{C_1, C_2, \dots, C_K\}$$

- ▶ The **Bernoulli** distribution with parameter $p \in [0, 1]$. The PMF of $X \in \{0, 1\}$ is given by

$$p_X(x) = \begin{cases} p, & \text{if } x = 1 \\ 1 - p, & \text{if } x = 0 \end{cases} = p^x(1 - p)^{1-x}$$

It can represent a coin toss when the coin has bias p where 1 denotes heads and 0 denotes tails.

- ▶ Other important distributions: Binomial, Geometric, Poisson, etc.
- ▶ The symbol “ \sim ” denotes “distributed as”, i.e. $X \sim \text{Ber}(p)$ means that X has a Bernoulli distribution with parameter p .

Expectation

The **expectation** of a random variable X is

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \, p_X(x).$$




Image source: Introduction to Probability for Data Science, Stanley H. Chan.

Expectation and its properties

For any function g , we have

$$\mathbb{E}[g(X)] = \sum_{x \in \mathcal{X}} g(x) p_X(x).$$

For any function g and h ,

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)].$$

For any constant c ,

$$\mathbb{E}[cX] = c \mathbb{E}[X].$$

For any constant c ,

$$\mathbb{E}[X + c] = \mathbb{E}[X] + c.$$

Moments and variance

The k -th **moment** of a random variable X is

$$\mathbb{E}[X^k] = \sum_{x \in \mathcal{X}} x^k p_X(x).$$

The **variance** of a random variable X is

$$\text{Var}(X) = \mathbb{E}[(X - \mu_X)^2],$$

where $\mu_X = \mathbb{E}[X]$. The **standard deviation** of X is $\sqrt{\text{Var}(X)}$.

Useful properties of the variance include:

- ▶ $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$
- ▶ $\text{Var}(cX) = c^2 \text{Var}(X)$
- ▶ $\text{Var}(X + c) = \text{Var}(X)$

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Probability density function



Image source:: Introduction to Probability for Data Science, Stanley H. Chan.

The **probability density function** (PDF) of a continuous random variable X is a function f_X , when integrated over an interval $[a, b]$, yields the probability of obtaining $\{x : a \leq x \leq b\}$:

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx.$$

For a PDF, we have:

$$(1) \quad f_X(x) \geq 0, \forall x \in \mathcal{X} \quad (2) \quad \int_{\mathcal{X}} f_X(x) dx = 1$$

Note: $f_X(x)$ is not the probability of $X = x$ since we can have $f_X(x) > 1$.

Some important continuous distributions

- ▶ Continuous **uniform** distribution on interval $[a, b]$. The PDF is given by

$$f_X(x) = \frac{1}{b-a} \quad (x \in [a, b]).$$

We write $X \sim \mathcal{U}[a, b]$.

- ▶ **Gaussian** or **Normal** distribution. With a location (mean) μ and scale (standard deviation) σ , the PDF is given by

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \quad (x \in \mathbb{R}).$$

We write $X \sim \mathcal{N}(\mu, \sigma^2)$.

Some important continuous distributions

- ▶ Continuous **uniform** distribution on interval $[a, b]$. The PDF is given by

$$f_X(x) = \frac{1}{b-a} \quad (x \in [a, b]).$$

We write $X \sim \mathcal{U}[a, b]$.

- ▶ **Gaussian** or **Normal** distribution. With a location (mean) μ and scale (standard deviation) σ , the PDF is given by

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \quad (x \in \mathbb{R}).$$

We write $X \sim \mathcal{N}(\mu, \sigma^2)$.

Expectation and its properties

The **expectation** of a continuous random variable X is given by

$$\mathbb{E}[X] = \int_{\mathcal{X}} x f_X(x) dx.$$

For any function g , we have

$$\mathbb{E}[g(X)] = \int_{\mathcal{X}} g(x) f_X(x) dx.$$

Let $I_A(X) = \begin{cases} 1, & X \in A \\ 0, & X \notin A \end{cases}$. Then, we have

$$\mathbb{E}[I_A(X)] = \int_{\mathcal{X}} I_A(x) f_X(x) dx = \int_A f_X(x) dx = \mathbb{P}(X \in A).$$

Expectation and its properties

The **expectation** of a continuous random variable X is given by

$$\mathbb{E}[X] = \int_{\mathcal{X}} x f_X(x) dx.$$

For any function g , we have

$$\mathbb{E}[g(X)] = \int_{\mathcal{X}} g(x) f_X(x) dx.$$

Let $I_A(X) = \begin{cases} 1, & X \in A \\ 0, & X \notin A \end{cases}$. Then, we have

$$\mathbb{E}[I_A(X)] = \int_{\mathcal{X}} I_A(x) f_X(x) dx = \int_A f_X(x) dx = \mathbb{P}(X \in A).$$

Expectation and its properties

The **expectation** of a continuous random variable X is given by

$$\mathbb{E}[X] = \int_{\mathcal{X}} x f_X(x) dx.$$

For any function g , we have

$$\mathbb{E}[g(X)] = \int_{\mathcal{X}} g(x) f_X(x) dx.$$

Let $I_A(X) = \begin{cases} 1, & X \in A \\ 0, & X \notin A \end{cases}$. Then, we have

$$\mathbb{E}[I_A(X)] = \int_{\mathcal{X}} I_A(x) f_X(x) dx = \int_A f_X(x) dx = \mathbb{P}(X \in A).$$

Moments and variance

The k -th **moment** of a continuous random variable X is

$$\mathbb{E}[X^k] = \int_{\mathcal{X}} x^k f_X(x) dx$$

The **variance** of a continuous random variable X is

$$\text{Var}(X) = \mathbb{E}[(X - \mu_X)^2] = \int_{\mathcal{X}} (x - \mu_X)^2 f_X(x) dx,$$

where $\mu_X = \mathbb{E}[X]$. The **standard deviation** of X is $\sqrt{\text{Var}(X)}$.

See the useful properties of the variance introduced previously.

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions



Conditional expectations

Random vectors (more than two variables)

Inference

More than one random variable?

- ▶ **Multivariate** random variables or **random vectors** are ubiquitous in modern data analysis.
- ▶ The uncertainty in the random vector is characterized by a **joint** PMF or PDF.



Important concepts

- ▶ Joint distribution
- ▶ Marginal distribution
- ▶ Independence
- ▶ Joint expectations
- ▶ Covariance and correlation
- ▶ Conditional distribution
- ▶ Conditional expectations

Joint distributions

- ▶ $f_X(x)$
- ▶ $f_{X_1, X_2}(x_1, x_2)$
- ▶ $f_{X_1, X_2, X_3}(x_1, x_2, x_3)$
- ▶ \dots
- ▶ $f_{X_1, \dots, X_n}(x_1, \dots, x_n)$
- ▶ We often just write $f_X(x)$ when the dimensionality is clear from context.

In the following, we will mainly focus on **bivariate random variables**.

Joint PMF

Let X and Y be two discrete random variables. The **joint PMF** of X and Y is defined as

$$p_{X,Y}(x, y) = \mathbb{P}(X = x \text{ and } Y = y).$$




Image source:: Introduction to Probability for Data Science, Stanley H. Chan.

For any $A \subseteq \mathcal{X} \times \mathcal{Y}$, we have

$$\mathbb{P}((X, Y) \in A) = \sum_{(x,y) \in A} p_{X,Y}(x, y).$$

Joint PMF

Let X and Y be two discrete random variables. The **joint PMF** of X and Y is defined as

$$p_{X,Y}(x, y) = \mathbb{P}(X = x \text{ and } Y = y).$$




Image source:: Introduction to Probability for Data Science, Stanley H. Chan.

For any $A \subseteq \mathcal{X} \times \mathcal{Y}$, we have

$$\mathbb{P}((X, Y) \in A) = \sum_{(x,y) \in A} p_{X,Y}(x, y).$$

Example

Let X be a coin flip, Y be a dice. Find the **joint PMF**.

The *sample space* of X is $\{0, 1\}$. The *sample space* of Y is $\{1, 2, 3, 4, 5, 6\}$. The joint PMF is

	Y					
	1	2	3	4	5	6
X = 0	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$
X = 1	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{12}$

Equivalently, we have

$$p_{X,Y}(x,y) = \frac{1}{12}, \quad x = 0, 1, \quad y = 1, 2, 3, 4, 5, 6.$$

Joint PDF

Let X and Y be two **continuous** random variables. The **joint PDF** of X and Y is a function $f_{X,Y}(x,y)$ that can be integrated to yield a probability:

$$\mathbb{P}((X, Y) \in A) = \int_A f_{X,Y}(x,y) dx \, dy,$$

for any $A \subseteq \mathcal{X} \times \mathcal{Y}$.




Image source:: Introduction to Probability for Data Science, Stanley H. Chan.


Marginal distribution

The **marginal PMFs** are defined as

$$p_X(x) = \sum_{y \in \mathcal{Y}} p_{X,Y}(x, y) \text{ and } p_Y(y) = \sum_{x \in \mathcal{X}} p_{X,Y}(x, y),$$

and the **marginal PDFs** are defined as

$$f_X(x) = \int_{\mathcal{Y}} f_{X,Y}(x, y) dy \text{ and } f_Y(y) = \int_{\mathcal{X}} f_{X,Y}(x, y) dx.$$



Independence

If two random variables X and Y are **independent**, then

$$p_{X,Y}(x,y) = p_X(x)p_Y(y), \quad \text{and } f_{X,Y}(x,y) = f_X(x)f_Y(y)$$

If a sequence of random variables X_1, \dots, X_N are **independent**, then their joint PDF (or joint PMF) can be **factorized**:

$$f_{X_1, \dots, X_n}(x_1, \dots, x_n) = \prod_{j=1}^n f_{X_j}(x_j)$$

Independent and Identically Distributed (i.i.d.)

A collection of random variables X_1, \dots, X_N are called **independent and identically distributed (i.i.d.)** if

1. All X_1, \dots, X_N are **independent**.
2. All X_1, \dots, X_N have the **same distribution**.

Joint expectations

Recall that the expectation of a discrete random variable X is given by

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x p_X(x).$$

How about the expectation for two variables?

Let X and Y be two *discrete* random variables. For any function g , the **joint expectation** is

$$\mathbb{E}[g(X, Y)] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} g(x, y) p_{X, Y}(x, y).$$

If X and Y are *continuous*, we have

$$\mathbb{E}[g(X, Y)] = \int_{\mathcal{X}} \int_{\mathcal{Y}} g(x, y) f_{X, Y}(x, y) dx dy.$$

Joint expectations

Let $g(X, Y) = XY$. If X and Y are **discrete**, we have

$$\mathbb{E}[XY] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} xy \, p_{X,Y}(x, y).$$

If X and Y are **continuous**, we have

$$\mathbb{E}[XY] = \int_{\mathcal{X}} \int_{\mathcal{Y}} xy \, f_{X,Y}(x, y) \, dx \, dy.$$

Covariance

Let X and Y be two random variables. Then the **covariance** of X and Y is

$$\begin{aligned}\text{Cov}(X, Y) &= \mathbb{E}[(X - \mu_X)(Y - \mu_Y)] \\ &= \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y],\end{aligned}$$

where $\mu_X = E[X]$ and $\mu_Y = E[Y]$.

Note that $\text{Cov}(X, X) = \text{Var}(X)$.

Useful properties

For any X and Y , we have

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y],$$

and

$$\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y] + 2\text{Cov}(X, Y).$$

If X and Y are **independent**, then

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

Correlation

Let X and Y be two random variables. The **correlation coefficient** is

$$\rho = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}},$$

where $-1 \leq \rho \leq 1$.

- ▶ When $X = Y$ (**fully correlated**), $\rho = 1$.
- ▶ When $X = -Y$ (**fully correlated**), $\rho = -1$.
- ▶ When X and Y are **uncorrelated** then $\rho = 0$.

Independence vs correlation

Consider the following two statements:

1. X and Y are independent;
2. $\text{Cov}(X, Y) = 0$.

We have

- ▶ $(1) \implies (2)$ (independence \implies uncorrelated)
- ▶ $(2) \not\implies (1)$ (uncorrelated $\not\implies$ independence)
- ▶ Independence is a **stronger** condition than correlation

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables




Conditional distributions

Conditional expectations




Random vectors (more than two variables)

Inference

Conditional distributions



Conditional distributions



Conditional distributions

Let X and Y be two **discrete** random variables. The **conditional PMF** of Y given X is

$$p_{Y|X}(y|x) = \frac{p_{Y,X}(y, x)}{p_X(x)}.$$

Let X and Y be two **continuous** random variables. The **conditional PDF** of Y given X is

$$f_{Y|X}(y|x) = \frac{f_{Y,X}(x, y)}{f_X(x)}.$$


Example

Consider **two coins** which can take values in $\{0, 1\}$. Let Y be the *sum* of the two coins, and X , the *value of the first coin*.

- ▶ What is $p_Y(y)$?
- ▶ What is $p_{Y|X}(y|x = 1)$?
- ▶ What is $p_{Y,X}(y, x)$??

Example

$p_Y(y)$:



$p_{Y|X}(y|x=1)$:




Image source:: Introduction to Probability for Data Science, Stanley H. Chan.

Conditional distributions

Let X and Y be two **discrete** random variables. For any $A \subseteq \mathcal{Y}$, we have

$$\mathbb{P}(Y \in A | X = x) = \sum_{y \in A} p_{Y|X}(y|x),$$

and

$$\mathbb{P}(Y \in A) = \sum_{x \in \mathcal{X}} \mathbb{P}(Y \in A | X = x) p_X(x).$$

Let X and Y be two **continuous** random variables. For any $A \subseteq \mathcal{Y}$, we have

$$\mathbb{P}(Y \in A | X = x) = \int_A f_{Y|X}(y|x) dy,$$

and

$$\mathbb{P}(Y \in A) = \int_{\mathcal{X}} \mathbb{P}(Y \in A | X = x) f_X(x) dx.$$

Conditional distributions

Let X and Y be two **discrete** random variables. For any $A \subseteq \mathcal{Y}$, we have

$$\mathbb{P}(Y \in A | X = x) = \sum_{y \in A} p_{Y|X}(y|x),$$

and

$$\mathbb{P}(Y \in A) = \sum_{x \in \mathcal{X}} \mathbb{P}(Y \in A | X = x) p_X(x).$$

Let X and Y be two **continuous** random variables. For any $A \subseteq \mathcal{Y}$, we have

$$\mathbb{P}(Y \in A | X = x) = \int_A f_{Y|X}(y|x) dy,$$

and

$$\mathbb{P}(Y \in A) = \int_{\mathcal{X}} \mathbb{P}(Y \in A | X = x) f_X(x) dx.$$

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables



Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Conditional expectations



Conditional expectations

For a **discrete** random variable Y , the **conditional expectation** of Y given X is

$$\mathbb{E}[Y|X = x] = \sum_{y \in \mathcal{Y}} y \, p_{Y|X}(y|x).$$

For a **continuous** random variable Y , the conditional expectation of Y given X is

$$\mathbb{E}[Y|X = x] = \int_{\mathcal{Y}} y \, f_{Y|X}(y|x) dy$$

The summation/integration is taken w.r.t. y , because $X = x$ is **given and fixed**.

Conditional expectations

For a **discrete** random variable Y , the **conditional expectation** of Y given X is

$$\mathbb{E}[Y|X = x] = \sum_{y \in \mathcal{Y}} y \, p_{Y|X}(y|x).$$

For a **continuous** random variable Y , the conditional expectation of Y given X is

$$\mathbb{E}[Y|X = x] = \int_{\mathcal{Y}} y \, f_{Y|X}(y|x) dy$$

The summation/integration is taken w.r.t. y , because $X = x$ is **given and fixed**.

Law of Total Expectation

The **law of total expectation**² is a decomposition rule which allows to decompose the computation of $\mathbb{E}[Y]$ into **conditional expectations** that are smaller/easier to compute.

$$\mathbb{E}[Y] = \sum_{x \in \mathcal{X}} \mathbb{E}[Y|X = x] p_X(x)$$

or

$$\mathbb{E}[Y] = \int_{\mathcal{X}} \mathbb{E}[Y|X = x] f_X(x) dx$$

²Also known as the **law of iterated expectations** and the **tower rule**.

Law of Total Expectation

The **law of total expectation** can also be written as

$$\mathbb{E}[Y] = \mathbb{E}_X[\mathbb{E}_{Y|X}[Y|X]].$$

Note the difference between

$$h(x) = \mathbb{E}_{Y|X}[Y|X = x], \quad (\text{A deterministic function in } x)$$

and

$$h(X) = \mathbb{E}_{Y|X}[Y|X]. \quad (\text{A function of the random variable } X)$$


Example

Suppose there are two classes of cars. Let $C \in \{1, 2\}$ be the **class** and $S \in \mathbb{R}$, the **speed**.

We know that

- ▶ $\mathbb{P}(C = 1) = p$
- ▶ When $C = 1$, $S \sim \mathcal{N}(\mu_1, \sigma_1^2)$
- ▶ When $C = 2$, $S \sim \mathcal{N}(\mu_2, \sigma_2^2)$

You see a car on the freeway, what is its **average speed**?



Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Random vectors

An image from a dataset can be represented by a **high-dimensional** vector.




Image source:: Introduction to Probability for Data Science, Stanley H. Chan.

Random vectors

Random vector:

$$X = \begin{pmatrix} X_1 \\ X_2 \\ \dots \\ X_n \end{pmatrix} \text{ and } x = \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix}$$

Joint PDF:

$$f_X(x) = f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n)$$

Probability:

$$\mathbb{P}(X \in A) = \int_A f_X(x) dx$$

Mean vector and covariance matrix

Let $X = (X_1, X_2, \dots, X_n)^T$ be a random vector. The **expectation** is

$$\mu = \mathbb{E}[X] = \begin{pmatrix} \mathbb{E}[X_1] \\ \mathbb{E}[X_2] \\ \dots \\ \mathbb{E}[X_n] \end{pmatrix}.$$

The **covariance** matrix is

$$\Sigma = \begin{pmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \dots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \dots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \dots & \text{Var}(X_n) \end{pmatrix},$$

which can be written in a more compact way as


$$\Sigma = \mathbb{E}[(X - \mu)(X - \mu)^T].$$

Diagonal covariance matrix


If the coordinates X_1, X_2, \dots, X_n are *uncorrelated*, the covariance matrix is a **diagonal** matrix:

$$\Sigma = \begin{pmatrix} \text{Var}(X_1) & 0 & \dots & 0 \\ 0 & \text{Var}(X_2) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \text{Var}(X_n) \end{pmatrix}$$


Bivariate Gaussian



$$(\mu, \Sigma) = \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 5 & 0 \\ 0 & 0.5 \end{bmatrix}$$



$$\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 & -0.5 \\ -0.5 & 1 \end{bmatrix}$$



$$\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 & 1.9 \\ 1.9 & 2 \end{bmatrix}$$

Multivariate Gaussian

The PDF of a d -dimensional **joint Gaussian** is given by

$$f_X(x) = \frac{1}{\sqrt{(2\pi)^d |\Sigma|}} \exp \left\{ -\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu) \right\}$$

Outline

Probability

Random variables

Discrete random variables

Continuous random variables

Multivariate random variables

Conditional distributions

Conditional expectations

Random vectors (more than two variables)

Inference

Estimators

A central concept of machine learning (or statistics) is to **learn (or estimate)** certain properties about some underlying (stochastic) process on the basis of samples (data).

Point estimation refers to calculating a single “best guess” of the value of an unknown quantity of interest, which could be a **parameter** or a **density function**. We typically use $\hat{\theta}$ to denote a point estimator for θ .

Given $X_1, X_2, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} p_X$, a (point) **estimator** is a function of the observed sample, i.e.

$$\hat{\theta} = T(X_1, X_2, \dots, X_n),$$

so that $\hat{\theta}$ is a *random variable*.

For example, the *sample mean* $\hat{\theta} = \bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ is an estimator for the expectation ($\theta = \mathbb{E}[X]$).

Properties of estimators

The **bias of an estimator** is given by

$$b(\hat{\theta}) = \mathbb{E}[\hat{\theta}] - \theta.$$

The **variance of an estimator** is given by

$$v(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \mathbb{E}[\hat{\theta}])^2].$$

The **standard errors of an estimator** is given by

$$se(\hat{\theta}) = \sqrt{v(\hat{\theta})},$$

i.e., its standard deviation.

The **sampling distribution** of an estimator is the probability distribution of the estimator.

Example - The sample mean

Let $X_1, X_2, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} p_X$, with $\mathbb{E}[X] = \mu_X$ and $\text{Var}(X) = \sigma_X^2$. The *sample mean* estimator is defined as

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i.$$

What are the bias and variance of \bar{X}_n ?

Since $\mathbb{E}[\bar{X}_n] = \mu_X$, \bar{X}_n is unbiased, i.e. the bias is equal to zero. Also, using the fact that $\text{Var}(\sum_{i=1}^n X_i) = \sum_{i=1}^n \sum_{j=1}^n \text{Cov}(X_i, X_j)$, we can show that $\text{Var}(\bar{X}_n) = \frac{\sigma_X^2}{n}$.

Example - The sample mean

Let $X_1, X_2, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} p_X$, with $\mathbb{E}[X] = \mu_X$ and $\text{Var}(X) = \sigma_X^2$. The *sample mean* estimator is defined as

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i.$$

What are the bias and variance of \bar{X}_n ?

Since $\mathbb{E}[\bar{X}_n] = \mu_X$, \bar{X}_n is unbiased, i.e. the bias is equal to zero. Also, using the fact that $\text{Var}(\sum_{i=1}^n X_i) = \sum_{i=1}^n \sum_{j=1}^n \text{Cov}(X_i, X_j)$, we can show that $\text{Var}(\bar{X}_n) = \frac{\sigma_X^2}{n}$.

More on the sample mean

- ▶ The variance of the average is **much smaller** than the variance of the individual random variables. This is one of the core principles of statistics and helps us learn various quantities reliably by making **repeated independent measurements**.
- ▶ Why independent measurements are **essential**? The extreme case of non-independence is when $X_1 = X_2 = \dots = X_n$, for which we have

$$\text{Var}(\bar{X}_n) = \sigma_X^2.$$

Inference

Let $y_1, y_2, \dots, y_n \stackrel{\text{i.i.d.}}{\sim} p_Y$. How can we estimate p_Y ?

- ▶ We often **assume** that the sample was generated from some (parametric) model.
- ▶ When we specify a model, we hope that it can provide a useful **approximation** to the data generation mechanism.
- ▶ The George Box quote is worth remembering in this context: “**all models are wrong, but some are useful**”.

Maximum likelihood estimation

Let us restrict ourselves to a set of possible distributions $p(y; \theta)$, described by a finite number of parameters $\theta \in \Theta$.

An example for $y \in \mathbb{R}$ is

$$\left\{ p(y; \mu; \sigma) = \frac{1}{2\sigma\sqrt{2\pi}} \exp \left\{ \frac{(y - \mu)^2}{\sigma^2} \right\} : \mu \in \mathbb{R}, \sigma > 0 \right\},$$

where $\theta = (\mu, \sigma)^T$, and, for $y \in \{0, 1\}$,

$$\{ p(y; \alpha) = \alpha^y (1 - \alpha)^{1-y} : 0 \leq \alpha \leq 1 \},$$

where $\theta = \alpha$.

The goal of maximum likelihood estimation is to select the distribution $p(y; \theta)$ that is **most likely** to have generated the sample y_1, y_2, \dots, y_n .

Maximum likelihood estimation

The **likelihood function** is defined as

$$\mathcal{L}(\theta) \equiv \mathcal{L}(\theta; y_1, y_2, \dots, y_n) \quad (1)$$

$$= p(y_1, y_2, \dots, y_n; \theta) \quad (2)$$

$$= \prod_{i=1}^n p(y_i; \theta). \quad (3)$$

The **maximum likelihood estimator**, or MLE – denoted by $\hat{\theta}$ – is the value of θ that maximizes $\mathcal{L}(\theta)$. Note that $\hat{\theta}$ also maximizes the **log-likelihood function** $\log \mathcal{L}(\theta)$.

We write

$$\hat{\theta} = \operatorname{argmax}_{\theta \in \Theta} \mathcal{L}(\theta) = \operatorname{argmax}_{\theta \in \Theta} \log \mathcal{L}(\theta),$$

where Θ is the parameter space.

Example

We observe y_1, \dots, y_n where $y_i \in \{0, 1\}$ with unknown PMF p_Y . If we assume

$$y_1, \dots, y_n \stackrel{\text{i.i.d.}}{\sim} p(y; \alpha),$$

where

$$p(y; \alpha) = \alpha^y (1 - \alpha)^{1-y}$$

with $0 \leq \alpha \leq 1$.

What is the maximum likelihood estimate $\hat{\alpha}$?

Example

The **likelihood function** is given by

$$\begin{aligned}\mathcal{L}(\alpha; y_1, \dots, y_n) &= p(y_1, \dots, y_n; \alpha) \\ &= \prod_{i=1}^n p(y_i; \alpha) \\ &= \prod_{i=1}^n \alpha^{y_i} (1 - \alpha)^{1-y_i} \\ &= \alpha^{\sum_{i=1}^n y_i} (1 - \alpha)^{\sum_{i=1}^n (1-y_i)},\end{aligned}$$

and the **log-likelihood function** is given by

$$\begin{aligned}\log \mathcal{L}(\alpha; y_1, \dots, y_n) &= \sum_{i=1}^n y_i \log(\alpha) + (1 - y_i) \log(1 - \alpha) \\ &= n\bar{y} \log(\alpha) + n(1 - \bar{y}) \log(1 - \alpha),\end{aligned}$$

where $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$.

Example

The first derivative of the log-likelihood is given by

$$(\log \mathcal{L})'(\alpha) = n\bar{y}\frac{1}{\alpha} - n(1 - \bar{y})\frac{1}{1 - \alpha}.$$

A necessary condition for a maximum is given by

$$(\log \mathcal{L})'(\alpha) = 0 \iff \hat{\alpha} = \bar{y}.$$

We can verify that it is indeed a maximum by checking that the second derivative of the log-likelihood at $\hat{\alpha}$ is indeed negative, i.e. $(\log \mathcal{L})''(\hat{\alpha}) < 0$.