

Summary Data on Acid Cropland and Lime Requirements in Malawi

GAIA

www.acidsoils.africa

Last Updated: November 8, 2024


Outlook

- 1 Introduction
- 2 Acidic Cropland
- 3 Additional Production
- 4 Yield Loss
- 5 Yield Response
- 6 Lime Requirement


Introduction

- The slide deck provides a **data-driven, ex-ante analysis** of soil acidity across **Malawi**, evaluating the scale of affected areas, potential crop yield impacts, and the expected benefits of lime application for proactive soil remediation.
- **Highlights:**
 - **Acidic Cropland:** Identifies regions with high acidity levels impacting crop yields.
 - **Yield Loss:** Quantifies productivity losses in key crops due to acidic soils.
 - **Lime Requirements:** Estimates the lime amounts needed to improve soil health in these areas.
 - **Economic Potential:** Highlights estimated increases in production and production value from lime application.
- Provide policymakers, farmers, and stakeholders with actionable insights to prioritize areas with high potential for productivity gains and economic return from soil remediation.


Top 10 Districts in Malawi with the highest acidic cropland



Top 10 Districts in Malawi with the highest acidic cropland (Percentage of total cropland)




Total crop area in the top-10 most acidic Districts, not segregated by acidic cropland



Total crop area in the top-10 most acidic Districts, not segregated by acidic cropland (table)

Districts	Bean	Groundnut	Maize	Other-Cereal	Other-Legume	Other-Non-food	Other-RTBs	Potato
Chitipa	6.05	4.73	46.09	1.22	7.25	10.64	5.47	5.50
Dedza	29.40	18.52	103.19	4.01	48.81	7.30	5.82	23.63
Karonga	36.59	4.24	38.04	0.80	1.45	4.36	13.71	1.16
Lilongwe	19.70	53.00	204.13	7.30	32.26	38.19	8.79	32.53
Mangochi	11.79	17.86	108.91	14.38	22.24	17.18	8.39	13.28
Mchinji	7.77	37.49	111.69	8.45	32.12	10.15	2.91	7.77
Mzimba	20.02	24.22	144.90	8.98	32.08	30.53	16.45	16.90
Nkhata Bay	0.48	0.17	142.57	0.12	0.28	3.75	25.48	0.17
Nkhotakota	49.01	5.37	71.21	0.25	2.43	13.76	24.52	0.69
Rumphi	0.62	0.67	52.61	0.31	0.76	1.57	0.55	0.34


Potential for additional production (tone) in top 10 acidic Districts in Malawi



Potential for additional production (1000 tone) in top 10 acidic Districts in Malawi(Table)

Districts	Bean	Groundnut	Maize	Other-Cereal	Other-Legume	Other-Non-food	Other-RTBs	Potato
Chitipa	0.11	0.07	0.07	-	0.08	-	0.01	0.09
Dedza	0.06	0.19	0.08	-	0.20	-	-	0.42
Karonga	0.40	0.06	0.07	-	0.03	-	-	0.03
Lilongwe	0.03	0.16	0.16	0.01	0.06	-	0.03	0.50
Mangochi	0.06	0.14	0.18	0.02	0.11	-	-	0.29
Mchinji	0.03	0.42	0.32	0.01	0.09	-	-	0.21
Mzimba	0.11	0.14	0.43	0.01	0.12	0.01	0.01	0.74
Nkhata Bay	0.04	0.01	14.40	-	0.01	-	0.11	0.03
Nkhotakota	0.93	0.13	0.18	-	0.04	-	-	-
Rumphi	-	-	0.53	-	-	-	-	-

Potential for additional production value (US\$) in top 10 acidic Districts in Malawi




Potential for additional production value (1000 US\$) in top 10 acidic Districts in Malawi(Table)


Districts	Bean	Groundnut	Maize	Other-Cereal	Other-Legume	Other-Non-food	Other-RTBs	Potato
Chitipa	81.8	45.7	20.1	0.6	40.1	-	3.6	37.6
Dedza	46.0	116.0	21.5	1.1	104.0	3.4	1.0	178.9
Karonga	289.1	39.0	20.4	0.3	13.4	-	0.1	14.5
Lilongwe	23.2	99.3	44.7	5.1	28.7	8.7	8.0	214.6
Mangochi	40.4	88.4	50.7	5.6	56.5	0.6	0.1	124.5
Mchinji	24.7	260.0	89.4	4.2	50.6	5.4	0.1	90.8
Mzimba	80.2	85.4	119.8	4.8	57.7	22.1	1.9	319.0
Nkhata Bay	27.6	9.2	4001.6	0.3	6.8	-	29.7	11.0
Nkhotakota	672.9	82.5	50.2	-	21.8	-	-	0.7
Rumphi	2.5	1.0	147.1	0.1	0.7	0.1	-	1.2

Yield loss due to soil acidity for Maize and Groundnut in Malawi

Maize Yield Loss (%)




Groundnut Yield Loss (%)




Yield response to lime application in Malawi


Maize Yield Response (t/ha)



Groundnut Yield Response (t/ha)



Lime requirements (t) for acid soil remediation in Malawi




Lime requirements (1000 t) for acid soil remediation in Malawi (table)


Districts	Bean	Groundnut	Maize	Other-Cereal	Other-Legume	Other-RTBs	Potato	Other-Non-food
Chitipa	0.65	0.45	0.09	0.01	0.53	-	0.02	-
Dedza	0.71	0.47	-	-	0.77	-	-	-
Karonga	6.16	0.64	0.38	0.02	0.25	-	0.04	-
Lilongwe	0.43	0.57	0.28	0.07	0.23	0.01	0.06	-
Mangochi	0.71	1.08	-	-	0.71	-	-	-
Mchinji	0.22	1.54	-	-	0.25	-	-	-
Mzimba	1.42	0.99	1.05	0.09	1.02	-	0.15	-
Nkhata Bay	0.37	0.13	36.50	-	0.12	-	-	-
Nkhotakota	6.99	0.57	0.41	-	0.12	-	-	-
Rumphi	0.01	0.01	2.12	0.01	0.01	-	-	0.01

Lime rate for Maize and Groundnut in Malawi

Maize Lime Rate (t/ha)



Groundnut Lime Rate (t/ha)



Main assumptions and parameters

- Acidic cropland is defined as having acidity saturation levels above 10% of the Effective Cation Exchange Capacity (ECEC).
- Crop yields are unaffected by soil acidity until a specific acidity saturation threshold Target Acidity Saturation(TAS) is reached; beyond this threshold, yields decrease linearly, eventually reaching zero at high acidity levels.
- The TAS that separates maximum yield from yield decline is used as a critical value for making liming recommendations.
- Actual yields in acid tropical soils are a fraction of the potential yields in non-acidic soils, determined by the estimated yield loss due to acidity.
- Economic benefits from liming are calculated by multiplying the additional yield after soil remediation by median crop prices from sub-Saharan Africa (2016-2020 FAOSTAT data). The cost of liming is calculated using estimated lime application rates and a fixed lime price of 100 US\$/ton.
- The benefits of liming are assessed only for the year of application due to a lack of robust, large-scale data on long-term effects in sub-Saharan Africa.

For details, visit www.acidsoils.africa.