

Summary Data on Acid Cropland and Lime Requirements in Uganda

GAIA


www.acidsoils.africa

Last Updated: October 22, 2024


Outlook

- 1 Introduction
- 2 Acidic Cropland
- 3 Yield Loss
- 4 Lime Requirement
- 5 Yield Response
- 6 Additional Production
- 7 ROI for Lime Application
- 8 Profitability of Lime Application


Top 10 Districts in Uganda with the highest acidic cropland



Top 10 Districts in Uganda with the highest acidic cropland (Percentage of total cropland)



Total crop area in the top-10 most acidic Districts, not segregated by acidic cropland





Uganda-Summary

└ Total crop area distribution (ha)


└ Total crop area in the top-10 most acidic Districts, not segregated by acidic cropland

Districts	Bean	Cassava	Maize	Other-Cereal	Other-Legume	Other-Non-food	Other-RTBs	Sweet potato
Apac	35.58	19.10	37.75	10.90	63.61	35.85	0.14	6.48
Bushenyi	20.96	3.83	103.15	5.63	3.99	3.03	1.55	14.17
Gulu	26.64	7.69	9.77	16.48	31.01	31.73	0.01	6.84
Iganga	16.87	9.12	33.92	2.44	14.07	24.55	-	23.80
Kamuli	31.03	15.13	42.46	5.51	10.28	24.15	0.01	33.02
Katakwi	8.35	8.69	154.86	21.04	23.40	11.69	-	10.80
Lira	25.88	11.97	27.96	28.65	74.52	41.15	0.06	7.30
Luwero	8.85	5.42	40.85	83.82	77.60	20.32	0.64	9.42
Mbarara	33.37	130.09	56.56	12.10	10.20	7.55	26.52	82.78
Soroti	132.17	10.06	13.46	15.17	13.57	9.97	-	10.72

Yield loss due to soil acidity for Maize and Bean in Uganda





Lime requirements (t) for acid soil remediation in Uganda



Lime requirements (t) for acid soil remediation in Uganda (table)


Districts	Bean	Cassava	Maize	Other-Cereal	Other-Legume	Other-Non-food	Other-RTBs	Sweet potato
Apac	50.82	0.34	41.65	11.71	76.51	4.47	0.16	7.07
Bushenyi	29.92	-	104.12	5.32	3.95	0.10	1.65	5.71
Gulu	33.20	-	4.96	6.42	34.04	0.78	-	3.37
Iganga	24.09	-	19.97	1.99	16.62	0.43	-	13.32
Kamuli	44.34	-	44.79	5.74	13.54	0.40	-	35.18
Katakwi	11.69	-	69.32	7.98	16.31	0.24	-	4.31
Lira	36.98	-	25.18	26.75	99.00	1.92	0.05	6.41
Luwero	9.17	-	11.61	31.31	81.01	1.03	0.32	1.28
Mbarara	42.49	-	22.82	1.73	11.35	-	14.81	0.25
Soroti	188.81	-	14.46	16.32	17.60	1.85	-	11.89

Lime rate for Maize and Bean in Uganda




Yield response to lime application in Uganda


Maize Yield Response (t/ha)




Bean Yield Response (t/ha)





Potential for additional production (tone) in top 10 acidic Districts in Uganda





Potential for additional production value (US\$) in top 10 acidic Districts in Uganda



Return on investment for lime application in Uganda



Profitability of lime application in Uganda



Main assumptions and parameters

- Acidic cropland is defined as having acidity saturation levels above 10% of the Effective Cation Exchange Capacity (ECEC).
- Crop yields are unaffected by soil acidity until a specific acidity saturation threshold Target Acidity Saturation(TAS) is reached; beyond this threshold, yields decrease linearly, eventually reaching zero at high acidity levels.
- The TAS that separates maximum yield from yield decline is used as a critical value for making liming recommendations.
- Actual yields in acid tropical soils are a fraction of the potential yields in non-acidic soils, determined by the estimated yield loss due to acidity.
- Economic benefits from liming are calculated by multiplying the additional yield after soil remediation by median crop prices from sub-Saharan Africa (2016-2020 FAOSTAT data). The cost of liming is calculated using estimated lime application rates and a fixed lime price of 100 US\$/ton.
- The benefits of liming are assessed only for the year of application due to a lack of robust, large-scale data on long-term effects in sub-Saharan Africa.

For details, visit www.acidsoils.africa.