

Infinite Flatland medium, Isotropic Point Source, Isotropic Scattering

Exponential Random Flight

This is code to accompany the book:

A Hitchhiker's Guide to Multiple Scattering

© 2015 Eugene d'Eon

www.eugenedeon.com


Path Setup

Put a file at `~/hitchhikerpath` with the path to your hitchhiker repo so that these worksheets can find the MC data from the C++ simulations for verification

```
In[1]:= SetDirectory[Import["~/hitchhikerpath"]]
```

```
Out[1]:= /Users/eug/Documents/research/hitchhikersscatter
```

Notation



α - single-scattering albedo

Σ_t - extinction coefficient

r - radial position coordinate in medium (distance from point source at origin)

$u = \cos \theta$ - direction cosine

Analytic solutions

Fluence: exact solution (1)

$$\text{In[2]:= infflatlandisopointisoscatter}\phi_{\text{exact1}}[r_, \Sigma t_, \alpha_] := \frac{\text{Exp}[-\Sigma t r]}{2 \text{ Pi } r} + \frac{\Sigma t}{2 \text{ Pi }} \text{NIntegrate}\left[\frac{\alpha z \text{ BesselJ}[0, r z \Sigma t]}{1 + z^2 - \alpha \sqrt{1 + z^2}}, \{z, 0, \text{Infinity}\}, \text{Method} \rightarrow \text{"LevinRule"}\right]$$

Fluence: exact solution (2)

$$\text{In[3]:= infflatlandisopointisoscatter}\phi_{\text{exact2}}[r_, \Sigma t_, \alpha_] := \frac{\Sigma t \alpha}{\text{Pi }} \frac{\text{BesselK}[0, r \Sigma t \sqrt{1 - \alpha^2}]}{\text{Pi }} + \frac{\Sigma t}{2 \text{ Pi }} \text{NIntegrate}\left[\frac{z \text{ BesselJ}[0, \Sigma t r z]}{\sqrt{1 + z^2} + \alpha}, \{z, 0, \text{Infinity}\}, \text{Method} \rightarrow \text{"LevinRule"}\right]$$

Fluence: exact solution (3)

$$\text{In[4]:= infflatlandisopointisoscatter}\phi_{\text{exact3a}}[r_, \Sigma t_, \alpha_] := \text{NIntegrate}\left[\frac{\Sigma t}{2 \text{ Pi }} \frac{k \text{ BesselJ}[0, k \Sigma t r]}{\sqrt{k^2 + 1} - \alpha}, \{k, 0, \text{Infinity}\}, \text{Method} \rightarrow \text{"LevinRule"}\right]$$

$$\text{In[5]:= bessellk}[n_, x_] := \sqrt{\frac{2}{\text{Pi } x}} \text{BesselK}[n + 1/2, x];$$

$$\text{infflatlandisopointisoscatter}\phi_{\text{exact3b}}[r_, \Sigma t_, \alpha_, M_] := \frac{\text{Exp}[-\Sigma t r]}{2 \text{ Pi } r} + \frac{\alpha \Sigma t}{2 \text{ Pi }} \text{BesselK}[0, \Sigma t \sqrt{1 - \alpha^2} r] + \frac{\Sigma t}{2 \text{ Pi }} \text{Sum}\left[\frac{\alpha^{2n} n!}{(2n)!} (2 \Sigma t r)^n \text{bessellk}[n - 1, \Sigma t r], \{n, 1, M\}\right]$$

Classical diffusion approximation

$$\text{In[7]:= infflatlandisopointisoscatter}\phi_{\text{Diffusion}}[r_, \Sigma t_, \alpha_] := \frac{\Sigma t \text{BesselK}[0, \sqrt{2 - 2\alpha} r \Sigma t]}{\pi}$$

Rigorous diffusion approximation

$$\text{In[8]:= infflatlandisopointisoscatter}\phi_{\text{RigorousDiffusion}}[r_, \Sigma t_, \alpha_] := \frac{\Sigma t \alpha \text{BesselK}[0, \sqrt{1 - \alpha^2} r \Sigma t]}{\pi}$$

Grosjean-style diffusion approximation

$$\text{In[9]:= infflatlandisopointisoscatter}\phi_{\text{Grosjean}}[r_, \Sigma t_, \alpha_] := \frac{\text{Exp}[-r \Sigma t]}{2 \text{ Pi } r} + \frac{\alpha \Sigma t}{(2 - \alpha) \text{ Pi }} \text{BesselK}\left[0, r \Sigma t \left(\sqrt{2} \frac{\sqrt{1 - \alpha}}{\sqrt{2 - \alpha}}\right)\right]$$

n-th scattered fluence

```
In[10]:= infflatlandisotropicpointscatter`phi[r_, Sigma t_, alpha_, n_] :=
  2^(1/2 (-1-n)) alpha^n r^(1/2 (-1+n)) Sigma t^(1+n/2) BesselK[1/2 (-1+n), r Sigma t]
  / (pi Gamma[1+n/2])
```

Moments

```
In[11]:= infflatlandisotropicpointscatter`phi_m[Sigma t_, alpha_, m_] := (1 - alpha^2)^(-1-m/2) Sigma t^(-1-m)
  (2^m alpha Gamma[1 + m/2]^2 + Gamma[1 + m] Hypergeometric2F1[-1/2, -m/2, 1/2, alpha^2])
```

```
In[12]:= infflatlandisotropicpointscatter`phi_m[Sigma t_, alpha_, n_, m_] :=
  2^m alpha^n Sigma t^(-1-m) Gamma[1 + m/2] Gamma[1/2 (1 + m + n)]
  / Gamma[1+n/2]
```

Angular phi integral

```
In[140]:= infflatlandisotropicpointscatter`Lintegral[r_, u_, Sigma t_, alpha_, phi_] :=
  alpha Sigma t / (2 Pi) NIntegrate[phi[Sqrt[r^2 + t^2 - 2 r t u], Sigma t, alpha] Exp[-Sigma t t], {t, 0, Infinity}]
```

Angular Classical diffusion approximation

```
In[138]:= infflatlandisotropicpointscatter`Ldiffusion[r_, u_, Sigma t_, alpha_] :=
  BesselK[0, Sqrt[2] Sqrt[(1 - alpha) Sigma t^2] r] / (4 Pi^2 1/(2 Sigma t)) +
  Sqrt[2] Sqrt[(1 - alpha) Sigma t^2] BesselK[1, Sqrt[2] Sqrt[(1 - alpha) Sigma t^2] r] / (2 Pi^2) u
```

load MC data

```
In[49]:= infflatlandisotropicpointscatter`ppoints[xs_, dr_, maxx_, Sigma t_] :=
  Table[{dr[i] - 0.5 dr, xs[[i]]/Sigma t}, {i, 1, Length[xs]}}[[1 ;; -2]]
```

```
In[110]:= infflatlandisotropicpointscatter`ppointsu[xs_, du_, Sigma t_] :=
  Table[{-1.0 + du[i] - 0.5 du, Sqrt[1 - (-1.0 + du[i] - 0.5 du)^2] xs[[i]]/(2 Sigma t)},
  {i, 1, Length[xs]}}[[1 ;; -1]]
```

```
In[51]:= infflatlandisotropicpointscatter`fs =
  FileNames["code/flatland/infiniteFlatland/Isotropicpointsource/data/
  infflatland_isotropicpoint_isotropicscatter*"];
```

```

In[52]:= infflatlandisopointisoscatter`index[x_] := Module[{data,  $\alpha$ ,  $\Sigma t$ },
  data = Import[x, "Table"];
   $\Sigma t$  = data[[1, 13]];
   $\alpha$  = data[[2, 3]];
  { $\alpha$ ,  $\Sigma t$ , data}};
infflatlandisopointisoscatter`simulations =
  infflatlandisopointisoscatter`index /@ infflatlandisopointisoscatter`fs;
infflatlandisopointisoscatter`alphas =
  Union[#[[1]] & /@ infflatlandisopointisoscatter`simulations]

Out[54]= {0.01, 0.1, 0.3, 0.5, 0.7, 0.8, 0.9, 0.95, 0.99, 0.999}

In[55]:= infflatlandisopointisoscatter`mutts =
  Union[#[[2]] & /@ infflatlandisopointisoscatter`simulations]

Out[55]= {1, 3}

In[56]:= infflatlandisopointisoscatter`numcollorders =
  infflatlandisopointisoscatter`simulations[[1]][[3]][[2, 15]];
infflatlandisopointisoscatter`maxr =
  infflatlandisopointisoscatter`simulations[[1]][[3]][[2, 7]];
infflatlandisopointisoscatter`dr =
  infflatlandisopointisoscatter`simulations[[1]][[3]][[2, 9]];
infflatlandisopointisoscatter`numr =
  Floor[infflatlandisopointisoscatter`maxr / infflatlandisopointisoscatter`dr];

```

Compare Deterministic and MC

Fluence - Exact solution (I) comparison to MC

```
In[21]:= Clear[alpha, Σt];
Manipulate[
  If[Length[inflatlandisotropicpointscatter`simulations] > 0,
    Module[{data, maxr, dr, pointsφ, plotpointsφ, logplotφ, plotφ, exactlpoints},
      data = SelectFirst[inflatlandisotropicpointscatter`simulations,
        #[[1]] == α && #[[2]] == Σt &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];


      pointsφ = data[[4]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = inflatlandisotropicpointscatter`ppoints[pointsφ, dr, maxr, Σt];

      exactlpoints =
        Quiet[{#[[1]], 2 Pi #[[1]] inflatlandisotropicpointscatter`φexactl[
          #[[1]], Σt, α]}] & /@plotpointsφ;

      plotφ = Quiet[Show[
        ListPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListPlot[exactlpoints, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      logplotφ = Quiet[Show[
        ListLogPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListLogPlot[exactlpoints, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 800],
        PlotLabel -> "Exact solution (I)\nInfinite Flatland, isotropic point
          source, isotropic scattering, fluence φ[r], α = "<>
          ToString[α]<> ", Σt = "<>ToString[Σt]]
      ]
    ,
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ]
, {{α, 0.95}, inflatlandisotropicpointscatter`alphas},
{Σt, inflatlandisotropicpointscatter`mutss}]
```

Out[22]=



Fluence - Exact solution (2) comparison to MC

```

In[23]:= Clear[alpha, Σt];
Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data, maxr, dr, pointsφ, plotpointsφ, logplotφ, plotφ, exact1points},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] == α && #[[2]] == Σt &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];

      pointsφ = data[[4]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = infflatlandisopointisoscatter`ppoints[pointsφ, dr, maxr, Σt];

      exact1points =
        Quiet[{#[[1]], 2 Pi #[[1]] infflatlandisopointisoscatter`φexact2[
          #[[1]], Σt, α]}] & /@ plotpointsφ;

      plotφ = Quiet[Show[
        ListPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      logplotφ = Quiet[Show[
        ListLogPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListLogPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 800],
        PlotLabel -> "Exact solution (2)\nInfinite Flatland, isotropic point
          source, isotropic scattering, fluence φ[r], α = "<>
          ToString[α]<> ", Σt = "<> ToString[Σt]]
      ]
    ],
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ],
  {{α, 0.99}, infflatlandisopointisoscatter`alphas},
  {{Σt, 3}, infflatlandisopointisoscatter`mutss}]

```

Out[24]=

α 0.99 +
v
 Σt 1 3
 \$Aborted

Fluence - Exact solution (3a) comparison to MC

```

In[25]:= Clear[alpha, Σt];
Manipulate[
  If[Length[infflatlandisotropicpointisoscatter`simulations] > 0,
    Module[{data, maxr, dr, pointsφ, plotpointsφ, logplotφ, plotφ, exact1points},
      data = SelectFirst[infflatlandisotropicpointisoscatter`simulations,
        #[[1]] == α && #[[2]] == Σt &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];

      pointsφ = data[[4]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = infflatlandisotropicpointisoscatter`ppoints[pointsφ, dr, maxr, Σt];


      exact1points =
        Quiet[{#[[1]], 2 Pi #[[1]] infflatlandisotropicpointisoscatter`φexact3a[
          #[[1]], Σt, α]}] & /@ plotpointsφ;

      plotφ = Quiet[Show[
        ListPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      logplotφ = Quiet[Show[
        ListLogPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListLogPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 800],
        PlotLabel -> "Exact solution (3a)\nInfinite Flatland, isotropic point
          source, isotropic scattering, fluence φ[r], α = "<>
          ToString[α] <> ", Σt = "<> ToString[Σt]]
      ]
    ],
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ],
  {{α, 0.99}, infflatlandisotropicpointisoscatter`alphas},
  {{Σt, 3}, infflatlandisotropicpointisoscatter`mutss}
]

```


α 0.99 **v** Σt 1 3

Out[26]=



Fluence - Exact solution (3b) comparison to MC

```

In[27]:= Clear[alpha,  $\Sigma_t$ ];
Manipulate[
  If[Length[infflatlandisotropicpointisoscatter`simulations] > 0,
    Module[{data, maxr, dr, points $\phi$ , plotpoints $\phi$ , logplot $\phi$ , plot $\phi$ , exact1points},
      data = SelectFirst[infflatlandisotropicpointisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma_t$  &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];

      points $\phi$  = data[[4]];

      (* divide by  $\Sigma_t$  to convert collision density into fluence *)
      plotpoints $\phi$  = infflatlandisotropicpointisoscatter`ppoints[points $\phi$ , dr, maxr,  $\Sigma_t$ ];

      exact1points =
        Quiet[{#[[1]], 2 Pi #[[1]] infflatlandisotropicpointisoscatter` $\phi$ exact3b[
          #[[1]],  $\Sigma_t$ ,  $\alpha$ , M]}] & /@plotpoints $\phi$ ;

      plot $\phi$  = Quiet[Show[
        ListPlot[plotpoints $\phi$ , PlotRange → All, PlotStyle → PointSize[.01]],
        ListPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r  $\phi$ [r]}, {r,}}
      ]];
      logplot $\phi$  = Quiet[Show[
        ListLogPlot[plotpoints $\phi$ , PlotRange → All, PlotStyle → PointSize[.01]],
        ListLogPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{2 Pi r  $\phi$ [r]}, {r,}}
      ]];
      Show[GraphicsGrid[{{plot $\phi$ , logplot $\phi$ }}, ImageSize → 800],
        PlotLabel -> "Exact solution (3b)\nInfinite Flatland, isotropic point
          source, isotropic scattering, fluence  $\phi$ [r],  $\alpha$  = "<>
          ToString[ $\alpha$ ] <> ",  $\Sigma_t$  = "<> ToString[ $\Sigma_t$ ]
        ]
      ,
      Text[
        "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
          ensure the data path is setup correctly."
      ]
    ]
  , {{ $\alpha$ , 0.7}, infflatlandisotropicpointisoscatter`alphas},
    {{ $\Sigma_t$ , 3}, infflatlandisotropicpointisoscatter`mutss}, {{M, 10}, Range[20]}]


```

α 0.7

Σ_t

M 10

Out[28]=



Fluence - Diffusion approximations (Classical and Grosjean) comparison to MC

```

In[29]:= Clear[alpha, Σt];
Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data, maxr, dr, pointsφ, plotpointsφ, logplotφ, plotφ, exact1points},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] == α && #[[2]] == Σt &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];

      pointsφ = data[[4]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = infflatlandisopointisoscatter`ppoints[pointsφ, dr, maxr, Σt];



      plotφ = Quiet[Show[
        ListPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        Plot[2 Pi r infflatlandisopointisoscatter`φGrosjean[r, Σt, α],
          {r, 0, maxr}, PlotRange → All],
        Plot[2 Pi r infflatlandisopointisoscatter`φDiffusion[r, Σt, α],
          {r, 0, maxr}, PlotRange → All, PlotStyle → Dashed],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      logplotφ = Quiet[Show[
        ListLogPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        LogPlot[2 Pi r infflatlandisopointisoscatter`φGrosjean[r, Σt, α],
          {r, 0, maxr}, PlotRange → All],
        LogPlot[2 Pi r infflatlandisopointisoscatter`φDiffusion[r, Σt, α],
          {r, 0, maxr}, PlotRange → All, PlotStyle → Dashed],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}}
      ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 1000],
        PlotLabel -> "Classical (dashed) and Grosjean Modified (thin)
          Diffusion Approximation\nInfinite Flatland, isotropic
          point source, isotropic scattering, fluence φ[r], α = "<>
          ToString[α]<> ", Σt = "<> ToString[Σt]]
    ]
  ,
  Text[
    "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
      ensure the data path is setup correctly."
  ]
]
, {{α, 0.5}, infflatlandisopointisoscatter`alphas},
{Σt, infflatlandisopointisoscatter`mutts}]

```

α 0.5 Σt

Classical (dashed) and Grosjean Modified (thin) Diffusion
Infinite Flatland, isotropic point source, isotropic scattering, fluen

Out[30]=



Fluence - Diffusion approximation (Rigorous) comparison to MC

```

In[31]:= Clear[alpha, Σt];
Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data, maxr, dr, pointsφ, plotpointsφ, logplotφ, plotφ, exact1points},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] == α && #[[2]] == Σt &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];

      pointsφ = data[[4]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = infflatlandisopointisoscatter`ppoints[pointsφ, dr, maxr, Σt];



      plotφ = Quiet[Show[
        ListPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        Plot[2 Pi r infflatlandisopointisoscatter`φRigorousDiffusion[r, Σt, α],
          {r, 0, maxr}, PlotRange → All],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}},
      ]];
      logplotφ = Quiet[Show[
        ListLogPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        LogPlot[2 Pi r infflatlandisopointisoscatter`φRigorousDiffusion[r, Σt, α],
          {r, 0, maxr}, PlotRange → All],
        Frame → True,
        FrameLabel -> {{2 Pi r φ[r]}, {r,}},
      ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 1000], PlotLabel ->
        "Rigorous Diffusion Approximation\nInfinite Flatland, isotropic
          point source, isotropic scattering, fluence φ[r], α = "<>
          ToString[α] <> ", Σt = "<> ToString[Σt]]
    ]
  ,
  Text[
    "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
      ensure the data path is setup correctly."
  ]
]
, {{α, 0.5}, infflatlandisopointisoscatter`alphas},
{Σt, infflatlandisopointisoscatter`mutts}]

```

α 0.5

Σt

Rigorous Diffusion Approximation
Infinite Flatland, isotropic point source, isotropic scattering, fluence



N-th order fluence / scalar flux

```

In[33]:= Clear[alpha,  $\Sigma t$ ];
Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data, maxr, dr, points $\phi$ ,
      plotpoints $\phi$ , logplot $\phi$ , plot $\phi$ , exactlpoints, numorders},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma t$  &][[3]];
      maxr = data[[2, 7]];
      dr = data[[2, 9]];
      numorders = data[[2, 15]];

      points $\phi$  = data[[9 + numorders + n + 1]];

      (* divide by  $\Sigma t$  to convert collision density into fluence *)
      plotpoints $\phi$  = infflatlandisopointisoscatter`ppoints[points $\phi$ , dr, maxr,  $\Sigma t$ ];

      plot $\phi$  = Quiet[Show[
        ListPlot[plotpoints $\phi$ , PlotRange → All, PlotStyle → PointSize[.01]],
        Plot[2 Pi r infflatlandisopointisoscatter` $\phi$ [r,  $\Sigma t$ ,  $\alpha$ , n],
          {r, 0, maxr}, PlotRange → All],
        Frame → True,
        FrameLabel -> {{2 Pi r  $\phi$ [r | n]}, {r,}},
        ]];
      logplot $\phi$  = Quiet[Show[
        ListLogPlot[plotpoints $\phi$ , PlotRange → All, PlotStyle → PointSize[.01]],
        LogPlot[2 Pi r infflatlandisopointisoscatter` $\phi$ [r,  $\Sigma t$ ,  $\alpha$ , n],
          {r, 0, maxr}, PlotRange → All],
        Frame → True,
        FrameLabel -> {{2 Pi r  $\phi$ [r | n]}, {r,}},
        ]];
      Show[GraphicsGrid[{{plot $\phi$ , logplot $\phi$ }}, ImageSize → 1000],
        PlotLabel -> "Infinite Flatland, isotropic point source, isotropic
          scattering, n-th scattered fluence  $\phi$ [r|n],  $\alpha$  = "<>
          ToString[ $\alpha$ ] <> ",  $\Sigma t$  = "<> ToString[ $\Sigma t$ ]]
      ]
    ],
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ],
  {{ $\alpha$ , 0.9}, infflatlandisopointisoscatter`alphas},
  { $\Sigma t$ , infflatlandisopointisoscatter`mut},
  {{n, 8}, Range[If[NumberQ[infflatlandisopointisoscatter`numcollorders],
    infflatlandisopointisoscatter`numcollorders, 1]]}]

```



α 0.9

Σt

n 8

Infinite Flatland, isotropic point source, isotropic scattering, n-th scatterer

Out[34]=



Compare moments of ϕ

```

In[35]:= Manipulate[
  If[Length[infflatlandisotropicpointisoscatter`simulations] > 0,
    Module[{data, nummoments,  $\phi$ moments, ks, analytic, j},
      data = SelectFirst[infflatlandisotropicpointisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma t$  &][[3]];
      nummoments = data[[2, 17]];
       $\phi$ moments = N[{ $\frac{\text{data}[[6]]}{\Sigma t}$ }]];
      ks = {Table[k, {k, 0, nummoments - 1}]}];
      analytic = Table[infflatlandisotropicpointisoscatter` $\phi m[\Sigma t, \alpha, k]$ , {k, ks}];
      j = Join[ks, analytic,  $\phi$ moments];
      TableForm[
        Join[{"k", "analytic", "MC"}, Transpose[j]]
      ]
    ],
  Text[
    "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
    ensure the data path is setup correctly."
  ]
],
{ $\alpha$ , 0.95}, infflatlandisotropicpointisoscatter`alphas},
{ $\Sigma t$ , 3}, infflatlandisotropicpointisoscatter`mutss}]

```

α
 Σt

k	analytic	MC
0	6.66667	6.66647
1	10.931	10.9305
2	29.6296	29.6046
3	111.733	111.425
4	539.918	536.417

Out[35]=

n-th collided moments of ϕ

```

In[36]:= Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data,  $\phi$ moments, ks, analytic, j, nummoments},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma t$  &][[3]];
      nummoments = data[[2, 17]];
       $\phi$ moments = N[ $\frac{\text{data}[[9 + n]]}{\Sigma t}$ ];
      ks = {Table[k, {k, 0, nummoments - 1}]};
      analytic =
        Table[Quiet[N[infflatlandisopointisoscatter` $\phi m[\Sigma t, \alpha, n, k]$ ]], {k, ks}];
      j = Join[ks, analytic,  $\phi$ moments];
      TableForm[
        Join[{{"k", "analytic", "MC"}}, Transpose[j]]
      ]
    ],
  Text[
    "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
    ensure the data path is setup correctly."
  ]
],
  {{ $\alpha$ , 0.8}, infflatlandisopointisoscatter`alphas},
  {{ $\Sigma t$ , 3}, infflatlandisopointisoscatter`mutss},
  {{n, 11}, Range[If[NumberQ[infflatlandisopointisoscatter`numcollorders],
    infflatlandisopointisoscatter`numcollorders, 1]]}]

```

Out[36]=

α	0.8	V
Σt	1	3
n	11	V

k	analytic	MC
0	0.0286331	0.0286669
1	0.0405845	0.0406517
2	0.076355	0.0765137
3	0.175866	0.176313
4	0.475098	0.476537

Angular distributions


```
In[149]:= Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data, numorders, pointsu, plotpointsu, du, r, dr},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma_t$  &][[3]];
      numorders = data[[2, 15]];
      du = data[[2, 11]];
      dr = data[[2, 9]];

      pointsu = data[[9 + 2 numorders + n]];

      r = dr * n - 0.5 dr;

      (* divide by  $\Sigma_t$  to convert collision density into fluence *)
      plotpointsu = infflatlandisopointisoscatter`ppointsu[pointsu, du,  $\Sigma_t$ ];
      Show[
        ListPlot[plotpointsu, PlotRange -> All,
          Frame -> True,
          FrameLabel -> {{2 Pi r L[r, u]}, {u}},
        Plot[2 Pi r infflatlandisopointisoscatter`Ldiffusion[r, u,  $\Sigma_t$ ,  $\alpha$ ],
          {u, -1, 1}, PlotRange -> All
        ],
        PlotLabel -> "Infinite Flatland, isotropic point source,
          isotropic scattering\nAngular distribution: classical
          diffusion approximation\n $2 \pi r L[r, u]$ , r = "<>
          ToString[r]<>" ,  $\alpha$  = "<>ToString[ $\alpha$ <>" ,  $\Sigma_t$  = "<>ToString[ $\Sigma_t$ 
      ]
    ],
    Text["Uh oh! Couldn't find MC data.
      Try to evaluate this entire notebook and ensure the data path is setup
      correctly."]
  ],
  {{ $\alpha$ , 0.99}, infflatlandisopointisoscatter`alphas},
  {{ $\Sigma_t$ , 1}, infflatlandisopointisoscatter`muts},
  {{n, 100}, Range[If[NumberQ[infflatlandisopointisoscatter`numr],
    infflatlandisopointisoscatter`numr, 1]]}]
```

Out[149]=



```

In[152]:= Manipulate[
  If[Length[infflatlandisopointisoscatter`simulations] > 0,
    Module[{data, numorders, pointsu, plotpointsu, du, r, dr},
      data = SelectFirst[infflatlandisopointisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma_t$  &][[3]];
      numorders = data[[2, 15]];
      du = data[[2, 11]];
      dr = data[[2, 9]];

      pointsu = data[[9 + 2 numorders + n]];

      r = dr * n - 0.5 dr;

      (* divide by  $\Sigma_t$  to convert collision density into fluence *)
      plotpointsu = infflatlandisopointisoscatter`ppointsu[pointsu, du,  $\Sigma_t$ ];
      Show[
        ListPlot[plotpointsu, PlotRange -> All,
          Frame -> True,
          FrameLabel -> {{2 Pi r L[r, u], }, {u, }},
          Plot[2 Pi r infflatlandisopointisoscatter`Lintegral[r, u,  $\Sigma_t$ ,  $\alpha$ ,
            infflatlandisopointisoscatter`phiGrosjean], {u, -1, 1}, PlotRange -> All],
          PlotLabel -> "Infinite Flatland, isotropic point source,
            isotropic scattering\nAngular distribution:
              Fluence integral Grosjean\n 2  $\pi$  r L[r,u], r = "<>
              ToString[r]<> ",  $\alpha$  = "<> ToString[ $\alpha$ <> ",  $\Sigma_t$  = "<> ToString[ $\Sigma_t$ ]
        ]
      ],
    Text["Uh oh! Couldn't find MC data.
      Try to evaluate this entire notebook and ensure the data path is setup
      correctly."]
  ]
, {{ $\alpha$ , 0.9}, infflatlandisopointisoscatter`alphas},
  {{ $\Sigma_t$ , 1}, infflatlandisopointisoscatter`mutss},
  {{n, 53}, Range[If[NumberQ[infflatlandisopointisoscatter`numr],
    infflatlandisopointisoscatter`numr, 1]]}]

```

Out[152]=

