

Infinite 3D medium, Isotropic Plane Source, Isotropic Scattering

Exponential Random Flight

This is code to accompany the book:

A Hitchhiker's Guide to Multiple Scattering

© 2016 Eugene d'Eon


www.eugenedeon.com

Path Setup

Put a file at `~/hitchhikerpath` with the path to your hitchhiker repo so that these worksheets can find the MC data from the C++ simulations for verification

```
In[1]:= SetDirectory[Import["~/hitchhikerpath"]]
```

Notation



α - single-scattering albedo

Σ_t - extinction coefficient

z - scalar position coordinate in medium (distance from plane source at origin)

$u = \cos \theta$ - direction cosine

Analytic solutions

Caseology quantities

$$\text{In[2]:= CaseN0}[c_ , v0_] := \frac{1}{2} c v0^3 \left(\frac{c}{v0^2 - 1} - \frac{1}{v0^2} \right)$$

$$\text{In[3]:= Casev0}[c_?NumericQ] := \text{FindRoot}\left[c v \text{ArcTanh}\left[\frac{1}{v}\right] - 1 == 0, \{v, 1.00000000001, 10^{10}\}, \text{Method} \rightarrow \text{"Brent"}\right][[1]][[2]]$$

$$\text{In[4]:= CaseN}[c_ , v_] := v \left(\text{Case}\lambda[v, c]^2 + \left(\frac{\pi c v}{2} \right)^2 \right)$$

$$\text{In[5]:= Case}\lambda[v_ , c_] := 1 - c v \text{ArcTanh}[v]$$

$$\text{In[6]:= Case}\psi0[u_ , v0_ , c_ , z_] := \frac{c}{2} \frac{v0}{v0 - \text{Sign}[z] u}$$

Rigorous diffusion approximation

$$\text{In[7]:= inf3Disoplaneisoscatter}\phi_{\text{rigorousDiffusion}}[z_ , \Sigma t_ , \alpha_] := \frac{1}{2} \frac{e^{-\text{Abs}[z] \Sigma t / \#}}{\text{CaseN0}[\alpha, \#]} \&[\text{Casev0}[\alpha]]$$

Fluence: exact solution

[Case and Zwiefel 1967]

$$\text{In[8]:= inf3Disoplaneisoscatter}\phi_{\text{exact}}[z_ , \Sigma t_ , \alpha_] := \text{inf3Disoplaneisoscatter}\phi_{\text{rigorousDiffusion}}[z, \Sigma t, \alpha] + \frac{1}{2} \text{NIntegrate}\left[\frac{e^{-\Sigma t \text{Abs}[z] / v}}{\text{CaseN}[\alpha, v]}, \{v, 0, 1\}\right]$$

Nth-scattered fluence

$$\text{In[9]:= inf3Disoplaneisoscatter}\phi_{\text{exact1}}[x_ , \Sigma t_ , \alpha_ , n_] := \frac{(\alpha \Sigma t)^n}{\pi} \text{NIntegrate}\left[\frac{\text{ArcTan}\left[\frac{z}{\Sigma t}\right]^{n+1} \text{Cos}[x z]}{z^{n+1}}, \{z, 0, \text{Infinity}\}, \text{Method} \rightarrow \text{"ExtrapolatingOscillatory"}\right]$$

$$\text{In[10]:= inf3Disoplaneisoscatter}\phi_{\text{exact2}}[x_ , \Sigma t_ , \alpha_ , n_] := \frac{\alpha^n}{\pi} \text{Chop}\left[\text{NIntegrate}\left[\frac{1}{2} 2^{-2-n} e^{-\text{Abs}[x] z \Sigma t} \left(\left(\frac{-\frac{1}{2} \pi + \text{Log}\left[\frac{z+1}{z-1}\right]}{z} \right)^{1+n} - \left(\frac{\frac{1}{2} \pi + \text{Log}\left[\frac{z+1}{z-1}\right]}{z} \right)^{1+n} \right), \{z, 1, \text{Infinity}\}\right]\right]$$

$$\text{In[11]:= inf3Disoplaneisoscatter}\phi_{\text{Gaussian}}[x_ , \Sigma t_ , \alpha_ , n_] := \frac{e^{-\frac{3 x^2 \Sigma t^2}{4 (1+n)}} \sqrt{\frac{3}{\pi}} \alpha^n}{2 \sqrt{1+n}}$$

Classical Diffusion Approximation

$$\text{In[12]:= inf3Disoplaneisoscatter}\phi\text{Diffusion}[x_, \Sigma t_, \alpha_] := \frac{3 e^{-\text{Abs}[x] \sqrt{3-3\alpha}} \Sigma t}{2 \sqrt{3-3\alpha}}$$

Grosjean Modified Diffusion Approximation

$$\text{In[13]:= inf3Disoplaneisoscatter}\phi\text{Grosjean}[x_, \Sigma t_, \alpha_] :=$$

$$-\frac{1}{2} \text{ExpIntegralEi}[-\text{Abs}[x] \Sigma t] + \frac{e^{-\text{Abs}[x] \sqrt{3 + \frac{3}{-2+\alpha}} \Sigma t} \sqrt{3 + \frac{3}{-2+\alpha}} \alpha}{2 (1 - \alpha)}$$

Radiance (Angular Flux)

$$\text{In[14]:= inf3Disoplaneisoscatter}\text{LrigorousDiffusion}[z_, u_, \Sigma t_, \alpha_] :=$$

$$\frac{1}{4 \text{Pi}} \text{Case}\psi 0[u, \#, \alpha, z] \frac{e^{-\text{Abs}[z] \Sigma t / \#}}{\text{CaseN} 0[\alpha, \#]} \&[\text{Case}v 0[\alpha]]$$

$$\text{In[15]:= inf3Disoplaneisoscatter}\text{Lexact}[z_, u_, \Sigma t_, \alpha_] :=$$

$$\text{inf3Disoplaneisoscatter}\text{LrigorousDiffusion}[z, u, \Sigma t, \alpha] +$$

$$\frac{1}{4 \text{Pi}} \left(\text{Case}\lambda [u, \alpha] \frac{e^{-\frac{\text{Abs}[z] \Sigma t}{u}}}{\text{CaseN}[\alpha, u]} \text{HeavisideTheta}[1 - u] \text{HeavisideTheta}[u] \right.$$

$$+ \text{NIntegrate}\left[\frac{e^{-\frac{\text{Abs}[z] \Sigma t}{v}}}{\text{CaseN}[\alpha, v]} \frac{\alpha}{2} \frac{v}{v - u} \right.$$

$$\left. , \{v, 0, u, 1\}, \text{Method} \rightarrow \text{"PrincipalValue"}, \text{PrecisionGoal} \rightarrow 5 \right]$$

load MC data

```

In[16]:= inf3Disoplaneisoscatter`ppoints[zs_, dz_, maxz_, Σt_] :=
  Table[{dz (i) - 0.5 dz - maxz, zs[[i]] / Σt}, {i, 1, Length[zs]}][[2 ;; -2]]

In[17]:= inf3Disoplaneisoscatter`ppointsu[xs_, du_, Σt_] :=
  Table[{-1.0 + du (i) - 0.5 du, xs[[i]] / (2 Σt)}, {i, 1, Length[xs]}][[1 ;; -1]]

In[18]:= inf3Disoplaneisoscatter`fs =
  FileNames["code/3D_medium/infinite3Dmedium/Isotropicplanesource/data/
  inf3D_isotropicplane_isotropicscatter*"];

In[19]:= inf3Disoplaneisoscatter`index[x_] := Module[{data, α, Σt},
  data = Import[x, "Table"];
  Σt = data[[1, 13]];
  α = data[[2, 3]];
  {α, Σt, data}];
inf3Disoplaneisoscatter`simulations =
  inf3Disoplaneisoscatter`index /@ inf3Disoplaneisoscatter`fs;
inf3Disoplaneisoscatter`alphas =
  Union[#[[1]] & /@ inf3Disoplaneisoscatter`simulations]

```

```
Out[21]:= {0.01, 0.1, 0.3, 0.5, 0.7, 0.8, 0.9, 0.95, 0.99, 0.999}
```

```
In[22]:= inf3Disoplaneisoscatter`mutts =  
         Union[#[[2]] & /@ inf3Disoplaneisoscatter`simulations]
```

```
Out[22]= {1, 3}
```

```
In[23]:= inf3Disoplaneisoscatter`numcollorders =  
         inf3Disoplaneisoscatter`simulations[[1]][[3]][[2, 13]];  
inf3Disoplaneisoscatter`maxz =  
         inf3Disoplaneisoscatter`simulations[[1]][[3]][[2, 5]];  
inf3Disoplaneisoscatter`dz =  
         inf3Disoplaneisoscatter`simulations[[1]][[3]][[2, 7]];  
inf3Disoplaneisoscatter`numz =  
         Floor[2 inf3Disoplaneisoscatter`maxz / inf3Disoplaneisoscatter`dz];
```

Compare Deterministic and MC

Fluence - Exact solution comparison to MC

```

In[26]:= Manipulate[
  If[Length[inf3Disoplaneisoscatter`simulations] > 0,
    Module[{data, maxz, dz, pointsφ, plotpointsφ, logplotφ, plotφ, exact1points},
      data = SelectFirst[
        inf3Disoplaneisoscatter`simulations, #[[1]] == α && #[[2]] == Σt &][[3]];
      maxz = data[[2, 5]];
      dz = data[[2, 7]];

      pointsφ = data[[4]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = inf3Disoplaneisoscatter`ppoints[pointsφ, dz, maxz, Σt];

      exact1points =
        Quiet[{#[[1]], inf3Disoplaneisoscatter`φexact[#[[1]], Σt, α]}] & /@
          plotpointsφ;



      numpoints = Length[plotpointsφ];
      skip = Floor[numpoints  $\frac{6}{7} \frac{1}{2}$ ];

      plotφ = Quiet[Show[
        (*ListPlot[exact1points[[skip;;-skip]], PlotRange→{0,6}, Joined→True], *)
        Plot[inf3Disoplaneisoscatter`φexact[z, Σt, α],
          {z, - $\frac{\text{maxz}}{7}$ ,  $\frac{\text{maxz}}{7}$ }, PlotRange → {0, 6}],
        Plot[inf3Disoplaneisoscatter`φrigorousDiffusion[z, Σt, α],
          {z, -maxz, maxz}, PlotRange → All, PlotStyle → {Red, Dashed}],
        ListPlot[plotpointsφ[[skip;;-skip]], PlotRange → All,
          PlotStyle → {Black, PointSize[.01]}],
        Frame → True,
        FrameLabel -> {{φ[z],}, {z,}}
      ]];
      logplotφ = Quiet[Show[
        ListLogPlot[exact1points, PlotRange → All, Joined → True],
        LogPlot[inf3Disoplaneisoscatter`φrigorousDiffusion[z, Σt, α],
          {z, -maxz, maxz}, PlotRange → All, PlotStyle → {Red, Dashed}],
        ListLogPlot[plotpointsφ, PlotRange → All,
          PlotStyle → {Black, PointSize[.01]}],
        Frame → True,
        FrameLabel -> {{φ[z],}, {z,}}
      ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 800], PlotLabel ->
        "Exact solution (thin), Rigorous Diffusion component (dashed)\nInfinite
          3D, isotropic plane source, isotropic scattering, fluence
          φ[z], α = "<>ToString[α]<>" , Σt = "<>ToString[Σt]]
    ]
  ],
  Text[
    "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
      ensure the data path is setup correctly."
  ]
],
  {α, 0.95}, inf3Disoplaneisoscatter`alphas},
  {Σt, 1}, inf3Disoplaneisoscatter`mutss}
]

```

α 0.95 Σ_t

Exact solution (thin), Rigorous Diffusion component (dashed)
Infinite 3D, isotropic plane source, isotropic scattering, fluence $\phi[z]$, $\alpha = 0.95$, $\Sigma_t = 1$



Out[26]=

Fluence - Diffusion approximations (Classical and Grosjean) comparison to MC

```

In[27]:= Manipulate[
  If[Length[inf3Disoplaneisoscatter`simulations] > 0,
    Module[{data, maxz, dz, points $\phi$ ,
      plotpoints $\phi$ , logplot $\phi$ , plot $\phi$ , exact1points, numpoints, skip},
      data = SelectFirst[inf3Disoplaneisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma t$  &][[3]];
      maxz = data[[2, 5]];
      dz = data[[2, 7]];

      points $\phi$  = data[[4]];

      (* divide by  $\Sigma t$  to convert collision density into fluence *)
      plotpoints $\phi$  = inf3Disoplaneisoscatter`ppoints[points $\phi$ , dz, maxz,  $\Sigma t$ ];



      numpoints = Length[plotpoints $\phi$ ];
      skip = Floor[numpoints  $\frac{6}{7} \frac{1}{2}$ ];

      plot $\phi$  = Quiet[Show[
        ListPlot[plotpoints $\phi$ [[skip ;; -skip]],
          PlotRange  $\rightarrow$  All, PlotStyle  $\rightarrow$  {PointSize[.01], Black}],
        Plot[inf3Disoplaneisoscatter` $\phi$ Grosjean[z,  $\Sigma t$ ,  $\alpha$ ],
          {z, -maxz, maxz}, PlotRange  $\rightarrow$  All],
        Plot[inf3Disoplaneisoscatter` $\phi$ Diffusion[z,  $\Sigma t$ ,  $\alpha$ ],
          {z, -maxz, maxz}, PlotRange  $\rightarrow$  All, PlotStyle  $\rightarrow$  {Red, Dashed}],
        ListPlot[plotpoints $\phi$ [[skip ;; -skip]], PlotRange  $\rightarrow$  All,
          PlotStyle  $\rightarrow$  {PointSize[.01], Black}],
        Frame  $\rightarrow$  True,
        FrameLabel  $\rightarrow$  {{ $\phi$ [z],}, {z,}}
      ]];
      logplot $\phi$  = Quiet[Show[
        ListLogPlot[plotpoints $\phi$ ,
          PlotRange  $\rightarrow$  All, PlotStyle  $\rightarrow$  {PointSize[.01], Black}],
        LogPlot[inf3Disoplaneisoscatter` $\phi$ Grosjean[z,  $\Sigma t$ ,  $\alpha$ ],
          {z, -maxz, maxz}, PlotRange  $\rightarrow$  All],
        LogPlot[inf3Disoplaneisoscatter` $\phi$ Diffusion[z,  $\Sigma t$ ,  $\alpha$ ],
          {z, -maxz, maxz}, PlotRange  $\rightarrow$  All, PlotStyle  $\rightarrow$  {Red, Dashed}],
        ListLogPlot[plotpoints $\phi$ , PlotRange  $\rightarrow$  All,
          PlotStyle  $\rightarrow$  {PointSize[.01], Black}],
        Frame  $\rightarrow$  True,
        FrameLabel  $\rightarrow$  {{ $\phi$ [z],}, {z,}}
      ]];
      Show[GraphicsGrid[{{plot $\phi$ , logplot $\phi$ }}, ImageSize  $\rightarrow$  800],
        PlotLabel  $\rightarrow$  "Classical (dashed) and Grosjean Modified (thin)
          Diffusion Approximation\nInfinite 3D, isotropic plane
          source, isotropic scattering, fluence  $\phi$ [z],  $\alpha$  = "<>
          ToString[ $\alpha$ ] <> ",  $\Sigma t$  = "<> ToString[ $\Sigma t$ ]]
      ]
    ],
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ]
, {{ $\alpha$ , 0.5}, inf3Disoplaneisoscatter`alphas},
  {{ $\Sigma t$ , 1}, inf3Disoplaneisoscatter`mutss}]

```


α 0.5 Σ_t

Classical (dashed) and Grosjean Modified (thin) Diffusion Approximation
Infinite 3D, isotropic plane source, isotropic scattering, fluence $\phi[z]$, $\alpha = 0.5$, $\Sigma_t = 1$



Out[27]=

N-th order fluence / scalar flux

N-th collided Fluence - Exact solution (I) comparison to MC

```
In[28]:= Manipulate[
  If[Length[inf3Disoplaneisoscatter`simulations] > 0,
    Module[{data, maxz, dz, points $\phi$ ,
      plotpoints $\phi$ , logplot $\phi$ , plot $\phi$ , exactlpoints, numorders},
      data = SelectFirst[inf3Disoplaneisoscatter`simulations,
        #[[1]] ==  $\alpha$  && #[[2]] ==  $\Sigma_t$  &][[3]];
      maxz = data[[2, 5]];
      dz = data[[2, 7]];

      numorders = data[[2, 13]];

      points $\phi$  = data[[9 + numorders + n + 1]];

      (* divide by  $\Sigma_t$  to convert collision density into fluence *)
      plotpoints $\phi$  = inf3Disoplaneisoscatter`ppoints[points $\phi$ , dz, maxz,  $\Sigma_t$ ];

      exactlpoints =
        Quiet[{#[[1]], inf3Disoplaneisoscatter` $\phi$ exactl[#[[1]],  $\Sigma_t$ ,  $\alpha$ , n]}] & /@
          plotpoints $\phi$ ;

      plot $\phi$  = Quiet[Show[
        ListPlot[plotpoints $\phi$ , PlotRange → All, PlotStyle → PointSize[.01]],
        ListPlot[exactlpoints, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel → {{ $\phi$ ["z|" <> ToString[n]],}, {z,}}
      ]];
      logplot $\phi$  = Quiet[Show[
        ListLogPlot[plotpoints $\phi$ , PlotRange → All, PlotStyle → PointSize[.01]],
        ListLogPlot[exactlpoints, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel → {{ $\phi$ ["z|" <> ToString[n]],}, {z,}}
      ]];
      Show[GraphicsGrid[{{plot $\phi$ , logplot $\phi$ }}, ImageSize → 800],
        PlotLabel → "Exact solution (1)\nInfinite 3D, isotropic plane source,
          isotropic scattering, n-th scattered fluence  $\phi$ [z|n],  $\alpha$  = " <>
          ToString[ $\alpha$ ] <> ",  $\Sigma_t$  = " <> ToString[ $\Sigma_t$ ]]
      ]
    ],
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ],
  {{ $\alpha$ , 0.99}, inf3Disoplaneisoscatter`alphas},
  {{ $\Sigma_t$ , 3}, inf3Disoplaneisoscatter`mutss},
  {{n, 13}, Range[If[NumberQ[inf3Disoplaneisoscatter`numcollorders],
    inf3Disoplaneisoscatter`numcollorders, 1]]}
]
```

Out[28]=

α

0.95

+

v

Σt

1

3

n

13

v

\$Aborted

N-th collided Fluence - Exact solution (2) comparison to MC

```

In[29]:= Manipulate[
  If[Length[inf3Disoplaneisoscatter`simulations] > 0,
    Module[{data, maxz, dz, pointsφ,
      plotpointsφ, logplotφ, plotφ, exact1points, numorders},
      data = SelectFirst[inf3Disoplaneisoscatter`simulations,
        #[[1]] == α && #[[2]] == Σt &][[3]];
      maxz = data[[2, 5]];
      dz = data[[2, 7]];

      numorders = data[[2, 13]];


      pointsφ = data[[9 + numorders + n + 1]];

      (* divide by Σt to convert collision density into fluence *)
      plotpointsφ = inf3Disoplaneisoscatter`ppoints[pointsφ, dz, maxz, Σt];

      exact1points =
        Quiet[#[[1]], inf3Disoplaneisoscatter`φexact2[#[[1]], Σt, α, n]] & /@
          plotpointsφ;

      plotφ = Quiet[Show[
        ListPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{φ["z" <> ToString[n]],}, {z,}},
        ]];
      logplotφ = Quiet[Show[
        ListLogPlot[plotpointsφ, PlotRange → All, PlotStyle → PointSize[.01]],
        ListLogPlot[exact1points, PlotRange → All, Joined → True],
        Frame → True,
        FrameLabel -> {{φ["z" <> ToString[n]],}, {z,}},
        ]];
      Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 800],
        PlotLabel -> "Exact solution (2)\nInfinite 3D, isotropic plane source,
          isotropic scattering, n-th scattered fluence φ[z|n], α = "<>
            ToString[α] <> ", Σt = "<> ToString[Σt]]
      ]
    ],
  ,
  Text[
    "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
      ensure the data path is setup correctly."
  ]
],
  {{α, 0.99}, inf3Disoplaneisoscatter`alphas},
  {{Σt, 3}, inf3Disoplaneisoscatter`mut},
  {{n, 13}, Range[If[NumberQ[inf3Disoplaneisoscatter`numcollorders],
    inf3Disoplaneisoscatter`numcollorders, 1]]}
]

```



N-th collided Fluence - Approximations

```
In[30]:= Manipulate[
  If[Length[inf3Disoplaneisoscatter`simulations] > 0,
    Module[{data, maxz, dz, pointsphi,
      plotpointsphi, logplotphi, plotphi, exactlpoints, numorders},
      data = SelectFirst[inf3Disoplaneisoscatter`simulations,
        #[[1]] == alpha && #[[2]] == Sigma t & [[3]]];
      maxz = data[[2, 5]];
      dz = data[[2, 7]];
      numorders = data[[2, 13]];

      pointsphi = data[[9 + numorders + n + 1]];

      (* divide by Sigma t to convert collision density into fluence *)
      plotpointsphi = inf3Disoplaneisoscatter`ppoints[pointsphi, dz, maxz, Sigma t];

      seriesclassical = alpha^n SeriesCoefficient[
        inf3Disoplaneisoscatter`phiDiffusion[z, Sigma t, C], {C, 0, n}];
      seriesG = alpha^n SeriesCoefficient[inf3Disoplaneisoscatter`phiGrosjean[z, Sigma t, C],
        {C, 0, n}];

      numpoints = Length[plotpointsphi];
      skip = Floor[numpoints * (6/7 - 1/2)];

      plotphi = Quiet[Show[
        Plot[inf3Disoplaneisoscatter`phiGaussian[z, Sigma t, alpha, n],
```

```

{z, - $\frac{\text{maxz}}{7}$ ,  $\frac{\text{maxz}}{7}$ }, PlotRange → All],
Plot[seriesclassical, {z, - $\frac{\text{maxz}}{7}$ ,  $\frac{\text{maxz}}{7}$ }, PlotRange → All,
PlotStyle → {Red, Dashed}],
Plot[seriesG, {z, - $\frac{\text{maxz}}{7}$ ,  $\frac{\text{maxz}}{7}$ }, PlotRange → All,
PlotStyle → {DotDashed, Purple}],
ListPlot[plotpointsφ[[skip ;; -skip]], PlotRange → All,
PlotStyle → {Black, PointSize[.01]}],
Frame → True,
FrameLabel -> {{φ["z" <> ToString[n]],}, {z,}}
];
logplotφ = Quiet[Show[
LogPlot[inf3Disoplaneisoscatter`φGaussian[z, Σt, α, n],
{z, -maxz, maxz}, PlotRange → {10-8, 1}],
LogPlot[seriesclassical, {z, -maxz, maxz},
PlotRange → {10-8, 1}, PlotStyle → {Red, Dashed}],
LogPlot[seriesG, {z, -maxz, maxz}, PlotRange → {10-8, 1},
PlotStyle → {DotDashed, Purple}],
ListLogPlot[plotpointsφ, PlotRange → {10-8, 1},
PlotStyle → {Black, PointSize[.01]}],
Frame → True,
FrameLabel -> {{φ["z" <> ToString[n]],}, {z,}}
];
Show[GraphicsGrid[{{plotφ, logplotφ}}, ImageSize → 1000],
PlotLabel -> "Approximate solutions, Gaussian (thin) Classical
Diffusion (Dashed) Grosjean (Dot-Dashed)\nInfinite
3D medium, isotropic planesource, isotropic
scattering, n-th scattered fluence φ[z|" <>
ToString[n] <> "], α = " <> ToString[α] <> ", Σt = " <> ToString[Σt]]
],
Text[
"Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
ensure the data path is setup correctly."
],
, {{α, 0.9}, inf3Disoplaneisoscatter`alphas},
{Σt, inf3Disoplaneisoscatter`mut},
{{n, 3}, Range[If[NumberQ[inf3Disoplaneisoscatter`numcollorders],
inf3Disoplaneisoscatter`numcollorders, 1]]}]

```


α 0.9

Σt

n 3

Approximate solutions, Gaussian (thin) Classical Diffusion (Dashed)
Infinite 3D medium, isotropic planesource, isotropic scattering, n-th scatter

Out[30]=



Angular Distributions

```

In[31]:= Manipulate[
  If[Length[inf3Disoplaneisoscatter`simulations] > 0,
    Module[{data, numorders, pointsu, plotpointsu, du, r, dz, maxz, zsim},
      data = SelectFirst[
        inf3Disoplaneisoscatter`simulations, #[[1]] ==  $\alpha$  & &#[[2]] ==  $\Sigma_t$  &][[3]];
      numorders = data[[2, 13]];
      du = data[[2, 9]];
      dz = data[[2, 7]];
      maxz = data[[2, 5]];

      pointsu = data[[9 + 2 numorders + n]];

      zsim = dz * n - 0.5 dz - maxz;

      plotpointsu = inf3Disoplaneisoscatter`ppointsu[pointsu, du,  $\Sigma_t$ ];
      pp = Show[
        ListPlot[plotpointsu, PlotRange → All, PlotStyle → Black,
          Frame → True,
          FrameLabel -> {{Pi L[zsim, u]}, {u}},
        Plot[Pi inf3Disoplaneisoscatter`LrigorousDiffusion[zsim, u,  $\Sigma_t$ ,  $\alpha$ ],
          {u, -1, 1}, PlotStyle → {Red, Dashed}
        ],
        Plot[Pi inf3Disoplaneisoscatter`Lexact[zsim, u,  $\Sigma_t$ ,  $\alpha$ ], {u, -1, 1}
        ],
        PlotLabel -> "Infinite 3D Medium, isotropic plane
          source, isotropic scattering\nAngular distribution:
          rigorous diffusion approximation\n $\pi$  L[z,u], z = "<>
          ToString[zsim]<> ",  $\alpha$  = "<> ToString[ $\alpha$ ]<> ",  $\Sigma_t$  = "<> ToString[ $\Sigma_t$ ]
        ]
      ],
    Text[
      "Uh oh! Couldn't find MC data. Try to evaluate this entire notebook and
        ensure the data path is setup correctly."
    ]
  ],
  {{ $\alpha$ , 0.9}, inf3Disoplaneisoscatter`alphas},
  {{ $\Sigma_t$ , 1}, inf3Disoplaneisoscatter`mutss}, {{n, 127}, Range[If[
    NumberQ[inf3Disoplaneisoscatter`numz], inf3Disoplaneisoscatter`numz, 1]]}]

```


Out[31]=

