

**TABLE 3.** Summary of relevant quantities by river stratum (area) for drift nets. Numbers in parentheses are 95% confidence intervals.

Stratum	Interviews	Effort Est.	Estimated Harvest			
			Chinook	Chum	Sockeye	Total
<b>Tuntutuliak ↔ Johnson R.</b>	46	150	2,000 (1,360 – 2,790)	510 (230 – 900)	720 (490 – 970)	<b>3,230</b> (2,290 – 4,350)
<b>Johnson R. ↔ Napaskiak</b>	69	104	1,330 (1,060 – 1,640)	200 (150 – 270)	480 (370 – 580)	<b>2,010</b> (1,650 – 2,380)
<b>Napaskiak ↔ Akiachak</b>	111	212	2,040 (1,650 – 2,480)	180 (60 – 350)	860 (670 – 1,070)	<b>3,070</b> (2,500 – 3,650)
<b>Akiachak ↔ Akiak</b>	0	45	430 (340 – 520)	40 (20 – 70)	180 (140 – 220)	<b>650</b> (530 – 770)
<b>All</b>	<b>226</b>	<b>511</b>	<b>5,810</b> (4,970 – 6,740)	<b>920</b> (610 – 1,330)	<b>2,230</b> (1,910 – 2,570)	<b>8,950</b> (7,830 – 10,230)

**TABLE 4.** Average (95% confidence limits) total salmon catch per trip and percent Chinook salmon, summarized for the areas above and below the confluence of the Johnson River with the Kuskokwim River. Quantities are derived from the strata- and species-specific harvest estimates, not the raw interview data.

Quantity	Proximity to Johnson R. Mouth	
	Downstream	Upstream
Total Catch/Trip	22 (15 – 29)	16 (14 – 18)
% Chinook Salmon	62% (55% – 68%)	66% (63% – 69%)

**FIGURE 1.** Distributions of relevant quantities from all completed trips using drift nets. The mean quantity by primary data source is shown in the top right; BBH = Bethel Boat Harbor (ONC), CBM = Other Villages (BSFA/KRITFC), FC = Bethel Area Fish Camps (ONC).

