

Example Large Multi-page Table

Produced by: Ben Staton

This example is adapted from Table 5 in Staton (2018) – a report documenting in-season harvest and effort estimates and associated methodology. That report was prepared with ‘bookdown’ using a custom template.

TABLE 1. Salmon harvest from drift nets from all five openers by species and geographic stratum¹. Numbers within parentheses are 95% confidence limits^{a,b}.

Species	Geographic Stratum ¹				Total
	A	B	C	D	
6/12/2018					
<i>Chinook</i>	1,910 (1,370-2,490)	540 (340-770)	1,600 (1,190-2,050)	540 (310-810)	4,590 (3,820-5,440)
<i>Chum</i>	1,250 (760-1,890)	140 (90-210)	220 (130-320)	50 (20-100)	1,670 (1,150-2,330)
<i>Sockeye</i>	90 (40-150)	60 (10-130)	90 (40-140)	20 (0-40)	250 (160-360)
<i>Total</i>	3,250 (2,290-4,310)	740 (500-1,030)	1,910 (1,440-2,400)	610 (360-900)	6,500 (5,370-7,730)
6/16/2018					
<i>Chinook</i>	1,570 (970-2,370)	1,220 (930-1,530)	1,890 (1,480-2,360)	650 (340-1,000)	5,330 (4,500-6,330)
<i>Chum</i>	860 (480-1,380)	490 (340-650)	980 (750-1,220)	340 (160-520)	2,670 (2,160-3,250)
<i>Sockeye</i>	100 (20-220)	90 (50-140)	240 (130-390)	10 (0-40)	430 (290-610)
<i>Total</i>	2,520 (1,600-3,770)	1,790 (1,370-2,200)	3,110 (2,520-3,770)	1,000 (760-1,280)	8,430 (7,250-9,810)
6/24/2018					
<i>Chinook</i>	1,450 (1,080-1,860)	1,040 (710-1,420)	2,500 (2,050-3,030)	890 (540-1,330)	5,890 (5,110-6,720)
<i>Chum</i>	2,920 (2,120-3,710)	1,120 (780-1,470)	3,320 (2,760-3,880)	1,480 (910-2,170)	8,820 (7,560-10,030)
<i>Sockeye</i>	1,650 (1,230-2,130)	450 (290-660)	1,130 (890-1,380)	230 (140-330)	3,460 (2,940-4,030)
<i>Total</i>	6,020 (4,740-7,310)	2,610 (1,820-3,430)	6,950 (5,970-7,970)	2,600 (1,820-3,560)	18,180 (16,230-20,200)
6/29/2018					
<i>Chinook</i>	520 (310-750)	710 (510-970)	1,130 (750-1,650)	600 (390-830)	2,970 (2,410-3,670)
<i>Chum</i>	5,790 (3,910-8,040)	5,110 (3,330-7,520)	5,530 (3,860-7,960)	1,210 (790-1,700)	17,640 (14,400-21,380)
<i>Sockeye</i>	2,440 (1,260-3,840)	1,680 (1,180-2,300)	2,550 (1,860-3,410)	530 (370-720)	7,200 (5,690-8,880)
<i>Total</i>	8,750 (6,190-11,740)	7,510 (5,400-10,190)	9,210 (6,890-12,410)	2,340 (1,930-2,840)	27,810 (23,560-32,390)

TABLE 1. Salmon harvest from drift nets from all five openers by species and geographic stratum¹. Numbers within parentheses are 95% confidence limits^{a,b}. (*continued*)

Species	A	B	C	D	Total
7/5/2018					
<i>Chinook</i>	230 (160-320)	290 (190-400)	340 (220-510)	100 (50-170)	970 (780-1,190)
<i>Chum</i>	3,660 (2,670-4,760)	4,180 (2,790-5,640)	3,150 (2,080-4,310)	1,040 (700-1,400)	12,030 (9,960-14,310)
<i>Sockeye</i>	2,540 (1,660-3,470)	3,060 (1,960-4,250)	4,490 (3,190-5,990)	460 (290-640)	10,550 (8,570-12,630)
<i>Total</i>	6,430 (4,660-8,400)	7,530 (5,140-10,080)	7,980 (6,230-10,080)	1,610 (1,270-2,000)	23,560 (20,090-27,490)
All Openers					
<i>Chinook</i>	5,680 (3,890-7,790)	3,800 (2,680-5,090)	7,460 (5,690-9,600)	2,780 (1,630-4,140)	19,750 (16,620-23,350)
<i>Chum</i>	14,480 (9,940-19,780)	11,040 (7,330-15,490)	13,200 (9,580-17,690)	4,120 (2,580-5,890)	42,830 (35,230-51,300)
<i>Sockeye</i>	6,820 (4,210-9,810)	5,340 (3,490-7,480)	8,500 (6,110-11,310)	1,250 (800-1,770)	21,890 (17,650-26,510)
<i>Total</i>	26,970 (19,480-35,530)	20,180 (14,230-26,930)	29,160 (23,050-36,630)	8,160 (6,140-10,580)	84,480 (72,500-97,620)

Note

All estimates rounded to the nearest 10 fish.

Totals were rounded to the nearest 10 fish after calculating the sum of non-rounded estimates. As a result, the total presented in this table may differ by approximately 10 – 20 fish from the total obtained by summing the rounded estimates.

Geographic Strata

¹ A = Below Johnson River, B = Johnson River to Napaskiak, C = Napaskiak to Akiachak, D = Akiachak to Akiak

Confidence Limits

^a Total means and 95% confidence limits within an opener were obtained via non-parametric bootstrap.

^b Total means and 95% confidence limits across openers are the sum of opener-specific estimates.