

Supplemental material for XYZ

1 Snow covered area and streamflow anomalies

To investigate the hydrologic ramifications of snow ablation, we obtained streamflow data from the USGS Sagehen creek gage which has long term records. The monthly average streamflow at Sagehen creek site peaked in March, two months earlier than normal (Fig. S1) and reached flows well above typical for March.




Figure S1: The monthly averaged streamflow anomalies in cubic-feet-per second (cfs) between 1954 to present for the USGS Sagehen Creek gage site. The historical mean, historical median, historical 1-standard deviation (shaded region), and WY2026 streamflow are depicted.

The synoptic pattern of snow covered area change also reveals the extent of the anomalous March. The SPIReS snow covered area (SCA) product shows the lowest snow covered area on record (Fig. S2). SCA was already very low leading up to March, but reached even lower levels by March 31. High elevation regions above 2250 m had nearly 100% SCA prior to the March event but those regions also declined substantially.




Figure S2: SPIReS SCA across the study region during the March heatwave. Watershed boundaries shown in black and SNOTEL monitoring sites indicated by magenta diamonds. **(a, b, c)** SCA (%) on March 2 and March 29, 2026, respectively, and the difference between the two dates. **(d)** Daily mean SCA (%) for each individual watershed and aggregated across all four basins. The 2000–2020 historical mean and ± 1 standard deviation are depicted. **(e)** Mean SCA as a function of elevation band (250 m bins) for March 2 (blue) and March 29 (red), shown alongside historical values for the same dates (dashed lines) with ± 1 standard deviation shading.

2 SUMMA details

This section describes assumptions implicit in SUMMA as well as provides additional details about the model forcing data preparation.

2.1 Model flux equations

The bulk expression for the H_s implemented in SUMMA is given by:

$$H_s = c_p \rho C_s u (T_{air} - T_{snow}), \quad (1)$$

where c_p is the specific heat capacity of dry air (constant), u is the horizontal wind speed, ρ is the dry air density, and C_s is the bulk transport coefficient. In this work, we use the SUMMA formulation for C_s , which models the stability correction as a function of the bulk Richardson number. We use the “Loisinv” stability correction function implemented in the SUMMA model and assume that $z_h = z_0$

$$C_h = \frac{k^2}{\ln(z/z_0)^2} \cdot f(\text{Ri}) \quad (2)$$

Where Ri is the bulk Richardson number, defined as

$$\text{Ri} = \frac{9.81(T_{air} - T_{snow})}{0.5(T_{air} + T_{snow})u^2}. \quad (3)$$

and $f(\text{Ri})$ is a function with range (0,1] that declines as a function of increasing Ri. An equivalent formulation for latent heat flux follows Eq. 1. The stability correction is reproduced and used in Fig. 12 of the main text. The $f(\text{Ri})$ equation used in that figure is given by $f(\text{Ri}) = (1 - 5 * \text{Ri})^2$.

2.2 Forcing data

We use bias corrected ERA5 data as forcing for SUMMA with several key steps implemented. ERA5 data was obtained from the Copernicus data store for the grid cell nearest CSSL. $SW \downarrow$ was compared between ERA and CSSL CNR4 radiometers for water year 2026. We developed a correction factor to account for potential shadowing by canopy adjacent to the site, using solar zenith angle (SZA) and solar azimuth angle (AZM) as predictors. These were computed from the python py-ephem package. The correction grid is depicted in Fig. S3. It shows the multiplicative correction between 0 and 1 for each SZA and AZM angle.




Figure S3: $SW \downarrow$ multiplicative correction factors for the ERA5 radiation data. The top plot shows the correction grid. The bottom plot shows the observed, uncorrected, and corrected data.

2.3 Additional SUMMA model outputs and model validation

Additional SUMMA outputs with relevance to interpreting the main text are displayed in this section, as well as additional model validation plots.

The snowpack accumulated substantial liquid water following the February ROS event, and snow-temperatures were nearly isothermal (Fig. S4). Cold-content generation began during March for a thin layer at the top of the snowpack, where liquid also refroze. A dense ice-layer formed during mid-February and remained buried under the late February snow and rain.

2.3.1 Canopy site

We validate the SUMMA model in several ways. First, we evaluated the canopy radiation parameterization against the CSSL sub-canopy Kipp and Zonene SPLite 2 radiometer located under the canopy. We find good correlation for March of 2026 for the UEB 2-stream parameterization and an LAI value of 1.5 (Fig S6), using the aforementioned $SW \downarrow$ forcing as the top boundary condition to the model.




Figure S4: A timeseries of modeled vertical cross-sections of snow temperature, snow liquid water content volumetric fraction, snow ice content volumetric fraction, and snow density (bottom) during the snow season of WY2026 for the open site at CSSL.




Figure S5: A comparison of sub-canopy downwelling radiation for different LAI values using the UEB-2stream shortwave radiation parameterization. The top plot depicts a hourly $SW_{\downarrow}$ two day period in early March. The bottom plot depicts hourly $SW_{\downarrow}$ during March. Observed radiation by the Kipp and Zonen radiometer is depicted in the blue solid line

2.3.2 Fluxes

CSSL hosts an experimental eddy-covariance instrument on a movable arm above the snow-pack (see Fig. 1 in the main text). Eddy covariance fluxes were processed using the Licor EddyPro v7.0.9 software with despiking, double coordinate rotation, and flux quality assignment. A qualitative comparison between the SUMMA and eddy-covariance fluxes of sensible and latent heat show reasonable agreement, though there were many low-quality flux measurements during the month of March. Key periods of sublimation during the first and latter portions of the the month show reasonable agreement.




Figure S6: A comparison of SUMMA and eddy-covariance (ECOR) measured fluxes at CSSL for sensible heat (H) and latent heat (LE). The top shows a timeseries for the month of March 2026. The bottom two plots show 1:1 plots between the ECOR and SUMMA estimated fluxes.

2.3.3 Albedo

SUMMA modelled albedo is compared against available observations from the CSSL CNR4 radiometers, as well as SPIReS remote sensing based albedo. Specifically we compare the




Figure S7: (a) SUMMA albedo evaluated against CSSL and SPIRES observations from 1 Jan 2026 to 1 April 2026. (b) Example maps of SPIReS snow albedo during water year 2026. CSSL is highlighted with the purple box. 500m elevation contours are depicted for reference.

3 Initial condition experiments

As we have shown, anomolous conditions also preceded the March 2026 ablation event. To interrogate the role of antecedent conditions, we performed initial condition experiments by altering the amount of ice, liquid water, and cold-content.

The first test (Test 1) evaluates the effects of initial liquid water content (Fig. S8.) It was constructed as follows. The first is the natural starting condition (liq=1). Then, we halved the amount of liquid water (liq=0.5) and increased the amount ice to keep the total SWE (the sum of liquid and ice water equivalent) approximately constant. The third experiment completely zeros out liquid water (liq=0) and likewise increases the amount of ice to keep SWE constant. For internal consistency in SUMMA, the layer temperature is also reduced in the zero-liquid water case (to prevent refreezing) which yields the higher starting CC for that case (Fig. S8a). The outcomes show that there is minimal sensitivity to the timing in the rate of SWE change; the all-ice case completely ablated about one-day later than the standard case.




Figure S8: Initial condition Test 1: Effects of liquid water content. (a) shows the hourly CC of the snowpack throughout the model run. (b) Depicts the total SWE, liquid water content, and ice-water equivalent integrated throughout the depth of the pack in units of water-equivalent (mm).

The second test (Test 2) evaluates the effects of initial snow mass (Fig. S9) on ΔSWE before and after the March heatwave. It is composed of four model runs and uses a dry starting snow case for all experiments. The initial dry-SWE amount is multiplied by 1,

1.5, 2, and 3. The snow height is also adjusted maintain an approximately uniform bulk density between the experiments (Fig. S9a). The surface temperature was assigned the same starting value and decays exponentially to 0 at the bottom of the snow layer. Fig. S9b may be interpreted as follows. First, it is important to recognize that ΔSWE is equivalent to ΔIce plus ΔLiquid integrated throughout the depth of the snowpack content over time (middle columns of Fig. S9b). These terms are also equivalent with the cumulative fluxes at the snowpack top/bottom boundaries, namely precipitation, sublimation/frost, and liquid water outflow through the bottom of the snowpack (rightmost columns of Fig. S9b).

The results of this experiment show that nearly the same amount of ice melts across all scenarios (ΔIce), save for the SWE1x case which is limited by the amount of ice available to ablate. However, ΔSWE for the SWEx3 experiment is -449 mm compared to a ΔSWE of -498 for the SWEx1.5 experiment. This discrepancy is due to the amount of liquid water retained in the snowpack (+70 vs +12 for the SWEx3 vs SWEx1.5 respectively). In this way, we can see that there is not always a 1:1 relationship between the snow energy balance and δSWE ; the residual holding capacity of snow liquid means that two snowpacks can undergo a nearly identical energy forcing, but the deeper snowpack may retain more SWE in the form of liquid water.




Figure S9: Initial condition Test 2: effects of dry snow amount. (a) Depicts the initial and final SWE and snow depth (starting on March 1 and ending on March 31) for the four experiments. (b) Depicts mass components of the change in SWE for the different experiments. Text lists ΔSWE , ΔIce , ΔLiq , and total Outflow, respectively.