# The Dark Sector, Wave Equations, and the Torsion Origin of Classical Fields

## Consequences of the Two-Twistor Framework: Part II

March 2026

---


## Abstract

Building on the two-twistor mass emergence framework (Part I), we derive five classes of results from the same axiom $F_2$. Both cones originate perpendicular to each other in the Fano plane, rotating in opposite directions. At 45-degree half-angle they contact and fragment; the fragmentation asymmetry tilts them to 30 degrees, where the photon mode becomes available. The tilt splits the time vector $t_2$, creating Cone B with its own φ′-circle and a complete dark sector with effective quantum scale $N' = N/2$ and universal fermion mass scaling $\sqrt{2}$. The dark electron mass $\sqrt{(2m_0^2/N)} = 105.67$ MeV coincides with the observed muon mass, identifying the second generation as the φ′-circle shadow of the first. The bottom quark mass follows from Fano geometry: $m_b = (7/3)m_\tau = 4148$ MeV (observed: 4180 MeV). The four Maxwell equations emerge from the contorsion tensor K on flat 5D spacetime: the homogeneous pair from the Bianchi identity $R(\Omega) = 0$, the inhomogeneous pair from the TEGR field equation, and the magnetic monopole absence from the circle topology of φ. The Einstein field equation $G_{\mu\nu} = 8\pi G T_{\mu\nu}$ is shown to be the torsion field equation rewritten in curvature variables, via the identity $R(\omega) = -D(K) - K \wedge K$, exact in the late-time limit. The 5D massless wave equation $\Box_5\Psi = 0$ separates on the φ-circle into Bessel modes, reducing to the massive Klein-Gordon equation per mode and to the Schrödinger equation in the non-relativistic limit. The quantum mechanical wave function is identified with the Fourier decomposition $e^{in\varphi}$ on $S^1$. Dark black holes have Chandrasekhar limit $0.7\ M_\odot$, form earlier than visible black holes, and provide natural seeds for supermassive black hole formation at high redshift. Eight gravitons reside on the shared boundary (graviton wall) at $\sigma = \Delta$ between sectors, yielding one universal gravitational interaction from the asymmetric torsion stress $\sin^2(30°) \neq \cos^2(30°)$. All results follow from the derivation chain in Part I with zero additional parameters.


---

## 1. Introduction

Part I established the framework: from $F_2$, through the Fano plane, two cones, and the twist at 45-degree contact, a constrained 5D metric emerges with mass formula $m_{app} = m_0\sqrt{(1 - n^2\varepsilon)}$. Seventeen predictions matched observation above 98.7%.

At 45-degree contact, fragmentation begins; the asymmetry energy tilts the cones to 30 degrees, splitting the time vector $t_2$ and creating $t_2'$ for Cone B. Part I focused on Cone A (the visible sector). This paper derives the consequences of Cone B (the dark sector), shows that classical field equations are components of the torsion tensor, and identifies the quantum mechanical wave function with Fourier modes on the φ-circle.

The results fall into five categories:

1. **Dark sector spectrum** — complete particle table from Cone B
2. **Three generations** — origin of the generation structure
3. **Maxwell equations** — all four from torsion on flat spacetime

4. **General relativity** — Einstein equation as the late-time limit of torsion
5. **Wave equation** — Schrödinger from $\Box_5\Psi = 0$

No new axioms, parameters, or assumptions are introduced.

---

# 2. The Dark Sector

## 2.1 Cone B Geometry

Both cones originate in the Fano plane, lying perpendicular to each other and rotating in opposite directions within the plane. As the Fano lines wrap around the cone axes, the half-angle increases. When the half-angle reaches 45 degrees, the two perpendicular cones contact and fragmentation begins. The energy of the fragmentation asymmetry tilts them to 30 degrees. At this angle, the photon mode becomes available: n = N with full precessional sweep on the φ-circle.

At the tilt, the time vector $t_2$ splits:

- $t_2$ remains with Cone A (visible sector), defining the φ-circle
- $t_2' = t_2 \sin(30°) = t_2/2$ departs at 30° with Cone B, defining the φ′-circle

The dark sector time-product is:

$\sigma' = t_1 \cdot t_2' = t_1 \cdot t_2/2 = \sigma/2$

The dark φ′-circle has the same topology as φ (both $S^1$) but different radius. The epoch parameter transforms as:

$\varepsilon' = \hbar^2 A^4/(4E^2\sigma'^2) = 4\varepsilon$

The effective quantum number scale:

$N' = N/2$

where $N \cdot \Delta = 254$ forces $N = 1.0858 \times 10^7$ (Part I). Both sectors share the same bare mass $m_0 = 246.22$ GeV (same cone energy) and the same fine structure constant $\alpha = 1/137$ (algebraic, from the prime skeleton $163 - 2 \times 13$, independent of N).

## 2.2 Universal √2 Mass Scaling

Since mass scales as $m \sim 1/\sqrt{\sigma}$ in the framework (Part I, §6.5), the ratio of dark to visible fermion masses is:

$m_{dark}/m_{visible} = \sqrt{(\sigma/\sigma')} = \sqrt{2}$

This is exact for all fermions whose mass formula depends on σ.

## 2.3 Dark Lepton Masses

Applying the lepton mass formulae (Part I, §7.3) with N → N/2:

| Lepton | Visible | Dark (N/2) | Ratio |
|---|---|---|---|
| e-type | 0.514 MeV | 0.727 MeV | √2 |

| Lepton | Visible | Dark (N/2) | Ratio |
|---|---|---|---|
| μ-type | 105.66 MeV | 149.39 MeV | √2 |
| τ-type | 1774 MeV | 2509 MeV | √2 |

## 2.4 Dark Quark Masses

| Quark | Visible | Dark (N/2) | Ratio |
|---|---|---|---|
| u-type | 2.22 MeV | 3.14 MeV | √2 |
| d-type | 4.71 MeV | 6.67 MeV | √2 |
| s-type | 95 MeV | 134.4 MeV | √2 |
| c-type | 1273 MeV | 1800 MeV | √2 |
| b-type | 4148 MeV | 5866 MeV | √2 |
| t-type | 174.1 GeV | 174.1 GeV | 1 |

The top quark is an exception: $m_t = m_0/\sqrt{2}$ comes from the cone projection angle (45°), which is the same for both cones.

## 2.5 Dark Bosons

The boson masses derive from the energy budget of each cone (Part I, §7.2). Since both cones carry 250 GeV with the same internal structure:

$m_{Z'} = m_Z$, $m_{W'} = m_W$, $m_{H'} = m_H$

The dark photon ($n = N/2$, fully precessing on φ') is massless.

## 2.6 The Dark Photon and Kinetic Mixing

The dark photon γ' is the massless gauge boson of the φ'-circle, in exact analogy to the visible photon on the φ-circle.

Kinetic mixing between γ and γ' arises from the shared graviton wall. The mixing parameter equals the electron gap:

$\varepsilon_{mix} = \Delta = 4\alpha^2/9 = 2.37 \times 10^{-5}$

This is an $\alpha^2$ effect: two electromagnetic vertices are required to cross the graviton wall (one on each side). The factor 4/9 comes from the dimensional ratio (Part I, §2).

The oscillation probability:

$P(\gamma \to \gamma') = \varepsilon^2/4 \approx 1.4 \times 10^{-10}$

## 2.7 Dark Matter Properties

| Property | Value | Origin |
|---|---|---|
| Fraction of total matter | cos(30°) = 86.6% | Time overlap integral |
| Self-interaction | Yes (dark α = 1/137) | Same algebraic origin |
| Dark hydrogen binding | 19.3 eV | Scaled from 13.6 eV by √2 |
| Dark Bohr radius | 0.372 Å | Scaled from 0.529 Å by 1/√2 |
| Cooling rate relative to visible | 4.1× faster | $\propto m_e^4$, dark m_e = √2 × visible |
| Dark-visible cross section | $\varepsilon^2 \approx 5.6 \times 10^{-10}$ | Kinetic mixing squared |

The dark sector is cold (m_dark_e = 0.727 MeV ≫ T_CMB), self-interacting (dark atoms and dark chemistry exist), and gravitationally coupled to the visible sector through the shared graviton wall.

---

# 3. Three Generations

## 3.1 The Problem

Why three generations of fermions? The Standard Model provides no explanation for the replication of (e, u, d) → (μ, c, s) → (τ, t, b).

## 3.2 The Solution: Two Circles

The framework contains two φ-circles: φ (visible, Cone A) and φ' (dark, Cone B). Three classes of states arise:

**Generation 1** — φ-circle eigenstates: e, u, d Quantum numbers near n ≈ N. Maximal precession on φ. Lightest.

**Generation 2** — φ'-circle shadows: μ, c, s The φ'-circle at N/2 produces states that project into the visible sector. The dark electron (n' = N/2 − 1/2 on φ') has mass:

$m'_e = \sqrt{(2m_0^2/N)} = 105.67$ MeV

This is the muon mass to 100.0% accuracy.

The muon is not a "heavy electron." It is the lightest dark lepton, observed in the visible sector through the φ-φ' overlap.

**Generation 3** — φ × φ' interference: τ, t, b Cross-coupling between the two circles. The tau mass involves Δn = 282, a large step combining both circle structures. The top quark projects from both cones equally ($m_t = m_0/\sqrt{2}$). The bottom quark mass encodes the Fano geometry directly.

## 3.3 The Bottom Quark from Fano Geometry

$m_b = (7/3) \times m_\tau$

where:

- 7 = number of points on the Fano plane PG(2,2)
- 3 = number of points per line = number of quark colors

Computed: 4148 MeV. Observed: 4180 MeV. Match: 99.2%.

This formula connects the heaviest quark–lepton pair in the third generation directly to the combinatorial structure of the seed geometry.

### 3.4 Why Exactly Three

Two circles produce exactly three classes: eigenstates of one, eigenstates of the other, and interference. A third circle would require a third cone, but only two cones fit the 90-degree quaternionic phase structure (Part I, §5.1). Three generations is a geometric consequence, not a free parameter.

---

## 4. The Graviton Wall

### 4.1 Shared Boundary

The graviton wall consists of 8 graviton states at $\sigma = \Delta$, the minimum of the effective potential. It sits on the boundary between the visible and dark sectors:

Visible ($\sigma > \Delta$) ← graviton wall ($\sigma = \Delta$) → Dark ($\sigma' > \Delta$)

The wall is not in either sector. It is the shared interface.

### 4.2 Asymmetric Torsion and Gravity

The visible sector contributes torsion stress proportional to $\cos^2(30°) = 3/4$, and the dark sector contributes $\sin^2(30°) = 1/4$. The ratio of matter content follows: 86.6% dark, 13.4% visible.

The graviton wall experiences unequal pressure from the two sides. This asymmetric stress is the contorsion $K^0{}_1$. If both sectors contributed equally (at 45°), $\sin^2 = \cos^2$ and the net torsion would vanish — no gravity.

Gravity exists because $\sin(30°) \neq \cos(30°)$.

### 4.3 One Gravity

Since there is one wall, there is one gravitational interaction. Both sectors contribute to and respond to the same torsion. This is why gravity is universal: visible matter, dark matter, and antimatter all gravitate identically.

The mass formula m_app depends on $n^2$ (not n): both +n (particle) and −n (antiparticle) yield the same gravitational mass. This is confirmed by the ALPHA experiment at CERN (2023), which measured normal gravitational acceleration for antihydrogen.

---

## 5. Dark Black Holes

### 5.1 Chandrasekhar Limit

The Chandrasekhar mass scales as $M_{Ch} \propto 1/m_n^2$, where $m_n$ is the constituent fermion mass. Since dark fermions are $\sqrt{2}$ heavier:

$M_{Ch}(\text{dark}) = M_{Ch}(\text{visible})/2 = 0.7\ M_\odot$

Similarly, the Tolman–Oppenheimer–Volkoff limit:

M_TOV(dark) ≈ 1.1 M⊙

Dark neutron stars occupy the range 0.7 – 1.1 M⊙. Dark black holes form from progenitors above ~1 M⊙.

## 5.2 Black Holes as Sector Bridges

At the core of any black hole, σ approaches Δ — the graviton wall. From the visible side, this is the event horizon interior. From the dark side, the same object appears as a dark black hole.

Every black hole is a two-sided object. One face toward each sector. The graviton wall at its center connects both. There is no singularity: the maximum torsion is finite (T_max = N/254 for visible, N/508 for dark).

## 5.3 Supermassive Black Hole Seeds

Supermassive black holes ($10^9$ M⊙) exist at redshift z > 7, within 800 Myr of the Big Bang. Visible-sector accretion is too slow to explain this.

The dark sector resolves the problem:

1. Dark cooling is 4.1× faster (rate ∝ $m_e^4$)
2. Dark stars form first
3. Dark Chandrasekhar limit (0.7 M⊙) is half the visible value
4. Dark matter provides 6.5× more fuel
5. Combined growth advantage: ~53× faster than visible-only models

Dark black holes form first and grow rapidly, creating the gravitational wells around which visible galaxies later assemble.

## 5.4 LIGO Predictions

| Observable | Dark sector value | Visible sector value |
|---|---|---|
| NS mass range | 0.7 – 1.1 M⊙ | 1.4 – 2.2 M⊙ |
| BH minimum mass | ~2.1 M⊙ | ~3 M⊙ |
| EM counterpart | None | Yes (usually) |
| Hawking radiation | Dark photons | Visible photons |

Events in the "mass gap" (2.2 – 5 M⊙) without electromagnetic counterparts are candidates for dark sector mergers. GW190814 (secondary mass 2.6 M⊙) is consistent with a dark neutron star.

Sub-solar compact objects, if detected in gravitational wave data, would have no formation channel in standard physics but are predicted here (dark Chandrasekhar limit = 0.7 M⊙).

## 5.5 Bullet Cluster

The dark-visible scattering cross section is suppressed by $\varepsilon^2 \approx 5.6 \times 10^{-10}$, making dark matter effectively collisionless with respect to visible matter while maintaining self-interaction within the dark sector. This is

consistent with the Bullet Cluster observation (1E 0657-56), where dark matter passes through the collision while baryonic gas is shock-heated.

---

# 6. Maxwell Equations from Torsion

## 6.1 The Connection

The 5D connection decomposes as:

$\Omega = \omega + K$

where ω is the Levi-Civita connection and K is the contorsion tensor. Teleparallelism requires vanishing curvature:

$R(\Omega) = 0$

All dynamics reside in the torsion $T^a = K^a_b \wedge \theta^b$.

## 6.2 Identification of the Electromagnetic Field

The Lie algebra so(1,4) has 10 generators. The contorsion K decomposes into irreducible components:

| Generator | Physical field | Count |
|---|---|---|
| $J_{01}$ | Gravitational (σ-φ sector) | 1 |
| $J_{0i}$ (i=2,3,4) | Electric field $E_i$ | 3 |
| $J_{1i}$ (i=2,3,4) | Magnetic field $B_i$ | 3 |
| $J_{ij}$ (i,j=2,3,4) | Spin connection | 3 |

The electromagnetic field tensor:

$F_{0i} = E_i = m_0\sigma K_{0i}$ $F_{ij} = \varepsilon_{ijk}B_k = m_0\sigma K_1[_{ij}]$

## 6.3 Homogeneous Equations: dF = 0

From $R(\Omega) = 0$, the Bianchi identity gives $D_\omega(K) = 0$. Since $F = m_0\sigma K_{EM}$ and $D(K_{EM}) = 0$:

$dF = 0$

In 3+1 components:

**∇·B = 0** — The magnetic field B resides in $K_{1i}$, which is the torsion component along the φ-direction. Since φ is topologically $S^1$ (a circle), and a circle has no boundary, there can be no magnetic source. The absence of magnetic monopoles is a topological theorem, not a postulate.

**∇×E = −∂B/∂t** — Faraday's law of induction. The projection of dF = 0 onto the (0,i,j) components. The rate of change of the magnetic flux through the φ-circle generates the electric circulation.

## 6.4 Inhomogeneous Equations: d⋆F = ⋆J

From the TEGR (Teleparallel Equivalent of General Relativity) field equation, varying the torsion Lagrangian L = T^a_{bc} S_a^{bc}/(4κ):

∂_μ(e S_a^{μν}) = eκ t_a^ν

For the electromagnetic generators ($J_{0i}$, $J_{1i}$), the superpotential S reduces to the Hodge dual ⋆F, and the source t reduces to the 4-current J^μ.

In 3+1 components:

**$\nabla\cdot\mathbf{E} = \rho/\varepsilon_0$** — Gauss's law. Charge density ρ is the torsion source density.

**$\nabla\times\mathbf{B} = \mu_0\mathbf{J} + (1/c^2)\partial\mathbf{E}/\partial t$** — Ampère's law with Maxwell's displacement current.

### 6.5 The Constants $\varepsilon_0$ and $\mu_0$

These are not fundamental. They are unit conversion factors derivable from the sole electromagnetic constant α = 1/137:

$e^2 = 4\pi\alpha\hbar c\ \varepsilon_0 = 1/(4\pi\alpha\hbar c)\ \mu_0 = 4\pi\alpha\hbar/c\ \varepsilon_0\mu_0 = 1/c^2$

Electromagnetism has zero free constants. α = 137 = 163 − 2×13 is determined by the prime skeleton of PG(5,2).

### 6.6 Charge Quantization

The charge quantum derives from the Fano plane structure. Each line of PG(2,2) contains 3 points. The fractional charges of quarks are:

- Electron: 1 (full point)
- Up quark: 2/3 (2 of 3 points on a Fano line)
- Down quark: −1/3 (1 of 3 points, opposite orientation)

### 6.7 Electromagnetic and Gravitational Waves

Both are torsion waves on a flat background (R = 0):

| **Property** | **EM wave** | **Gravitational wave** |
|---|---|---|
| Torsion component | $K_{0i}$, $K_{1i}$ (spatial) | $K_{01}$ (temporal) |
| φ-circle mode | Fourier modes | Cross modes σ-φ |
| Quantum | Photon (n = N, spin 1) | Graviton (δn = 1, spin 2) |
| Speed | c | c |
| Polarizations | 2 | 2 |
| Source | Charge (n on φ) | Mass (torsion asymmetry) |

---

## 7. General Relativity as the Late-Time Limit

### 7.1 The Torsion-Curvature Equivalence

In the teleparallel framework, the fundamental variables are the contorsion tensor K and the vierbein $e^a_\mu$. The connection decomposes as $\Omega = \omega + K$, with vanishing total curvature:

$$R(\Omega) = R(\omega) + D_\omega(K) + K \wedge K = 0$$

This is an identity — not a field equation. It means:

$$\mathbf{R(\omega) = -D_\omega(K) - K \wedge K}$$

The Levi-Civita curvature $R(\omega)$ — which is what general relativity calls "spacetime curvature" — is identically equal to a quadratic expression in the torsion. Curvature and torsion are not independent. They are two descriptions of the same geometric content.

## 7.2 The Einstein Field Equation

The TEGR field equation (from varying the torsion Lagrangian) is:

$$\partial_\mu(e\, S_a{}^{\mu\nu}) = e\kappa(t_a{}^\nu + T_a{}^\nu)$$

where S is the superpotential, t is the gravitational energy-momentum pseudo-tensor, and T is the matter energy-momentum tensor. Using the identity $R(\omega) = -D_\omega(K) - K \wedge K$ to rewrite the left side:

$$\mathbf{G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G\, T_{\mu\nu}}$$

The Einstein equation is not an independent postulate. It is the torsion field equation rewritten in curvature variables. The cosmological constant $\Lambda$ arises from the vacuum torsion — the residual K at $\sigma \to \infty$.

## 7.3 What Changes

In general relativity, the metric $g_{\mu\nu}$ is the dynamical variable and curvature $R(\omega)$ encodes the physics. In the torsion framework:

| GR quantity | Torsion origin | Status |
|---|---|---|
| $g_{\mu\nu}$ | Flat 5D metric (given, not dynamical) | Background |
| $R(\omega)$ | $-D_\omega(K) - K \wedge K$ | Derived, not fundamental |
| $\Gamma^\mu{}_{\nu\rho}$ (Christoffel) | $\omega + K$ projected to 4D | Effective connection |
| $G_{\mu\nu} = 8\pi G T_{\mu\nu}$ | TEGR field equation in curvature variables | Rewriting |
| $\Lambda$ | Vacuum torsion at $\sigma \to \infty$ | Computed: $(4/3)m_0^4/N^8$ |
| Gravitational waves | $K_{01}$ oscillation on flat background | Torsion wave |
| Singularities | Do not exist ($T_{max} = N/254$, finite) | Resolved |
| Black hole interior | $\sigma \to \Delta$ (graviton wall), two-sided | Finite |

The curvature description becomes exact in the limit $\sigma \to \infty$ (late cosmic epoch), where K becomes small and the quadratic term $K \wedge K$ is negligible. In this limit, $R(\omega) \approx -D_\omega(K)$, which is the linearized Einstein equation.

At early times, strong fields, or small scales, the full torsion structure (with finite N and discrete n) is required. General relativity is the macroscopic, late-time, continuous approximation of the discrete torsion dynamics.

### 7.4 The Gravitational Constant

Newton's constant G is not a free parameter. It is determined by the contorsion at the graviton wall:

$G = A^2/(8\pi m_0^2) \times$ (geometric factors from $K_{01}$)

where $A = H_0 c = 6.54 \times 10^{-10}$ m/s$^2$ is the acceleration scale from the two-time constraint (Part I, §6.1). The weakness of gravity ($G \sim 10^{-38}$ in natural units) follows from the smallness of $A/m_0$ — the ratio of the cosmological acceleration to the mass scale.

---

## 8. The Wave Equation and Quantum Mechanics

### 7.1 The 5D Massless Equation

On the constrained metric (Part I, §6.2):

$ds^2 = (A^2/2)(d\sigma^2 + \sigma^2 d\varphi^2) - dx^2 - dy^2 - dz^2$

The massless Klein-Gordon equation $\Box_5\Psi = 0$ expands to:

$(2/A^2)[\sigma^{-1}\partial_\sigma(\sigma\partial_\sigma\Psi)] + (2/A^2\sigma^2)\partial^2_\varphi\Psi - \nabla^2\Psi = 0$

### 7.2 Separation on the Circle

Since φ is periodic ($S^1$), we expand in Fourier modes:

$\Psi(\sigma, \varphi, x^\mu) = \Sigma_n \psi_n(\sigma, x^\mu) \cdot e^{in\varphi}$

with n integer (bosons) or half-integer (fermions), the latter from the $Pin_{+,-,-}$ double cover ($\varphi \to \varphi + 4\pi$ identification).

Substituting $e^{in\varphi}$ replaces $\partial^2_\varphi \to -n^2$. The equation for each mode becomes:

$(2/A^2)\partial^2_\sigma\psi_n + (2/A^2\sigma)\partial_\sigma\psi_n - \nabla^2\psi_n = (2n^2/A^2\sigma^2)\psi_n$

### 7.3 The 4D Massive Equation

The right side is the mass term:

$m_n^2 = 2n^2\hbar^2/(A^2\sigma^2)$

Each Fourier mode satisfies the massive Klein-Gordon equation:

$(\Box_4 + m_n^2)\psi_n = 0$

Mass originates from the φ-circle. A 5D massless field decomposes into a tower of 4D massive modes indexed by n. This is the Kaluza-Klein mechanism applied to the twist geometry.

### 7.4 The Radial Equation

The σ-dependent part R(σ) satisfies:

$\sigma^2 R'' + \sigma R' + (u^2 - n^2)R = 0$

where u = kAσ/√2. This is **Bessel's equation** of order n.

Solutions: R(σ) = J_n(kAσ/√2)

The full wave function:

Ψ = J_n(kAσ/√2) · e^{inφ} · e^{ik□·x□}

## 7.5 Special Cases

**n = 0** (classical limit): $J_0$ is smooth, non-oscillatory at small argument. This gives classical wave propagation — sound waves, fluid dynamics. No quantum effects.

**n = 1/2** (lightest fermion): J_{1/2}(z) = √(2/πz) sin(z). The Bessel function reduces to a sinusoidal oscillation. This is the electron's radial profile.

**n = N** (photon): J_N for large N and argument comparable to N gives the classical electromagnetic wave. The transition from quantum to classical happens when n and the argument are both large (Debye asymptotic regime).

## 7.6 Non-Relativistic Limit: The Schrödinger Equation

For E ≈ m_n $c^2$ + (small kinetic energy), write:

ψ_n(σ, x□) = u_n(x□) · e^{−im_n$c^2$σ/ℏ}

Substituting into the massive Klein-Gordon equation and keeping leading-order terms:

**iℏ ∂u_n/∂σ = −(ℏ²/2m_n)∇²u_n**

This is the Schrödinger equation, with σ playing the role of time and m_n the particle mass.

Adding external fields from contorsion components:

- V_grav = m_n Φ_grav from $K_{01}$ (gravitational contorsion)
- V_EM = qΦ_EM from $K_{0i}$ (electromagnetic contorsion)

The full Schrödinger equation with potentials:

**iℏ ∂ψ/∂t = [−ℏ²/(2m)∇² + V_grav + V_EM]ψ**

Both potentials arise from the same torsion structure.

## 7.7 Why Quantum Mechanics Has Its Properties

The following features of quantum mechanics are structural consequences of the φ-circle:

| QM feature | Origin in framework |
|---|---|
| ψ is complex | e^{inφ} is complex; imaginary part from circle |
| Quantized states | Fourier modes on $S^1$ are discrete |
| | ψ |

| QM feature | Origin in framework |
|---|---|
| $\Delta x \cdot \Delta p \geq \hbar/2$ | $\varepsilon + \omega + P = 0$ prevents simultaneous diagonalization |
| Superposition | Fourier completeness on $S^1$ |
| Entanglement | Shared φ-coordinate from shared $t_2$ |

---

## 9. The Correspondence Table

The Faraday induction law $\nabla \times E = -\partial B/\partial t$, the σ-evolution on the φ-circle, and the Bockstein bridge $F_2 \to \mathbb{Z} \to \mathbb{Q} \to \mathbb{R}$ share a common mathematical structure. The following table collects the correspondences:

| Mathematical structure | Physical realization | Algebraic origin |
|---|---|---|
| φ is $S^1$ (circle) | $\nabla \cdot B = 0$ (no monopole) | Circle has no boundary |
| $dF = 0$ (Bianchi) | Faraday induction | $R(\Omega) = 0$ on flat space |
| $d\star F = \star J$ (TEGR) | Gauss + Ampère | Torsion field equation |
| $e^{in\varphi}$ on $S^1$ | Wave function ψ | Fourier mode on circle |
| $\Box_5 \Psi = 0$ | Schrödinger equation | 5D → 4D reduction |
| $K_{0i}$ oscillation | EM wave | Spatial torsion mode |
| $K_{01}$ oscillation | Gravitational wave | Temporal torsion mode |
| σ-evolution through $S^1$ | $\oint E \cdot dl = -d\Phi_B/d\sigma$ | Stokes theorem on circle |
| Lenz resistance of $S^1$ | $m_{app} = m_0\sqrt{(1-n^2\varepsilon)}$ | Inertia from angular momentum conservation |
| Bockstein: $F_2 \to \mathbb{R}$ | Discrete → continuous | Short exact sequence |

The resistance of the φ-circle to changes in σ (angular momentum conservation of the Fourier mode $e^{in\varphi}$) is the mechanism that produces apparent mass. A mode with n = N (photon) precesses maximally and offers no resistance — zero mass. A mode with small n (top quark) precesses minimally and resists maximally — large mass.

---

## 10. Experimental Signatures

### 9.1 Near-Term (2025–2030)

| Test | Prediction | Experiment |
|---|---|---|
| Sub-solar compact object in GW | Exists (dark $M_{Ch} = 0.7\ M_\odot$) | LIGO/Virgo/KAGRA |
| Mass gap events without EM | Dark mergers | LIGO O4/O5 |
| Dark photon at $\varepsilon = 2.4\times10^{-5}$ | Massless | Belle II, LDMX |
| 30° entanglement anomaly | $\delta C(30°) \sim \Delta$ | Tabletop Bell test |

| Test | Prediction | Experiment |
|---|---|---|
| DM direct detection at ~1 GeV | Dark proton | SuperCDMS, NEWS-G |

### 9.2 Mid-Term (2030–2040)

| Test | Prediction | Experiment |
|---|---|---|
| Higgs self-coupling | $\lambda = 0.1305$ (vs SM 0.129) | HL-LHC |
| Neutron spin-gravity coupling | Torsion correction O(n/N) | Improved COW |
| Photon oscillation | $\varepsilon^4 \approx 3.2\times10^{-19}$ | Light-through-wall |

### 9.3 Decisive Test

Detection of a sub-solar compact object in gravitational wave data. Standard physics provides no formation channel for compact objects below ~1 $M_\odot$. The framework predicts them (dark Chandrasekhar = 0.7 $M_\odot$). If found, no alternative explanation exists within conventional astrophysics.

---

## 11. Summary of New Results

| # | Result | Section | Match |
|---|---|---|---|
| 1 | Dark sector with N' = N/2 | §2 | structural |
| 2 | Universal √2 mass scaling | §2.2 | exact |
| 3 | Dark electron = muon (105.67 MeV) | §3.2 | 100.0% |
| 4 | $m_b = (7/3)m_\tau = 4148$ MeV | §3.3 | 99.2% |
| 5 | Three generations from two circles | §3.4 | structural |
| 6 | Graviton wall shared | §4.1 | structural |
| 7 | Gravity from $\sin^2(30°) \neq \cos^2(30°)$ | §4.2 | structural |
| 8 | Dark Chandrasekhar = 0.7 $M_\odot$ | §5.1 | prediction |
| 9 | SMBH seeds from dark BHs | §5.3 | consistent |
| 10 | $\nabla\cdot B = 0$ from $S^1$ topology | §6.3 | exact |
| 11 | All 4 Maxwell eqs from torsion | §6 | derived |
| 12 | $\Box_5\Psi = 0 \rightarrow$ Schrödinger | §8.6 | derived |
| 13 | $\psi = e^{in\varphi}$ (wave function identity) | §8.2 | structural |
| 14 | Dark photon $\varepsilon = 2.37\times10^{-5}$ | §2.6 | prediction |
| 15 | Bullet Cluster consistency | §5.5 | consistent |
| 16 | $G_{\mu\nu} = 8\pi G T_{\mu\nu}$ from torsion | §7.2 | derived |

| # | Result | Section | Match |
|---|---|---|---|
| 17 | Λ from vacuum torsion | §7.3 | 99.6% (Part I) |
| 18 | No singularities (T_max finite) | §7.3 | structural |

Combined with Part I: 32 numerical predictions, 26+ resolved problems, zero free parameters beyond $F_2$.

---

## 12. Conclusion

The derivation chain $F_2$ → Fano → Cones → Twist → Torsion, established in Part I, extends naturally to the dark sector, the classical field equations, and the wave equation. No additional structure is needed.

The contorsion tensor K on flat 5D spacetime contains all interactions:

- $K_{01}$: gravity
- $K_{0i}$: electric field
- $K_{1i}$: magnetic field
- $J_{ij}$: spin

The wave equation $\Box_5\Psi = 0$ contains all dynamics:

- n = 0: classical wave propagation
- n = ½: fermion (electron, quark)
- n = N: photon (classical EM limit)
- Bessel J_n: radial profile
- Schrödinger: non-relativistic limit

General relativity itself is not independent: the Einstein equation $G_{\mu\nu} = 8\pi G T_{\mu\nu}$ is the torsion field equation rewritten in curvature variables, exact in the late-time limit $\sigma \to \infty$. The cosmological constant, Newton's constant, and the absence of singularities all follow from the torsion structure.

The electromagnetic field, gravitational field, quantum wave function, and dark matter all emerge as aspects of one geometric structure: torsion modes on a flat five-dimensional spacetime with two time dimensions, forced into existence by the algebraic identity $\varepsilon + \omega + P = 0$ over $F_2$.

---

## References

[1–12] As in Part I.

13. Weitzenböck, R. "Invariantentheorie." Noordhoff, Groningen (1923).
14. Aldrovandi, R. and Pereira, J.G. "Teleparallel Gravity." Springer (2013).
15. Cho, Y.M. "Einstein Lagrangian as the translational Yang-Mills Lagrangian." Phys. Rev. D 14, 2521 (1976).
16. Clowe, D. et al. "A direct empirical proof of the existence of dark matter." Astrophys. J. 648, L109 (2006).
17. Anderson, E.K. et al. "Observation of the effect of gravity on the motion of antimatter." Nature 621, 716 (2023).
18. Abbott, R. et al. "GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object." Astrophys. J. Lett. 896, L44 (2020).
19. Kaluza, T. "Zum Unitätsproblem der Physik." Sitzungsber. Preuss. Akad. Wiss. Berlin, 966 (1921).

---

## Accompanying Computations (in repository root)

1. compute_dark_sector.js — Dark fermion mass spectrum
2. compute_dark_sector_v2.js — Complete visible + dark particle table
3. compute_dark_bh.js — Dark black hole properties and LIGO predictions
4. compute_maxwell.js — Maxwell equations from torsion derivation
5. compute_schrodinger.js — 5D wave equation to Schrödinger reduction
6. compute_experiments.js — Experimental proposals with sensitivity estimates

---

Copyright 2026. All rights reserved.

**Author:** Szőke Barna