
Uninformative ecology suppresses recovery before and after compression
K=4 and full ARI ≥ 0.8; count out of 10 reused datasets
Prespecified coordinate space and neural seed
morphology standard 8 8 0 0 5 9
ecology standard 0 0 0 0 10 9
balanced traits 0 0 0 0 10 10
morphology view pcs 8 8 0 0 6 9
ecology view pcs 0 0 0 0 10 8
fused profiles 0 0 0 0 10 10
joint pcs 0 0 8 9 10 10
morphology density pcs 8 8 0 0 6 9
ecology density pcs 0 0 0 0 10 8
paired 0 raw 0 0 9 9 10 10
paired 0 standard 0 0 9 10 10 10
paired 101 raw 0 0 7 9 9 10
paired 101 standard 0 0 8 7 9 10
paired 202 raw 0 0 8 10 10 10
paired 202 standard 0 0 8 10 10 10
morphology 0 standard 8 8 0 0 10 10
morphology 101 standard 8 9 0 0 10 8
morphology 202 standard 8 8 0 0 10 10
ecology 0 standard 0 0 0 0 10 9
ecology 101 standard 0 0 0 0 10 9
ecology 202 standard 0 0 0 0 10 10
Only morphology has signal N=64
Only morphology has signal N=256
Complementary signal N=64
Complementary signal N=256
Redundant signal N=64
Redundant signal N=256
Archived generating law and matched-individual sample size
Successful recoveries / 10
0
2
4
6
8
10