ZOO603T-PRINCIPLES OF FISH BIOLOGY
MCQs OF FINAL TERM PAST PAPERS
COMPILED BY: AYESHA BUTT, HAFIZAH SAMEEN, SABA ALI

Shared by Students:

1. Herbivory is the one of most important processes in maintaining ecological balance on Mesoamerican
2. Coprophage which is a feeding strategy involving the consumption of feces.
3. Parrot fish or butterfly fish graze on coral reefs.
4. An adult Menhaden is capable of staining as much *1-2 gallons* of water per minutes.
5. Abdominal dropsy of carps is caused is *Bacterial disease*
6. Limnology is the study of structural and functional interrelationships of organisms of inland waters as they are affected by their dynamic, physical, chemical and biotic environments.
7. Advantage of cages it *reduced the fish mortilty*
8. Father of ichthyology; Peter Artedi
9. Hagfishes belong to which class. Myxini
10. Comparison of salinity of brackish and fresh water ;brackish water salinity is 0.5-30ppt while fresh water salinity is less than 0.5.
11. Claris batrachus is scientific name on (mangur)
12. Fish belong to the phylum (chordata)
13. Scientific name of tiger prawn(penaeus monodon)
14. Bivalves have (bilateral symmetry) square
15. Crustacean are characterized by(armor like thick or thin shell)
16. Migration from sea to fresh water (diadromous)
17. *700* species, elasmobranchs
18. There are currently over *1,100* species of fish that are at risk of extinction.
19. Eurythermal* means can tolerate wide range of temperature.
20. *electric eel* has powerful electric organs
21. Consumption of fish reduces risk of *autoimmune diseases.*
22. Marine ecosystem have a *more* substantial salt content and greater biodiversity in comparison to the fresh water.
23. The aim of the law conservation is to enforce the ..... *International fishes.*
24. .How brackish water are challenging for fishes? *Due to fluctuating salanity*
25. Snakeheads consume *plankton, aquatic insects, and mollusks* during early life stages.
26. Over 20% from areas of upwelling that make up___of ocean surface area? 0.1%
27. Roughly ___ of species inhabit freshwater environments, ranging from desert springs to high altitude streams? : 40%
28. Osteichthyes is the largest class of vertebrates
29. Carp belongs to class Actinopterygii
30. Which is non living abiotic component....water

31. Mariculture refers to.... branch of aquaculture that focuses on cultivation of marine organisms.
32. The scientific name of sardine is *_Sardina pilchardus._*
33. Why a square mesh panel is often included in an otherwise diamond net? *A diamond-shaped mesh catches fish more easily.*
34. First jawed fishes appeared in the geologic record about *100 million* years after jawless fish came about. 5. Muller gave classification of fishes in *1844.*
35. Lampreys* generally inhabit shallow coastal regions and freshwater environments.
36. Strainer fish selects food by *size.*
37. What does the term 'Methaline aquaculture' refer to? *Rearing of aquatic organisms in water containing high salt concentration.*
38. Romer* classified fishes into three classes.
39. Hagfish have *six or ten* barbels around the mouth and a single nostril. *(The available options were: i.6 or 7 ii.7 or 8 iii.8 or 9 iv.6 or 10)*
40. *Cannibalism* is the act of consuming another individual of the same species as fo
41. Cichlids and poecilids are example of *secondary freshwater fish.*
42. Fringe lips are used for *sucking.*
43. Overfishing:* when the number of fish caught is greater than the rate at which the fish reproduce leading to a fall in fish numbers in an area.
44. Which species of voracious fish is found in Africa? *Hemichromis*
45. Which gonadotropin hormones are secreted by pituitary gland that influences the maturation of gonads and spawning in fishes? *FSH & LH
46. Osteichthyes are *bony fishes.*
47. The water body moving in one direction is known as *lotic freshwater.*
48. Fish oil is extracted from *liver.*
49. One mcqs related to body of flatfish
50. *Placoid Scales* are found in sharks and rays.
51. For *Poikilotherm* physiological processes like the development of the embryo, the progress rate is determined by temperature.
52. Lateral line system consists of *100* receptors.
53. frogs* are used to make bait to catch game fish.
54. Ray finned fish *Actinopterygii*
55. The number of individuals and total biomass *decreases*
56. Which of the following is not included in fish management? *fish marketing*
57. *1100* species are at the risk of extinction
58. Cat fish is named *because it has cat's whisker like barbels*
59. Bombay duck is captured by using a characteristic bag-net known as *dol* that is operated against tidal current
60. Cephalopods are *molluscs*
61. Lateral line *Detect water movement and pressure changes*

62. Lung fish *Dipnoi*
63. Scales in lung fish *Cosmoid*
64. Difference in ponds and river *Ponds stagnant water and lakes flowing water*
65. Which one is not an example of carnivore?
66. Productivity in aquaculture is _High productivity
67. Desirable range of alkanity *20-250
68. What is lobester? crustacean
69. L.S. Berg classified fishes in to *seven* classes and separated lampreys and hagfishes into a separate class.
70. Proportion *(40–60%)..* Of the mass of a fish's muscle is made up of locomotory muscle.
71. Characteristics of Osteichthyes is the *presence of bony skeleton.*
72. Scientific name of Bombay-duck is *Harpodon nehereus*
73. Cephalopods belongs to phylum *mollusca* .
74. Which is not a chondrostean *lamprey*.
75. Class MYXINI have mouth with *four* pairs of tentacles.
76. The head shield provided some lift though, *and would have made the* *cephalaspidomorphi* *better swimmers than most of their contemporaries.*
77. Which is not a characteristic of holocephali ..... *Body is covered with scales
78. In the anatomy of fish , which layer gives rise to scales... *Dermis
79. Bottom feeders are fish that inhabit the lower level of the aquarium o feed off the bottom are *bottom dwellers.
80. Marine fishery deals with the *fishery aspects of the sea water or ocean.*
81. Cephalospidomorph were, like most contemporary *fishes, very well armored
82. The gonadotropin hormone secreted by... (pituitary gland)
83. Fish flesh is an excellent source of protein and Vitamin (A+D)
84. Muller's classification (1844) .
85. Cephalopod phylum mollusc
86. Type of ecosystem 2 1-aquatic ecosystem 2-terrestrial ecosystem
87. Rohu fish is an example of Marginal feeders.
88. Driftnets or gill nets banned in 198
89. Ecology derived from two words oikas and logos
90. influences the maturation of gonads and spawning in fishes.pituitary gland
91. Some deep sea male argb (ceration) Fish are mandatory parasites of female fish of some species.
92. The Characiformes constitute more than 30% of the total fish yield in the Amazon basin, whereas the catfishes are of minor importance .
93. The Detritivores living in the beneath region. These organisms often called bottom feeder.
94. Many shark species are pelagic, meaning that they spend much of their lives swimming throughout the open ocean (pelagic zone).
95. The gill rot (branchyomyces) of carps involves the attack of Saprolegnia on the gills of carps.

96. The number of individuals and total biomass decreases at successive trophic levels as Organisms increase in size.
97. The fish flesh is an excellent source of protein, has very little fat, carries a good amount of minerals and vitamins A and D and rich in iodine
98. The water examines for global fisheries employment are 1.75 time higher than the food and agriculture organization estimates.
99. The freshwater system is mainly divided into 3 types based on range lotic, lentic and wetland.
100. The freshwater ecosystem is an aquatic ecosystem that includes lakes, ponds, rivers, streams and wetlands
101. During summers, the temperature of the freshwater ecosystem generally ranges from 30-71f degrees Fahrenheit.
102. Whereas during winters, the temperature ranges from 35-45f degrees Fahrenheit
103. Amiais commonly referred to as the Dog fish or bowfin.
104. The second group of actinopterygians (neopterygii) flourished in the jurassic period and succeeded most chondrosteans.
105. Paddlefishes are large, freshwater chondrosteans.
106. They feed by sieving through large gill rakers or by mucus secreted from the epibranchial glands.
107. (lepomus macrochirus) often eat the larvae of dipteran insects from the bottom of lakes.
108. Members of this subclass are characterized by having five to seven pairs of g
109. The circulatory, or blood vascular, system consists of the heart, the arteries, the capillaries, and the veins.
110. Fish that eat large animals as food are called predatory fish.
111. Roughly …….f species inhabit freshwater environments, ranging from desert springs to high altitude streams. 40%
112. Most of the earth surface (around 70%) is covered with water, but not all the water is freshwater.
113. 80% of available fish stock fully exploited, overexploited, or depleted/recovering belongs to…… overfishing
114. Herbivores also eat algae that grow on rocks or pebbles by mouth using wide lips under their snout.
115. presented the classification of fishes inin particular and chordata in general in a linear order. nelson (2006)
116. Scientific name for shovelnose sturgeons (scaphirhynchus platorynchus)
117. Fish consuming or craving large quantities of food are called? Answer: Voracious

118. ___is an active, voracious predator of fish and invertebrates? Answer: Shark

. Fish consume decaying plants and animal matter as food? Answer: Detrivores

119. Detritivores are ___of all species? Answer: 10%

120. . ___fishes form an important part of the ichthyomass in the Amazon basin? Answer: Detritivorous

121. Most immature adult fish are usually? Answer: Plankton eater

122. This type of fish eats certain selected foods? Answer: Stenophagic

123. This type of fish eats a variety of foods/ Answer: Euryphagic

124. ___typically live in the upper layers of the acquaculture and consume food from the surface? Answer: Surface feeders

125. ___Prefers algae and other submerged vegetation? Answer: Marginal feeders

126. ___Often live in the middle layer of the aquarium? Answer: Middle feeders

127. They eat bottom food such as rotten food? Answer: Bottom feeders

128. Fish can be divided into___categories based on the type of prey and swallowing? Answer: 5

129. The Southeast Asian archer fish preys on___by spitting on nearby trees? Answer: Insects

130. ___Often eat the larvae of dipteran insects from the bottom of lakes? Answer: Bluegills

131. Deep sea eel is also? Answer: Parasitic

132. Female fish obligatorily responds by developing a ___ through which males can absorb nutrients? Answer: Fleshy papillae

133. In this case, the fish swallow the living creatures that live in the water? Answer: Strainers

134. Sometimes ducklings and small birds are also seen in the belly of___? Answer: goosefish

135. . ___change their eating habits as the seasons change? Answer: Bermuda angelfish

136. Even small rats,snakes and tortoises can be found in the stomachs of ___/ Answer: Pike Lepisosteus or Amia

137. The___is an excellent source of protein? Answer: Fish flesh

138. Larvivorous fish eat? Answer: Larvae of mosquito

139. Some fish like the___ are of zoological importance because of their discontinuous distribution and anatomical features? Answer: Lungfish

140. The fish waste after the extraction of oil, is used as? Answer: Fertilizers

141. It is a sticky product, obtained from the skin of the cod and is used as gum? Answer: Fish Glue

142. It is a gelatinous substance, obtained from the air bladder of perches? Answer: Isinglass

143. Indian salmons and___are used in the preparation of special cement and in the clarification of wine and beer? Answer: Catfish

144. Global trade in fish over___billion in 2008? Answer: $102

145. The silvery boney scales of ___are used in the manufacture of artificial pearls especially in___? Answer: Cypnmids+ France

146. Blue revolution refers to? Answer: Fish production

147. Green revolution refers to? Answer: For producing enough food for all

148. ___Deals with the fishery aspects of waters other than marine water? Answer: Inland fishery

149. Mixed farming is called? Answer: Composite fish farming

150. Marine fisheries are conducted in all the oceans and seas of the world, including? Answer: Bays and estuarie

151. . ___Migrates from the sea to the river for breeding? Answer: Hilsa

152. Abdominal dropsy of carps caused by? Answer: Aeromonas punctuata

153. Furunculosis of salmon and trout is caused by? Answer: Aeromonas salmonicida

154. VHS stands for? Answer: Viral haemorrhagic septicaemia

155. Over 20% from areas of upwelling that make up___of ocean surface area? Answer: 0.1%

156. MSY stands for? Answer: Reduction in maximum sustainable yield

157. Bycatch may be upto ___times more than the intended catch? Answer: 8 times

158. It refers to the rate of biomass production? Answer: Productivity

159. The top-soil is the major site for? Answer: Decomposition

160. An ecosystem can be small as an? Answer: Ocean

161. Hagfish are deep-sea? Answer: Bottom dwellers

162. Most species of ___ inhabit marine environments? Answer: Chondrichthyes

163. For example, in many regions, such as ___ the fishing industry is a major source of employment? Answer: Alaska

164. Some organisms are specialists and others are generalist in terms of? Morphology

MID TERM PAST MCQs+ final TERM

1. Father of ichthyology – Peter Artedi.

2. Systematics is the study of the diversity of fish and the evolutionary relationships between these organisms.

3. L.S. Berg classified fishes into seven classes and separated lampreys and hagfishes into a separate class.

4. Romer classified fishes into three classes.

5. J.S. Nelson classified fishes into five classes and raised Sarcopterygii and Actinopterygii to class level.

6. Agnatha do not have an identifiable stomach or any appendages.

7. Gnathostome diversity comprises roughly 60,000 species (99% of all living vertebrates).

8. Subclass Elasmobranchii contains about 700 species.

9. Ostichthys is the largest class of vertebrates.

10. Lobe-finned fish have muscular lobes associated with their fins and usually use lungs for gas exchange.

11. Beaklike mouth is used to graze on small algae growing on hard surfaces.

12. A protrusible mouth allows a fish to extend its reach when attempting to snatch prey or food particles.

13. Three dorsal fins are used for locomotion.

14. The caudal (tail) fin is the primary fin that gives forward propulsion and speed.

15. Placoid scales are found in sharks and rays.

16. Placoid scales are made of a flattened base with a spine protruding towards the rear of the fish.

17. Sharks have placoid scales.

18. Carp belong to the class Actinopterygii (ray-finned fishes).

19. Paddlefishes are large, freshwater chondrosteans.

20. Example of omnivorous fish: Goldfish.

21. Larvivorous fish feed on immature stages of mosquitoes.

22. Herbivory is one of the most important processes in maintaining ecological balance on coral reefs and the Mesoamerican Reef.

23. A feeding strategy involving the consumption of feces: Coprophagy.

24. Less voracious fish species in Africa: Hemichromis.

25. Snakeheads consume plankton, aquatic insects, and mollusks during early life stages.

26. Parrot fish or butterfly fish graze on coral reefs.

27. Cannibalism is the act of consuming another individual of the same species as food.

28. Overfishing occurs when fish are caught at a rate greater than reproduction.

29. Which species of voracious fish is found in Africa? Hemichromis.

30. First jawed fishes appeared in the geologic record about 100 million years after jawless fish came about.

31. The lateral line system consists of about 100 or more sensory organs.

32. Lateral lines can detect water movement and pressure changes; in some species, they contain electroreceptors.

33. Fish belong to the phylum Chordata.

34. Fish class MYXINI has four pairs of tentacles.

35. Bivalves possess bilateral symmetry.

36. Cephalopods belong to the phylum Mollusca.

37. Lungfish belong to the subclass Dipnoi.

38. Scales in lungfish are cosmoid.

39. Hagfish are classified under family Myxinidae.

40. Catfishes are ray-finned fish named for their barbels resembling cat whiskers.

41. Fringe lips are used for sucking food.

42. The anatomy layer that gives rise to fish scales is the dermis.

43. Fishes contain 6–7 gill openings.

44. Snake-like locomotion is aided by trunked dorsal fins.

45. Physoclistic condition occurs when the bladder loses its connection with the digestive tract.

46. Eurythermal fishes can tolerate a wide range of temperatures.

47. Euryhaline fishes can tolerate a wide range of salinity.

48. Brackish water challenges fishes due to fluctuating salinity.

49. The water body moving in one direction is called a lotic freshwater ecosystem.

50. Carp is a freshwater species.

51. Bombay duck is captured by using a bag-net known as dol against tidal currents.

52. Fish oil is extracted from the liver.

53. Fish flesh is an excellent source of protein and vitamins A & D.

54. Global trade in fish was over $102 billion in 2008.

55. Blue revolution refers to increased fish production.

56. Green revolution refers to increased agricultural productivity.

57. Inland fishery deals with the fishery aspects of waters other than marine water.

58. Composite fish farming refers to mixed farming of fish species.

59. Marine fisheries are conducted in all the oceans and seas, including bays and estuaries.

60. Hilsa migrates from the sea to rivers for breeding (anadromous migration).

61. Abdominal dropsy in carps is caused by Aeromonas punctuata.

62. Furunculosis of salmon and trout is caused by Aeromonas salmonicida.

63. VHS stands for Viral Haemorrhagic Septicaemia.

64. Shark is about 15m in length.

65. Hagfish are typically about 50 cm in length..

66. Gnathostome diversity accounts for 99% of all living vertebrates, including humans.

67. Tetrodotoxic poisoning occurs following the ingestion of pufferfish, boxfish, or porcupine fish.

68. Swedish naturalist Peter Artedi is known as the father of ichthyology.

69. Romer gave classification of Osteichthyes, Chondrichthyes, and Placodermi.

70. Dorsal fins assist in rolling and sharp turns.