

In sampling without replacement, an element can be chosen:

Select the correct option

☐	More than once
☐	Less than once
☐	only once
☐	Difficult to tell

Click to S

Question # 3 of 10 (Start time: 03:05:58 PM, 21 July 2023)

Total Marks

Who introduced the basic concept of ranked set sampling?

Select the correct option

[Reload Math Equations](#)

☒	McIntyre (1952)
☐	Muhante (1952)
☐	John (1952)
☐	Isaki (1952)

Question # 4 of 10 (Start time: 03:06:33 PM, 21 July 2023)

Total Marks: 1

When the population size is what size, it is recommended to use the entire population instead of a sample?

Select the correct option

 Reload Math Equations

☐	100 or greater
☒	100 or less
☐	1000 or greater
☐	1000 or less

Question # 6 of 10 (Start time: 03:07:50 PM, 21 July 2023)

Total Marks

If the respondents do not provide the required information to the researcher, then it is known as

Select the correct option

 Reload Math Equations

- | | |
|----------------------------------|-------------------------|
| ☐ | none of the above |
| ☐ | both of the above |
| ☒ | problem of non-response |
| ☐ | non-sampling error |

Click to Save Answer & Move to Next Question

Question # 7 of 10 (Start time: 03:08:24 PM, 21 July 2023)

Total Marks: 1

When Grant estimated the population of England:

Select the correct option

 Reload Math Equations

☒	1662
☐	1962
☐	1952
☐	1667

STA632 – Sampling Techniques (Quiz 4)

Question # 9 of 10 (**Start time: 03:08:45 PM, 21 July 2023**)

Problem of non-response

Select the correct option

☐	none
☐	can be solved 
☐	no solution
☐	no meaning

Question # 10 of 10 (Start time: 03:10:23 PM, 21 July 2023)

Total Marks: 1

What is the R command to find out mean of x?

Select the correct option

[Reload Math Equations](#)

☒	mean(x)
☐	mean=x

The term used to define the smallest element in sample selection is:

Select the correct option

Rebad Math Equations

- | | |
|-----------------------|------------------|
| ☐ | population unit |
| ☐ | statistical unit |
| ☐ | respondent |
| ☐ | sampling unit |

Click to Save Answer & Move to Next Question



Question # 2 of 10 (Start time: 02:53:03 PM, 21 July 2023)

Total Marks: 1

Horvitz and Thompson introduced an unbiased estimator in:

Select the correct option

 Rebad Math Equations

☐	1952
☐	1652
☐	1967
☐	1697

[Click to Save Answer & Move to Next Question](#)

Question #1 of 10 (Start time: 02:51:36 PM, 21 July 2023)

Total Marks:

What is the value of combined type ratio estimator with $\bar{y}_s = 18.5$, $\bar{x}_s = 80.25$ and $\bar{x} = 75.32$?

Select the correct option

 Rebad Math Equations

☐	16.323
☐	19.364
☒	17.363
☐	15.632

[Click to Save Answer & Move to Next Question](#)

Question # 3 of 10 (Start time: 02:44:24 PM, 21 July 2023)

Total Marks:

What is the value of combined type ratio estimator with $\bar{y}_x = 15.5$, $\bar{x}_y = 70.25$ and $\bar{x} = 65.32$?

Select the correct option

 Reload Math Equations

☐	14.50	14.50
☐	13.50	

[Click to Save Answer & Move to Next Question](#)

Question # 6 of 10 (Start time: 02:47:28 PM, 21 July 2023)

Total Marks: 1

Auxiliary variable is also called:

Select the correct option

 Reload Math Equations

- | | |
|-----------------------|----------------------|
| ☐ | dependent Variable |
| ☐ | independent Variable |
| ☐ | sampling Variable |
| ☐ | supporting variable |
- supporting variable ✓

[Click to Save Answer & Move to Next Question](#)

Which of the following sampling methods is the best way to select a group of people for a study if you are interested in making statements about the larger population?

Select the correct option

 Reload Math Equations



Non-Random sampling



Random sampling

