

STA641 - Statistical Packages and its Applications

Question # 8 of 8 (Start time: 11:31:01 PM, 09 January

There are basically ... types of errors

Select the correct option



☐	Four
☐	Two
☐	Five
☐	Three

A large, thick green checkmark is drawn over the 'Two' option, indicating it is the correct answer.

Question # 7 of 8 (Start time: 11:29:28 PM, 09 January 2024)

Formula which is used to multiply a matrix is

Select the correct option

☒	=MMULT(array1,array2)
☐	=MMULT(array+array2)
☐	=MMULT(array*array2)
☐	=MUL(array1,array2)

STA841 - Statistical Packages and its Applications (Quiz 3)

Question # 1 of 2 (Start time: 11:23:11 PM, 09 January 2024)

If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

- | | |
|-----------------------|--------------|
| ☐ | F-test |
| ☐ | t-test |
| ☐ | Z-test |
| ☐ | Both a and b |
- 

Question # 6 of 8 (Start time: 11:28:37 PM, 09 Ja

=NORM.S.DIST function. Returns the

Select the correct option

☒	Normal distribution
☐	Standard normal distribution
☐	Chi distribution
☐	Poisson distribution

Question # 8 of 8 (Start time: 11:10:35 PM, 09

=NORM.S.DIST function. Returns the

Select the correct option

- | | |
|-----------------------|------------------------------|
| ☐ | Poisson distribution |
| ☐ | Standard normal distribution |
| ☐ | Normal distribution |
| ☐ | Chi distribution |
- 

TA641 - Statistical Packages and its Applications (

Question # 5 of 8 (Start time: 11:08:09 PM, 09 January

Test statistics for Z-test is

Select the correct option



$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$



$$Z = \frac{\bar{x} - x}{\frac{\sigma}{\sqrt{n}}}$$



$$Z = \frac{\mu - \bar{x}}{\frac{\sigma}{\sqrt{n}}}$$



$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$



STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 4 of 8 (Start time: 11:06:41 PM, 09 January 2024)

Rejecting the null hypothesis when it's actually true is

Select the correct option

☐ Level of significance

☒ Type I error

☐ Power of a test

☐ Type II error

Formula which is used to multiply a matrix is

Select the correct option

☐	=MUL(array1, array2)
☐	=MMULT(array1, array2)
☐	=MMULT(array*array2)
☐	=MMULT(array+array2)




Question # 3 of 8 (Start time: 11:05:56 PM, 09 January 2024)

Function to generate random numbers from normal distribution is

Select the correct option

- | | |
|----------------------------------|---|
| ☐ | =NORM.INV(rand(), standard deviation, mean) |
| ☐ | =NORM.INV(rand(), mean, median) |
| ☒ | =NORM.INV(rand(), mean, standard deviation) |
| ☐ | =NORM.S.DIST(rand(), mean, variance) |



Z-test and T-test are used for

Select the correct option

☐ Three and more than three means

☒ Singles or two mean testing



Question # 1 of 8 (Start time: 11:03:27 PM, 09 January 2024)

When p value is less than alpha then we will

Select the correct option

☐	Don't Reject the Null Hypothesis
☒	Reject the Null Hypothesis

Question # 3 of 8 (Start time: 10:49:41 PM, 09 January 202

Null hypothesis is denoted by

Select the correct option

H_0



H_1



If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

☐ T-test

☐ Both a and b

☐ F-test

☐ Z-test

Function to generate random numbers from normal distribution is

Select the correct option



=NORM.S.DIST(rand(), mean, variance)



=NORM.INV(rand(), standard deviation, mean)



=NORM.INV(rand(), mean, median)



=NORM.INV(rand(), mean, standard deviation)




Question # 1 of 8 (Start time: 10:47:01 PM, 09 January 2024)

F test is used for

Select the correct option

☐ Singles or two mean testing

☒ Three and more than three means testing



Function to generate random numbers from normal distribution is

Select the correct option



=NORM.S.DIST(rand(), mean, variance)



=NORM.INV(rand(), standard deviation, mean)



=NORM.INV(rand(), mean, median)



=NORM.INV(rand(), mean, standard deviation)



If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

☐	T-test
☐	Both a and b
☐	F-test
☐	Z-test

Test statistics for Z-test is

Select the correct option



$$Z = \frac{\bar{x} + \mu}{\frac{\sigma}{\sqrt{n}}}$$



$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$



$$Z = \frac{\mu - \bar{x}}{\frac{\sigma}{\sqrt{n}}}$$



$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$



Question # 8 of 8 (start time: 10:16:35 PM, 09 January 2024)

Total

If Many of the Observation lies on the line or closest to the line

Select the correct option



Reload Math Equations

☒ Follows the Normal distribution



☐ Doesn't follows the Normal distribution

When p value is less then alpha then we will

Select the correct option

☐ Don't Reject the Null Hypothesis

☒ Reject the Null Hypothesis



STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 5 of 8 (Start time: 10:14:34 PM, 09 January 2024)

Function to generate random numbers from normal distribution is

Select the correct option

- | | | |
|----------------------------------|--|---|
| ☐ | =NORMINV(rand(), mean, median) | |
| ☐ | =NORMSDIST(rand(), mean, variance) | |
| ☒ | =NORMINV(rand(), mean, standard deviation) |  |
| ☐ | =NORMINV(rand(), standard deviation, mean) | |

Question # 4 of 8 (start time: 10:13:55 PM, 09 January 2024)

Standard normal distribution has a mean ... and a standard deviation ...

Select the correct option

- | | |
|----------------------------------|-----------|
| ☐ | One, Two |
| ☐ | One, Zero |
| ☒ | Zero, One |
| ☐ | Zero, Two |
- 

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 3 of 8 (**start time: 10:13:08 PM, 09 January 2024**)

If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

- | | |
|----------------------------------|--------------|
| ☒ | T-test |
| ☐ | Z- test |
| ☐ | F-test |
| ☐ | Both a and b |

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 2 of 8 (**start time: 10:12:09 PM, 09 January 2024**)

P- value for the two tailed test is

Select the correct option

☐ Two tailed Test: P-value = $1 - NORM.S.DIST(|z_{\alpha}|, TRUE)$

☒ Two tailed Test: P-value = $2(1 - NORM.S.DIST(|z_{\alpha}|, TRUE))$



STA641 – Statistical Packages and its Applications (Quiz 3)

Question # 1 of 8 (start time: 10:08:01 PM, 09 January 2024)

F test is used for

select the correct option

☒ Three and more than three means testing

☐ Singles or two mean testing

Question # 8 of 8 (start time: 09:47:16 PM, 09 January 2024)

Do not rejecting the null hypothesis when it's actually false is

Select the correct option

- ☐ Type I error
- ☒ Type II error
- ☐ Power of a test
- ☐ Level of significance

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 7 of 8 (Start time: 09:45:38 PM, 09 January 2024)

Test statistics for Z-test is

Select the correct option

☐	$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$
☐	$Z = \frac{\bar{x} + \mu}{\frac{\sigma}{\sqrt{n}}}$
☒	$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$

Question # 5 of 8 (start time: 09:44:25 PM, 09 January 2024)

If the data is not normal, we will go for the

Select the correct option

☐	Parametric Procedures
☐	Chi Square
☐	Rayleigh distribution
☐	Non-Parametric Procedures

Question # 4 of 8 (Start time: 09:43:48 PM, 09 January 2024)

Z-test and T-test are used for

Select the correct option

☒	Singles or two mean testing
☐	Three and more than three means

Question # 3 of 8 (start time: 09:43:02 PM, 09 January 2024)

If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

- | | |
|----------------------------------|--------------|
| ☐ | Both a and b |
| ☒ | T-test |
| ☐ | F-test |
| ☐ | Z- test |

Question # 1 of 8 (Start time: 09:43:28 PM, 09 January 2024)

Formula which is used to multiply a matrix is

Select the correct option

☒ =MMULT(array1,array2)

☐ =MUL(array1,array2)

☐ =MMULT(array*array2)

☐ =MMULT(array+array2)

Question # 7 of 8 (Start time: 08:41:12 PM, 09 January 2024)

F test is used for

Select the correct option

☒ Three and more than three means testing

☐ Singles or two mean testing

Question # 4 of 8 (**start time: 08:38:49 PM, 09 January 2024**)

=NORM.S.DIST function. Returns the

select the correct option

- | | |
|-----------------------|------------------------------|
| ☐ | Poisson distribution |
| ☐ | Standard normal distribution |
| ☐ | Chi distribution |
| ☐ | Normal distribution |



Function to generate random numbers from normal distribution is

Select the correct option

☐	=NORM.INV(rand(), mean, median)
☐	=NORM.S.DIST(rand(), mean, variance)
☐	=NORM.INV(rand(), standard deviation, mean)
☐	=NORM.INV(rand(), mean, standard deviation)



F test is used for

Select the correct option

☐ Singles or two mean testing

☐ Three and more than three means testing



Question # 5 of 8 (Start time: 07:57:58 PM, 09 January 2024)

If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

☐	Both a and b
☐	Z- test
☐	F-test
☐	T-test



7:56



Quiz

vulms.vu.edu.pk

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 4 of 8 (**Start time: 07:56:53 PM, 09 January 2024**)

P- value for the left tail test is

Select the correct option

☐	Left tailed Test: P-value = $1 - \text{NORM.S.DIST}(z_{\alpha}, \text{TRUE})$
☐	Left tailed Test: P-value = $\text{NORM.S.DIST}(z_{\alpha}, \text{TRUE})$





Quiz

vulms.vu.edu.pk



STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 3 of 8 (**Start time: 07:55:55 PM, 09 January 2024**)

Function to generate random numbers from normal distribution is

Select the correct option



=NORM.INV(rand(), mean, standard deviation)



=NORM.INV(rand(), standard deviation, mean)



=NORM.INV(rand(), mean, median)



=NORM.S.DIST(rand(), mean, variance)

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 1 of 8 (**Start time: 07:54:23 PM, 09 January 2024**)

If Many of the Observation lies on the line or closest to the line

Select the correct option

☒ Follows the Normal distribution



☐ Doesn't follows the Normal distribution

P-value for the two tailed test is

Select the correct option

Two tailed Test P-value = $2(1 - \text{NORM.S.DIST}(|z_o|, \text{TRUE}))$



Two tailed Test P-value = $1 - \text{NORM.S.DIST}(|z_o|, \text{TRUE})$



Question # 6 of 8 (Start time: 07:51:40 PM, 09 January 2024)

F test is used for

Select the correct option

Singles or two mean testing



Three and more than three means testing



Question # 5 of 8 (Start time: 07:51:03 PM, 09 Jan)

Formula which is used to multiply a matrix is

Select the correct option

☐	=MUL(array1, array2)
☐	=MMULT{array*1array2}
☐	=MMULT{array+1array2}
☐	=MMULT(array1, array2)



Question # 4 of 8 (**Start time: 07:50:27 PM, 09 January 2024**)

Rejecting the null hypothesis when it's actually true is

Select the correct option

<input data-bbox="15 963 103 1041" type="radio"/>	Power of a test
<input data-bbox="15 1209 103 1288" type="radio"/>	Type II error
<input data-bbox="15 1456 103 1534" type="radio"/>	Type I error
<input data-bbox="15 1702 103 1780" type="radio"/>	Level of significance




Question # 1 of 8 (Start time: 07:48:48 PM, 09 January 2024)

=NORM.S.DIST function. Returns the

Select the correct option

☐	Chi distribution
☐	Normal distribution
☐	Poisson distribution
☐	Standard normal distribution



STA541 - Statistical Packages and its Applications (Quiz 3)

Question # 7 of 8 (start time: 07:47:22 PM, 09 January 2024)

standard normal distribution has a mean ... and a standard deviation ...

Select the correct option

☐	Zero, Two
☐	One, Two
☐	One, Zero
☐	Zero, One



P- value for the left tail test is

Select the correct option



Left tailed Test: P-value = $\text{NORM.S.DIST}(x_{\alpha}, \text{TRUE})$



Left tailed Test: P-value = $1 - \text{NORM.S.DIST}(x_{\alpha}, \text{TRUE})$

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 1 of 8 (Start time: 07:41:58 PM, 09 January 2024)

Test statistics for Z-test is

select the correct option



$$z = \frac{\bar{x} - \mu_0}{\sqrt{\frac{s^2}{n}}}$$



$$z = \frac{\bar{x} - \mu_0}{\sqrt{\frac{\sigma^2}{n}}}$$



$$z = \frac{\bar{x} - \mu_0}{\sqrt{\frac{s^2}{n-1}}}$$



$$z = \frac{\bar{x} - \mu_0}{\sqrt{\frac{\sigma^2}{n-1}}}$$

Question # 7 of 8 (Start time: 07:40:06 PM, 09 January 2024)

If Many of the Observation lies on the line or closest to the line

Select the correct option

☐ Doesn't follows the Normal distribution

☒ Follows the Normal distribution




Question # 6 of 8 (Start time: 07:39:29 PM, 09 January 2024)

=NORM.S.DIST function. Returns the

Select the correct option

☐	Chi distribution
☐	Poisson distribution
☐	Normal distribution
☐	Standard normal distribution



Question # 3 of 8 (Start time: 07:37:37 PM, 09 January 2024)

F test is used for

Select the correct option

☐ Singles or two mean testing

☒ Three and more than three means testing




Statistics: Concepts and its Applications (Quiz 3)

Question # 2 of 8 (Start time: 07:36:37 PM, 09 January 2024)

A ____ can only be used if the population standard deviation is known

Select the correct option

☒ Z- test



☐ T-test

STA647 - Statistical Packages and its Applications (Quiz 3)

Question # 1 of 8 (**start time: 07:35:52 PM, 09 January 2024**)

Formula which is used to multiply a matrix is

Select the correct option

☐	=MUL(array1, array2)
☐	=MMULT(array1, array2)
☐	=MMULT(array*array2)
☐	=MMULT(array+array2)

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 8 of 8 (**Start time: 07:34:39 PM, 09 January 2024**)

If the sample size is large ($n > 30$) and the standard deviation is unknown then we will apply

Select the correct option

Z- test



F- test

F test is used for



Select the correct option

☐	Singles or two mean testing
☐	Three and more than three means testing



Question # 6 of 8 (**Start time: 07:33:43 PM, 09 January 2024**)

There are basically types of errors

Select the correct option


- | | |
|-----------------------|-------|
| ☐ | Five |
| ☐ | Three |
| ☐ | Four |
| ☐ | Two |
- 

Question # 5 of 8 (Start time: 07:33:04 PM, 09 January)

When p value is less than alpha then we will

Select the correct option

☒ Reject the Null Hypothesis



☐ Don't Reject the Null Hypothesis

STA641 – Statistical Packages and its Applications (Quiz 3)

Question # 4 of 8 (**Start time: 07:32:34 PM, 09 January 2024**)

Standard normal distribution has a mean ... and a standard deviation

Select the correct option

☒	Zero, One
☐	One, Two
☐	Zero, Two
☐	One, Zero

Question # 3 of 8 (Start time: 07:31:19 PM, 09 January 2024)

P- value for the left tail test is

Select the correct option

☐	Left tailed Test: P-value = $1 - \text{NORM.S.DIST}(z_{\alpha}, \text{TRUE})$
☒	Left tailed Test: P-value = $\text{NORM.S.DIST}(z_{\alpha}, \text{TRUE})$



Question # 2 of 8 (**Start time: 07:30:49 PM, 09 January 2024**

If the data is not normal, we will go for the

Select the correct option

☐	Parametric Procedures
☐	Non-Parametric Procedures
☐	Rayleigh distribution
☐	Chi Square



Question # 1 of 8 (Start time: 07:30:13 PM, 09 January 202

Rejecting the null hypothesis when it's actually true is

Select the correct option

☐	Type I error	
☐	Power of a test	
☐	Type II error	
☐	Level of significance	

Question # 8 of 8 (Start time: 07:28:16 PM, 09 January 202

Test statistics for Z-test is


Select the correct option



$$z = \frac{\bar{x} + \mu}{\sqrt{\frac{\sigma^2}{n}}}$$



$$z = \frac{\bar{x} - \mu}{\sqrt{\frac{\sigma^2}{n}}}$$



$$z = \frac{\bar{x} - \mu}{\sqrt{\frac{\sigma^2}{n}}}$$



$$z = \frac{\bar{x} - \mu}{\sqrt{\frac{\sigma^2}{n}}}$$

Question # 6 of 8 (Start time: 07:26:12 PM, 09 January 20

=NORM.S.DIST function. Returns the

Select the correct option

☐	Poisson distribution
☒	Normal distribution
☐	Chi distribution
☐	Standard normal distribution

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 2 of 8 (**Start time: 07:23:14 PM, 09 January 2024**)

Function to generate random numbers from normal distribution is

Select the correct option



=NORM.INV(rand(), standard deviation, mean)



=NORM.INV(rand(), mean, median)



=NORM.INV(rand(), mean, standard deviation)



=NORM.S.DIST(rand(), mean, variance)

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 1 of 8 (**start time: 07:22:28 PM, 09 January 2024**)

If the data is not normal, we will go for the

Select the correct option

☐

Chi Square

☐

Non-Parametric Procedures

☐

Parametric Procedures

☐

Rayleigh distribution

Question # 7 of 8 (**Start time: 07:16:25 PM**)

=NORM.S.DIST function. Returns the

Select the correct option

☐	Standard normal distribution
☐	Poisson distribution
☐	Normal distribution
☐	Chi distribution

Question # 6 of 8 (Start time: 07:15:55 PM, 09 Jan

Formula which is used to multiply a matrix is

Select the correct option


☐	=MMULT{array*array2}
☐	=MMULT(array1, array2)
☐	=MUL(array1, array2)
☐	=MMULT{array+array2}



If the data is not normal, we will go for the

Select the correct option

☐	Parametric Procedures
☐	Chi Square
☐	Rayleigh distribution
☐	Non-Parametric Procedures



QQ plot is used to check

Select the correct option

☐	Mean
☒	Normality



STA641 – Statistical Packages and its Applications (Quiz 3)

Question # 3 of 8 (**Start time: 07:14:26 PM, 09 January 2024**)

A ____ can only be used if the population standard deviation is known

Select the correct option

☐ T-test

☒ Z- test



Question # 2 of 8 (**Start time: 07:13:50 PM, 09 January 2024**)

Standard normal distribution has a mean and a standard deviation

Select the correct option

- | | |
|----------------------------------|-----------|
| ☒ | Zero, One |
| ☐ | One, Zero |
| ☐ | One, Two |
| ☐ | Zero, Two |

Question # 1 of 8 (**Start time: 07:13:09 PM, 09 January 2024**)

Alternative hypothesis is denoted by

Select the correct option

☐	H_0
☐	H_1



Question # 8 of 8 (start time: 07:11:04 PM, 09 January 2024)

Total Marks: 1

if the sample size is large ($n > 30$) and the standard deviation is unknown then we will apply

Select the correct option

 Reload Math Equations☐ Z- test☐ F-test

Question # 7 of 8 (Start time: 07:10:12 PM, 09 January 2024)

If the sample size is less than 30 and the standard deviation is unknown then we will apply

Select the correct option

☐	F-test
☐	Both a and b
☐	Z- test
☐	T- test



STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 6 of 8 (**start time: 07:09:32 PM, 09 January 2024**)

QQ plot is used to check

Select the correct option

☒	Normality
☐	Mean



STA641 – Statistical Packages and its Applications (Quiz 3)

Question # 5 of 8 (start time: 07:08:22 PM, 09 January 2024)

Test statistics for Z-test is

Select the correct option

☐	$Z = \frac{\mu - \bar{x}}{\frac{\sigma_{\bar{x}}}{\sqrt{n}}}$
☒	$Z = \frac{\bar{x} - \mu}{\frac{\sigma_{\bar{x}}}{\sqrt{n}}}$
☐	$Z = \frac{\bar{x} + \mu}{\frac{\sigma_{\bar{x}}}{\sqrt{n}}}$
☐	$Z = \frac{\bar{x} - x}{\frac{\sigma_{\bar{x}}}{\sqrt{n}}}$

STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 4 of 8 (Start time: 07:07:30 PM, 09 January 2024)

Formula which is used to multiply a matrix is

Select the correct option

☒ =MMULT(array1, array2)



☐ =MMULT{array+1array2}

☐ =MMULT{array*1array2}

☐ =MUL(array1, array2)

Question # 3 of 8 (start time: 07:06:44 PM, 09 January 2024)

A can only be used if the population standard deviation is known

Select the correct option

☐	T-test
☒	Z- test



STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 2 of 8 (**start time: 07:05:31 PM, 09 January 2024**)

Function to generate random numbers from normal distribution is

Select the correct option

☐ =NORM.INV(rand(), mean, standard deviation)

☐ =NORM.S.DIST(rand(), mean, variance)

☐ =NORM.INV(rand(), mean, median)

☐ =NORM.INV(rand(), standard deviation, mean)




Question # 1 of 8 (start time: 07:03:48 PM, 09 January

If the data is not normal, we will go for the

Select the correct option

☐	Parametric Procedures
☐	Rayleigh distribution
☐	Non-Parametric Procedures
☐	Chi square



STA641 - Statistical Packages and its Applications (Quiz 3)

Question # 4 of 8 (Start time: 04:09:15 PM, 09 January 2024)

A can only be used if the population standard deviation is known

► Select the correct option

T-test



Z- test

