STA301 - Statistics and Probability (Quiz#3) Quiz Start

Question # 1 of 10 (Start time: 01:04:22 PM, 23 July 2023)

In Binomial distribution the sample size is considered to be sufficiently large, if both np and nq are greater than or equal to

Select the correct option

- 10
- 5
- 15
- 3

Click to Save Answ

Question # 4 of 10 (Start time: 01:07:52 PM, 23 July 2023)

Which of the following is a major difference between the binomial and the hyper geometric distributions?

Select the correct option

- The sum of the outcomes can be greater than 1 for the hyper geometric.
- The number of trials changes in the hyper geometric distribution.
- The probability of a success changes from trial to trial in the hyper geometric distribution.
- The outcomes cannot be whole numbers in the hyper geometric distribution.

Click to Save Answe

STA301 - Statistics and Probability (Quiz#3)

Question # 5 of 10 (Start time: 01:40:10 PM, 23 July 2023)

In repeating sampling, which of the following possess desirable property of being Unbiased

Select the correct option

- All of above
- Sample proportion
- Sample median
- Sample Mean

Question # 2 of 10 (Start time: 01:21:45 PM, 23 July 2023) Total Marks: 1

"A point estimate plus/minus a few times the standard error of that estimate". This statement represents:

Select the correct option

- ○ Critical region
- ○ Critical value
- ○ Acceptance region
- ○ Confidence interval

Click to Save Answer & Move to Next Question

Question # 7 of 10 (Start time: 01:27:36 PM, 23 July 2023) **Total Marks:** 1

The Maximum Likelihood Estimators (MLE) are ..... and ..... but not necessarily .....

Select the correct option

- ○ Consistent, efficient, unbiased
- ○ Unbiased, efficient, consistent
- ○ Unbiased, consistent, efficient
- ○ Consistent, unbiased, efficient

Click to Save Answer & Move to Next Question

Question # 4 of 10 (Start time: 01:24:00 PM, 23 July 2023) Total Marks: 1

What will be the value of C.V if S.D = 0.5 and mean = 1.

Select the correct option

- 45%
- 50%
- 30%
- 10%

Click to Save Answer & Move to Next Question

STA301 - Statistics and Probability (Quiz#3)

Quiz Start Time:

Question # 9 of 10 (Start time: 01:46:11 PM, 23 July 2023)

The range of the binomial distribution is:

Select the correct option

- 0, 1, 2, ... n
- 1,2, ... n
- 0, 1, 2, ... 100
- 0, 1, 2, ... x

Click to Save Answer & Mov

1:12 PM 39%

Quiz
vulms.vu.edu.pk

λ is a positive _____ equal to the expected number of occurrences that occur during the given interval.

Select the correct option

- Complex number
- Transcendental number
- Irrational number
- Real number

Click to Save

1:12 PM 39%

Quiz
vulms.vu.edu.pk

λ is a positive _____ equal to the expected number of occurrences that occur during the given interval.

Select the correct option

- Complex number
- Transcendental number
- Irrational number
- Real number

Click to Save

vulms.vu.edu.pk

STA301 - Statistics and Probability (Quiz#3)

Question # 2 of 10 (Start time: 01:36:42 PM, 23 July 2023)

If Var(T1) >Var(T2) ,where T1 and T2 are two unbiased estimators ,then:

Select the correct option

- T2 is more consistent
- T2 is more efficient
- T1 is more consistent
- T1 is more efficient

What is the mean and variance of binomial distribution $b(x; n, p)$?

Select the correct option

○	np and pq
○	xp and npq
○	np and npq
○	Zero and one

Question # 9 of 10 (Start time: 01:30:01 PM, 23 July 2023) Total Marks: 1

Which of the following is NOT applicable to a Poisson distribution?

Select the correct option

- IF P = 0.5 & n = 19
- IF P = 0.03 & n = 500
- IF P = 0.01 & n = 200
- IF P = 0.02 & n = 300

Click to Save Answer & Move to Next Question

An estimator is always a:

Select the correct option

- Parameter
- Statistic and Random variable
- Statistic
- Random variable

Type here to search

Question # 9 of 10 (Start time: 01:30:01 PM, 23 July 2023) Total Marks: 1

Which of the following is NOT applicable to a Poisson distribution?

Select the correct option

- IF P = 0.5 & n = 19
- IF P = 0.03 & n = 500
- IF P = 0.01 & n = 200
- IF P = 0.02 & n = 300

Click to Save Answer & Move to Next Question

Gmail YouTube Maps YouTube WhatsApp Learning Managem... New Tab www.mgt603 quizz... https://chat.openai... Google Account

A randomly selected sample of 400 students at a university with 15-week semesters was asked whether or not they think the semester should be shortened to 14 weeks.46% of the 400 students surveyed answered "yes". Which one of the following statements about the number 46% is correct?

Select the correct option

- It is a sample statistic
- It is a population parameter
- It is a margin of error
- It is a standard error

Type here to search

Question # 10 of 10 (Start time: 01:32:00 PM, 23 July 2023) Total Marks: 1

According to emperical rule, how much data lies between $\mu - 3\sigma$ and $\mu + 3\sigma$?

Select the correct option

Reload Math Equations

- 99.73%
- 95.44%
- 50%
- 68.26%

Click to Save Answer & Move to Next Question

What are the numbers of parameter of Hyper geometric distribution ?

Select the correct option

- 2
- 5
- 3
- 4

1:10 PM 39%

Quiz
vulms.vu.edu.pk

BC210422860: AQSA

STA301 - Statistics and Probability (Quiz#3) Quiz

Question # 5 of 10 (Start time: 01:10:07 PM, 23 July 2023)

The limitation of point estimation is that:

Select the correct option

- There is no way of ascertaining how close the point estimate to parameter?
- All of these
- It is not calculated for small sample
- It is not calculated for non-normal distribution

Click to Save

Question # 6 of 10 (Start time: 01:26:05 PM, 23 July 2023) **Total Marks: 1**

The relative efficiency of T1 compared to T2 where:

Select the correct option

- ○ T2 is unbiased estimator
- ○ both T1 and T2 are unbiased estimators
- ○ both T1 and T2 are biased estimators
- ○ T1 is unbiased estimator

Click to Save Answer & Move to Next Question

STA301 - Statistics and Probability (Quiz#3) Quiz start Ti

Question # 2 of 10 (Start time: 01:05:08 PM, 23 July 2023)

Accept H0/H0 is false, is the _____

Select the correct option

- Type I error
- Type II error
- Correct decision
- None of the above

Click to Save Answer

STA301 - Statistics and Probability (Quiz#3)

Masters

Question # 1 of 10 (Start time: 11:18:11 AM, 23 July 2023)

Which type of data is used in the binomial distribution?

Select the correct option

- ○ Both discrete and continuous
- ○ Qualitative data only
- ○ Discrete data only
- ◉ Continues data only

correct

In the method of moments, how many equations are required for finding two unknown population parameters?

pg#257

Select the correct option

○	2 correct
○	3
○	4
○	1

STA301 - Statistics and Probability (Quiz#3)

Question # 4 of 10 (Start time: 11:22:23 AM, 23 July 2023)

pg#223

According to emperical rule, how much data lies between $\mu - 2\sigma$ and $\mu + 2\sigma$?

Select the correct option

- 50%
- 95.44% correct
- 99.73%
- 68.26%

Uniform distribution is a:

Select the correct option

◉	Continuous Distribution
○	Mixed
○	None of the above
○	Discrete distribution

correct

Normal approximation to the binomial distribution is used when:

Select the correct option

○	None of the above
○	np>5
○	Both of the above
○	nq>5

correct

Question # 2 of 10 (Start time: 11:19:46 AM, 23 July 2023) **Total Mark**

In hyper geometric distribution; N = number of units in the population, n = number of units in the sample and k = number of successes in the population, then Variance of hyper geometric distribution is defined as:

pg#214

Select the correct option

○	n.k/N ((N-k)/N)((N-n)/(N-1))
○	n.k/N ((N-k)/N)((N-n)/(N-k))
○	n.k/N ((N-k)/n)((N-n)/(N-1)) correct
○	n.k/N ((N-1)/N)((N-n)/(N-1))

If the second moment ratio is less than 3 the distribution will be:

pg#223

Select the correct option

○	Leptokurtic
○	Mesokurtic
○	None of these
○	Platykurtic

correct

In sampling from a large population with $\sigma = 60$. The size of the sample used is 100, the value of the standard error will be:

$$S = \frac{\sigma}{\sqrt{n}}$$

Select the correct option

○	6 correct
○	12
○	4
○	45

A random sample of size n = 25 is drawn from a normal population with population mean = 50 and sigma = 10 then mean of the sampling distribution of the sample mean is:

Select the correct option

- 45
- 25
- 50 correct
- 2

The rang of hypergeometric distribution is:

Select the correct option

- ○ 0 to k
- ○ 0 to N
- ○ 0 to n or k (whichever is less)
- ○ 0 to n

correct