paired t test is applicabl...

All Images Books News Videos Finan

اردو میں In English

A paired t-test is used to compare two population means where you have two samples in which **observations in one sample can be paired with observations in the other sample**.

https://www.statstutor.ac.uk › ... PDF

	P										
DF	0.995	0.975	0.2	0.1	0.05	0.025	0.02	0.01	0.005	0.002	0.001
1	.0004	.00016	1.642	2.706	3.841	5.024	5.412	6.635	7.879	9.55	10.828
2	0.01	0.0506	3.219	4.605	5.991	7.378	7.824	9.21	10.597	12.429	13.816
3	0.0717	0.216	4.642	6.251	7.815	9.348	9.837	11.345	12.838	14.796	16.266
4	0.207	0.484	5.989	7.779	9.488	11.143	11.668	13.277	14.86	16.924	18.467
5	0.412	0.831	7.289	9.236	11.07	12.833	13.388	15.086	16.75	18.907	20.515
6	0.676	1.237	8.558	10.645	12.592	14.449	15.033	16.812	18.548	20.791	22.458
7	0.989	1.69	9.803	12.017	14.067	16.013	16.622	18.475	20.278	22.601	24.322
8	1.344	2.18	11.03	13.362	15.507	17.535	18.168	20.09	21.955	24.352	26.124
9	1.735	2.7	12.242	14.684	16.919	19.023	19.679	21.666	23.589	26.056	27.877
10	2.156	3.247	13.442	15.987	18.307	20.483	21.161	23.209	25.188	27.722	29.588
11	2.603	3.816	14.631	17.275	19.675	21.92	22.618	24.725	26.757	29.354	31.264
12	3.074	4.404	15.812	18.549	21.026	23.337	24.054	26.217	28.3	30.957	32.909
13	3.565	5.009	16.985	19.812	22.362	24.736	25.472	27.688	29.819	32.535	34.528
14	4.075	5.629	18.151	21.064	23.685	26.119	26.873	29.141	31.319	34.091	36.123
15	4.601	6.262	19.311	22.307	24.996	27.488	28.259	30.578	32.801	35.628	37.697
16	5.142	6.908	20.465	23.542	26.296	28.845	29.633	32	34.267	37.146	39.252
17	5.697	7.564	21.615	24.769	27.587	30.191	30.995	33.409	35.718	38.648	40.79
18	6.265	8.231	22.76	25.989	28.869	31.526	32.346	34.805	37.156	40.136	42.312
19	6.844	8.907	23.9	27.204	30.144	32.852	33.687	36.191	38.582	41.61	43.82
20	7.434	9.591	25.038	28.412	31.41	34.17	35.02	37.566	39.997	43.072	45.315

What does β mean in statistics?

Beta (β) refers to **the probability of Type II error in a statistical hypothesis test**. Frequently, the power of a test, equal to 1−β rather than β itself, is referred to as a measure of quality for a hypothesis test.

What does β mean in statistics?

Beta (β) refers to **the probability of Type II error in a statistical hypothesis test**. Frequently, the power of a test, equal to 1−β rather than β itself, is referred to as a measure of quality for a hypothesis test.

Ufone ZONG 80% 10:52 AM

Your Results for: "Chapter 4: Multiple Choice Questions"

Site Title: Statistics in Psychology 1e
Book Title: Statistics in Psychology
Book Author: Steele et al
Location on Site: Student Resources > Chapter 4: Chi-square analysis > Chapter 4: Multiple Choice Questions
Date/Time Submitted: February 6, 2023 at 5:50 AM (UTC/GMT)

---

1. CORRECT By which other name is the chi-square goodness of fit test known?

Your Answer: One-sample chi-square

---

2. CORRECT What type of data do you need for a chi-square test?

Your Answer: Categorical

---

3. INCORRECT How many variables do you need to run a one-sample chi-square analysis?

Your Answer: At least 5
Correct Answer: You should only have 1

Incorrect! You should only have 1

---

4. CORRECT What is the purpose of a goodness-of-fit test?

Your Answer: To assesses whether the central tendency, variability and distribution of sam

---

5. CORRECT How many cases need to appear in one category for chi-square?

Your Answer: 5

---

6. INCORRECT How can you deal with low expected values?

Your Answer: (blank)

---

7. INCORRECT In which section of the SPSS output will you find the chi-square analysis?

Your Answer: (blank)

---

8. INCORRECT What does a significant result in a chi-square test imply?

4:16 PM
vulms.vu.edu.pk/
15

Question # 6 of 10 (Start time: 04:16:55 PM, 06 February 2023)

What does a significant result in a chi-square test imply?

Select the correct option

- ○ It implies that the sample is not representative of the population
- ○ That homogeneity of variance has not been established
- ○ That there is a significant difference between the three categorical variables included in the
- ○ That there is a significant positive relationship

Clic

The one-sample t-test is a statistical hypothesis test used to determine whether an unknown population mean is different from a specific value.

What other name is used for a contingency table?

Select the correct option

- A cross-classification table
- A histogram.
- None of the above
- An ANOVA table.

5:05 4G 46%

‘Mean of the Chi-...are distribution is’

Copy text Listen Translate Se



اردو میں In English

A chi-square distribution is a continuous probability distribution. The shape of a chi-square distribution depends on its degrees of freedom, k. The mean of a chi-square distribution is **equal to its degrees of freedom (k)** and the variance is 2k.

20-May-2022

https://www.scribbr.com › statistics

Chi-Square (X²) Distributions | Definition & Examples - Scribbr

About featured snippets Feedback

People also ask

Is the mean of a chi-square distribution 0?

Did you find these results useful? Yes No

43% 5:09 pm

vulms.vu.edu.pk/

BC200410038: ABDUL LATIF

Time Left 89 sec(s)

STAT406 - Statistical Inference Methods (Quiz 3)

Quiz Start Time: 05:00 PM, 06 February 2023

Question # 10 of 10 (Start time: 05:09:33 PM, 06 February 2023)

Total Marks: 1

The meaning of α is

Select the correct option

- All of the above
- Probability of Type I error
- Significant level
- Probability of rejecting the null hypothesis given its true

Click to Save Answer & Move to Next Question

5:35 4G 41%

with chi square test the effect of t

19-Feb-2020 — There are three different ways to calculate effect size for a Chi-Square test.

BMJ
https://www.bmj.com › publications

8. The Chi squared tests

The χ²tests The distribution of a categorical variable in a sample often needs to be compared with the distribution of a categorical variable in another ...

Investopedia
https://www.investopedia.com › terms

Chi-Square (x2) Statistic: What It Is, Examples, How and When to Use the ...

A chi-square (χ2) statistic is a test that is used to measure how expectations compare to actual observed data or model results.

What Is a Chi-square Test Used for?

Who Uses Chi-Square Analysis?

sphweb.bumc.bu.edu
https://sphweb.bumc.bu.edu › otlt

Hypothesis Testing - Chi Squared Test - SPH - Boston University

We find the critical value in a table of probabilities for the

Discover Snapshot Search Collections

5:35 4G 40%

Google H

range of coefficient of assosiatior

All Images Videos News Books Maps

Showing results for range of coefficient of *association* test

Search instead for range of coefficient of assosiation test

اردو میں In English

That said, a coefficient of association is independent of its measurement scale. These coefficients typically range **between 0 and 1**, where 0 is no relationship and 1 is a perfect relationship. However, some measures of association range from -1 to 1, where -1 indicates a perfect inverse relationship.

https://www.statisticshowto.com › c...

Coefficient of Association: Definition, Types, Examples - Statistics How To

About featured snippets Feedback

People also ask

Discover Snapshot Search Collections

5:37 4G 40%

Google H

what would b chi square p 0.05 si

All Images Videos News Books Maps

اردو میں In English

It is the Asymptotic Significance, or p- value, of the chi-square we've just run in SPSS. This value determines the statistical significance of the relationship we've just tested. In all tests of significance, if $p < 0.05$, we can say that **there is a statistically significant relationship between the two variables**.

https://cdn.southampton.ac.uk › ... PDF

Chi Square Test - University of Southampton

About featured snippets Feedback

People also ask

What does P 0.05 mean in chi-square?

What if p-value is greater than 0.05 in chi square test?

What does a P value of .05 represent?

Discover Snapshot Search Collections

5:48 4G 37%
wps.pearsoned.co.uk

CORRECT How many variables do you need to run a one-sample chi-square analysis?

Your Answer: You should only have 1

CORRECT What is the purpose of a goodness-of-fit test?

Your Answer: To assesses whether the central tendency, variability and distribution of sample is different fro

---

CORRECT How many cases need to appear in one category for chi-square?

Your Answer: 5

CORRECT How can you deal with low expected values?

Your Answer: You can increase your sample size or combine categories

---

CORRECT In which section of the SPSS output will you find the chi-square analysis?

Your Answer: Test statistics box

INCORRECT What does a significant result in a chi-square test imply?

Your Answer: That there is a significant difference between the three categorical variables included in the
Correct Answer: It implies that the sample is not representative of the population

Incorrect! It implies that the sample is not representative of the population

---

CORRECT What symbol is used to represent chi-square?

Your Answer: χ^2

CORRECT What would a chi-square significance value of P > 0.05 suggest?

Your Answer: That there is no significant difference between the sample and the population

---

CORRECT By which other name is a chi-square contingency table analysis known?

Your Answer: Chi-square test for independence

CORRECT Chi-square test for independence assesses which of the following?

Your Answer: It assesses whether there is a relationship between two categorical variables

---

INCORRECT Which tests could be used if your expected cases were fewer than 5?

Your Answer: Paired or independent t-tests

5:49 4G 37%

CORRECT In which section of the SPSS output will you find the chi-square analysis?

Your Answer: Test statistics box

INCORRECT What does a significant result in a chi-square test imply?

Your Answer: That there is a significant difference between the three categorical variables included in the a
Correct Answer: It implies that the sample is not representative of the population

Incorrect! It implies that the sample is not representative of the population

---

CORRECT What symbol is used to represent chi-square?

Your Answer: χ^2

CORRECT What would a chi-square significance value of P > 0.05 suggest?

Your Answer: That there is no significant difference between the sample and the population

---

CORRECT By which other name is a chi-square contingency table analysis known?

Your Answer: Chi-square test for independence

CORRECT Chi-square test for independence assesses which of the following?

Your Answer: It assesses whether there is a relationship between two categorical variables

---

INCORRECT Which tests could be used if your expected cases were fewer than 5?

Your Answer: Paired or independent t-tests
Correct Answer: Fisher's Exact Probability Test or Yates' Continuity Correction

Incorrect! You could use Fisher's Exact Probability Test or Yates' Continuity Correction

---

CORRECT Which row would you normally consult to find your chi-square results in a Chi-square test for independence al

Your Answer: Pearson chi-square

CORRECT What is an effect size?

Your Answer: The magnitude of the relationship between variables

---

E-mail Your Results

My name is (first last):		
E-mail my results to:		
	E-mail address:	**Send as:**
☐ Me		Text
☐ Instructor		Text
☐ TA		Text
☐ Other		Text

5:52 4G 36%

‘The meaning of a ...iven its true Click’

Copy text Listen Translate Se

Accepting a false null...

Answer · 0 votes

1. The option which best represents the correct answer ...
More

error B. level of sig

Answer · 0 votes

The power of a test
a "true positive" an
More

Feedback

People also ask

What is the probability of rejecting the null hypothesis called?

The probability of committing a type I error (rejecting the null hypothesis when it is actually true) is called **α (alpha)** the other name for this is the level of statistical significance.

https://www.ncbi.nlm.nih.gov › pmc

Hypothesis testing, type I and type II errors - PMC - NCBI

More results

What does it mean to reject the null hypothesis?

What is the significance of null hypothesis?

Did you find these results useful? Yes No

Ufone ZONG 80% 10:51 AM

Your Answer: 5

How can you deal with low expected values?

Your Answer: (blank)

In which section of the SPSS output will you find the chi-square analysis?

Your Answer: (blank)

What does a significant result in a chi-square test imply?

Your Answer: That there is a significant difference between the three categorical variables included in
Correct Answer: It implies that the sample is not representative of the population

Incorrect! It implies that the sample is not representative of the population

---

What symbol is used to represent chi-square?

Your Answer: χ^2

What would a chi-square significance value of P > 0.05 suggest?

Your Answer: That there is no significant difference between the sample and the population

---

By which other name is a chi-square contingency table analysis known?

Your Answer: One-sample chi-square
Correct Answer: Chi-square test for independence

Incorrect! This is a chi-square test for independence

Chi-square test for independence assesses which of the following?

Your Answer: It assesses whether there is a significant difference between two categorical variables
Correct Answer: It assesses whether there is a relationship between two categorical variables

Incorrect! It assesses whether there is a relationship between two categorical variables

---

Which tests could be used if your expected cases were fewer than 5?

DATE: _____ M T W T F S

Now, taking 'ln'

$$\ln e^{3t} = \ln \frac{8}{5}$$

$$3t = \ln \frac{8}{5}$$

$$t = \frac{1}{3} \ln \frac{8}{5}$$

$$\boxed{t = 0.157}$$ Ans

Now

$$d^2x = -\frac{5}{2} e^{-t} + \frac{8}{2} e^{-4t}$$

$$\Rightarrow -\frac{5}{3}e^{-t} + \frac{8}{3}e^{-4t} = 0$$

$$e^{-t}\left(-\frac{5}{3} + \frac{8}{3}e^{-3t}\right) = 0$$

$$-\frac{5}{3} + \frac{8}{3}e^{-3t} = 0$$

$$\frac{8}{3}e^{-3t} = \frac{5}{3}$$

$$e^{-3t} = \frac{5}{3} \times \frac{3}{8}$$

$$e^{-3t} = \frac{5}{8}$$

$$\boxed{e^{3t} = \frac{8}{5}}$$

$$\frac{d^2x}{dt^2}$$

Taking agai

$$\frac{d^2x}{dt^2} =$$

When $t = 0$

$$\frac{d^2x}{dt^2} = \frac{5}{3}$$

$$\boxed{\frac{d^2x}{dt^2} = \frac{5}{3}}$$

$$\frac{dt}{3} \qquad 3$$

$$x'(t) = 0 \Rightarrow -\frac{5}{3}e^{-t} + \frac{8}{3}e^{-4t} = 0$$

$$e^{3t} = \frac{8}{5} \Rightarrow t = \frac{1}{3}\ln\frac{8}{5}$$

$$t = 0.157$$

to two
motion if the w
3 ft / sec.

Solution

Since

Therefore, by Hoo

© Copyright Virtual University of Pakistan

8:35 3 K/s 72

Quiz
vulms.vu.edu.pk

chi-square distribution is _____skewed.

Select the correct option

- symmetrical.
- positively.
- none of these.
- negatively.

Click to Save Answer &

A matrix A is defined as

$$A = \begin{bmatrix} a & b \\ \\ c & d \end{bmatrix}, k \text{ is a constant then}$$

$$Ak = kA$$

$$Ak \neq kA$$

$$A = kA$$

$$-Ak = Ak$$

9:32 4G 74%
Done



Text editor
Edit image
Fill Form
Annotation

9:32 4G 74%
Done





Text editor
Edit image
Fill Form
Annotation

9:32 4G 73%
Done



Variance of the Chi-square distribution is:

Select the correct option

- 1
- n
- 2
- 2n

Text editor
Edit image
Fill Form
Annotation

9:32 4G 73%
Done



The chi-square distribution contains _____ parameters.

Select the correct option

- two.
- four.
- three.
- one.

Text editor
Edit image
Fill Form
Annotation

9:32 4G 73%
Done

Question # 5 of 10 (Start time: 02:05:30 PM, 06 February 2023)

In coefficient of association, when Q = 0, the attributes are:

Select the correct option

- none of these.
- Independent.
- dependent.
- vanished.

Text editor Edit image Fill Form Annotation

9:32 4G 73%
Done



Question # 6 of 10 (Start time: 02:06:10 PM, 06 February 2023)

The null hypothesis in the chi-square test states that

Select the correct option

- The rows and columns in the table are associated
- The rows and columns in the table are not associated ✓
- The means are equal
- The means are not equal

Text editor Edit image Fill Form Annotation

9:32 4G 73%
Done



Text editor
Edit image
Fill Form
Annotation

9:32 4G 73%
Done



Text editor
Edit image
Fill Form
Annotation

9:33 4G 73%
Done



Text editor
Edit image
Fill Form
Annotation

9:33 4G 73%
Done
4:16 PM
vulms.vu.edu.pk/
15

STAT406 - Statistical Inference Methods (Quiz 3)

Question # 7 of 10 (Start time: 04:17:34 PM, 06 February 2023)

With the Chi-square test, the effects of the test are calculated as

Select the correct option

- the average of the marginal totals
- n - K
- (r - 1) (c - 1)
- differences between observed and expected frequencies ✓

Text editor Edit image Fill Form Annotation


9:33
4G
73%
Done


4:16 PM
vulms.vu.edu.pk/
15
Question # 6 of 10 (Start time: 04:16:55 PM, 06 February 2023)
What does a significant result in a chi-square test imply?
Select the correct option
It implies that the sample is not representative of the population
That homogeneity of variance has not been established
That there is a significant difference between the three categorical variables included in the
That there is a significant positive relationship


4:15 PM
Text editor
Edit image
Fill Form
Annotation

9:33 4G 73%
Done



Text editor
Edit image
Fill Form
Annotation

9:33 4G 73%
Done

4:13 PM 16.2 K/s 72

STAT406 - Statistical Inference Methods (Quiz 3)

Question # 3 of 10 (Start time: 04:14:21 PM, 06 February 2023)

How many cases need to appear in one category for chi-square?

Select the correct option

- 6
- 3
- 2
- 5 ✓



Text editor Edit image Fill Form Annotation


9:33
4G
73%
Done


4:24
New tab
Question # 4 of 10 (Start time: 04:28:09 PM, 06 February 2023)
In analyzing the results of an experiment involving seven paired samples, tabulated t should be obtained for:
Select the correct option
12 degrees of freedom
6 degrees of freedom
13 degrees of freedom
14 degrees of freedom
Click to Save Answer


Text editor
Edit image
Fill Form
Annotation

9:33 4G 73%
Done

4G G 4:25 55
New tab

STAT406 - Statistical Inference Methods (Quiz 3) Quiz start Time: 04:24

Question # 5 of 10 (Start time: 04:29:14 PM, 06 February 2023)

The null hypothesis in the chi-square test states that

Select the correct option

- The means are equal
- The rows and columns in the table are associated
- The means are not equal
- The rows and columns in the table are not associated ✓

Click to Save Answer & Mov

Text editor Edit image Fill Form Annotation


9:34
73%
Done
4:26
New tab
The degrees of freedom for a t-test with sample size 11 is:
Select the correct option
11
8
10
9
Click to
Text editor
Edit image
Fill Form
Annotation


9:34
73%
Done
4:28
New tab
Question # 7 of 10 (Start time: 04:32:31 PM, 06 February 2023)
Which symbol is used to represent chi-square test?
Select the correct option
T
χ2
P
R
Click to Save Answer & Mov
Text editor
Edit image
Fill Form
Annotation

9:34 4G 73%
Done

Question # 2 of 10 (Start time: 04:29:10 PM, 06 February 2023)

In coefficient of association, when Q = 0, the attributes are:

Select the correct option

- none of these.
- Independent. ✓
- dependent.
- vanished.

Click to Sa

Text editor Edit image Fill Form Annotation

9:34 4G 73%
Done

Question # 3 of 10 (Start time: 04:29:40 PM, 06 February 2023)

The chi-square test is applied to:

Select the correct option

- Ratio Scale
- Interval scale
- Categorical Data ✓
- Ordinal scale

Click to Sc

Text editor Edit image Fill Form Annotation

9:34 4G 73%
Done

Question # 3 of 10 (Start time: 04:29:16 PM, 06 February 2023)

The meaning of β is

Select the correct option

- Probability of rejecting a null hypothesis when its true
- The probability of type II error
- Significance level
- All of the above

Text editor Edit image Fill Form Annotation

9:34 4G 73%
Done

Question # 4 of 10 (Start time: 04:30:06 PM, 06 February 2023)

Chi square distribution tends to normal distribution as the number of degrees freedom approaches to:

Select the correct option

- 0
- $-\infty$
- none of these.
- ∞ ✓

Click to Save Answer

Text editor Edit image Fill Form Annotation

9:34 4G 73%
Done

Question # 5 of 10 (Start time: 04:31:12 PM, 06 February 2023)

If we are considering one attribute such as A, then number of class frequencies:

Select the correct option

- 4
- 2
- 1
- 3 ✓

Click to Save Answer

Text editor Edit image Fill Form Annotation

9:35 4G 73%
Done

For testing of independence, parameters used in chi-square distribution is observed value and:

Select the correct option

- Expected value ✓
- Estimated value
- Mean value
- Error value

Click to Save Answer

Text editor Edit image Fill Form Annotation

9:35 4G 73%
Done

Question # 8 of 10 (Start time: 04:32:37 PM, 06 February 2023)

chi-square distribution is _____skewed.

Select the correct option

- none of these.
- positively.
- symmetrical.
- negatively.

Click to Save Answer &

Text editor Edit image Fill Form Annotation

9:35 4G 73%
Done

Question # 9 of 10 (Start time: 04:33:08 PM, 06 February 2023)

As the degrees of freedom increase (and especially when the degrees of freedom are more than 90), the graph of th looks more and more _____.

Select the correct option

- skewed left
- asymmetrical
- Symmetrical
- skewed right ✓

Click to Save Answer

Question # 8 of 10 (Start time: 04:33:07 PM, 06 February 2023)

Text editor Edit image Fill Form Annotation


9:35
4G
73%
Done


4:29
54
New tab
Question # 9 of 10 (Start time: 04:33:53 PM, 06 February 2023)
The degrees of freedom for a t-test with sample size 10 is:
Select the correct option
7
9
10
8
Click to Save Ans


Text editor
Edit image
Fill Form
Annotation


9:35
4G 73%
Done
4:30
New tab
Question # 10 of 10 (Start time: 04:34:41 PM, 06 February 2023)
Which of the following values cannot occur in a chi-square distribution?
Select the correct option
0.62
100
384
-2.45
Click to Save Answer & Mov
Text editor
Edit image
Fill Form
Annotation

9:35 4G 73%
Done

Question # 4 of 10 (Start time: 04:36:22 PM, 06 February 2023)

A single-sample t ratio is considered statistically significant only when

Select the correct option

- the null hypothesis is rejected. ✓
- there is no Type I error.
- the standard error is quite large.
- the population values are known.

Click to Save Answer

Text editor Edit image Fill Form Annotation

9:35 4G 73%
Done

Question # 6 of 10 (Start time: 04:36:58 PM, 06 February 2023)

The value of χ^2 with 1 degree of freedom is equal to:

Select the correct option

- 3.84 ✓
- 5.33
- 4.91
- 2.33

Click to Save Answer &

Text editor Edit image Fill Form Annotation

9:35 4G 73%
Done

Question # 1 of 10 (Start time: 04:34:41 PM, 06 February 2023)

The degrees of freedom for a t-test with sample size 'n' is:

Select the correct option

- n+1
- n-1
- n-2
- n-3

Click to Save Answer & M

Text editor Edit image Fill Form Annotation



Question # 10 of 10 (Start time: 04:33:35 PM, 06 February 2023)

While applying chi-square test to a contingency table of 4 rows and 4 columns, the degrees of freedom would be:

Select the correct option

- 1
- 4
- 8
- 9 (●)

Click to Save Answer & Move to



9:36 4G 73%
Done

Mean of the Chi-square distribution is:

Select the correct option

- 2n
- 1
- n
- 2

Text editor
Edit image
Fill Form
Annotation

9:36 4G 73%
Done

In analyzing the results of an experiment involving seven paired samples, tabulated t should be obtained for:

Select the correct option

- 13 degrees of freedom
- 14 degrees of freedom
- 12 degrees of freedom
- 6 degrees of freedom



Text editor
Edit image
Fill Form
Annotation

9:36 4G 73%
Done

The purpose of a goodness-of-fit test is to:

Select the correct option

- To assesses whether the central tendency, variability and distribution of sample is different from that of the population
- To extract factors from the data
- To identify significant effects
- To assess whether several categorical variables are related

Text editor
Edit image
Fill Form
Annotation

9:36 4G 73%
Done



Text editor
Edit image
Fill Form
Annotation

9:36 4G 73%
Done

If we are considering one attribute such as A, then number of class frequencies:

Select the correct option

- 2
- 1
- 3
- 4

Text editor
Edit image
Fill Form
Annotation

9:37 4G 73%
Done

Range of coefficient of association lies between:

Select the correct option

- -1 to +1
- 0 to 1
- -1 to 0
- 0 to 2

62

Text editor
Edit image
Fill Form
Annotation

9:37 4G 73%
Done

For a test of the relationship between two variables using chi-square, the null hypothesis is stated as

Select the correct option

- Chi-square is greater than zero
- Chi-square is less than zero
- Chi-square is equal to zero ✓
- Chi-square is equal to 1

64

Text editor Edit image Fill Form Annotation

9:37 4G 73%
Done

By which other name is the chi-square goodness of fit test known?

Select the correct option

- Chi-square Anova
- None of these
- One-sample chi-square
- Directional chi-square

Text editor
Edit image
Fill Form
Annotation

9:37 4G 73%
Done



Text editor
Edit image
Fill Form
Annotation

9:37 4G 73%
Done





Text editor
Edit image
Fill Form
Annotation


9:37
72%
Done
5:13
66%
Quiz
vulms.vu.edu.pk
Question # 4 of 10 (Start time: 05:13:48 PM, 06 February 2023)
How can you deal with low expected values?
Select the correct option
You can add more of the same number
You can transform your data
You can increase your sample size or combine categories
You can exclude outliers
Click to Save Answer & Move to Next
Text editor
Edit image
Fill Form
Annotation

9:37 4G 72%
Done



Text editor
Edit image
Fill Form
Annotation


9:38
4G 72%
Done
5:16
66%
Quiz
vulms.vu.edu.pk
What would a chi-square significance value of P > 0.05 suggest?
Select the correct option
That there is no significant difference between the sample and the population
That there is a significant relationship between the sample and the population
That there is a significant relationship between the sample and the population
That there is no significant difference between time 1 and time 2
Click to Save Answer & Move
Text editor
Edit image
Fill Form
Annotation

9:38 4G 72%
Done



Text editor
Edit image
Fill Form
Annotation

9:38 4G 72%
Done



Text editor
Edit image
Fill Form
Annotation

9:38 4G 72%
Done
Quiz
vulms.vu.edu.pk

Question # 1 of 10 (Start time: 05:36:59 PM, 06 February 2023)

chi-square distribution is _____skewed.

Select the correct option

- ○ symmetrical.
- ○ none of these.
- ○ negatively.
- ○ positively. ✓

Text editor
Edit image
Fill Form
Annotation

9:38 4G 72%
Done



5:38
Text editor
Edit image
Fill Form
Annotation

9:38 4G 72%
Done
Quiz
vulms.vu.edu.pk

BC180406297: MUHAMMAD HASEEB

STAT406 - Statistical Inference Methods (Quiz 3)

Question # 2 of 10 (Start time: 05:38:00 PM, 06 February 2023)

Paired t-test is applicable when the observations in the two samples are:

Select the correct option

- ○ All of the above
- ○ Paired ✓
- ○ Equal in number
- ○ Correlation

Text editor Edit image Fill Form Annotation

9:39 4G 72%
Done

STAT406 - Statistical Inference Methods (Quiz 3) Quiz Start Time: 06:

Question # 2 of 10 (Start time: 06:40:47 PM, 06 February 2023)

With the Chi-square test, the effects of the test are calculated as

Select the correct option

- ○ the average of the marginal totals
- ○ n - K
- ○ differences between observed and expected frequencies ✓
- ○ (r - 1) (c - 1)

Text editor Edit image Fill Form Annotation


9:39
4G 72%
Done
BC200408292: FOUZIA RIAZ
STAT406 - Statistical Inference Methods (Quiz 3)
Quiz Start Time: 06:
Question # 1 of 10 (Start time: 06:39:45 PM, 06 February 2023)
How can you deal with low expected values?
Select the correct option
You can add more of the same number
You can increase your sample size or combine categories
You can exclude outliers
You can transform your data
Click to Save Answer & M


6:38
VoLTE
90
vulms.vu.edu.pk/
BC200410084: ASRA AMIN
Time Left
86
sec(s)
PM, 06 February 2023
Total Marks:
Text editor
Edit image
Fill Form
Annotation