File by MAM MEHWISH 03184148783
Paid LMS Handling Available

We also provide Past papers for Mids. and Finals

Cross Check your answers before submitting directly from the file. We are not responsible for 100% correction as human error exists

Question 1:

In testing H0: μ = 100 against H1: μ ≤ 100 at the 10% level of significance, H0 is rejected if:

Options:

A. The p-value is greater than 0.10

B. 100 is contained in the 90% confidence interval

C. The p-value is less than 0.10

D. The value of the test statistic is in the acceptance region

Correct Option: C. The p-value is less than 0.10

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

Question 2:

The variance of the Hypergeometric distribution is:

Options:

A. pq{nk(N-k)(N-n)}/N*N(N-1)

B. npq{nk(N-n)}/N(N-1)

Correct Option: A. pq{nk(N-k)(N-n)}/N*N(N-1)

Question 3:

Alpha is the probability of:

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

Options:

A. Type I error

B. All of these

C. Committing Type I error

D. Rejecting H0 when H0 is true

Correct Option: A. Type I error

Question 4:

Which of the following is a major difference between the binomial and the hypergeometric distributions?

Options:

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

A. The outcomes cannot be whole numbers in the hypergeometric distribution.

B. The probability of a success changes from trial to trial in the hypergeometric distribution.

C. The number of trials changes in the hypergeometric distribution.

D. The sum of the outcomes can be greater than 1 for the hypergeometric.

Correct Option: B. The probability of a success changes from trial to trial in the hypergeometric distribution.

Question 5:

If the first moment ratio is equal to 0, then the distribution will be:

Options:

A. Symmetrical

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

B. Negatively Skewed

C. Positively Skewed

D. Leptokurtic

Correct Option: A. Symmetrical

Question 6:

If p is very small and n is considerably large then we shall apply the:

Options:

A. Binomial distribution

B. Exponential distribution

C. Poisson distribution

D. Hypergeometric distribution

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

Correct Option: C. Poisson distribution

Question 7:

In a normal distribution, how much area lies between $\mu \pm 2\sigma$?

Options:

A. 75%

B. 95.45%

C. 94%

D. 85%

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

Correct Option: B. 95.45%

Question 8:

A standard deviation obtained from the sampling distribution of sample statistics is known as:

Options:

A. Standard error

B. Universal error

C. Sampling error

D. Minimum error

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

Correct Option: A. Standard error

Question 9:

We can apply the method of Maximum Likelihood on:

Options:

A. Discrete as well as continuous variables

B. Continuous variables only

C. Qualitative variables only

D. Discrete variables only

Correct Option: A. Discrete as well as continuous variables

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

Question 10:

In a binomial experiment, the probability of success changes on:

Options:

A. 10th trial

B. Second trial

C. Each trial

D. 30th trial

Correct Option: C. Each trial

File by MAM MEHWISH 03184148783
Paid LMS Handling Available

File by MAM MEHWISH 03184148783
Paid LMS Handling Available