For More Visit VU Answer

MTH301 Final Term Papers By Waqar (File 1)

Question No : 1 of 52 Marks: 1 (Budgeted Time 1 Min)

Which one of the following is correct Wallis Sine formula when n is odd and $n \geq 3$?

Answer (Please select your correct option)

VuAnswers.com

- $\int_0^{\frac{\pi}{2}} \sin^n x\,dx = \frac{\pi}{2}\frac{(n-1)}{n}\frac{(n-3)}{(n-2)}\frac{(n-5)}{(n-4)}\cdots\cdots\frac{5}{6}\frac{3}{4}\frac{1}{2}$
- $\int_0^{\frac{\pi}{2}} \sin^n x\,dx = \frac{\pi}{2}\frac{(n)}{(n-1)}\frac{(n-2)}{(n-3)}\frac{(n-4)}{(n-5)}\cdots\cdots\frac{6}{5}\frac{4}{3}\frac{2}{1}$
- $\int_0^{\frac{\pi}{2}} \sin^n x\,dx = \frac{(n-1)}{n}\frac{(n-3)}{(n-2)}\frac{(n-5)}{(n-4)}\cdots\cdots\frac{6}{7}\frac{4}{5}\frac{2}{3}$ correct
- $\int_0^{\frac{\pi}{2}} \sin^n x\,dx = \frac{(n)}{(n-1)}\frac{(n-2)}{(n-3)}\frac{(n-4)}{(n-5)}\cdots\cdots\frac{6}{5}\frac{4}{3}\frac{2}{1}$

Made by: Waqar Siddhu

Question No : 2 of 52 Marks: 1 (Budgeted Time 1 Min)

What is the period of a periodic function defined by $f(x) = 4\cos 3x$?

Answer (Please select your correct option)

VuAnswers.com

- $\frac{\pi}{4}$
- $\frac{\pi}{3}$
- $\frac{2\pi}{3}$ correct
- π

Made by: Waqar Siddhu





Question No : 3 of 52 Marks: 1 (Budgeted Time 1 Min)

Match the following periodic function with its graph.

$$f(x) = \begin{cases} 2-x & 0 < x < 3 \\ -1 & 3 < x < 5 \end{cases}$$

Answer (Please select your correct option)

VuAnswers.com

-  correct
- 
- 
- 


Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 4 of 52 Marks: 1 (Budgeted Time 1 Min)

What is the period of periodic function whose graph is as below?



Answer (Please select your correct option)

VuAnswers.com

- 2
- 5
- 6


correct

- 8

Made by: Waqar Siddhu

Question No : 5 of 52 Marks: 1 (Budgeted Time 1 Min)

What is the period of periodic function whose graph is as below?



Answer (Please select your correct option)

VuAnswers.com

- 0
- 4
- 6
- 8

correct

Made by: Waqar Siddhu

Question No : 6 of 52 Marks: 1 (Budgeted Time 1 Min)

Which of the following condition must be satisfied for a vector field $\vec{F}$ to be a conservative vector field?

Answer (Please select your correct option)

VuAnswers.com

- Line integral of $\vec{F}$ along a curve, depends only on the endpoints of that curve, not on the particular route taken.
- Divergence of $\vec{F}$ should be zero
- Gradient of $\vec{F}$ should be zero.
- $\vec{F} = 0$

correct

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 7 of 52 Marks: 1 (Budgeted Time 1 Min)

What is the period of periodic function whose graph is as below?



Answer (Please select your correct option)

VuAnswers.com

- π
- $-\pi$ correct
- 2π
- -2π

Made by: Waqar Siddhu

Question No : 8 of 52 Marks: 1 (Budgeted Time 1 Min)

Let L denotes the Laplace Transform.
According to First Shift Theorem, if $L(F(t)) = f(s)$ then which of the following equation holds?
s *and* a are constants.

Answer (Please select your correct option)

VuAnswers.com

- $L(e^{-at}F(t)) = f(s+a)$
- $L(e^{-at}F(t)) = f(s)$
- $L(e^{-at}F(t)) = f(a)$
- $L(e^{-at}F(t)) = f(s-a)$

Made by: Waqar Siddhu

Question No : 9 of 52 Marks: 1 (Budgeted Time 1 Min)

The function $f(x) = x^2 \cos 2x$ is ----------

Answer (Please select your correct option)

VuAnswers.com

- Neither even nor odd
- Odd function
- Even function correct

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 10 of 52 Marks: 1 (Budgeted Time 1 Min)

The graph of an even function is symmetrical about ----------

Answer (Please select your correct option)

VuAnswers.com

- x-axis
- y-axis

correct

- origin

Made by: Waqar Siddhu

Question No : 11 of 52 Marks: 1 (Budgeted Time 1 Min)

Which of the following is Laplace inverse transform of the function $f(s)$ defined by $f(s) = \frac{3}{s-2} - \frac{2}{s}$?

Answer (Please select your correct option)

VuAnswers.com

- $3te^{2t} - 2$
- $3e^{2t} - 2t$
- $3e^{2t} - 2$

correct

- None of these.

Made by: Waqar Siddhu

Question No : 12 of 52 Marks: 1 (Budgeted Time 1 Min)

What is laplace transform of the function $F(t)$ if $F(t) = \sin 3t$?

Answer (Please select your correct option)

VuAnswers.com

- $L(\sin 3t) = \frac{3}{s^2+9}$

correct

- $L(\sin 3t) = \frac{s}{s^2+9}$
- $L(\sin 3t) = \frac{1}{s-3}$
- $L(\sin 3t) = \frac{3!}{s^4}$

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 13 of 52 Marks: 1 (Budgeted Time 1 Min)

If L denotes laplace transform then
$L(te^{5t}) =$

Answer (Please select your correct option)

VuAnswers.com

- $L(te^{5t}) = \frac{1}{s^2-5}$
- $L(te^{5t}) = \frac{1}{s^2+5}$
- $L(te^{5t}) = \frac{1}{(s+5)^2}$
- $L(te^{5t}) = \frac{1}{(s-5)^2}$ correct

Made by: Waqar Siddhu

Question No : 14 of 52 Marks: 1 (Budgeted Time 1 Min)

What is $L(-6)$ if L denotes Laplace Transform?

Answer (Please select your correct option)

VuAnswers.com

- $L(-6) = \frac{1}{s+6}$
- $L(-6) = \frac{-6}{s}$ correct
- $L(-6) = \frac{s}{s^2+36}$
- $L(-6) = \frac{-6}{s^2+36}$

Made by: Waqar Siddhu

Question No : 15 of 52 Marks: 1 (Budgeted Time 1 Min)

A vector field is a vector each of whose components is a scalar field

Answer (Please select your correct option)

VuAnswers.com


- True correct
- False

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 16 of 52 Marks: 1 (Budgeted Time 1 Min)




Answer (Please select your correct option)

VuAnswers.com

- An odd function
- An even function
- Neither even nor odd

Made by: Waqar Siddhu

Question No : 16 of 52 Marks: 1 (Budgeted Time 1 Min)



Answer (Please select your correct option)

VuAnswers.com

- An odd function
- An even function
- Neither even nor odd

Made by: Waqar Siddhu

Question No : 16 of 52 Marks: 1 (Budgeted Time 1 Min)



The graph of "saw tooth wave" given above is ----------

Answer (Please select your correct option)

VuAnswers.com

- An odd function
- An even function
- Neither even nor odd

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 17 of 52 Marks: 1 (Budgeted Time 1 Min)

All axes are positive in ---------- octant.

Answer (Please select your correct option)

VuAnswers.com

- First — correct
- Second
- Fourth
- Eighth

Made by: Waqar Siddhu

Question No : 18 of 52 Marks: 1 (Budgeted Time 1 Min)

The angles which a line makes with positive x ,y and z-axes are known as ----------

Answer (Please select your correct option)

VuAnswers.com

- Direction cosines
- Direction ratios
- Direction angles — correct

Made by: Waqar Siddhu

Question No : 19 of 52 Marks: 1 (Budgeted Time 1 Min)

If α, β *and* γ are the direction angles of a line then
$\cos^2\alpha+\cos^2\beta+\cos^2\gamma=$

Answer (Please select your correct option)

VuAnswers.com

- 0
- 1 — correct
- 2
- T

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 20 of 52 Marks: 1 (Budgeted Time 1 Min)

Intersection of two non parallel planes is a . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- Plane
- Point
- Straight line

correct

- None of these.

Made by: Waqar Siddhu

Question No : 21 of 52 Marks: 1 (Budgeted Time 1 Min)

$\lim_{(x,y)\to(0,1)} \frac{xy^2}{x^2+y^2} =$

Answer (Please select your correct option)

VuAnswers.com

- ∞
- 0

correct

- 1
- 0.5

Made by: Waqar Siddhu

Question No : 22 of 52 Marks: 1 (Budgeted Time 1 Min)

The function $z = \frac{1}{\sqrt{x+y}}$ is discontinuous at origin because at the point (0,0), it . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- has a jump
- has a hole
- approaches towards infinity
- approaches towards zero

correct

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 23 of 52 Marks: 1 (Budgeted Time 1 Min)

Gradient of a scalar function always results in a . . . . . function.

Answer (Please select your correct option)

VuAnswers.com

- Scalar
- Continuous
- Vector

correct

- Constant

Made by: Waqar Siddhu

Question No : 24 of 52 Marks: 1 (Budgeted Time 1 Min)

At the critical point (x_0, y_0), if $D = f_{xx}(x_0,y_0) f_{yy}(x_0,y_0) - f_{xy}^2(x_0,y_0) < 0$ then f has a . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- Relative maximum at (x_0, y_0)
- Relative minimum at (x_0, y_0)

correct

- Saddle point at (x_0, y_0)
- No conclusion can be drawn

Made by: Waqar Siddhu

Question No : 25 of 52 Marks: 1 (Budgeted Time 1 Min)

The volume of parallelepiped with dimensions x, y, z is . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- $V = x^2y^2z^2$
- $V = x + y + z$
- $V = \sqrt{xyz}$
- $V = xyz$

correct

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 26 of 52 Marks: 1 (Budgeted Time 1 Min)

If $R = \{(x,y) : 0 \le x \le 2 \text{ and } 1 \le y \le 4\}$, then $\iint_R (6x^2 + 4xy^3)dA =$ . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- $\int_1^4 \int_0^2 (6x^2 + 4xy^3)dydx$
- $\int_0^2 \int_1^4 (6x^2 + 4xy^3)dxdy$
- $\int_1^4 \int_0^2 (6x^2 + 4xy^3)dxdy$ correct
- $\int_2^4 \int_0^1 (6x^2 + 4xy^3)dxdy$

Made by: Waqar Siddhu

Question No : 27 of 52 Marks: 1 (Budgeted Time 1 Min)

Polar co-ordinates of a point are $\left(7, \frac{-\pi}{4}\right)$. Which of the following is another possible polar co-ordinates representation of this point?

Answer (Please select your correct option)

VuAnswers.com

- $\left(7, \frac{3\pi}{4}\right)$ correct
- $\left(-7, \frac{3\pi}{4}\right)$
- $\left(-7, \frac{-\pi}{4}\right)$
- $\left(7, \frac{-3\pi}{4}\right)$

Made by: Waqar Siddhu

Question No : 28 of 52 Marks: 1 (Budgeted Time 1 Min)

Polar co-ordinates of a point are $\left(-2, \frac{-3\pi}{2}\right)$. Which of the following is another possible polar co-ordinates representation of this point?

Answer (Please select your correct option)

VuAnswers.com

- $\left(-2, \frac{\pi}{2}\right)$ correct
- $\left(-2, \frac{\pi}{4}\right)$
- $\left(-2, \frac{3\pi}{4}\right)$
- $\left(-2, \frac{\pi}{3}\right)$

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 29 of 52 Marks: 1 (Budgeted Time 1 Min)

If the equation of a curve, in polar co-ordinates, remains unchanged after replacing (r, θ) by $(r, \pi - \theta)$ then the curve is said to be symmetric about . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- y-axis
- Pole
- Initial line
- origin

correct

Made by: Waqar Siddhu

Question No : 30 of 52 Marks: 1 (Budgeted Time 1 Min)

In polar coordinate system, x- axis is also called . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- Polar axis
- Pole
- Imaginary axis
- None of these

correct

Made by: Waqar Siddhu

Question No : 31 of 52 Marks: 1 (Budgeted Time 1 Min)

The point $p(0, \theta)$ in polar coordinate system lies on . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- Polar axis
- y-axis
- Pole
- None of these

correct

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 32 of 52 Marks: 1 (Budgeted Time 1 Min)

Given the integral $\iint_R f(x,y)dxdy$, after converting to polar coordinates, it will become . . . . . where $a \le \theta \le b$ and $c \le r \le d$.

Answer (Please select your correct option)

VuAnswers.com

- $\int_a^b \int_c^d f(r,\theta) r dr d\theta$
- $\int_a^b \int_c^d f(r,\theta) dr d\theta$ correct
- $\int_a^b \int_c^d f(r,\theta) r d\theta dr$
- $\int_a^b \int_c^d f(r,\theta) d\theta dr$

Made by: Waqar Siddhu

Question No : 33 of 52 Marks: 1 (Budgeted Time 1 Min)

The parametric equations that correspond to the vector equation $\vec{r}(t) = \sin^2 t\,\hat{i} + (1-\cos 2t)\,\hat{j}$ are . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- $x = \sin^2 t \;,\quad y = 1-\cos 2t \;,\; z = 0$ correct
- $y = \sin^2 t \;,\quad x = 1-\cos 2t \;,\; z = 0$
- $x = \sin^2 t \;,\quad y = 1-\cos 2t \;,\; z = 1$
- $x = \sin^2 t \;,\quad y = \cos 2t \;,\; z = 1$

Made by: Waqar Siddhu

Question No : 34 of 52 Marks: 1 (Budgeted Time 1 Min)

If $\vec{R}(t)$ is the anti-derivative of a given vector valued function $\vec{r}(t)$ i.e., $\vec{R}'(t) = \vec{r}(t)$ then $\int_a^b \vec{r}(t)dt =$ . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- $\vec{R}(a) + \vec{R}(b)$
- $\vec{R}(b) - \vec{R}(a) + c$ where c is a non-zero constant.
- $\vec{R}(a).\vec{R}(b)$
- $\vec{R}(b) - \vec{R}(a)$

Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 35 of 52 Marks: 1 (Budgeted Time 1 Min)

For the given vector valued functions $\vec{r}_1(t)$ and $\vec{r}_2(t)$, $\frac{d}{dt}[\vec{r}_1(t)\times\vec{r}_2(t)]$ = ... where "×" and "•" represent the cross and dot product respectively.

Answer (Please select your correct option)

VuAnswers.com

- $\vec{r}_1 \cdot \frac{d\vec{r}_2}{dt} + \frac{d\vec{r}_1}{dt} \cdot \vec{r}_2$
- $\vec{r}_1 \times \frac{d\vec{r}_2}{dt} + \frac{d\vec{r}_1}{dt} \times \vec{r}_2$
- $\frac{d\vec{r}_2}{dt} \times \vec{r}_1 + \vec{r}_2 \times \frac{d\vec{r}_1}{dt}$
- $\frac{d\vec{r}_2}{dt} \times \vec{r}_1 + \frac{d\vec{r}_1}{dt} \times \vec{r}_2$

Made by: Waqar Siddhu

Question No : 36 of 52 Marks: 1 (Budgeted Time 1 Min)

A vector valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ is continuous if . . . . . .

Answer (Please select your correct option)

VuAnswers.com

- Atleast one of its components is continuous.
- All of its components are necessarily differentiable.

correct

- All of its components are continuous.
- Limit exists for all of its components.

Made by: Waqar Siddhu

Question No : 37 of 52 Marks: 1 (Budgeted Time 1 Min)

Given a vector valued function $\vec{r}(t) = \frac{1}{(t-3)}\hat{i} + e^t\hat{j}$ and its anti-derivative $\vec{R}(t) = \ln(t-3)\hat{i} + e^t\hat{j}$, then $\int \vec{r}(t)dt$ = ...

Answer (Please select your correct option)

VuAnswers.com

- $\ln(t-3)\hat{i} + e^t\hat{j} + c$
- $(t-3)\hat{i} + \frac{e^t}{2}\hat{j} + c$
- $(t-3)^{-1}\hat{i} + \frac{e^t}{2}\hat{j} + c$
- $\frac{1}{(t-3)}\hat{i} + e^t\hat{j}$

correc Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer

Question No : 38 of 52 Marks: 1 (Budgeted Time 1 Min)

For any two vector valued functions $\vec{r}_1(t)$ and $\vec{r}_2(t)$, $\frac{d}{dt}[\vec{r}_1(t).\vec{r}_2(t)] =$ ..... where "×" and "." represent the cross and dot product respectively.

Answer (Please select your correct option)

VuAnswers.com

- $\vec{r}_1 . \frac{d\vec{r}_2}{dt} - \frac{d\vec{r}_1}{dt} . \vec{r}_2$
- $\vec{r}_1 . \frac{d\vec{r}_2}{dt} + \frac{d\vec{r}_1}{dt} . \vec{r}_2$
- $\vec{r}_1 \times \frac{d\vec{r}_2}{dt} + \frac{d\vec{r}_1}{dt} \times \vec{r}_2$
- $\vec{r}_1 . \frac{d\vec{r}_2}{dt} - \frac{d\vec{r}_1}{dt} \times \vec{r}_2$ correct

Made by: Waqar Siddhu

Question No : 39 of 52 Marks: 1 (Budgeted Time 1 Min)

A single curve can be represented by ..... vector valued function(s).

Answer (Please select your correct option)

VuAnswers.com

- Two
- Infinitely many
- Single correct
- Three

Made by: Waqar Siddhu

Question No : 40 of 52 Marks: 1 (Budgeted Time 1 Min)

A function is said to be smooth if it's derivative is ..... on any value of its domain.

Answer (Please select your correct option)

VuAnswers.com

- continuous and non zero
- piecewise continuous
- defined and non zero
- differentiable correct

Made by: Waqar Siddhu

For More Visit VU Answer


For More Visit VU Answer

Question No : 41 of 52 Marks: 2 (Budgeted Time 4 Min)

Use Wallis sine formula to evaluate $\int_0^{\frac{\pi}{2}} \sin^5 x \, dx$

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu


Question No : 42 of 52 Marks: 2 (Budgeted Time 4 Min)

Prove whether the following function is even, odd or neither.

$f(x) = x^3 + x^2$

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 43 of 52 Marks: 2 (Budgeted Time 4 Min)

Consider the point $\left(\frac{\sqrt{3}}{2}, \sqrt{3}, \frac{1}{2}\right)$ in rectangular coordinate system. Find the value of "ρ" in spherical coordinates.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

For More Visit VU Answer


For More Visit VU Answer

Question No : 44 of 52 Marks: 2 (Budgeted Time 4 Min)

Write down the expression for the arc- length of the curve represented by the vector valued function $\vec{r}(t) = \cos t^2 \hat{i} + \sin t^2 \hat{j}$ where $0 \le t \le \frac{\pi}{2}$. (Do not evaluate the expression).

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 45 of 52 Marks: 3 (Budgeted Time 6 Min)

Use Wallis cosine formula to evaluate $\int_0^{\frac{\pi}{2}} \left(\cos^3 x + \cos^4 x\right) dx$

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu


Question No : 46 of 52 Marks: 3 (Budgeted Time 6 Min)

Prove whether the following function is even, odd or neither.

$f(x) = \sin^2 x \cos 3x$

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

For More Visit VU Answer


For More Visit VU Answer

Question No : 47 of 52 Marks: 3 (Budgeted Time 6 Min)

Find the region where the function $f(x, y) = \sqrt{9 - x^2 - y^2}$ is continuous.

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 48 of 52 Marks: 3 (Budgeted Time 6 Min)

What is the arc-length of the curve $\vec{r}(t) = (4 + 3t)\,\hat{i} + (2 - 2t)\,\hat{j}$ when $3 \le t \le 4$?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu


Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

Determine the Fourier co-efficient b_n of the periodic function defined below:

$f(x) = 2x + 1 \qquad 0 < x < 2$

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

For More Visit VU Answer

For More Visit VU Answer


Question No : 50 of 52 Marks: 5 (Budgeted Time 10 Min)

Determine whether the following vector field $\vec{F}$ is conservative or not.

$\vec{F}(x, y, z) = (3x + y)\hat{i} + xy^2z\,\hat{j} + xz^2\hat{k}$

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 51 of 52 Marks: 5 (Budgeted Time 10 Min)

Using second partial derivative test show that the function $f(x, y) = xy(24 - x - y)$ is maximum at $x = 8, y = 8$.

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 52 of 52 Marks: 5 (Budgeted Time 10 Min)

Let $\vec{r}(t) = t^2\,\hat{i} + t\,\hat{j} + (t^2 - 5)\,\hat{k}$. Find t, such that $\vec{r}(t)$ and $\vec{r}'(t)$ are perpendicular to each other.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

For More Visit VU Answer