Question # 1 of 10 (Start time: 03:57:06 PM, 29 June 2022) Total Marks: 1

After reversing the order of limits of $\int_0^{\frac{1}{2}} \int_{2x}^{1} e^{y^2} dy\, dx$, we get _____.

Select the correct option

Reload Math Equations

- $\int_0^1 \int_{2x}^{\frac{1}{2}} e^{y^2} dx\, dy$
- $\int_0^1 \int_0^{\frac{y}{2}} e^{y^2} dx\, dy$
- $\int_0^1 \int_{\frac{1}{2}}^{2x} e^{y^2} dx\, dy$
- $\int_0^1 \int_{\frac{y}{2}}^{0} e^{y^2} dx\, dy$

Click to Save Answer & Move to Next Question

Question # 9 of 10 (Start time: 02:49:09 PM, 29 June 2022)

...ression for the double integral for the volume of a solid bounded above by $z=8-x-y$ and below by rectangle $R=0 \leq x \leq 4,\ 5 \leq y \leq 8$ is

...t the correct option

- $V=\int_5^8 \int_0^4 (8-x-y)dxdy$
- $V=\int_0^4 \int_5^8 (8-x-y)dxdy$
- $V=\int_5^8 \int_0^4 (8-x-y)dx$
- $V=\int_5^8 \int_0^4 (8-x-y)dydx$

Question # 4 of 10 (Start time: 03:29:15 PM, 29 June 2022)

If the value of the integral $\int_0^3 \int_0^1 (x+1)dydx$ =15/2 , then what will be the value of the integral $\int_0^1 \int_0^3 (x+1)dxdy$?

Select the correct option

- 0
- 15/2
- -15/2
- 9/2

Click to Save Answer & Mo

BC200417213: MUHAMMAD USMAN JAMEEL

Time Lef

MTH201:Quiz#2

Quiz Start Time: 03:26 PM, 2

Question # 7 of 10 (Start time: 03:34:06 PM, 29 June 2022)

T

$$\int_0^1 \int_1^2 dx\, dy = _____.$$

Select the correct option

Reload Ma

- 0
- 2
- 3
- 1



Click to Save Answer & Move to Next Que

Quiz Start Tim

rt time: 03:35:54 PM, 29 June 2022)

The equation of the line segment between the points $(0, -5)$ and $(4, 0)$ where slopeis$5/4$ is

n

$y = (5/4)x - 5$

$y = (-5/4)x + 5$

$y = (-5/4)x - 5$

$y = (5/4)x + 5$

Click to Save Answer & Mo

rt time: 03:35:54 PM, 29 June 2022)

The equation of the line segment between the points $(0, -5)$ and $(4, 0)$ where slopeis$5/4$ is

n

$$y = (5/4)x - 5$$

$$y = (-5/4)x + 5$$

$$y = (-5/4)x - 5$$

$$y = (5/4)x + 5$$

Click to Save Answer & Mo

MTH201:Quiz#2 Quiz Start Time: 04:00 PM, 29 June 202

Question # 3 of 10 (Start time: 04:01:43 PM, 29 June 2022) Total Marks

If the value of the integral $\int_0^3 \int_0^1 (x+1)dydx$ =15/2 , then what will be the value of the integral $\int_0^1 \int_0^3 (x+1)dxdy$?

Select the correct option

Reload Math Equation

- 0
- -15/2
- 15/2
- 9/2

Click to Save Answer & Move to Next Question

MTH201:Quiz#2 Quiz Start Time: 04:00 PM, 29 June 202

Question # 3 of 10 (Start time: 04:01:43 PM, 29 June 2022) Total Marks

If the value of the integral $\int_0^3 \int_0^1 (x+1)dydx$ =15/2 , then what will be the value of the integral $\int_0^1 \int_0^3 (x+1)dxdy$?

Select the correct option

Reload Math Equation

- 0
- -15/2
- 15/2
- 9/2

Click to Save Answer & Move to Next Question

MTH201:Quiz#2 Quiz Start Time: 04:00 PM, 29 June 202

Question # 3 of 10 (Start time: 04:01:43 PM, 29 June 2022) Total Marks

If the value of the integral $\int_0^3 \int_0^1 (x+1)dydx$ =15/2 , then what will be the value of the integral $\int_0^1 \int_0^3 (x+1)dxdy$?

Select the correct option

Reload Math Equation

- 0
- -15/2
- 15/2
- 9/2



Click to Save Answer & Move to Next Question

Question # 2 of 10 (Start time: 04:00:21 PM, 29 June 2022) Total Marks

Reversing the order of integration $\int_0^1 \int_{4x}^4 e^{-y^2} dy dx =$

Select the correct option

Reload Math Equation

- $\int_0^4 \int_0^{3y} e^{-y^2} dx dy$
- $\int_0^4 \int_0^{\frac{y}{4}} e^{-y^2} dx dy$
- $\int_0^4 \int_0^{\frac{y}{2}} e^{-y^2} dx dy$
- $\int_0^4 \int_0^{2y} e^{-y^2} dx dy$

Click to Save Answer & Move to Next Question

MTH201:Quiz#2

Quiz Start Time: 03:43 PM, 29 Jun

Question # 9 of 10 (Start time: 03:57:13 PM, 29 June 2022)

Total M

$$\int_0^1 \int_1^2 dx\, dy = _____.$$

Select the correct option

Reload Math Eq

- 1
- 3
- 0
- 2

Click to Save Answer & Move to Next Question

MTH201:Quiz#2

Quiz Start Time: 03:43 PM, 29 June 2022

Question # 10 of 10 (Start time: 03:58:00 PM, 29 June 2022)

Total Marks: 1

If the region is bounded below and above by the horizontal lines $y = c$ and $y = d$ and is bounded on the left and right by the continuous curves, $x = h_1(y)$ and $x = h_2(y)$, satisfying $h_1(y) \leq h_2(y)$, for $c \leq y \leq d$. Then $\iint_R f(x,y)\,dx\,dy =$ _____.

Select the correct option

Reload Math Equations

- $\int_c^d f(x,y)\,dx\,dy$
- $\int_c^d \int_{h_1(y)}^{h_2(y)} f(x,y)\,dx\,dy$
- $\int f(x,y)\,dy\,dx$
- $\int_{h_1(y)}^{h_2(y)} f(x,y)\,dx\,dy$

Click to Save Answer & Move to Next Question

3:48 75
quiz.vu.edu.pk/Qu 9

If ------- on R, then $\iint_R f(x, y)\, dA \geq 0.$

orrect option

Reload Ma

$f(x, y) \geq 0$

$f(x, y) = 0$

$f(x, y) \leq 0$

$f(x, y) < 0$

For each fixed $x; a \leqslant x \leqslant b$, the function $f(x, y)$ is a function ofyalone, then $A(x)$ the area of this function on the interval $c \leqslant y \leqslant d$ then the volume is

orrect option

$$V(S) = \int_c^d \left(\int_a^b f(x, y) dx \right) dy$$

$$V(S) = \int_a^b \left(\int_c^d f(x, y) dx \right) dy$$


$$V(S) = \int_c^d \left(\int_a^b f(x, y) dx \right) dy$$

$$V(S) = \int_a^b \left(\int_c^d f(x, y) dy \right) dx$$

$$\int_0^1 \int_0^{\ln y} x \, dy \, dx = \text{----------}$$

$$\frac{\ln x}{2}$$

$$\frac{\ln y}{2}$$




$$\frac{(\ln x)^2}{2}$$

$$\ln y$$

Quiz - Google Chro...

The equation of the line segment between the points $(0, -5)$ and $(4, 0)$ where slopeis$5/4$ is

$y = (5/4)x - 5$



$y = (5/4)x + 5$


$y = (-5/4)x - 5$

$y = (-5/4)x + 5$

The critical point of the given function $f(x, y) = 4xy - 3x + 2y + 10$ where $f_x(x, y) = 4y - 3$and $f_y(x, y) = 4x + 2$is

$(\frac{3}{4}, \frac{1}{2})$

$(\frac{1}{2}, \frac{3}{4})$

$(\frac{3}{4}, \frac{-1}{2})$ 


$(\frac{-1}{2}, \frac{3}{4})$

Question # 9 of 10 (Start time: 02:49:09 PM, 29 June 2022)

...ression for the double integral for the volume of a solid bounded above by $z=8-x-y$ and below by rectangle $R=0 \leq x \leq 4,\ 5 \leq y \leq 8$ =

...t the correct option

- $V=\int_5^8 \int_0^4 (8-x-y)dxdy$
- $V=\int_0^4 \int_5^8 (8-x-y)dxdy$
- $V=\int_5^8 \int_0^4 (8-x-y)dx$
- $V=\int_5^8 \int_0^4 (8-x-y)dydx$



Quiz - Google Chro...

Which of the following is a correct integral?

$$\iint_R f(x, y)\, dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y)\, dx\, dy$$

$$\iint_R f(x, y)\, dA = \int_c^d \int_{h_1(x)}^{h_2(x)} f(x, y)\, dx\, dy$$



$$\iint_R f(x, y)\, dA = \int_{g_1(x)}^{g_2(x)} \int_{h_1(y)}^{h_2(y)} f(x, y)\, dy\, dx$$

$$\iint_R f(x, y)\, dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y)\, dy\, dx$$

Reversing the order of integration $\int_{0}^{2}\int_{\frac{y}{2}}^{1} e^{x^2}\,dx\,dy =$

Reload Ma

$\int_{0}^{1}\int_{0}^{x^2} e^{x^2}\,dy\,dx$

$\int_{0}^{1}\int_{\frac{x}{2}}^{x} e^{x^2}\,dy\,dx$

$\int_{0}^{1}\int_{0}^{x} e^{x^2}\,dy\,dx$

$\int_{0}^{1}\int_{0}^{2x} e^{x^2}\,dy\,dx$




After reversing the order of limits of $\int_0^3 \int_{x^2}^9 y \cos x \, dy \, dx$, we get - - -

$$\int_0^3 \int_0^{\sqrt{x}} y \cos x \, dx dy$$

$$\int_0^9 \int_0^{\sqrt{y}} y \cos x \, dx dy$$

$$\int_{x^2}^9 \int_0^3 y \cos x \, dx dy$$

$$\int_{}^{9} \int_{}^{\sqrt{y}} y \cos x \, dy dx$$



SAMSUNG


$$\int_0^{10} (x^3y + y^2)\ dz = \text{-------}$$

$$10\,(x^3y + y^2)$$

$$500y^2 + \frac{1000}{3}$$

$$10\left(x^3y^2 + \frac{y^3}{3}\right)$$

$$50x^3 + \frac{1000}{3}$$



SAMSUNG

MTH201:Quiz#2 Quiz Start Time: 02:00 PM, 29 June 2022

Question # 10 of 10 (Start time: 02:08:53 PM, 29 June 2022) Total Marks: 1

If the region is bounded the left and right by vertical lines $x = a$ and $x = b$ and is bounded below and above by curves, $y = g_1(x)$ and $y = g_2(x)$, where $g_1(x) \leqslant g_2(x)$ for $a \leqslant x \leqslant b$. Then $\iint_R f(x,y)\,dA =$ _____.

Select the correct option

Reload Math Equations

- $\int f(x,y)\,dy\,dx$
- $\int_a^b \int_{g_1(x)}^{g_2(x)} f(x,y)\,dy\,dx$
- $\int_{g_1(x)}^{g_2(x)} f(x,y)\,dy\,dx$
- $\int_a^b f(x,y)\,dx\,dy$

Click to Save Answer & Move to Next Question

MTH201:Quiz#2

Quiz Start Time: 02:11 PM, 29 June 2022

Question # 1 of 10 (Start time: 02:11:22 PM, 29 June 2022)

Total Marks: 1

$$\int_0^1 \int_0^{\ln y} x \, dy \, dx = - - - - - - - - - -$$

Select the correct option

Reload Math Equations

- $\frac{(\ln x)^2}{2}$
- $\ln y$
- $\frac{\ln x}{2}$
- $\frac{\ln y}{2}$



Click to Save Answer & Move to Next Question

MTH201:Quiz#2

Quiz Start Time: 02:11 PM, 29 June 2022

Question # 2 of 10 (Start time: 02:11:53 PM, 29 June 2022)

Total Marks: 1

The critical point of $f(x) = 2x^2$ is $x =$ _____.

Select the correct option

Reload Math Equations

- 2
- 0
- 1
- -1

Click to Save Answer & Move to Next Question

Question # 3 of 10 (Start time: 02:12:36 PM, 29 June 2022) Total Marks: 1

The critical points of the given function $f(x) = x^3 + x^2 - x + 10$
where $f_x(x) = 3x^2 + 2x - 1$are

Select the correct option

Reload Math Equations

- $x = \frac{1}{3}, 1$
- $x = \frac{1}{3}, -1$
- $x = \frac{-1}{3}, -1$
- $x = \frac{-1}{3}, 1$

Click to Save Answer & Move to Next Question

Question # 4 of 10 (Start time: 02:13:20 PM, 29 June 2022) Total Marks: 1

The absolute extrema of $f(x) = 2x$ is _____.

Select the correct option

Reload Math Equations

- $(0, 0)$
- $(0, 1)$
- $(0, -1)$
- None

Click to Save Answer & Move to Next Question

MTH201:Quiz#2 Quiz Start Time: 02:11 PM, 29 June 2022

Question # 5 of 10 (Start time: 02:13:44 PM, 29 June 2022) Total Marks: 1

$$\text{The} \int_0^1 2xe^{x^2} dx =$$

Select the correct option

Reload Math Equations

- $e + 1$
- e^{x^2}
- e
- $e - 1$



Click to Save Answer & Move to Next Question

Question # 6 of 10 (Start time: 02:22:52 PM, 29 June 2022) Total Marks: 1

Expression for the double integral for the volume of a solid bounded above by z=8-x-y and below by rectangle R=$0 \leqslant x \leqslant 4,\ 5 \leqslant y \leqslant 8$ is.....

Select the correct option

Reload Math Equations

- $V = \int_5^8 \int_0^4 (8 - x - y)dydx$
- $V = \int_0^4 \int_5^8 (8 - x - y)dxdy$
- $V = \int_5^8 \int_0^4 (8 - x - y)dx$
- $V = \int_5^8 \int_0^4 (8 - x - y)dxdy$

sec(s)

MTH201:Quiz#2

Quiz Start Time: 02:11 PM, 29 June 2022

Question # 10 of 10 (Start time: 02:17:19 PM, 29 June 2022)

Total Marks: 1

The volume of a parallelopiped with dimensions $x,\ y,\ z$ is . . . . . .

Select the correct option

Reload Math Equations

- $V = x + y + z$
- $V = x\,y\,z$
- $V = \sqrt{x\,y\,z}$
- $V = x^2\,y^2\,z^2$

Click to Save Answer & Move to Next Question

BC190407302: MUHAMMAD SADDAM LIAQAT

Time Left 87 sec(s)

MTH201:Quiz#2

Quiz Start Time: 01:50 PM, 29 June 2022

Question # 5 of 10 (Start time: 01:55:46 PM, 29 June 2022)

Total Marks: 1

Reversing the order of integration $\int_0^1 \int_{4x}^4 e^{-y^2} dy dx =$

Select the correct option

Reload Math Equations

- $\int_0^4 \int_0^{\frac{y}{2}} e^{-y^2} dx dy$
- $\int_0^4 \int_0^{2y} e^{-y^2} dx dy$
- $\int_0^4 \int_0^{3y} e^{-y^2} dx dy$
- $\int_0^4 \int_0^{\frac{y}{4}} e^{-y^2} dx dy$



Click to Save Answer & Move to Next Question

BC190407302: MUHAMMAD SADDAM LIAQAT

Time Left 88 sec(s)

MTH201:Quiz#2

Quiz Start Time: 01:50 PM, 29 June 2022

Question # 6 of 10 (Start time: 01:56:51 PM, 29 June 2022)

Total Marks: 1

After reversing the order of limits of $\int_{0}^{\frac{1}{2}} \int_{2x}^{1} e^{y^2}\,dy\,dx$, we get _____.

Select the correct option

Reload Math Equations

- $\int_{0}^{1} \int_{\frac{1}{2}}^{0} e^{y^2}\,dx\,dy$
- $\int_{0}^{1} \int_{\frac{1}{2}}^{2x} e^{y^2}\,dx\,dy$
- $\int_{0}^{1} \int_{2x}^{\frac{1}{2}} e^{y^2}\,dx\,dy$
- $\int_{0}^{1} \int_{0}^{\frac{x}{2}} e^{y^2}\,dx\,dy$



Click to Save Answer & Move to Next Question

BC190407302: MUHAMMAD SADDAM LIAQAT

Time Left 88 sec(s)

MTH201:Quiz#2

Quiz Start Time: 01:50 PM, 29 June 2022

Question # 7 of 10 (Start time: 01:57:46 PM, 29 June 2022)

Total Marks: 1

$$\int_0^{\ln 3} \int_0^{\ln 5} e^{u+v}\, du\, dv = \text{- - - - - - - - - - -}$$

Select the correct option

Reload Math Equations

- 15
- 8
- 5
- 3

Click to Save Answer & Move to Next Question

BC190407302: MUHAMMAD SADDAM LIAQAT

Time Left 89 sec(s)

MTH201:Quiz#2

Quiz Start Time: 01:50 PM, 29 June 2022

Question # 8 of 10 (Start time: 01:58:23 PM, 29 June 2022)

Total Marks: 1

Let x be the length, width and height of a rectangular box. The area of its roof will be . . . . . .

Select the correct option

Reload Math Equations

- $A = x$
- $A = 2x$
- $A = x^2$
- $A = x + x + x$

Click to Save Answer & Move to Next Question

Question # 9 of 10 (Start time: 01:58:57 PM, 29 June 2022) Total Marks: 1

The expression $\int_a^b f(x,y)dx$ is a function of .....

Select the correct option

Reload Math Equations

- x
- y
- None of these
- both x and y

Click to Save Answer & Move to Next Question

Question # 10 of 10 (Start time: 01:59:29 PM, 29 June 2022) Total Marks: 1

For each fixed $x; a \leqslant x \leqslant b$, the function $f(x, y)$ is a function ofyalone, then the area of this function on the interval $c \leqslant y \leqslant d$is

Select the correct option

Reload Math Equations

- $A(x) = \int_a^b f(x, y)dx$
- $A(x) = \int_c^d f(x, y)dy$
- $A(x) = \int_a^b f(x, y)dy$
- $A(x) = \int_c^d f(x, y)dx$

Click to Save Answer & Move to Next Question

me: 02:01:00 PM, 29 June 2022)

Total Ma

Which of the following is a correct integral?

Reload Math Equa

$$\iint_R f(x,\, y)\, dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x,\, y)\ dy\, dx$$

$$\iint_R f(x,\, y)\, dA = \int_c^d \int_{h_1(x)}^{h_2(x)} f(x,\, y)\ dx\, dy$$



$$\iint_R f(x,\, y)\, dA = \int_{g_1(x)}^{g_2(x)} \int_{h_1(y)}^{h_2(y)} f(x,\, y)\ dy\, dx$$

$$\iint_R f(x,\, y)\, dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x,\, y)\ dx\, dy$$

Click to Save Answer & Move to Next Question

2:15 PM

Solution

$$\int_0^1 2x\,e^{x^2}\,dx$$

Solution

Show Steps

$$\int_0^1 2xe^{x^2}\,dx = e-1 \quad \text{(Decimal: } 1.71828...\text{)}$$

Steps

$$\int_0^1 2xe^{x^2}\,dx$$

Take the constant out: $\int a\cdot f(x)\,dx = a\cdot\int f(x)\,dx$

$$= 2\cdot\int_0^1 xe^{x^2}\,dx$$

Apply u – substitution — *Show Steps*

$$= 2\cdot\int_0^1 \frac{e^u}{2}\,du$$

Take the constant out: $\int a\cdot f(x)\,dx = a\cdot\int f(x)\,dx$

$$= 2\cdot\frac{1}{2}\cdot\int_0^1 e^u\,du$$

Use the common integral: $\int e^u\,du = e^u$

$$= 2\cdot\frac{1}{2}\left[e^u\right]_0^1$$

COOL OMOJI and LOVE OMOJI

Question # 2 of 10 (Start time: 02:01:52 PM, 29 June 2022) Total Mar

Let x, y, z be the length, width and height of a rectangular box respectively. The area of bottom will be . . . . . .

Select the correct option

Reload Math Equati

- $A = yz$
- $A = xyz$
- $A = xy$
- $A = xz$

Click to Save Answer & Move to Next Question

MTH201:Quiz#2 Quiz Start Time: 02:00 PM, 29 June 2022

Question # 3 of 10 (Start time: 02:02:23 PM, 29 June 2022) Total Marks: 1

$$\int_0^{\ln 6}\int_0^{\ln 3} f(x,\,y)\,dx\,dy \;=\; \int_0^{\ln 3}\int_0^{\ln 6} f(x,\,y)\,dy\,dx$$

Select the correct option

Reload Math Equations

- True
- False

Click to Save Answer & Move to Next Question

2:12 PM

Solution

$$\int_0^1 \int_0^{\ln y} x \, dydx$$

Solving $\int_0^1 \int_0^{\ln(y)} xdxdy$

Solution

Show Steps

$$\int_0^1 \int_0^{\ln(y)} xdxdy = 1$$

Steps

$$\int_0^1 \int_0^{\ln(y)} xdxdy$$

$$\int_0^{\ln(y)} xdx = \frac{\ln^2(y)}{2}$$ Show Steps

$$= \int_0^1 \left(\frac{\ln^2(y)}{2}\right) dy$$

$$\int_0^1 \left(\frac{\ln^2(y)}{2}\right) dy = 1$$ Show Steps

$$= 1$$

Got a different answer? Check if it's correct

Enter your answer

MTH201:Quiz#2 Quiz Start Time: 02:00 PM, 29 June 2022

Question # 4 of 10 (Start time: 02:02:58 PM, 29 June 2022) Total Marks: 1

After reversing the order of limits of $\int_0^3 \int_{x^2}^9 y \cos x \, dy \, dx$, we get ------

Select the correct option

Reload Math Equations

- $\int_0^9 \int_0^{\sqrt{y}} y \cos x \, dx dy$
- $\int_0^9 \int_0^{\sqrt{y}} y \cos x \, dy dx$
- $\int_0^3 \int_0^{\sqrt{x}} y \cos x \, dx dy$
- $\int_{x^2}^6 \int_0^3 y \cos x \, dx dy$

Click to Save Answer & Move to Next Question

Question # 6 of 10 (Start time: 02:04:49 PM, 29 June 2022) Total Marks: 1

After reversing the order of limits of $\int_{0}^{\frac{1}{2}} \int_{2x}^{1} e^{y^2}\, dy\, dx$, we get _____.

Select the correct option

Reload Math Equations

- $\int_{0}^{1} \int_{2x}^{\frac{1}{2}} e^{y^2}\, dx\, dy$
- $\int_{0}^{1} \int_{\frac{1}{2}}^{2x} e^{y^2}\, dx\, dy$
- $\int_{0}^{1} \int_{0}^{\frac{y}{2}} e^{y^2}\, dx\, dy$
- $\int_{0}^{1} \int_{\frac{y}{2}}^{0} e^{y^2}\, dx\, dy$

Click to Save Answer & Move to Next Question

Question # 7 of 10 (Start time: 02:05:53 PM, 29 June 2022) Total Marks: 1

For each fixed x; $a \leqslant x \leqslant b$, the function $f(x, y)$ is a function ofyalone,
then $A(x)$ the area of this function on the interval $c \leqslant y \leqslant d$ then the volume is

Select the correct option

Reload Math Equations

- $V(S) = \int_c^d \left(\int_a^b f(x, y)dx \right) dy$
- $V(S) = \int_a^b \left(\int_c^d f(x, y)dx \right) dy$
- $V(S) = \int_c^d \left(\int_a^b f(x, y)dx \right) dy$
- $V(S) = \int_a^b \left(\int_c^d f(x, y)dy \right) dx$

Click to Save Answer & Move to Next Question

Question # 8 of 10 (Start time: 02:06:55 PM, 29 June 2022) Total Marks: 1

The critical points of the given function $f(x) = x^3 + x^2 - x + 10$ where $f_x(x) = 3x^2 + 2x - 1$are

Select the correct option

Reload Math Equations

- $x = \frac{1}{3}, 1$
- $x = \frac{-1}{3}, 1$
- $x = \frac{-1}{3}, -1$
- $x = \frac{1}{3}, -1$



Click to Save Answer & Move to Next Question

Let x be the length, width and height of a rectangular box. The area of its roof will be . . . . . .

Select the correct option

Reload Math Equations

- $A = x + x + x$
- $A = x$
- $A = x^2$
- $A = 2x$



Click to Save Answer & Move to Next Question

Question # 10 of 10 (Start time: 01:35:05 PM, 29 June 2022) Total Marks: 1

The order of integration in the iterated integral does not matter.

Select the correct option

- False
- True



Click to Save Answer & Move to Next Question

$$\int_0^2 \int_0^4 10\,dx\,dy \;=\; - - - - - - -$$

Select the correct option

- 80
- 20
- 60
- 40

Click to Save Answer & Move

The integral to find the volume under the surface $z = 5x^3 + 5x^2y$ and the rectangle $\{(x, y) : 0 \leqslant x \leqslant 3, 0 \leqslant y \leqslant 2\}$ is - - - - - - -

Select the correct option

Reload Math Equations

- $\int_0^2 \int_0^3 (5x^3 + 5x^2y)\, dx\, dy$
- $\int_0^3 \int_0^2 (5x^3 + 5x^2y)\, dx\, dy$
- $\int_0^2 \int_0^3 (5x^3 + 5x^2y)\, dydx$
- $\int_0^2 \int_0^2 (5x^3 + 5x^2y)\, dydx$

Click to Save Answer & Move to Next Question

Question # 4 of 10 (Start time: 01:45:33 PM, 29 June 2022)

$$\int_0^1 \int_0^1 dx\, dy = _____.$$

Select the correct option

- 3
- 0
- 1
- 2



Click to Save Answer & M

Question # 10 of 10 (Start time: 01:45:50 PM, 29 June 2022) Total Marks: 1

For each fixed $y; c \leq y \leq d$, the function $f(x, y)$ is a function of x alone, then the area of this function on the interval $a \leq x \leq b$ is

Select the correct option

Reload Math Equations

- $A(y) = \int_c^d f(x, y)dy$
- $A(y) = \int_a^b f(x, y)dy$
- $A(y) = \int_a^b f(x, y)dx$
- $A(y) = \int_c^d f(x, y)dx$

Click to Save Answer & Move to Next Question

The critical point(s) of $f(x) = 3x - 2$ is(are) $x =$ _____.

Select the correct option

- $0, -1$
- $0, 0$
- None
- $0, 1$

Click to Save

After reversing the order of limits of $\int_0^2 \int_x^2 f(x, y)\, dy\, dx$, we get – – – – – – –

Select the correct option

Reload Math Equations

- $\int_y^2 \int_x^2 f(x, y)\, dx\, dy$
- $\int_0^2 \int_0^y f(x, y)\, dx\, dy$
- $\int_x^2 \int_0^y f(x, y)\, dy\, dx$
- $\int_0^2 \int_x^2 f(x, y)\, dx\, dy$

Click to Save Answer & Move to Next Question

The expression $\int_a^b f(x,y)dy$ is a function of .....

Select the correct option

- None of these
- y
- Both x and y
- x



Click to Sa

Question # 7 of 10 (Start time: 01:48:44 PM, 29 June 2022)

For each fixed $x; a \leqslant x \leqslant b$, the function $f(x, y)$ is a function of$y$alone, then $A(x)$ the area of this function on the interval $c \leqslant y \leqslant d$ then the volume is

Select the correct option

- $V(S) = \int_a^b \left(\int_c^d f(x, y) dy \right) dx$
- $V(S) = \int_c^d \left(\int_a^b f(x, y) dx \right) dy$
- $V(S) = \int_c^d \left(\int_a^b f(x, y) dx \right) dy$
- $V(S) = \int_a^b \left(\int_c^d f(x, y) dx \right) dy$

Click to Save Answer & Move t

The critical points of the given function $f(x) = x^3 + x^2 - x + 10$ where $f_x(x) = 3x^2 + 2x - 1$ are _____.

Select the correct option

Reload Math Equations

- $x = \frac{1}{3}, 1$
- $x = \frac{1}{3}, -1$
- $x = \frac{-1}{3}, -1$
- $x = \frac{-1}{3}, 1$

Click to Save Answer & Move to Next Question

For each fixed $y; c \leqslant y \leqslant d$, the function $f(x, y)$ is a function of x alone, then $A(y)$ the area of this function on the interval $a \leqslant x \leqslant b$ then the volume is

$$\mathrm{V(S)} = \int_{c}^{d} \left(\int_{a}^{b} \mathrm{f(x,y)dx} \right) \mathrm{dy}$$

Select the correct option

Reload Math Eq

- $V(S) = \int_{a}^{b} \left(\int_{c}^{d} f(x, y)dx \right) dy$
- $V(S) = \int_{c}^{d} \left(\int_{a}^{b} f(x, y)dx \right) dy$
- $V(S) = \int_{a}^{b} \left(\int_{c}^{d} f(x, y)dy \right) dx$
- $V(S) = \int_{c}^{d} \left(\int_{a}^{b} f(x, y)dx \right) dy$

Click to Save Answer & Move to Next Question

Question # 10 of 10 (Start time: 01:50:59 PM, 29 June 2022)

X,Y,Z are three positive numbers whose sum is 27 and such that the sum of their squares is as small as possible, find the numbers.

Select the correct option

- X=12, Y=6, Z=9
- X=9, Y=12, Z=6
- X=9, Y=9, Z=9
- X=3, Y=9, Z=15

Click to Save Answer & Move to Next Question

Question # 8 of 10 (Start time: 01:50:25 PM, 29 June 2022)

If $------$ on R, then $\iint_R f(x, y)\, dA \geqslant 0.$

Select the correct option

- $f(x, y) = 0$
- $f(x, y) \geqslant 0$
- $f(x, y) \leqslant 0$
- $f(x, y) < 0$

Click to Save Answer & Move t

Quiz#2

Quiz Start Time: 01:41 PM

10 of 10 (Start time: 01:53:11 PM, 29 June 2022)

The double integral of two variables x and y over the certain region R of the xy - plane can be expressed as _____.

correct option

Reload

$$\iint_R f(x,y)\,dx\,dy$$

$$\int_R f(x,y)\,dx\,dy$$

$$\iint_R f(x,y)\,dy$$

$$\iint_R f(x,y)\,dx$$

Click to Save Answer & Move to Next