MTH501

Quiz no 2

28 Dec 2025

SOLVED BY


IQRA AZHAR

SERVICES

Assignment || Quiz ||GDB ||Project ||
Seminar ||Online Book Shop ||
LMS Hndling Services Available

CONTACT 03305516507

MTH501 QUIZ 2 BY IQRA AZHAR



MTH501 - MTH501 - Linear Algebra (Quiz 02)

Quiz ‹Back

#	Quiz Title	Start Date	End Date	Total Marks	Quiz Status	Submit Status	Result
1	Quiz 01	Nov 06, 2025 12:00 AM	Nov 07, 2025 11:59 PM	10	Closed	Submitted Submit Date: Nov 07,2025 02:48 PM	10
2	Quiz 02	Dec 26, 2025 12:00 AM	Dec 28, 2025 11:59 PM	10	Open	Submitted Submit Date: Dec 26,2025 10:40 AM	
3	Quiz 03	Jan 15, 2026 12:00 AM	Jan 16, 2026 11:59 PM	10	-	-	
4	Quiz 04	Jan 18, 2026 12:00 AM	Jan 19, 2026 11:59 PM	10	-	-	

Note: To take Quiz in Chrome browser, It is mandatory to Install/Enable VU Quiz Firewall Extension.

The rank of a 3 x 3 matrix C (= AB), found by multiplying a non-zero column matrix A of size 3 x 1 and a non-zero row matrix B of size 1 x 3, is

0

2

3

1

Cylindrical coordinate system is also called as_____

Spherical coordinate system.

Space coordinate system.

Circular coordinate system.

Rectangular coordinate system.

MTH501 QUIZ 2 BY IQRA AZHAR

In two coordinate system, \[{\left({\mathop P\limits_{C \leftarrow B} } \right)^{ - 1}}\] is the matrix that converts

B-coordinates into B-coordinates

B-coordinates into C-coordinates

C-coordinates into B-coordinates

C-coordinates into C-coordinates

The pivot columns of a matrix \[A\] form a basis for ----------

[RowA]

[ColA]

The rank of the matrix (m × n) where m<n cannot be more than?

m

n

The vectors spaces - C(-infty, +infty) are ----------

Finite-dimensional

Infinite-dimensional

The number of parameters in the solution set of Ax = 0 if A is a 4× 4 matrix of rank 0 then nullity (A) =?

0

4

If A is an m x n, matrix, then

Rank (A) = the number of leading variables in the solution of Ax = 0

Nullity (A) = the number of parameters in the general solution of Ax = 0

True

False

The dimension of the zero vector space \[\left\{ 0 \right\}\] is defined to be

None of these

[\infty\]

0

1

MTH501 QUIZ 2 BY IQRA AZHAR

If Coordinates vector of the polynomial \[p = {a_0} + {a_1}x + {a_2}{x_2}\] Relative to the basis S = \[\{ 1,x,{x^2}\} \,\,\,\] for \[{p_2}\] then linear combination of the basis set S is

[P]_s = (a_0 + a_1 + a_2)

[P]_s = [a_0 + a_1 + a_2]

An indexed set of vectors \[{v_1,v_2,…,v_p}\] in \[V\] is said to be linearly independent if the vector equation

\[{c_1}{v_1} + {c_2}{v_2} + \ldots + {c_p}{v_p} \ne 0\]

\[{c_1}{v_1} + {c_2}{v_2} + \ldots + {c_p}{v_p} = 1\]

c_1v_1}+ c_2v_2 + \ldots + {c_p}{v_p} =0\]

None of the above.

Any set of linearly independent vectors cannot contain the ---------- vector.

Identity

None of these

zero

Unit

If A is any matrix, then rank (A)=rank(A^t)

False

True

If A is an m x n, matrix, then

Rank (A) = the number of leading variables in the solution of Ax = 0

Nullity (A) = the number of parameters in the general solution of Ax = 0

False

True

Every finite dimensional vector space contains a/an

Basis

Empty set

Infinite set

None of the above

If two matrices A and B are row equivalent, then their row spaces are the same.

MTH501 QUIZ 2 BY IQRA AZHAR

true

False

If vector space is not spanned by a finite set of vectors, then V is said to be --------

Finite-dimensional

Infinite-dimensional

The set of unit vectors \[{i,j,k}\]form a basis of

\[R^4\]

\[R^2\]


R^3

\[R\]

If two matrices A and B are row equivalent, then their row spaces are the.

Different

Same



MTH501 QUIZ 2 BY IQRA AZHAR

Youtube

Vu zone with iqra azhar

Whatsapp

03305516507

 Allah Hafiz 