

MC170201745: Shakeel Ahmad

Time Left 73 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 9 of 10 (Start time: 07:54:42 PM, 16 January 2018)

Total Marks: 1

The limit

$$\lim_{x \rightarrow 1} \sqrt{x} - 1.$$

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|-----------------|
| ☐ | 1. |
| ☒ | 0. |
| ☐ | does not exist. |
| ☐ | -1. |

By google

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 31 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 8 of 10 (Start time: 02:13:27 PM, 28 November 2017)

Total Marks: 1

The set of terms of $\{s_{n_k}\}$ is contained in the set of terms of $\{s_n\}$ implies

Select the correct option

Reload Math Equations

- ☐ $\sup\{s_n\} = \sup\{s_{n_k}\}.$
- ☐ $\sup\{s_n\} \leq \sup\{s_{n_k}\}.$
- ☒ $\sup\{s_n\} \geq \sup\{s_{n_k}\}.$
- ☐ $\{s_n\}$ is convergent.

page 38

MC170202886: Aqsa Zunaira

Time Left 62 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 2 of 10 (Start time: 10:03:30 PM, 06 February 2018)

Total Marks: 1

The Mean Value Theorem says, suppose that f is differentiable on $[a, b]$, $f'(a) \neq f'(b)$, and μ is between $f'(a)$ and $f'(b)$.

Select the correct option

Reload Math Equations

Then $f'(c) \neq \mu$ for some c in (a, b) .Then $f'(c) = \mu$ for some c in (a, b) .

page 111

Then $f'(c) = 0$ for some c in (a, b) .Then $f'(c)$ may be 0 or μ for some c in (a, b) .

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 74 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 1 of 10 (Start time: 07:47:22 PM, 16 January 2018)

Total Marks: 1

The limit $\lim_{x \rightarrow \infty} x^2 - x$.

is
Select the correct option

Reload Math Equations

- ☒ does not exist.
- ☐ 2.
- ☐ 0.
- ☐ 6.

by google uncle

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 35 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 1 of 10 (Start time: 02:06:09 PM, 28 November 2017)

Total Marks: 1

Which statement(s) is(are) true about the following sequence $s_0 = 1$ and $s_n = 1 - e^{-s_{n-1}}$

Select the correct option

Reload Math Equations

- ☐ $0 < s_n \leq 1$ for all n .
- ☐ $s_{n+1} - s_n = -(e^{-s_n} - e^{-s_{n-1}})$.
- ☐ Sequence is convergent.
- ☒ All of the above.

page 33

not sure

MC170202886: Aqsa Zunaira

Time Left 55
sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 9 of 10 (Start time: 10:05:45 PM, 06 February 2018)

Total Marks: 1

If $f : I \rightarrow \mathbb{R}$ has a derivative at $c \in I$, then f is

Out of sylubs

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|-------------------------|
| ☒ | continuous at c . |
| ☐ | discontinuous at c . |
| ☐ | undefined at c . |
| ☐ | having one sided limit. |

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 30 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 7 of 10 (Start time: 02:12:09 PM, 28 November 2017)

Total Marks: 1

The series $\sum_{n=0}^{\infty} r^n$, $-1 < r < 1$, is

Select the correct option

Reload Math Equations

- ☒ divergent.
- ☐ convergent and converges to $\frac{1}{1+r^2}$.
- ☐ convergent and converges to 5.
- ☒ convergent and converges to $\frac{1}{1-r}$.

page 43

MC170201745: Shakool Ahmad

Time Left 40 sec(s)

MTH621:Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 2 of 10 (Start time: 02:07:17 PM, 28 November 2017)

Total Marks: 1

If the sequence is increasing, then it

Select the correct option

- ☐ converges to its supremum.
- ☐ diverges.
- ☒ may converges to its supremum.
- ☐ is bounded.

✓ by google

MC170201745: Shakool Ahmad

Time Left 53 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 4 of 10 (Start time: 02:09:40 PM, 28 November 2017)

Total Marks: 1

Identify the identical sequences.

Select the correct option

Reload Math Equations

- ☐ $\left\{ \frac{1}{n^2-2} \right\}_3^\infty$ and $\left\{ \frac{1}{n} \right\}_1^\infty$.
- ☐ $\left\{ \frac{1}{n} \right\}_3^\infty$ and $\left\{ \frac{1}{n} \right\}_1^\infty$.
- ☐ $\left\{ \frac{1}{n-2} \right\}_3^\infty$ and $\left\{ \frac{1}{n^2} \right\}_1^\infty$.
- ☒ $\left\{ \frac{1}{n-2} \right\}_3^\infty$ and $\left\{ \frac{1}{n} \right\}_1^\infty$.

page 29

MC170201745: Shakeel Ahmad

Time Left 63 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 09:37 PM, 06 February 2018

Question # 10 of 10 (Start time: 09:42:41 PM, 06 February 2018)

Total Marks: 1

$\lim_{x \rightarrow 0^+} x \log x$, has the following indeterminate form

Select the correct option

Reload Math Equations

 $\infty \times \infty$. $(0)(\infty)$. $\frac{0}{0}$. 0^∞ .

page 117

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 70 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 4 of 10 (Start time: 10:04:12 PM, 06 February 2018)

Total Marks: 1

Rolle's Theorem says. suppose that f is continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) and $f(a) = f(b)$.

Select the correct option

Reload Math Equations

page 111

Then $f'(c) = 0$ for some c in the open interval (a, b) .Then $f'(c) \neq 0$ for some c in the open interval (a, b) .Then $f'(c)$ exists for some c in the open interval (a, b) .Then $f'(c) = 0$ for some c in the open interval (a, ∞) .

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 80 sec(s)

MTH621: Quiz 4 - Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 5 of 10 (Start time: 10:04:36 PM, 06 February 2018)

Total Marks: 1

Generalized Mean Value Theorem says. If f and g are continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) , then

Select the correct option

Reload Math Equations

for some c in (a, b) .

$$[g(b) - g(a)]f'(c) = [f(b) - f(a)]g'(c)$$

for some c in (a, b) .

$$[g(b) - g(a)] = [f(b) - f(a)]g'(c)$$

 $g'(x) = 0$ for all x in (a, b) . $g'(x) = f'(x)$ for all x in (a, b) .

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 77 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 5 of 10 (Start time: 09:47:26 PM, 18 January 2018)

Total Marks: 1

For the function defined as $f(x) = cx$, for every $\epsilon > 0$ the formal definition of limit of function ensures

Select the correct option

Reload Math Equations

- ☐ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } 0 < \delta < \epsilon.$
- ☒ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } 0 < \delta < \epsilon/|c|.$
- ☐ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } -\epsilon < \delta < \epsilon.$
- ☐ $|f(x) - c| < \epsilon, \quad |x - x_0| < \delta, \text{ where } 0 < \delta < \epsilon.$

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 82 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 2 of 10 (Start time: 10:32:00 PM, 06 February 2018)

Total Marks: 1

If $f : I \rightarrow \mathbb{R}$ has a derivative at $c \in I$, then f is

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|-------------------------|
| ☒ | continuous at c. |
| ☐ | discontinuous at c. |
| ☐ | undefined at c. |
| ☐ | having one sided limit. |

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 27
sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 6 of 10 (Start time: 07:51:32 PM, 16 January 2018)

Total Marks: 1

The $\lim_{x \rightarrow 0} \frac{4-4\cos x - 2\sin^2 x}{x^4}$ is

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|----------------|
| ☐ | $\frac{1}{4}$ |
| ☐ | $-\frac{1}{2}$ |
| ☒ | $\frac{1}{2}$ |
| ☐ | 2 |

page 116



Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 70
sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 10 of 10 (Start time: 10:06:24 PM, 06 February 2018)

Total Marks: 1

The function $h(x) = -x^3$ is

Select the correct option

Reload Math Equations

is undefined on $(-\infty, \infty)$.decreasing on $(-\infty, \infty)$.increasing on $(-\infty, \infty)$.positive on $(-\infty, \infty)$.

Click to Save Answer & Move to Next Question

page 97

MC170201745: Shakeel Ahmad

Time Left 51 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 09:37 PM, 06 February 2018

Question # 8 of 10 (Start time: 09:41:16 PM, 06 February 2018)

Total Marks: 1

The inverse of the given function $f(x) = 2x + 4$, $0 \leq x \leq 2$, is

Select the correct option



$$f^{-1}(y) = \frac{(y-4)}{2}$$

out



$$f^{-1}(y) = \frac{(y-2)}{4}$$



$$f^{-1}(y) = \frac{(x-2)}{4}$$



$$f^{-1}(y) = \frac{(y-4)}{4}$$

Click to Save Answer & Move to Next Question



MC170201745: Shakeel Ahmad

Time Left

52

sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 09:37 PM, 06 February 2018

Question # 7 of 10 (Start time: 09:40:26 PM, 06 February 2018)

Total Marks: 1

The inverse of the function $f(x) = x^2$, is

Select the correct option

Reload Math Equations



$$f^{-1}(y) = y^2$$



$$f^{-1}(y) = \sqrt{y}.$$



$$f^{-1}(y) = \frac{1}{x^3}$$



$$f^{-1}(y) = \frac{1}{y^2}$$

page 101

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 54 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 5 of 10 (Start time: 02:10:35 PM, 28 November 2017)

Total Marks: 1

Which statement(s) is(are) true about the following sequence $s_0 = 1$ and $s_n = 1 - e^{-s_{n-1}}$

Select the correct option

Reload Math Equations

- ☐ $0 < s_n \leq 1$ for all n .
- ☐ $s_{n+1} - s_n = -(e^{-s_n} - e^{-s_{n-1}})$.
- ☐ Sequence is convergent.
- ☒ All of the above.

MC170202886: Aqsa Zunaira

Time Left 83 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 3 of 10 (Start time: 10:04:02 PM, 06 February 2018)

Total Marks: 1

The inverse of the function $f(x) = x^2$, is

Select the correct option

Reload Math Equations



$$f^{-1}(y) = y^2$$



$$f^{-1}(y) = \sqrt{y}.$$



$$f^{-1}(y) = \frac{1}{x^3}$$



$$f^{-1}(y) = \frac{1}{y^2}$$

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 55 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 6 of 10 (Start time: 07:51:32 PM, 16 January 2018)

Total Marks: 1

The $\lim_{x \rightarrow 0} \frac{4-4\cos x - 2\sin^2 x}{x^4}$ is

Select the correct option

Reload Math Equations

- | | | |
|----------------------------------|----------------|----------|
| ☐ | $\frac{1}{4}$ | rep |
| ☐ | $-\frac{1}{2}$ | |
| ☒ | $\frac{1}{2}$ | page 116 |
| ☐ | 2 | |

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 47 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 3 of 10 (Start time: 02:08:24 PM, 28 November 2017)

Total Marks: 1

Which of the following given sets is compact subset of $\mathbb{R}$.

Select the correct option

Reload Math Equations

- ☐ (1,5]
- ☐ {1,2,4}
- ☒ [0,1]
- ☐ (2,3)

page 28

MC170201745: Shakeel Ahmad

Time Left 42 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 3 of 10 (Start time: 07:48:48 PM, 16 January 2018)

Total Marks: 1

For the function $\frac{|x|}{x}$, identify the false statement(s)

out out

Select the correct option

Reload Math Equations



$$\lim_{x \rightarrow 0^+} \frac{|x|}{x} = 1$$



$$\lim_{x \rightarrow 0^-} \frac{|x|}{x} = -1.$$



$$\lim_{x \rightarrow 0} \frac{|x|}{x} \text{ doesn't exist.}$$



$$\lim_{x \rightarrow 0} \frac{|x|}{x} = 1.$$



Click to Save Answer & Move to Next Question



MC170202886: Aqsa Zunaira

Time Left 81 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 7 of 10 (Start time: 09:48:03 PM, 18 January 2018)

Total Marks: 1

The value of the limit

$$\lim_{x \rightarrow 0^+} x \log x,$$

Select the correct option

Reload Math Equations

☒	0
☐	1
☐	2
☐	3

page 117

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 63 sec(s)

MTH621: Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 9 of 10 (Start time: 09:48:57 PM, 18 January 2018)

Total Marks: 1

For the function $f(x) = x \sin \frac{1}{x}$, $x \neq 0$, which statement is true

Select the correct option

Reload Math Equations



$$\lim_{x \rightarrow 1/\pi} f(x) = 0.$$



The function is not defined at

$$x = 0.$$



$$\lim_{x \rightarrow 0} f(x) = 1.$$



$$\lim_{x \rightarrow 0} f(x) = 0.$$

page 78

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 50 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 4 of 10 (Start time: 07:49:47 PM, 16 January 2018)

Total Marks: 1

For the function

$$f(x) = x \sin \frac{1}{x}, \quad x \neq 0,$$

Select the correct option

Reload Math Equations



$$\lim_{x \rightarrow 1/\pi} f(x) = 0.$$

The function is not defined at $x = 0$.

$$\lim_{x \rightarrow 0} f(x) = 1.$$



$$\lim_{x \rightarrow 0} f(x) = 0.$$

rep

Click to Save Answer & Move to Next Question

Time Left 62 sec(s)

Quiz Start Time: 09:46 PM, 18 January 2018

Total Marks: 1

$$\begin{aligned} x^3, & \quad x \leq 0, \\ x^2 \sin \frac{1}{x}, & \quad x > 0, \end{aligned}$$

[Reload Math Equations](#)

☒	0	page 108
☐	2	
☐	1	
☐	-1	

[Click to Save Answer & Move to Next Question](#)

MC170200890: Awwal Khan

Time Left 70 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 7 of 10 (Start time: 10:33:35 PM, 06 February 2018)

Total Marks: 1

The inverse of the given function $f(x) = 2x + 4, 0 \leq x \leq 2$, is

Select the correct option

out



$$f^{-1}(y) = \frac{(y-4)}{2}$$



$$f^{-1}(y) = \frac{(y-2)}{4}$$



$$f^{-1}(y) = \frac{(x-2)}{4}$$



$$f^{-1}(y) = \frac{(y-4)}{4}$$

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 79
sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 3 of 10 (Start time: 10:32:11 PM, 06 February 2018)

Total Marks: 1

The function $g(x) = x^2$ is

Select the correct option

Reload Math Equations

decreasing on $[0, \infty)$.increasing on $[0, \infty)$.having negative values on $[0, \infty)$.undefined on $[0, \infty)$.

Click to Save Answer & Move to Next Question

page 97

MC170202886: Aqsa Zunaira

Time Left 83 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 6 of 10 (Start time: 09:47:44 PM, 18 January 2018)

Total Marks: 1

If $\lim_{x \rightarrow 0} f(x) = 10$ and $\lim_{x \rightarrow 0} g(x) = -2$ then $\lim_{x \rightarrow 0} \frac{f(x)}{g(x)}$ is

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|-----|
| ☐ | -10 |
| ☐ | 5 |
| ☐ | 0 |
| ☒ | -5 |

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 81 sec(s)

MTH621: Quiz 4 - Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 6 of 10 (Start time: 10:33:22 PM, 06 February 2018)

Total Marks: 1

Generalized Mean Value Theorem says. If f and g are continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) , then

Select the correct option

Reload Math Equations

page 112

for some c in (a, b) .

$$[g(b) - g(a)]f'(c) = [f(b) - f(a)]g'(c)$$

for some c in (a, b) .

$$[g(b) - g(a)] = [f(b) - f(a)]g'(c)$$

 $g'(x) = 0$ for all x in (a, b) . $g'(x) = f'(x)$ for all x in (a, b) .

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 84 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 7 of 10 (Start time: 10:05:25 PM, 06 February 2018)

Total Marks: 1

The derivative of the function x^n is

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|---------------------|
| ☐ | x^{n-1} |
| ☒ | nx^{n-1} |
| ☐ | $\frac{n}{x^{n-1}}$ |
| ☐ | nx^n |

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 40 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 7 of 10 (Start time: 07:52:40 PM, 16 January 2018)

Total Marks: 1

For the function defined as

$$f(x) = cx,$$

for every $\epsilon > 0$ the formal definition ensures

Select the correct option

Reload Math Equations

- ☐ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } 0 < \delta < \epsilon.$
- ☒ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } 0 < \delta < \epsilon/|c|.$
- ☐ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } -\epsilon < \delta < \epsilon.$
- ☐ $|f(x) - cx_0| < \epsilon, \quad |x - x_0| < \delta, \text{ where } 0 < \delta < \epsilon.$

page 78



Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 80
sec(s)

MTH621: Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 4 of 10 (Start time: 09:47:08 PM, 18 January 2018)

Total Marks: 1

If $\lim_{x \rightarrow 0} f(x) = 10$ and $\lim_{x \rightarrow 0} g(x) = -2$ then $\lim_{x \rightarrow 0} \left(\frac{f(x)}{g(x)} \right)$ is

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|------|
| ☐ | -10. |
| ☐ | 5. |
| ☐ | 0. |
| ☒ | -5. |
- 

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 67 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 5 of 10 (Start time: 10:32:56 PM, 06 February 2018)

Total Marks: 1

The function $f(x) = \begin{cases} x, & 0 \leq x < 1, \\ 2, & 1 \leq x \leq 2, \end{cases}$ is

Select the correct option

Reload Math Equations

- | | | |
|----------------------------------|------------------------------|----------------|
| ☒ | is nondecreasing on | $I = [0, 2]$. |
| ☐ | is nonincreasing on | $I = [0, 2]$. |
| ☐ | is having negative values on | $I = [0, 2]$. |
| ☐ | is undefined on | $I = [0, 2]$. |

page 97

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 61 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 6 of 10 (Start time: 10:04:50 PM, 06 February 2018)

Total Marks: 1

The function $f(x) = \begin{cases} x, & 0 \leq x < 1, \\ 2, & 1 \leq x \leq 2, \end{cases}$ is

Select the correct option

Reload Math Equations

page 97



is nondecreasing on

$$I = [0, 2] .$$



is nonincreasing on

$$I = [0, 2] .$$



is having negative values on

$$I = [0, 2] .$$



is undefined on

$$I = [0, 2] .$$

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 84 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 9 of 10 (Start time: 10:34:12 PM, 06 February 2018)

Total Marks: 1

The inverse of the function $f(x) = x^2$, is

Select the correct option

Reload Math Equations



$$f^{-1}(y) = y^2$$



$$f^{-1}(y) = \sqrt{y}.$$

page 101



$$f^{-1}(y) = \frac{1}{x^3}$$



$$f^{-1}(y) = \frac{1}{y^2}$$

Click to Save Answer & Move to Next Question



MC170202886: Aqsa Zunaira

Time Left 69 sec(s)

MTH621: Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 1 of 10 (Start time: 09:46:10 PM, 18 January 2018)

Total Marks: 1

If $f(x) = \log x$ and $g(x) = \frac{1}{1-x^2}$, then $f \circ g$ is

Select the correct option

Reload Math Equations



$$(f \circ g)(x) = \log \frac{1}{1-x^2}.$$



$$(f \circ g)(x) = \log \frac{x}{1-x^2}.$$



$$(f \circ g)(x) = \log \frac{1}{x^2}.$$



$$(f \circ g)(x) = \log \frac{-1}{1-x^2}.$$

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 64 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 4 of 10 (Start time: 10:32:26 PM, 06 February 2018)

Total Marks: 1

Rolle's Theorem says. suppose that f is continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) and $f(a) = f(b)$.

Select the correct option

Reload Math Equations

page 111

Then $f'(c) = 0$ for some c in the open interval (a, b) .Then $f'(c) \neq 0$ for some c in the open interval (a, b) .Then $f'(c)$ exists for some c in the open interval (a, b) .Then $f'(c) = 0$ for some c in the open interval (a, ∞) .

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 38 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 9 of 10 (Start time: 02:14:42 PM, 28 November 2017)

Total Marks: 1

Every Cauchy sequence has a

Select the correct option

- ☒ convergent subsequence.
- ☐ increasing subsequence.
- ☐ decreasing subsequence.
- ☐ positive subsequence.

by google uncle

MC170201745: Shakeel Ahmad

Time Left 61 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 09:37 PM, 06 February 2018

Question # 9 of 10 (Start time: 09:41:59 PM, 06 February 2018)

Total Marks: 1

Generalized Mean Value Theorem says. If f and g are continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) , then

Select the correct option

Reload Math Equations

page 112

for some c in (a, b) .

$$[g(b) - g(a)]f'(c) = [f(b) - f(a)]g'(c)$$

for some c in (a, b) .

$$[g(b) - g(a)] = [f(b) - f(a)]g'(c)$$

 $g'(x) = 0$ for all x in (a, b) . $g'(x) = f'(x)$ for all x in (a, b) .

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 82 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 2 of 10 (Start time: 09:46:36 PM, 18 January 2018)

Total Marks: 1

The $\lim_{x \rightarrow 2} (3x - 5) =$

Select the correct option

Reload Math Equations

☐	0.
☒	1.
☐	-1.
☐	-2.

page 78

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 56 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 6 of 10 (Start time: 02:11:19 PM, 28 November 2017)

Total Marks: 1

$\limsup_{n \rightarrow \infty} s_n = \infty$ if

Select the correct option

Reload Math Equations

- ☒ $\{s_n\}$ is not bounded above.
- ☐ $\{s_n\}$ has unique limit.
- ☐ $\{s_n\}$ is convergent.
- ☐ $\{s_n\}$ has limit inferior.

page 40

MC170202886: Aqsa Zunaira

Time Left 78 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 09:46 PM, 18 January 2018

Question # 10 of 10 (Start time: 09:49:37 PM, 18 January 2018)

Total Marks: 1

The value of $x + \lim_{x \rightarrow 0^+} \frac{|x|}{x} =$

3rd chptr

Select the correct option

Reload Math Equations



$x - 1.$



$x + 1.$



$x.$



$x + 1/x.$

Click to Save Answer & Move to Next Question



MC170201745: Shakeel Ahmad

Time Left 77 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 5 of 10 (Start time: 07:51:09 PM, 16 January 2018)

Total Marks: 1

For the function $f(x) = x \sin \frac{1}{x}$, $x \neq 0$, which statement is true

Select the correct option

Reload Math Equations

- ☐ $\lim_{x \rightarrow 1/\pi} f(x) = 0.$
- ☐ The function is not defined at $x = 0.$
- ☐ $\lim_{x \rightarrow 0} f(x) = 1.$
- ☒ $\lim_{x \rightarrow 0} f(x) = 0.$

page 78

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 84 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 1 of 10 (Start time: 10:31:47 PM, 06 February 2018)

Total Marks: 1

$\lim_{x \rightarrow 0^+} x \log x$, has the following indeterminate form

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|--------------------------|
| ☐ | $\infty \times \infty$. |
| ☒ | $(0)(\infty)$. |
| ☐ | $\frac{0}{0}$. |
| ☐ | 0^∞ . |

page 117

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 82 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 8 of 10 (Start time: 10:05:33 PM, 06 February 2018)

Total Marks: 1

$\lim_{x \rightarrow 0^+} x \log x$, has the following indeterminate form

Select the correct option

Reload Math Equations

- | | |
|----------------------------------|--------------------------|
| ☐ | $\infty \times \infty$. |
| ☒ | $0)(\infty)$. |
| ☐ | $\frac{0}{0}$. |
| ☐ | 0^∞ . |

page 117

Click to Save Answer & Move to Next Question

MC170202886: Aqsa Zunaira

Time Left 68 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:03 PM, 06 February 2018

Question # 1 of 10 (Start time: 10:03:05 PM, 06 February 2018)

Total Marks: 1

If a function f is continuous on the closed interval then f attains its

Select the correct option

Reload Math Equations

page 110



extreme values in the closed interval.



extreme values at one point in the closed interval.



may or may not attain extreme values in the closed interval.



derivative in the closed interval.

Click to Save Answer & Move to Next Question

MC170200890: Awwal Khan

Time Left 80 sec(s)

MTH621:Quiz 4 -Lectures 30 to 37

Quiz Start Time: 10:31 PM, 06 February 2018

Question # 8 of 10 (Start time: 10:33:58 PM, 06 February 2018)

Total Marks: 1

If a function f is continuous on the closed interval then f attains its

Select the correct option

Reload Math Equations

page 110



extreme values in the closed interval.



extreme values at one point in the closed interval.



may or may not attain extreme values in the closed interval.



derivative in the closed interval.

Click to Save Answer & Move to Next Question

MC170201745: Shakeel Ahmad

Time Left 76 sec(s)

MTH621:Quiz 3 - Lecture 23 to 27

Quiz Start Time: 07:47 PM, 16 January 2018

Question # 8 of 10 (Start time: 07:53:43 PM, 16 January 2018)

Total Marks: 1

The value of $x + \lim_{x \rightarrow 0^+} \frac{|x|}{x} =$

Reload Math Equations

Select the correct option



$x - 1.$



$x + 1.$



$x.$



$x + 1/x.$

Chptr 3

Click to Save Answer & Move to Next Question

MC170201745: Shakool Ahmad

Time Left 38 sec(s)

MTH621: Quiz 2

Quiz Start Time: 02:06 PM, 28 November 2017

Question # 10 of 10 (Start time: 02:15:53 PM, 28 November 2017)

Total Marks: 1

The limit of the sequence

$$s_n = \frac{1}{n} + \frac{2(1 + 3/n)}{1 + 1/n}.$$

Select the correct option

Reload Math Equations

- ☐ 1
- ☐ 4
- ☐ 3
- ☒ 2

page 35 and 36