



VU NET



DOWNLOAD PAST PAPERS

vunet.blogspot.com

The probability of drawing a 'white' ball from a bag containing 4 red, 8 black and 3 white balls is:

Answer (Please select your correct option)

☐

0

☐ $\frac{3}{15}$

correct answer solve
by hadi

☐ $\frac{1}{12}$ ☐ $\frac{1}{2}$

Made By: Waqar Siddhu



If the probability of one event does not affected by the occurrence of another event then both events are:

Answer (Please select your correct option)

☐ Dependent

☐ Not Mutually Exclusive

☐ Mutually Exclusive

☐ Independent

correct answer solve
by hadi

Made By: Waqar Siddhu



A die is rolled. What is the probability of getting a number that is an even number and greater than 2.

Answer (Please select your correct option)

☐ $1/2$

☐ $1/3$

☐ $2/3$

☐ $5/6$

correct answer solve
by hadi

Made By: Waqar Siddhu



The collection of all the elements under study is called:

Answer (Please select your correct option)



Population



Sample

correct answer solve
by hadi



Data



Registration

Made By: Waqar Siddhu



The mode value from raw data can be obtained by the help of

Answer (Please select your correct option)

- ☐ Dot plot
- ☐ Stem and leaf plot
- ☐ Bar chart
- ☐ None of these



A student solved 25 questions from first 50 questions of a book . The probability that he will solve the remaining all questions is:

Answer (Please select your correct option)

☐ 0.25☐ 0.5

correct answer solve
by hadi

☐ 1☐ 0



The difference between the largest and the smallest data values is called the

Answer (Please select your correct option)



Variance



Interquartile range



Range

correct answer solve
by hadi



Coefficient of variation

Made By: Waqar Siddhu



Which one is the formula of mid- quartile range:

Answer (Please select your correct option)

☐ $Q_3 - Q_1$

☐ $\frac{Q_3 - Q_1}{2}$

☐ $\frac{Q_1 - Q_3}{2}$

☐ $\frac{Q_1 + Q_3}{2}$

correct answer solve
by hadi



For the given data 2, 3, 7, 0, -8; G. M will be:

Answer (Please select your correct option)

☐ Negative

☐ Positive

☐ Zero

correct answer solve
by hadi

☐ Undefined



Mean deviation is least when it is calculated from _____.

Answer (Please select your correct option)

☐

Mean

☐

Mode

☐

Median

☐

Geometric mean



If X and Y are independent variables then $\text{Var}(X-Y) =$ _____

Answer (Please select your correct option)

☐ $\text{Var}(X) - \text{Var}(Y)$

correct answer solve
by hadi

☐ $\text{Var}(X) + \text{Var}(Y)$

☐ $\text{Var}(X+Y)$

☐ $\text{Var}(X) \times \text{Var}(Y)$



Rang measures the dispersion around the _____:

Answer (Please select your correct option)



Extreme values



Minimum value



Maximum value



Mid-range value



The variable plotted on the horizontal or X-axis in a scatter diagram is called the:

Answer (Please select your correct option)



Scatter variable

correct answer solve
by hadi



Independent variable



Dependent variable



Correlation variable

Made By: Waqar Siddhu



In a scatter diagram we plot:

Answer (Please select your correct option)

☐ X and Y

☐ X and $\hat{Y}$

☐ Y and $\hat{Y}$

☐ $\hat{Y}$ and X



If $P(A \cap B) = P(A|B) \times P(B)$ then A and B will be:

Answer (Please select your correct option)

☐

Independent

correct answer solve
by hadi

☐

Mutually Exclusive

☐

Dependent

☐

Equally Likely

Made By: Waqar Siddhu



Which one of the following is the class frequency?

Answer (Please select your correct option)

☐

The number of observations in each class

correct answer solve
by hadi

☐

The difference between consecutive lower class limits

☐

Always contains at least 5 observations

☐

Usually a multiple of the lower limit of the first class

Made By: Waqar Siddhu



How to convert a frequency distribution to a relative frequency distribution?

Answer (Please select your correct option)

☐ Find the difference between consecutive lower class limits

☐ Find the difference between consecutive lower limits

correct answer solve
by hadi...check it

☐ Count the number of observations in the class

☐ Divide the class frequency by the total number of observations



In statistics, we deal with:

Answer (Please select your correct option)

- ☐ Individuals
- ☐ Isolated items
- ☐ Aggregates of facts
- ☐ Qualitative data

correct answer solve
by hadi

Made By: Waqar Siddhu



Pie chart consists of

Answer (Please select your correct option)

☐ A rectangular

☐ A triangle

☐ A square

☐ A circle

correct answer solve
by hadi

Made By: Waqar Siddhu



When $Q_1 = 2$ and $Q_3 = 4$, what is the value of Median, if the distribution is symmetrical:

Answer (Please select your correct option)

☐ 1

☐ 2

☐ 3

☐ 4

correct answer solve
by hadi



For two mutually exclusive events A and B, $P(A) = 0.2$ and $P(B) = 0.4$, then $P(A \cup B)$ is:

Answer (Please select your correct option)

☐ 0.8

☐ 0.2

☐ 0.6

☐ 0.5

correct answer solve
by hadi



When two coins are tossed the probability of at most one head is:

Answer (Please select your correct option)



1/4



2/4



3/4

correct answer solve
by hadi



4/4



A value which has half of the observations above it and half of the observations below it; is known as.....

Answer (Please select your correct option)



Rang



Median

correct answer solve
by hadi



Mean



Mode



In a set of 20 values all the values are 10, what will be median?

Answer (Please select your correct option)

☐ 2☐ 5☒ 10☐ 20

correct answer solve
by hadi

Made By: Waqar Siddhu



Which one of the following measure is not based on all the observations?

Answer (Please select your correct option)



Arithmetic Mean



Geometric Mean



Harmonic Mean



Mode

correct answer solve
by hadi

Made By: Waqar Siddhu



The collection of all the elements under study is called:

Answer (Please select your correct option)



Population



Sample

correct answer solve
by hadi



Data



Registration

Made By: Waqar Siddhu



A five-number summary consists of :

Answer (Please select your correct option)



X_0 , Q_1 , Median, Q_3 , and X_m

correct answer solve
by hadi



X_m , Q_1 , Mean, Q_3 , and X_0



X_m , Q_1 , Mode, Q_3 , and X_0



X_0 , Q_1 , Median, Q_2 , and X_m



For what condition of k , at least $1 - 1/k^2$ of the data-values fall within k standard deviations of the mean:

Answer (Please select your correct option)



Greater than 1

correct answer solve
by hadi



Less than 1



Greater or equal to 1



Less or equal to 1



Which of the following can never be taken as the probability of an event?

Answer (Please select your correct option)

☐ 1

☐ 0

☐ 0.5

☐ -0.5

correct answer solve
by hadi

Made By: Waqar Siddhu



If E is an impossible event, then $P(E)$ is:

Answer (Please select your correct option)

☐ 1

☐ 0.15

☐ 0

☐ 0.5

correct answer solve
by hadi

Made By: Waqar Siddhu



The average which is defined as the reciprocal of the arithmetic mean of the reciprocals of the values is called:

Answer (Please select your correct option)



Geometric Mean



Harmonic Mean

correct answer solve
by hadi



Mode



Median

Made By: Waqar Siddhu



An automobile is running, during the first 60 Km, at the rate of 10 Km/hr. During the second 60 Km at the rate of 30Km/hr, while during the third 60 Km its speed was 40 Km/hr. What method is more appropriate to calculate the average speed?

Answer (Please select your correct option)

☐ Median

☐ Arithmetic mean

☐ Harmonic mean

☐ Geometric mean

correct answer solve
by hadi

Made By: Waqar Siddhu



What is the standard deviation of the following values?

7, 7, 7, 7, 7, 7, 7

Answer (Please select your correct option)

7



1



49



0



correct answer solve
by hadi

Made By: Waqar Siddhu



Which one of the following is a meso-kurtic curve?

Answer (Please select your correct option)

☐

Negatively skewed

☐

Positively skewed

☐

J-shaped

☐

Normal

correct answer solve
by hadi

Made By: Waqar Siddhu



Men tend to marry women who are slightly younger than themselves. Suppose that every man married a woman who was exactly 5 years younger than themselves. Which of the following is correct?

Answer (Please select your correct option)

☐ The correlation coefficient is -5

☐ The correlation coefficient is 5

☐ The correlation coefficient is 1

☐ The correlation coefficient is 0



In a multiplication theorem $P(A \cap B)$ equals:

Answer (Please select your correct option)



$$P(A) P(B)$$



$$P(A) + P(B)$$



$$P(A) \times P(B|A)$$

correct answer solve
by hadi



$$P(B|A) \times P(B)$$



Which one of the following is the class frequency?

Answer (Please select your correct option)

☐

The number of observations in each class

correct answer solve
by hadi

☐

The difference between consecutive lower class limits

☐

Always contains at least 5 observations

☐

Usually a multiple of the lower limit of the first class

Made By: Waqar Siddhu



Which of the following comes first to make frequency distribution.

Answer (Please select your correct option)

☐ Number of Groups

☐ Class interval

☐ Range

correct answer solve
by hadi

☐ Tally marks

Made By: Waqar Siddhu



Data obtained from the Ministry of food, is an example of:

Answer (Please select your correct option)

☐ Primary data

☐ Secondary data

☐ Qualitative data

☐ None of these

correct answer solve
by hadi



Total angle of a Pie Chart is:

Answer (Please select your correct option)



90°



180°



270°



360°

correct answer solve
by hadi

Made By: Waqar Siddhu



The probability of drawing a 'white' ball from a bag containing 4 red, 8 black and 3 white balls is:

Answer (Please select your correct option)

☐

0

☐ $\frac{3}{15}$ correct☐ $\frac{1}{12}$ ☐ $\frac{1}{2}$



For two mutually exclusive events A and B, $P(A) = 0.2$ and $P(B) = 0.4$, then $P(A \cup B)$ is:

Answer (Please select your correct option)

☐ 0.8

☐ 0.2

☒ 0.6

correct

☐ 0.5



Which one is the formula of range:

Answer (Please select your correct option)

☐ $x_m - x_0$ correct

☐ $x_0 - x_m$

☐ $\frac{x_0 - x_m}{2}$

☐ $\frac{x_0 + x_m}{2}$



If X and Y are independent variables then $\text{Var}(X-Y) = \underline{\hspace{2cm}}$

Answer (Please select your correct option)

☐ $\text{Var}(X) - \text{Var}(Y)$

correct

☐ $\text{Var}(X) + \text{Var}(Y)$

☐ $\text{Var}(X+Y)$

☐ $\text{Var}(X) \times \text{Var}(Y)$



If a player well shuffles the pack of 52 playing card, then the probability of a black card from 52 playing cards is:

Answer (Please select your correct option)



$1/52$



$13/52$



$4/52$



$26/52$

correct

Made By: Waqar Siddhu



Pearson's coefficient of skewness is not possible if standard deviation is:

Answer (Please select your correct option)

☐ 1

☐ 1.5

☐ 0

☐ 3



Men tend to marry women who are slightly younger than themselves. Suppose that every man married a woman who was exactly 5 years younger than themselves. Which of the following is correct?

Answer (Please select your correct option)



The correlation coefficient is -5



The correlation coefficient is 5



The correlation coefficient is 1



The correlation coefficient is 0



What is the difference between permutation and combination?

Answer (Please select your correct option)

☐ In permutation order is important and in combination order is not important

correct

☐ In permutation order is not important and in combination order is important

☐ Only combination is based on the classical definition of probability

☐ Only permutation is based on the classical definition of probability

Made By: Waqar Siddhu



If data is categorized about any characteristic, then which measurement scale should be used?

Answer (Please select your correct option)

☐ Ordinal

☐ Interval

☒ Nominal

correct

☐ Ratio



Which of the following scale has true zero point?

agr not true zero ho
to then interval
hoga

Answer (Please select your correct option)

Ratio Scale



correct

Interval scale



Nominal scale



Ordinal scale



Made By: Waqar Siddhu



In a simple bar chart:

Answer (Please select your correct option)

☐ Width of bar is meaningful

☐ Height of bar is meaningful

☐ Height of bar is not meaningful

☐ Height and width both are meaningful

correct



What is the name of following curve?



Answer (Please select your correct option)

☐ Positively skewed curve

correct answer solve
by hadi

☐ Symmetrical curve

☐ Normal curve

☐ Negatively skewed curve



If $A = \{1, 2, 3, 4, 5, 10\}$ and $B = \{1, 3, 5\}$ then $B \subset A$ means:

Answer (Please select your correct option)



A is less than B



A is contained in B



B is contained in A.

correct



B is less than A



The graph of cumulative frequency distribution is called:

Answer (Please select your correct option)

☐ Frequency polygon

☒ Ogive

☐ Frequency curve

☐ Histogram

correct



Given the series 1,2,1,1,2,2,2,2,3,4,5,3,2,3,1,4,2,3. Which one of the following is mode of the given seires:

Answer (Please select your correct option)

☐ 4

☐ 3
3

☐ 2
3
2

☐ 1
3
2*



Which one of the following is equal to the 2nd quartile:

Answer (Please select your correct option)

 P_{33}  D_3 

Median



Mode

correct



The variance of a sample of 81 observations equal to 64. The standard deviation of the sample is equal to:

Answer (Please select your correct option)

☐ 256☐ 4096☐ 8☐ 6,561



Under which condition standard deviation could assume a negative value:

Answer (Please select your correct option)



When more than half of the data values are negative

correct answer solve
by hadi



If all the data values are same



The standard deviation cannot be negative



When all the data values are negative



In a set of 20 values all the values are 2, what will be the Geometric Mean?

Answer (Please select your correct option)

☐ 2

correct

☐ 5

☐ 10

☐ 20



In a set of 10 values all the values are 5, what will be the D_5 ?

Answer (Please select your correct option)

☐ 2

☒ 5

☐ 2
5
10

☐ 20

correct



When three coins are tossed simultaneously, P (3 Heads) is:

Answer (Please select your correct option)

☐ $\frac{3}{8}$

correct

☐ $\frac{1}{8}$

☐ $\frac{4}{8}$

☐ $\frac{2}{8}$



The distribution is mesokurtic if the moment ratio i.e. b_2 is:

Answer (Please select your correct option)



Equal to 0



Equal to 3

correct



Less than 3



Greater than 3



A set is any well-defined collection of

Answer (Please select your correct option)



Positive Objects



Negative Objects



Same Objects



Distinct Objects

correct

Made By: Waqar Siddhu



If an observation in the data set is negative, then the geometric mean is:

Answer (Please select your correct option)

☐ Positive

☐ negative

☐ Zero

☐ undetermined



Mean deviation is least when it is calculated from _____.

Answer (Please select your correct option)



Mean



Mode



Median



Geometric mean



If a player well shuffles the pack of 52 playing card, then the probability of a black card from 52 playing cards is:

Answer (Please select your correct option)



1/52



13/52



4/52



26/52

correct

Made By: Waqar Siddhu



For a positively skewed distribution m_3 will be:

Answer (Please select your correct option)

☐ Positive

correct

☐ Negative

☐ Zero

☐ 1



Which of the following value could not represent a coefficient of correlation?

Answer (Please select your correct option)

☐

$$r = 0.99$$

☐

$$r = 1.09$$

☐

$$r = -0.73$$

☐

$$r = -1$$



In a scatter diagram we plot:

Answer (Please select your correct option)



X and Y

correct answer solve
by hadi



X and $\hat{Y}$



Y and $\hat{Y}$



$\hat{Y}$ and X



The conditional probability is $P(A|B)$ is given by formula:

Answer (Please select your correct option)

☐

$$\frac{P(A \cup B)}{P(A)}$$

☐

$$\frac{P(A \cap B)}{P(B)}$$

correct

☐

$$\frac{P(A \cap B)}{P(A)}$$

☐

$$\frac{P(A \cup B)}{P(B)}$$



If a curve has a longer tail to the right, it is called :

Answer (Please select your correct option)

☐ Positively skewed

correct

☐ Negatively skewed

☐ J-shaped

☐ Symmetric



Which one of the following is the class frequency?

Answer (Please select your correct option)

☐

The number of observations in each class

correct

☐

The difference between consecutive lower class limits

☐

Always contains at least 5 observations

☐

Usually a multiple of the lower limit of the first class



Which of the following scale has not true zero point?

Answer (Please select your correct option)



Nominal scale



Ordinal scale



Interval scale



Ratio scale

correct

Made By: Waqar Siddhu

What is the name of following curve?



Answer (Please select your correct option)

☐ Positively skewed curve

correct

☐ Symmetrical curve

☐ Normal curve

☐ Negatively skewed curve



A set that contains all possible outcomes of a system is known as

Answer (Please select your correct option)

☐

Finite Set

☐

Infinite Set

☐

Universal Set

☐

None of these

correct

Made By: Waqar Siddhu



The graph of cumulative frequency distribution is called:

Answer (Please select your correct option)

☐ Frequency polygon

☒ Ogive

☐ Frequency curve

☐ Histogram

correct



Mean is affected by the change of

Answer (Please select your correct option)



Origin Only



Scale Only

correct answer solve
by hadi



Origin and Scale



Not affected



What will be the mode for the data 2,3,4,5,4,6,4?

Answer (Please select your correct option)

☐ 2

☐ 3

☒ 4

☐ 5

correct



In a set of 20 values all the values are 5, what will be the mean?

Answer (Please select your correct option)

4



5



10



20



correct

Made By: Waqar Siddhu



It is easy to find the median when data is.....

Answer (Please select your correct option)

☐ Arranged

correct

☐ Discrete

☐ Attributed

☐ Continuous



A letter is chosen at random from the word STATISTICS. The probability of getting a vowel is:

Answer (Please select your correct option)



3/10

correct answer solve
by hadi



4/10



5/10



6/10

Made By: Waqar Siddhu



When two coins are tossed simultaneously, P (one head) is:

Answer (Please select your correct option)

☐ $\frac{1}{4}$

☐ $\frac{1}{2}$

☐ $\frac{3}{4}$

☐ 1

correct answer solve
by hadi

Made By: Waqar Siddhu



A value which has half of the observations above it and half of the observations below it, is known as.....

Answer (Please select your correct option)



Rang



Median

correct answer solve
by hadi



Mean



Mode

Made By: Waqar Siddhu



In a set of 20 values all the values are 2, what will be the Geometric Mean?

Answer (Please select your correct option)

☐ 2

correct answer solve
by hadi

☐ 5

☐ 10

☐ 20

Made By: Waqar Siddhu



Which of the following measure of dispersion is least affected by the extreme values in a data?

Answer (Please select your correct option)



Range

correct answer solve
by hadi



Quartile deviation



Mean absolute deviation



Standard deviation



Which is appropriate average for finding the average speed of a car:

Answer (Please select your correct option)

Mean



Geometric mean



Harmonic mean



Weighted mean



correct answer solve
by hadi

Made By: Waqar Siddhu



What will be the value of the total amount of dispersion obtained by summing the $(X - \bar{X})$?

Answer (Please select your correct option)

- ☐ 0 correct answer solve by hadi
- ☐ -1
- ☐ 1
- ☐ 100



Rang measures the dispersion around the _____:

Answer (Please select your correct option)



Extreme values

correct answer solve
by hadi



Minimum value



Maximum value



Mid-range value



The deviation of a distribution from symmetry is called:

Answer (Please select your correct option)



Kurtosis



Skewness



Dispersion

correct answer solve
by hadi



Flatness

Made By: Waqar Siddhu



For the independent events A and B if $P(A) = 0.25$, $P(B) = 0.40$ then $P(A \cap B) = \dots\dots\dots$

Answer (Please select your correct option)

☐ 0.65☐ 0.1

correct answer solve
by hadi

☐ 0.50☐ 0.15



The variable plotted on the vertical or Y-axis in a scatter diagram is called the:

Answer (Please select your correct option)

☐ Scatter variable

correct answer solve
by hadi

☐ Independent variable

☐ Dependent variable

☐ Correlation variable



In a multiplication theorem $P(A \cap B)$ equals:

Answer (Please select your correct option)



$P(A) P(B)$



$P(A) + P(B)$



$P(A) \times P(B|A)$

correct answer solve
by hadi



$P(B|A) \times P(B)$

Made By: Waqar Siddhu



For two independent events A and B, if $P(A) = 0.2$ and $P(B) = 0.4$, then what is $P(A \cap B)$?

Answer (Please select your correct option)



0.06



0.08



0.02



0.04

correct answer solve
by hadi

Made By: Waqar Siddhu



The level of satisfaction for a certain facility which Govt. is providing is the example of

Answer (Please select your correct option)

☐

Constant

☐

Qualitative variable

correct answer solve
by hadi

☐

Variable

☐

Quantitative variable

Made By: Waqar Siddhu



Quantitative variable is further divided into:

Answer (Please select your correct option)



Continuous variable



Discrete variable

correct answer solve
by hadi



Continuous & Discrete variable



None of the above



If you connect the mid-points of rectangles in a histogram by a series of lines that also touches the x-axis from both ends, what will you get?

Answer (Please select your correct option)



Ogive



Frequency polygon



Frequency curve

correct answer solve
by hadi



Historigram

Made By: Waqar Siddhu



Which type of data is collected in population census?

Answer (Please select your correct option)



Grouped data

correct answer solve
by hadi,,but check
it



Secondary data



Primary data



Array data

Made By: Waqar Siddhu



Data obtained from the Bureau of Statistics, is an example of:

Answer (Please select your correct option)



None of these



Primary data



Secondary data

correct answer solve
by hadi



Qualitative data

Made By: Waqar Siddhu



A set that contains all possible outcomes of a system is known as

Answer (Please select your correct option)



Finite Set



Infinite Set



Universal Set

correct answer solve
by hadi



None of these

Made By: Waqar Siddhu



In positively skewed distribution, which of the following relationship exists:

Answer (Please select your correct option)



The mean, median, and mode are all equal



The mean is larger than the median



The median is larger than the mean



The standard deviation must be larger than the mean or the median

The probability of drawing a 'white' ball from a bag containing 4 red, 8 black and 3 white balls is:

Answer (Please select your correct option)

☐

0

☒ $\frac{3}{15}$

correct

☐ $\frac{1}{12}$ ☐ $\frac{1}{2}$



If the probability of one event does not affected by the occurrence of another event then both events are:

Answer (Please select your correct option)



Dependent



Not Mutually Exclusive



Mutually Exclusive



Independent

correct

Made By: Waqar Siddhu



A die is rolled. What is the probability of getting a number that is an even number and greater than 2.

Answer (Please select your correct option)

☐ $1/2$ ☐ $1/3$ ☒ $2/3$ ☐ $5/6$

correct



The collection of all the elements under study is called:

Answer (Please select your correct option)



Population



Sample



Data



Registration

correct

Made By: Waqar Siddhu



The mode value from raw data can be obtained by the help of

Answer (Please select your correct option)



Dot plot



Stem and leaf plot



Bar chart



None of these



A student solved 25 questions from first 50 questions of a book . The probability that he will solve the remaining all questions is:

Answer (Please select your correct option)



0.25



0.5

correct



1



0



The difference between the largest and the smallest data values is called the

Answer (Please select your correct option)



Variance



Interquartile range



Range



Coefficient of variation

correct

Made By: Waqar Siddhu

Which one is the formula of mid- quartile range:

Answer (Please select your correct option)

☐ $Q_3 - Q_1$

☐ $\frac{Q_3 - Q_1}{2}$

☐ $\frac{Q_1 - Q_3}{2}$

☐ $\frac{Q_1 + Q_3}{2}$

correct



For the given data 2, 3, 7, 0, -8; G. M will be:

Answer (Please select your correct option)



Negative



Positive



Zero



Undefined

correct

Made By: Waqar Siddhu



Mean deviation is least when it is calculated from _____.

Answer (Please select your correct option)

☐ Mean

☐ Mode

☐ Median

☐ Geometric mean

Made By: Waqar Siddhu

If X and Y are independent variables then $\text{Var}(X-Y) =$ _____

Answer (Please select your correct option)

☐

$\text{Var}(X) - \text{Var}(Y)$

correct

☐

$\text{Var}(X) + \text{Var}(Y)$

☐

$\text{Var}(X+Y)$

☐

$\text{Var}(X) \times \text{Var}(Y)$

Made By: Waqar Siddhu



Rang measures the dispersion around the _____:

Answer (Please select your correct option)



Extreme values



Minimum value



Maximum value



Mid-range value

correct

Made By: Waqar Siddhu



The variable plotted on the horizontal or X-axis in a scatter diagram is called the:

Answer (Please select your correct option)

☐ Scatter variable

correct

☐ Independent variable

☐ Dependent variable

☐ Correlation variable

In a scatter diagram we plot:

Answer (Please select your correct option)

☐ X and Y

☐ X and $\hat{Y}$

☐ Y and $\hat{Y}$

☐ $\hat{Y}$ and X



If $P(A \cap B) = P(A|B) \times P(B)$ then A and B will be:

Answer (Please select your correct option)

☐ Independent

correct

☐ Mutually Exclusive

☐ Dependent

☐ Equally Likely

Which one of the following is the class frequency?

Answer (Please select your correct option)

☐

The number of observations in each class

correct

☐

The difference between consecutive lower class limits

☐

Always contains at least 5 observations

☐

Usually a multiple of the lower limit of the first class

Made By: Waqar Siddhu



How to convert a frequency distribution to a relative frequency distribution?

Answer (Please select your correct option)



Find the difference between consecutive lower class limits



Find the difference between consecutive lower limits



Count the number of observations in the class



Divide the class frequency by the total number of observations



In statistics, we deal with:

Answer (Please select your correct option)



Individuals



Isolated items



Aggregates of facts



Qualitative data

correct

Made By: Waqar Siddhu



Pie chart consists of

Answer (Please select your correct option)



A rectangular



A triangle



A square



A circle

correct

Made By: Waqar Siddhu



When $Q_1 = 2$ and $Q_3 = 4$, what is the value of Median, if the distribution is symmetrical:

Answer (Please select your correct option)

☐ 1

☐ 2

☒ 3

☐ 4

correct



A letter is chosen at random from the word STATISTICS. The probability of getting a vowel is:

Answer (Please select your correct option)



3/10

correct



4/10



5/10



6/10

Made By: Waqar Siddhu



A letter is chosen at random from the word STATISTICS. The probability of getting a vowel is:

Answer (Please select your correct option)



3/10

correct



4/10



5/10



6/10

Made By: Waqar Siddhu



When two coins are tossed simultaneously, P (one head) is:

Answer (Please select your correct option)

☐ $\frac{1}{4}$

☐ $\frac{1}{2}$

☐ $\frac{3}{4}$

☐ 1

correct

Made By: Waqar Siddhu



When two coins are tossed simultaneously, P (one head) is:

Answer (Please select your correct option)

☐ $\frac{1}{4}$

☐ $\frac{1}{2}$

☐ $\frac{3}{4}$

☐ 1

correct

Made By: Waqar Siddhu



A value which has half of the observations above it and half of the observations below it; is known as.....

Answer (Please select your correct option)



Rang



Median

correct



Mean



Mode

Made By: Waqar Siddhu



A value which has half of the observations above it and half of the observations below it; is known as.....

Answer (Please select your correct option)



Rang



Median

correct



Mean



Mode

Made By: Waqar Siddhu



In a set of 20 values all the values are 2, what will be the Geometric Mean?

Answer (Please select your correct option)

☒ 2

correct

☐ 5

☐ 10

☐ 20

Made By: Waqar Siddhu



In a set of 20 values all the values are 2, what will be the Geometric Mean?

Answer (Please select your correct option)

☒ 2

correct

☐ 5

☐ 10

☐ 20

Made By: Waqar Siddhu



Which of the following measure of dispersion is least affected by the extreme values in a data?

Answer (Please select your correct option)

Range



correct

Quartile deviation



Mean absolute deviation



Standard deviation



Made By: Waqar Siddhu



Which of the following measure of dispersion is least affected by the extreme values in a data?

Answer (Please select your correct option)

Range



correct

Quartile deviation



Mean absolute deviation



Standard deviation



Made By: Waqar Siddhu



Which is appropriate average for finding the average speed of a car:

Answer (Please select your correct option)

Mean



Geometric mean



Harmonic mean



Weighted mean



correct

Made By: Waqar Siddhu



Which is appropriate average for finding the average speed of a car:

Answer (Please select your correct option)

Mean



Geometric mean



Harmonic mean



Weighted mean



correct

Made By: Waqar Siddhu



What will be the value of the total amount of dispersion obtained by summing the $(X - \bar{X})$?

Answer (Please select your correct option)

☐ 0

correct

☐ -1

☐ 1

☐ 100



What will be the value of the total amount of dispersion obtained by summing the $(X - \bar{X})$?

Answer (Please select your correct option)

☐ 0

correct

☐ -1

☐ 1

☐ 100



Rang measures the dispersion around the _____:

Answer (Please select your correct option)



Extreme values

correct



Minimum value



Maximum value



Mid-range value

Made By: Waqar Siddhu



Rang measures the dispersion around the _____:

Answer (Please select your correct option)



Extreme values

correct



Minimum value



Maximum value



Mid-range value

Made By: Waqar Siddhu



The deviation of a distribution from symmetry is called:

Answer (Please select your correct option)



Kurtosis



Skewness



Dispersion



Flatness

correct

Made By: Waqar Siddhu



The deviation of a distribution from symmetry is called:

Answer (Please select your correct option)



Kurtosis



Skewness



Dispersion



Flatness

correct

Made By: Waqar Siddhu



For the independent events A and B if $P(A) = 0.25$, $P(B) = 0.40$ then $P(A \cap B) = \dots\dots\dots$

Answer (Please select your correct option)



0.65



0.1



0.50



0.15

correct

Made By: Waqar Siddhu



For the independent events A and B if $P(A) = 0.25$, $P(B) = 0.40$ then $P(A \cap B) = \dots\dots\dots$

Answer (Please select your correct option)

☐ 0.65

☐ 0.1

☐ 0.50

☐ 0.15

correct



The variable plotted on the vertical or Y-axis in a scatter diagram is called the:

Answer (Please select your correct option)

☐ Scatter variable

correct

☐ Independent variable

☐ Dependent variable

☐ Correlation variable

Made By: Waqar Siddhu



The variable plotted on the vertical or Y-axis in a scatter diagram is called the:

Answer (Please select your correct option)

☐ Scatter variable

correct

☐ Independent variable

☐ Dependent variable

☐ Correlation variable

Made By: Waqar Siddhu



In a multiplication theorem $P(A \cap B)$ equals:

Answer (Please select your correct option)



$P(A) P(B)$



$P(A) + P(B)$



$P(A) \times P(B|A)$



$P(B|A) \times P(B)$

correct

Made By: Waqar Siddhu



In a multiplication theorem $P(A \cap B)$ equals:

Answer (Please select your correct option)



$P(A) P(B)$



$P(A) + P(B)$



$P(A) \times P(B|A)$



$P(B|A) \times P(B)$

correct

Made By: Waqar Siddhu



For two independent events A and B, if $P(A) = 0.2$ and $P(B) = 0.4$, then what is $P(A \cap B)$?

Answer (Please select your correct option)

☐

0.06

☐

0.08

☐

0.02

☐

0.04

correct

Made By: Waqar Siddhu



For two independent events A and B, if $P(A) = 0.2$ and $P(B) = 0.4$, then what is $P(A \cap B)$?

Answer (Please select your correct option)

☐

0.06

☐

0.08

☐

0.02

☐

0.04

correct

Made By: Waqar Siddhu



The level of satisfaction for a certain facility which Govt. is providing is the example of:

Answer (Please select your correct option)

☐

Constant

☐

Qualitative variable

correct

☐

Variable

☐

Quantitative variable

Made By: Waqar Siddhu



The level of satisfaction for a certain facility which Govt. is providing is the example of:

Answer (Please select your correct option)

☐

Constant

☐

Qualitative variable

correct

☐

Variable

☐

Quantitative variable

Made By: Waqar Siddhu



Quantitative variable is further divided into:

Answer (Please select your correct option)

☐ Continuous variable

☒ Discrete variable

☐ Continuous & Discrete variable

☐ None of the above

correct



Quantitative variable is further divided into:

Answer (Please select your correct option)

☐ Continuous variable

☒ Discrete variable

☐ Continuous & Discrete variable

☐ None of the above

correct



If you connect the mid-points of rectangles in a histogram by a series of lines that also touches the x-axis from both ends, what will you get?

Answer (Please select your correct option)



Ogive



Frequency polygon



Frequency curve

correct answer solve
by hadi



Historigram

Made By: Waqar Siddhu



If you connect the mid-points of rectangles in a histogram by a series of lines that also touches the x-axis from both ends, what will you get?

Answer (Please select your correct option)



Ogive



Frequency polygon



Frequency curve



Historigram

Made By: Waqar Siddhu



Which type of data is collected in population census?

Answer (Please select your correct option)



Grouped data



Secondary data



Primary data



Array data

Made By: Waqar Siddhu



Which type of data is collected in population census?

Answer (Please select your correct option)

☐ Grouped data



☐ Secondary data

☐ Primary data

☐ Array data



Data obtained from the Bureau of Statistics, is an example of:

Answer (Please select your correct option)

☐ None of these

☐ Primary data

☒ Secondary data

☐ Qualitative data

correct

Made By: Waqar Siddhu



Data obtained from the Bureau of Statistics, is an example of:

Answer (Please select your correct option)



None of these



Primary data



Secondary data



Qualitative data

correct

Made By: Waqar Siddhu



A set that contains all possible outcomes of a system is known as

Answer (Please select your correct option)

☐

Finite Set

☐

Infinite Set

☒

Universal Set

☐

None of these

correct

Made By: Waqar Siddhu



A set that contains all possible outcomes of a system is known as

Answer (Please select your correct option)

☐

Finite Set

☐

Infinite Set

☒

Universal Set

☐

None of these

correct

Made By: Waqar Siddhu



In positively skewed distribution, which of the following relationship exists:

Answer (Please select your correct option)



The mean, median, and mode are all equal



The mean is larger than the median



The median is larger than the mean



The standard deviation must be larger than the mean or the median



In positively skewed distribution, which of the following relationship exists:

Answer (Please select your correct option)



The mean, median, and mode are all equal



The mean is larger than the median



The median is larger than the mean



The standard deviation must be larger than the mean or the median



For two mutually exclusive events A and B, $P(A) = 0.2$ and $P(B) = 0.4$, then $P(A \cup B)$ is:

Answer (Please select your correct option)

☐

0.8

☐

0.2

☒

0.6

correct

☐

0.5

Made By: Waqar Siddhu



When two coins are tossed the probability of at most one head is:

Answer (Please select your correct option)

☐ $1/4$

☐ $2/4$

☐ $3/4$

☐ $4/4$

correct

Made By: Waqar Siddhu



A value which has half of the observations above it and half of the observations below it; is known as.....

Answer (Please select your correct option)



Rang



Median



Mean



Mode

correct

Made By: Waqar Siddhu



In a set of 20 values all the values are 10, what will be median?

Answer (Please select your correct option)

☐ 2

☐ 5

☐ 10

☐ 20

correct

Made By: Waqar Siddhu



Which one of the following measure is not based on all the observations?

Answer (Please select your correct option)



Arithmetic Mean



Geometric Mean



Harmonic Mean



Mode

correct

Made By: Waqar Siddhu



The collection of all the elements under study is called:

Answer (Please select your correct option)



Population



Sample



Data



Registration

correct

Made By: Waqar Siddhu



A five-number summary consists of :

Answer (Please select your correct option)

☐

X_0 , Q_1 , Median, Q_3 , and X_m

correct

☐

X_m , Q_1 , Mean, Q_3 , and X_0

☐

X_m , Q_1 , Mode, Q_3 , and X_0

☐

X_0 , Q_1 , Median, Q_2 , and X_m



For what condition of k , at least $1 - 1/k^2$ of the data-values fall within k standard deviations of the mean:

Answer (Please select your correct option)



Greater than 1

correct



Less than 1



Greater or equal to 1



Less or equal to 1

Made By: Waqar Siddhu



Which of the following can never be taken as the probability of an event?

Answer (Please select your correct option)

☐ 1

☒ 0

☐ 0.5

☐ -0.5

correct



If E is an impossible event, then $P(E)$ is:

Answer (Please select your correct option)

☐

1

☐

0.15

☒

0

☐

0.5

correct

Made By: Waqar Siddhu



The average which is defined as the reciprocal of the arithmetic mean of the reciprocals of the values is called:

Answer (Please select your correct option)



Geometric Mean



Harmonic Mean

correct



Mode



Median

Made By: Waqar Siddhu



An automobile is running, during the first 60 Km, at the rate of 10 Km/hr. During the second 60 Km at the rate of 30Km/hr, while during the third 60 Km its speed was 40 Km/hr. What method is more appropriate to calculate the average speed?

Answer (Please select your correct option)

☐ Median

☐ Arithmetic mean

☒ Harmonic mean

☐ Geometric mean

correct



What is the standard deviation of the following values?

7, 7, 7, 7, 7, 7, 7

Answer (Please select your correct option)

7

☐

1

☐

49

☐

0

☐

correct

Made By: Waqar Siddhu



Which one of the following is a meso-kurtic curve?

Answer (Please select your correct option)



Negatively skewed



Positively skewed



J-shaped



Normal

correct

Made By: Waqar Siddhu



Men tend to marry women who are slightly younger than themselves. Suppose that every man married a woman who was exactly 5 years younger than themselves. Which of the following is correct?

Answer (Please select your correct option)



The correlation coefficient is -5



The correlation coefficient is 5



The correlation coefficient is 1



The correlation coefficient is 0



In a multiplication theorem $P(A \cap B)$ equals:

Answer (Please select your correct option)

☐

$P(A) P(B)$

☐

$P(A) + P(B)$

☐

$P(A) \times P(B|A)$

☐

$P(B|A) \times P(B)$

correct

Made By: Waqar Siddhu



Which one of the following is the class frequency?

Answer (Please select your correct option)

☐

The number of observations in each class

correct

☐

The difference between consecutive lower class limits

☐

Always contains at least 5 observations

☐

Usually a multiple of the lower limit of the first class

Made By: Waqar Siddhu



Which of the following comes first to make frequency distribution.

Answer (Please select your correct option)



Number of Groups



Class interval



Range



Tally marks

correct

Made By: Waqar Siddhu



Data obtained from the Ministry of food, is an example of:

Answer (Please select your correct option)

☐ Primary data

☒ Secondary data

☐ Qualitative data

☐ None of these

correct



Total angle of a Pie Chart is:

Answer (Please select your correct option)



90°



180°



270°



360°

correct

Made By: Waqar Siddhu