Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐

Date:

Topic 68

Continuity and open sets:-

Continuity of a function depends on the topology of the domain space and the range space i.e., "The collection of open subsets of both sets."

The more rich space on domain side provides more room from continuity and more rich space on the range side ~~restriction~~ restricts more a function to be continuous.

In our previous three topics we have seen three things

(i) Constant function $f: X \to Y$ is always continuous.

(ii) Any function

$g: X \longrightarrow$ Indiscrete Space

is continuous.

(iii) Any function.

$h:$ Discrete Space $\longrightarrow Y$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

is continuous.

Let (X, T_x) be strictly finer than (X, T_x'). Then identity function

$$I:(X, T_x') \to (X, T_x)$$

is not continuous.

Proof:- $I : X \to X$

$I(x) = x \quad \forall x$

$T_x' \subset T_x \quad \exists \ U \in T_x$

such that $U \notin T_x'$

Now let $U \in T_x$

$I(U) = U \notin T_x'$

$\Rightarrow I:(X, T_x') \to (X, T_x)$ is not continuous.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 69

Examples: Continuity and open Sets

Let us see some examples to under-stand the role of open collection in continuity of a function.

Example:-

let $X = \{a, b, c, d\}$.

$T_X = \{\phi, \{b\}, X\}$

$Y = \{1, 2, 3, 4\}$

$T_Y = \{\phi, \{2\}, \{1, 2\}, \{2, 3, 4\}, Y\}$

$f : X \to Y$ defined as

$f(a) = 2$

$f(b) = 4$

$f(c) = 2$

$f(d) = 3$

Now f is not continuous

$\because f^{-1}(\{1, 2\}) = \{a, c\} \notin T_X$

$T_X' = \{\phi, \{b\}, \{a, c\}, \{a, b, c\}, X\}$

$g : X \to Y$

$T_Y = \{\phi, \{2\}, \{1, 2\}, \{2, 3, 4\}, Y\}$

$g(a) = 2$.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$g(b) = 4$

$g(c) = 2$

$g(d) = 3$

g is not continuous
($T_x' - T_y$ continuous)

Example 2:-

~~$X = \mathbb{R}_m$~~ $X = \mathbb{R}_u$, $Y = \mathbb{R}_u$

$f : \mathbb{R}_u \to \mathbb{R}_u$

$f(x) = x + 1$

$f(x)$ is continuous.

$g : \mathbb{R}_u \to \mathbb{R}_l$

$g(x) = x + 1$

g is not ~~constant~~ continuous.

$\because [0,2)$ open $\subset \mathbb{R}_l = Y$

$g^{-1}([0,2)) = [-1,1) \notin \mathbb{R}_u = X.$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Date:

Topic 70:-
Continuity and Compasiability

Proposition 1:-

Let $f:(X,T_x)\to(Y,T_y)$ be a continuous function and T_x' is finer than T_x. Then $f:(X,T_x')\to(Y,T_y)$ is continuous.

Proposition 2:-

Let $g:(X:T_x)\to(Y:T_y)$ is not continuous and T_x'' is weaker than T_x. Then $g:(X,T_x'')\to(Y,T_y)$ is not continuous.

Proof:- (Proposition 1):-

$f:(X,T_x)\xrightarrow{\text{cont.}}(Y,T_y)$

So $\forall\ U\in T_y \Rightarrow f^{-1}(U)\in T_x$

Given that $T_x\subset T_x'$

$\Rightarrow f^{-1}(U)\in T_x'$

$\Rightarrow f:(X,T_x')\to(Y,T_y)$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

is continuous.

Proposition 2:-

$g:(x,\tau_x) \xrightarrow{\text{Not cont.}} (Y,\tau_Y)$

This means that

$\exists \cdot u \in \tau_y$ such that

$g^{-1}(u) \notin \tau_x$

Now,

$\tau_x'' \subset \tau_x$

$\Rightarrow g^{-1}(u) \notin \tau_x''$

$\Rightarrow g:(x,\tau_x'') \to (Y,\tau_Y)$ is

~~cons~~ not continuous.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date: _____

Topic 71:- Continuity and Basis

Proposition:-

A function

$$f : (X, T_X) \to (Y, T_Y)$$

is continuous iff the inverse image of each member of a basis B_Y for T_Y is an open subset of X.

$$\forall\ B_Y \in \mathcal{B}_Y \Rightarrow f^{-1}(B_Y) \in T_X$$

Proof:-

Given $\forall\ B_Y \in \mathcal{B}_Y$

$$\Rightarrow f^{-1}(B_Y) \in T_X$$

let $U \in T_Y$

$$f^{-1}(U) \in T_X$$

Now $\because B_Y$ is a basis for T_Y so,

$\forall\ U \in T_Y$, $U = \bigcup_i B_{Y_i}$, $B_{Y_i} \in \mathcal{B}_Y$

$$f^{-1}(U) = f^{-1}\left(\bigcup_i B_{Y_i}\right)$$

$$= \bigcup_i \left(f^{-1}(B_{Y_i})\right)$$

Now, $f^{-1}(B_{Y_i}) \in T_X\ \forall i$

and T_X is a topology.

$\Rightarrow f$ is continuous.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Let

$f : X \to Y$ is continuous

$\Rightarrow \forall \cdot u \in T_y \Rightarrow f^{-1}(u) \in T_x$

But we know that

$B_y \subset T_y$

$\Rightarrow \forall \quad B_y \subset B_y \subset T_y$

$\Rightarrow f^{-1}(B_y) \in T_x$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Topic 72:- Continuity and subbasis

Theorem :-

A function $f:(X,T_X)\to(Y,T_Y)$ is continuous iff the inverse image of each member of a subbasis S_Y for T_Y is an open subset of X.

Proof :-

Given $\forall\ S_Y \in S_Y$

$\Rightarrow f^{-1}(S_Y) \in T_X$

let $U \in T_Y$

$\Rightarrow f^{-1}(U) \in T_X$

$\because S_Y$ is a subbasis for T_Y

$\Rightarrow \forall\ U \in T_Y,\ U = \bigcup_i \left(\bigcap_{j=1}^{n_i} S_{Yij}\right)$

$f^{-1}(U) = f^{-1}\left(\bigcup_i \left(\bigcap_{j=1} S_{Yij}\right)\right)$

$\Rightarrow f^{-1}(U)$ is open in X.

$\Rightarrow f$ is continuous function

let $f:X\to Y$ is continuous

$\Rightarrow \forall\ U \in T_Y$

$\Rightarrow f^{-1}(U) \in T_X$

$S_Y \subset T_Y$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$\Rightarrow \forall \quad S_Y \in \mathcal{S}_Y$

$f^{-1}(S_Y) \in T_X.$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐

Date:

Topic 73
Absolute Value Function

In this topic we will discuss the continuity of Absolute Value function.

$f : \mathbb{R} \longrightarrow \mathbb{R}$

$$f(x) = |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Show that $f : \mathbb{R}_u \longrightarrow \mathbb{R}_u$ is continuous when $f(x) = |x|$

Sol:-

Let $(a,b) \in \mathbb{R}_u = (\mathbb{R}, \mathcal{T}_u)$



Now

$$f^{-1}((a,b)) = \begin{cases} \phi & \text{if } a < b \leq 0 \\ (-b,b) & \text{if } a < 0 < b \\ (-b,-a) \cup (a,b) & \text{if } 0 \leq a < b \end{cases}$$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 74
Composition of Continuous Function:-

In this topic we will prove the fact

"Composition of two continuous functions is a continuous function."

$$f : X \longrightarrow Y$$

$$g : Y \longrightarrow Z$$

$$h(x) = gof(x) = g(f(x))$$

$$h : X \longrightarrow Z$$

Proposition:-

Let $f : X \to Y$

$g : Y \to Z$ are two continuous function. Then

$gof : X \longrightarrow Z$ is continuous.

Proof:-

Let $U \in T_Z$

$(gof)^{-1}(U) = f^{-1}(g^{-1}(U)) \in T_X$

$\Rightarrow (gof)^{-1}(U)$ is open in X.

gof is a continuous function $X \to Z$.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 75

Continuous Function:

Another Definition :-

Proposition :-

A function

$f : (X, T_X) \to (Y, T_Y)$

is continuous iff the inverse image of every closed subset of Y under is closed in X.

Proof :-

f is a continuous function from $X \to Y$.

$\forall\ U \in T_Y \Rightarrow f^{-1}(U) \in T_X$.

Let $A \underset{\text{closed}}{\subset} Y \Rightarrow A \underset{\text{open}}{\subset} Y$ $(A^c \in T_Y)$

$\therefore$ f is continuous.

$\Rightarrow f^{-1}(A^c)$ is open in X.

But $f^{-1}(A^c) = (f^{-1}(A))^c \to$ open in X.

$\Rightarrow (f^{-1}(A)^c)^c$ is closed in X.

$\Rightarrow f^{-1}(A)$ is closed in X.

(Given) We have $\forall\ A \subset Y \Rightarrow f^{-1}(A) \underset{\text{closed}}{\subset} X$

Let $U \underset{\text{open}}{\subset} Y \Rightarrow U^c \underset{\text{closed}}{\subset} Y$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$f^{-1}(U^c) = (f^{-1}(U))^c$

$\because f^{-1}(U) \subseteq_{open} X$ or $f^{-1}(U) \in T_X$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐

Date:

Topic 76
Restriction of Function

Defination:-

Let $f: X \to Y$ be a function and $A \subset X$.

The restriction of f to A is a function $f_A : A \to Y$ defined as

$f_A(a) = f(a) \quad \forall \quad a \in A$

Proposition:-

Let $f: X \to Y$ be a continuous function. Then $f_A : A \to Y$ is continuous. when $A \subset X$.

Proof:-

For any $B \subset Y$ $\quad f_A^{-1}(B) = A \cap f^{-1}(B)$

Let $U \in T_Y$

$\Rightarrow f_A^{-1}(U) = A \cap f^{-1}(U)$

Given f is continuous $\forall U$

$f^{-1}(U) \in T_X$.

$\Rightarrow f_A^{-1}(U) \in T_A$.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 77:- Continuous Function and Arbitrary Closeness

Recall closure point:-

$\bar{A} = A \cup A'$

A point that is arbitrary close to a set A.

Continuity preserve arbitrary closeness.

Theorem :-

A function $f:(X,T_X) \to (Y,T_Y)$ is continuous iff for every subset A of X.

$$f(\bar{A}) = \overline{f(A)}$$

Proof :-

Let $f: X \to Y$ is continuous.

Let $A \subset X$.

Now we know that $f(A) \subset \overline{f(A)}$

So

$$A \subset f^{-1}(f(A)) \subset f^{-1}(\overline{f(A)}) \to (1)$$

But $\overline{f(A)}$ is closed in Y.

$\Rightarrow f^{-1}(\overline{f(A)})$ is closed in X.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$\because f$ is continuous

Now ① $\Rightarrow A \subset f^{-1}(\overline{f(A)})$

i.e., $A \subset \bar{A} \subset f^{-1}(\overline{f(A)})$

$\Rightarrow f(\bar{A}) \subset f(f^{-1}(\overline{f(A)})) = \overline{f(A)}$

$\Rightarrow f(\bar{A}) \subset \overline{f(A)}$

For the other side

let $f(\bar{A}) \subset \overline{f(A)}$

let $B \overset{\text{closed}}{\subset} Y$

let $f^{-1}(B) = A$

we have to show that A is closed in X. i.e., $\bar{A} = A$

we know that

$f(\bar{A}) \subset \overline{f(A)}$

$\Rightarrow f(\overline{f^{-1}(B)}) \subset \overline{f(f^{-1}(B))} = \bar{B} = B$

$\Rightarrow f(\bar{A}) \subset B$

$\Rightarrow f^{-1}(f(\bar{A})) \subset f^{-1}(B)$

$\Rightarrow \bar{A} \subset f^{-1}(B) = A$

$\bar{A} \subset A \quad , \quad A \subset \bar{A}$

$\Rightarrow \bar{A} = A$

$\Rightarrow$ A is closed in X.

$\Leftrightarrow$ f is continuous.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 78
Continuity at a point

We have discussed continuity in global sense ~~i.e~~ i.e., continuity on a set.

Now let us discuss locally i.e., continuity at a point.

Depination:-

Let $f:(X,\mathcal{T}_X)\to(Y,\mathcal{T}_Y)$ be a function. f is continuous at a point $x\in X$ iff for every $U_{f(x)}$ (open set containing $f(x)$) There exist V_x (open set of X containing x) such that

$$V_x\subset f^{-1}(U_{f(x)}) \quad \text{or}$$

$$f(V_x)\subset U_{f(x)}.$$

For a given $x\in X$.

Fin $f(x)$.

Collect all the open subsets of Y containing $f(x)$ i.e.,

$$U_{f(x)}=\{U_{f(x)} \mid f(x)\in U_{f(x)}\}\subset\mathcal{T}_Y$$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

For each element of $U_{f(x)}$ (say $U_{f(x)}$) check is $\exists$ some V_x such that $V_x \subset f^{-1}(U_{f(x)})$

If your answer is yes for all $U_{f(x)}$. Then you can say f is continuous at x.

But if $\exists$ some $U_{f(x)}$ such that $\nexists U_x$ such that $V_x \subset f^{-1}(U_{f(x)})$. Then you say that f is not continuous at x.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date: _____

Topic 79
Examples: Continuity at a point:

Let us see some ~~eg~~ examples of continuity at a point.

Example 1:-

Let $X = \{a, b, c, d\}$ with

$\tau_x = \{\phi, \{a\}, \{b\}, \{a, b\}, \{b, c, d\}, X\}$

and $Y = \{1, 2, 3, 4\}$ with

$\tau_Y = \{\phi, \{2\}, \{1, 2\}, \{2, 3, 4\}, Y\}$



f is continuous at 'a'

$\forall\ U_3 \in \tau_Y\ \exists\ V_a \in \tau_X$

such that $V_a \subset f^{-1}(U_3)$

$U_3 = \{\{2, 3, 4\}, Y\}$

$f^{-1}\{2, 3, 4\} = X$ open in X.

f is not continuous at 'c'

$U_2 = \{\{2\}, \{1, 2\}, \{2, 3, 4\}, Y\}$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$f^{-1}(\{2\}) = \{c\} \neq V_c$

such that

$V_c \subset \{c\}$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date: _____

Topic 80:- Continuity at every and continuous Function

Theorem :-

A function $f: X \to Y$ is continuous iff it is continuous at every point of X.

Proof :-

Let $f: X \to Y$ is continuous

Let $x \in X$

Now take $U_{f(x)} \in T_Y$

(open subset of containing $f(x)$)

$\therefore$ f is continuous. so,

$f^{-1}(U_{f(x)}) \in T_X$

So, $V_x = f^{-1}(U_{f(x)}) \subset f^{-1}(U_{f(x)})$

$\Rightarrow$ f is continuous at x.

For the other side

$f: X \to Y$ is continuous at every $x \in X$. i.e., $\forall\ x \in X$ and

$\forall\ U_{f(x)} \in T_Y$ $\exists$ $V_x \in T_X$

such that $V_x \subset f^{-1}(U_{f(x)})$

$\therefore$ V_x $f'(U_{f(x)})$ can be written

Scanned by CamScanner

Scanned with CamScanner

CamScanner

as the union of such V_x

i.e.,

$$f'(U_{f(x)}) = \cup \{ V_p \mid p \in f'(U_{f(x)}) \}$$

$\Rightarrow f'(U_{f(x)})$ is open in X.

$\Rightarrow f$ is continuous function from (X, T_x) to (Y, T_y).

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 81:- Topology induced by function:-

Defination:-

Let $f: X \to Y$ be a function and T_Y be a topology on Y. The class of subsets of X.

$T_{Xf} = \{U = f^{-1}(V) \mid V \in T_Y\}$

forms a topology on X.

This topology is called "Topology on X induced by f."

Task:-

Show that T_{Xf} is a topology on X and also show that

$f: (X, T_X) \to (Y, T_Y)$

is continuous.

Proof:-

$\phi \in T_{Xf}$ $\because \phi = f^{-1}(\phi)$, $\phi \in T_Y$

$X \in T_{Xf}$ $\because X = f^{-1}(Y)$, $Y \in T_Y$

Let $U_i \in T_{Xf}$ $i \in I$

$U_i = f^{-1}(V_i)$ $(V_i \in T_Y)$ $\forall$ $i \in I$

$\bigcup_i U_i = \bigcup_i (f^{-1}(V_i)) = f^{-1}(\bigcup_i (V_i)) \in T_Y$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$\Rightarrow \forall\, U_i \in T_{x_f}$

let $U_1, U_2 \in T_{x_f}$

$\Rightarrow \exists\ V_1, V_2 \in T_Y$

such that

$U_1 = f^{-1}(V_1)$,

$U_2 = f^{-1}(V_2)$

$U_1 \cap U_2 = f^{-1}(V_1) \cap f^{-1}(V_2)$

$= f^{-1}(V_1 \cap V_2)$

$\Rightarrow U_1 \cap U_2 \in T_{x_f}$

$f : (X, T_{x_f}) \longrightarrow (Y, T_Y)$

f is continuous iff $\forall\ V \in T_Y$

$\Rightarrow f^{-1}(V) \in T_{x_f}$

T_{x_f} so, f is if continuous.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 82
Examples: Topology Induced by Function:-

Example 1:-

Let $X = \{a, b, c, d\}$, $Y = \{1, 2, 3\}$

$T_Y = \{\phi, \{2\}, \{1, 2\}, \{2, 3\}, Y\}$

$f : X \to Y$ defined as

$f(a) = 1$, $f(b) = 2$

$f(c) = 2$, $f(d) = 3$

$T_{X_f} = \{\phi, \{b, c\}, \{a, b, c\}, \{b, c, d\}, X\}$

Example 2:-

$g : X \to Y$ (X, Y are same as in ex. 1)

$g(a) = 1$, $g(b) = 1$

$g(c) = 3$, $g(d) = 2$

$\therefore T_{X_g} = \{\phi, \{d\}, \{a, b, d\}, \{c, d\}, X\}$

Question:-

What is T_X on X such that $f : X \to Y$, $g : X \to Y$ are both continuous.

Answer:-

The topology with the subbasis

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

$\mathcal{S}_{X_{f,g}} = \{f^{-1}(V) \mid V \in T_Y\} \cup \{g^{-1}(V) \mid V \in T_Y\}$

The topology generated by $\mathcal{S}_{X_{f,g}}$ (subbasis) is induced by f and g.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Date:

Topic 83
Topology on real line and linear function

In this topic we will show that the topology on real line induced by the set of all linear functions from real line to real line with lower limit topology is discrete.

$f : X \to Y$

$g : X \to Y$

$\mathcal{T}_Y$ is a topology on Y.

Then $\mathcal{T}_{X_{f,g}}$ has basis

$$\mathcal{S}_{X_{f,g}} = \{f^{-1}(v) \mid v \in \mathcal{T}_Y\} \cup \{g^{-1}(v) \mid v \in \mathcal{T}_Y\}$$

Let

$f : \mathbb{R} \to \mathbb{R}$

$g : \mathbb{R} \to \mathbb{R}$

$f(x) = x - p$, $g(x) = p - x$ $\quad p \in \mathbb{R}$.

$A \subset \mathbb{R}$, $A \in \mathcal{T}_{\mathbb{R}}$

$A = [0, 1)$

$f^{-1}([0,1)) = [p, p+1)$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

$$g^{-1}([0,1)) = (p-1, p]$$

$$f^{-1}([0,1)) \cap g^{-1}([0,1)) = [p, p+1) \cap (p-1, p].$$

$$= \{p\}$$

$\{p\} \in \mathcal{B}_x$

$\Rightarrow$ The topology on $\mathbb{R}$ induced by set of all linear function is discrete.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

Topic 84
Open Mapping

Defination:-

A map

$f:(X,T_X) \longrightarrow (Y,T_Y)$

is called an open mapping iff the image of each open subset of X is open in Y.

$\forall\ U \in T_X,\ f(U) \in T_Y$

Example:-

$X = \{1,2,3,4\}$ and with

$T_X = \{\phi, \{1\}, \{2\}, \{1,2\}, X\}$

$Y = \{a,b,c,d\}$ with

$T_Y = \{\phi, \{a\}, \{b\}, \{a,b\}, \{b,c,d\}, Y\}$

f



f is open map.

$f(\phi) = \phi$

$f(\{1\}) = \{b\}$

$f(\{2\}) = \{a\}$

$f(\{1,2\}) = \{a,b\}$

$f(X) = \{a,b\}$

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date: _____

Topic 85
Open Mapping and Basis

Proposition:-

A map

$f:(X,\mathcal{T}_X)\longrightarrow(Y,\mathcal{T}_Y)$

is an open map iff the image of each element of a basis $\mathcal{B}_X$ for $\mathcal{T}_X$ is an open subset of Y.

Proof:-

Let f is open

$\Rightarrow$ The image of every element of $\mathcal{T}_X$ is an element of $\mathcal{T}_Y$.

Now $B_X \subset \mathcal{T}_X$

$\Rightarrow f(B_X) \in \mathcal{T}_Y$

$B_X \in \mathcal{B}_X$.

For other side

Let $U \in \mathcal{T}_X$

$\mathcal{B}_X$ is a basis of $\mathcal{T}_X$.

So $U = \bigcup_i B_i \qquad B_i \in \mathcal{B}_X$

Now $f(U) = f(\bigcup_i B_i) = \bigcup_i f(B_i)$

Now $f(B_i)$ is open in Y.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Mon☐ Tues☐ Wed☐ Thurs☐ Fri☐ Sat☐ Date:

$\Rightarrow \bigcup_i f(B_i)$ is open in Y.

So $f(U)''$ is open in Y.

$\Rightarrow$ f is an open map.

Scanned by CamScanner

Scanned with CamScanner

CamScanner

Closed Mapping

Def:
A map

$$f : (X, \mathcal{T}_X) \to (Y, \mathcal{T}_Y)$$

is called a closed map iff the image of each closed subset of X is closed in Y.

Closed Mapping

Def:
A map

$$f : (X, \mathcal{T}_X) \to (Y, \mathcal{T}_Y)$$

is called a closed map iff the image of each closed subset of X is closed in Y.

$\forall\ U \underset{\text{closed}}{\subset} X \Rightarrow f(U) \underset{\text{closed}}{\subset} Y$

Closed Mapping

Ex: $X = \{1,2,3\}$, $\mathcal{T}_X = \{\emptyset, \{1\}, \{1,2\}, X\}$

$Y = \{a,b,c\}$, $\mathcal{T}_Y = \{\emptyset, \{a,b\}, \{b\}, Y\}$



$\mathcal{C}_X = \{X, \{2,3\}, \{3\}, \emptyset\}$

$\mathcal{C}_Y = \{Y, \{c\}, \{a,c\}, \emptyset\}$

$f(\emptyset) = \emptyset$ ✓

$f(\{3\}) = \{c\}$ ✓ | $f(\{2,3\}) = \{a,c\}$ ✓ | $f(X) = \{a,c\}$ ✓

Topology

Projection Mapping

Projection Mapping

Projection mapping is an example of mapping that is not closed in general.

But let us see first "What is projection map?"

Projection Mapping

Projection Map:
Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be two topological spaces. The maps

Product top:

$$\begin{aligned} \pi_1 : (X \times Y, \mathcal{T}_{X\times Y}) &\to (X, \mathcal{T}_X) \\ (x,y) &\mapsto x \end{aligned}$$

$$\begin{aligned} \pi_2 : (X \times Y, \mathcal{T}_{X\times Y}) &\to (Y, \mathcal{T}_Y) \\ (x,y) &\mapsto y \end{aligned}$$

are called projection maps, where $\mathcal{T}_{X\times Y}$ is product topology on $X \times Y$. π_1 is first projection and π_2 is second projection.

Projection Mapping

Fact:
Projection map may or may not be closed.
Justification:
Consider the first projection

$$\pi_1 : (\mathbb{R} \times \mathbb{R}, \mathcal{T}_{\mathbb{R}\times\mathbb{R}}) \to (\mathbb{R}, \mathcal{T}_u)$$

The set $A = \{(x,y) |\ y = \tan x\ ,\ \frac{-\pi}{2} < x < \frac{\pi}{2}\}$ is closed in $\mathbb{R} \times \mathbb{R}$ but

$$\pi_1(A) = \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$$

Projection Mapping

Fact:
Projection map may or may not be closed.
Justification:
Consider the first projection

$$\pi_1 : (\mathbb{R} \times \mathbb{R}, \mathcal{T}_{\mathbb{R}\times\mathbb{R}}) \to (\mathbb{R}, \mathcal{T}_u)$$

The set $A = \{(x,y) |\ y = \tan x\ ,\ \frac{-\pi}{2} < x < \frac{\pi}{2}\}$ is closed in $\mathbb{R} \times \mathbb{R}$ but

$$\pi_1(A) = \left(\frac{-\pi}{2}, \frac{\pi}{2}\right) \subset \mathbb{R}.$$

Not closed

Topology

Continuity and Open Mapping

Continuity and Open Mapping

We have studied

- Continuous Mapping
- Open Mapping
- Closed Mapping

Questions:

1. Is a continuous map an open map?
2. Is an open map a continuous map?

Continuity and Open Mapping

Example 1:
Let $X = \{a, b, c, d\}$ with $\mathcal{T}_X = \{\emptyset, \{b, c\}, \{a, b, c\}, X\}$
and
$Y = \{1, 2, 3\}$ with $\mathcal{T}_Y = \{\emptyset, \{1\}, \{1, 2\}, Y\}$.

Consider the following map

$$f : X \to Y$$

defined as $f(x) = 2$ for all $x \in X$.

Continuity and Open Mapping

Example 1:
Let $X = \{a, b, c, d\}$ with $\mathcal{T}_X = \{\emptyset, \{b, c\}, \{a, b, c\}, X\}$
and
$Y = \{1, 2, 3\}$ with $\mathcal{T}_Y = \{\emptyset, \{1\}, \{1, 2\}, Y\}$.

Consider the following map

$$f : X \to Y$$

Continuous.

$f(\{b,c\}) = \{2\}$ ∩ Not open in Y

defined as $f(x) = 2$ for all $x \in X$.

Continuity and Open Mapping

Example 2:
Let $X = \mathbb{R}$ with usual topology and $Y = \mathbb{R}$ with lower limit topology.

Consider the identity map

$$f : X \to Y$$
$$x \mapsto x$$

$[a, b) \subset \mathbb{R}_l$

$f^{-1}[a,b) = [a,b) \subset \mathbb{R}_u$

f is not cont.

$(a,b) \in \mathcal{B}_u \quad f((a,b)) = (a,b) \in \mathcal{T}_l$

Continuity and Open Mapping

Answers:

1. Continuous map may or may not be open map.
2. Open map may or may not be continuous map.

Topology

Openness and Restriction Mapping

Openness and Restriction...

Recall the restriction mapping and the fact about the continuity of restriction mapping.
Now the question is

"Can we prove the same kind of result for restriction of an open mapping?".

Openness and Restriction...

Example: Let $X = \{a, b, c, d\}$ with $\mathcal{T}_X = \{\emptyset, \{a\}, \{a, b\}, \{a, b, c\}, X\}$ and $Y = \{1, 2, 3\}$ with $\mathcal{T}_Y = \{\emptyset, \{1\}, \{1, 2\}, Y\}$.

Consider the map $f : X \to Y$ defined as

$$f(a) = 1,\ f(b) = 2,\ f(c) = 3,\ f(d) = 2$$

Let $A = \{b, c, d\} \subset X$ with relative topology $\mathcal{T}_A = \{\emptyset, \{b\}, \{b, c\}, A\}$. Now the restriction map $f_A : A \to Y$ is defined as

$$f_A(b) = 2,\ f_A(c) = 3,\ f_A(d) = 2$$

Openness and Restriction...

Now f is an open map but f_A is not an open map. Because $\{b\}$ is an open subset of A but

$$f_A(\{b\}) = \{2\} \underset{\text{NotOpen}}{\subset} Y$$

Remark:

Restriction of an open map may or may not be an open map.

KHALIL SAHIR
MC200400708

Topology

Homeomorphism and Homeomorphic Spaces

Homeomorphism...

- f is one-one.
- f is onto

Homeomorphism:
A map $f : X \to Y$ is called a homeomorphism between two topological spaces $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ iff

1. f is bijective.
2. f is continuous.
3. $f^{-1} : Y \to X$ is continuous.

Homeomorphism...

$\forall\ U \underset{\text{open}}{\subset} Y$

$f^{-1}(U) \underset{\text{open}}{\subset} X$

Homeomorphism:
A map $f : X \to Y$ is called a homeomorphism between two topological spaces $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ iff

1. f is bijective.
2. f is continuous.
3. $f^{-1} : Y \to X$ is continuous.

Homeomorphism...

$\forall\ V \underset{\text{open}}{\subset} X$

$\left(f^{-1}\right)^{-1}(V) \underset{\text{open}}{\subset} Y$

Homeomorphism:
A map $f : X \to Y$ is called a homeomorphism between two topological spaces $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ iff

1. f is bijective.
2. f is continuous.
3. $f^{-1} : Y \to X$ is continuous.

Homeomorphism...

Homeomorphic Spaces:
Two topological spaces $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ are called homeomorphic iff there exists a homeomorphism $f : X \to Y$. We also say such spaces topologically equivalent spaces.

$X \simeq Y$

Topology

Examples:
Homeomorphism

Examples: Homeomorphism

Let us see some examples of homeomorphism and homeomorphic spaces.

Examples: Homeomorphism

Example 1: Consider
$X = \{a, b, c, d\}$ with $T_X = \{\emptyset, \{a, b\}, \{c, d\}, X\}$
$Y = \{1, 2, 3, 4\}$ with $T_Y = \{\emptyset, \{2, 3\}, \{1, 4\}, Y\}$
and the map $f : X \to Y$ defined as

$$f(a) = 2,\ f(b) = 3,\ f(c) = 4,\ f(d) = 1$$

$f^{-1}(\{2,3\}) = \{a,b\} \subseteq_{open} X$ | $f^{-1}(Y) = X \subseteq_{open} X$.
$f^{-1}(\{1,4\}) = \{c,d\} \subseteq_{open} X$

Examples: Homeomorphism

Example 1: Consider
$X = \{a, b, c, d\}$ with $T_X = \{\emptyset, \{a, b\}, \{c, d\}, X\}$
$Y = \{1, 2, 3, 4\}$ with $T_Y = \{\emptyset, \{2, 3\}, \{1, 4\}, Y\}$
and the map $f : X \to Y$ defined as

$$f(a) = 2,\ f(b) = 3,\ f(c) = 4,\ f(d) = 1$$

Now $f^{-1} : Y \to X$
$f^{-1}(1) = d$, $f^{-1}(2) = a$, $f^{-1}(3) = b$, $f^{-1}(4) = c$
bij. + Cont + f^{-1} Cont.
f^{-1} is Cont.

Examples: Homeomorphism

Example 2: Consider
$X = [0, 1]$, $Y = [0, 2]$ as subspaces of $(\mathbb{R}, \mathcal{T}_u)$
and the map $f : X \to Y$ defined as

$$f(x) = 2x \quad \forall x \in X$$

- f is bijection.
- f is Cont.
- $f^{-1} : Y \to X$ $\quad f^{-1}(y) = \frac{y}{2}$
  So f^{-1} is Cont.

$X = [0,1]$ ⫋ $Y = [0,2]$

Examples: Homeomorphism

Example 3: Consider
$X = (\frac{-\pi}{2}, \frac{\pi}{2})$ as subspace of $(\mathbb{R}, \mathcal{T}_u)$ and $Y = \mathbb{R}$ with usual topology and the map $f : X \to Y$ defined as

$$f(x) = \tan x \quad \forall x \in X$$

- f is bij. ✓
- f is Cont. ✓
- f^{-1} is Cont ✓

$X = (\frac{-\pi}{2}, \frac{\pi}{2})$ ⫋ $Y = \mathbb{R}$

Topology

Bicontinuity and Open Mapping

Bicontinuity and Open Mapping

$f : X \to Y$
$\forall\ U \subseteq_{open} Y$, $f^{-1}(U) \subseteq_{open} X$

$f^{-1} : Y \to X$
$\forall\ V \subseteq_{open} X$, $(f^{-1})^{-1}(V) = f(V) \subseteq_{open} Y$
open mapping

Recall homeomorphism.
A bijective map $f : X \to Y$ such that f and f^{-1} both continuous or in other words f is bicontinuous.

Bicontinuity and Open Mapping

A map $f : X \to Y$ is a homeomorphism between two topological spaces $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ iff

1. f is bijective.
2. f is continuous.
3. f is open map.

Topology

Bicontinuity and Bijection

Bicontinuity and Bijection

Question:
Is there any relation between bicontinuity and bijection?

More specifically.

- Does bijection imply bicontinuity?
- Does bicontinuity imply bijection?

Bicontinuity and Bijection

Example 1: Consider
$X = \{a, b, c, d\}$ with $T_X = \{\emptyset, \{a\}, \{c, d\}, \{a, c, d\}, X\}$
$Y = \{1, 2, 3, 4\}$ with $T_Y = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, Y\}$
and the map $f : X \to Y$ defined as

$$f(a) = 1,\ f(b) = 2,\ f(c) = 3,\ f(d) = 4$$

Bicontinuity and Bijection

Example 1: Consider
$X = \{a, b, c, d\}$ with $T_X = \{\emptyset, \{a\}, \{c, d\}, \{a, c, d\}, X\}$
$Y = \{1, 2, 3, 4\}$ with $T_Y = \{\emptyset, \{1\}, \{2\}, \boxed{\{1, 2\}}, Y\}$
and the map $f : X \to Y$ defined as

$$f(a) = 1,\ f(b) = 2,\ f(c) = 3,\ f(d) = 4$$

f is bijective. but f is not cont.

$f^{-1}(\{1,2\}) = \{a, b\} \subset X$ (Not open)

Bijection $\not\Rightarrow$ Bicont.

Bicontinuity and Bijection

Example 2: Consider
$X = \{a, b, c, d\}$ with $T_X = \{\emptyset, X\}$
$Y = \{1, 2, 3, 4\}$ with $T_Y = \{\emptyset, \{1\}, Y\}$
and the map $f : X \to Y$ defined as

$$f(a) = 1,\ f(b) = 1,\ f(c) = 1,\ f(d) = 1$$

Bicontinuity and Bijection

Example 2: Consider
$X = \{a, b, c, d\}$ with $T_X = \{\emptyset, X\}$
$Y = \{1, 2, 3, 4\}$ with $T_Y = \{\emptyset, \{1\}, Y\}$
and the map $f : X \to Y$ defined as

$$f(a) = 1,\ f(b) = 1,\ f(c) = 1,\ f(d) = 1$$

f is (cont + open) $\not\Rightarrow$ Bijective.

Topology

Homeomorphism as Equivalence Relation

Homeomorphism...

Equivalence Relation:
A binary relation $\sim$ on a set X is said to be an equivalence relation iff $\sim$ satisfies the following three properties for all $a, b, c \in X$

1. Reflexive: $a \sim a$
2. Symmetric: $a \sim b$ iff $b \sim a$
3. Transitive: If $a \sim b$ and $b \sim c$ then $a \sim c$

Homeomorphism...

Proposition:
Homeomorphism of topological spaces is an equivalence relation.

Proof: • $X \approx X$ $\quad Id: X \to X$, $x \mapsto x$

• Let $X \approx Y$ $\quad f: X \to Y$ Hom. Bijectn + Bicont.

$f^{-1}: Y \to X$ Bijectn + Cont + f is cont

$\Rightarrow X \approx Y \Leftrightarrow Y \approx X$.

Homeomorphism...

• $X \approx Y$, $Y \approx Z$

$f: X \to Y$ Bijectn + Bicont. | $g: Y \to Z$ Bijective + Bicont.

$g \circ f: X \to Z$

Bijective. + $g \circ f$ is bicont.

$\to g \circ f$ is cont

$f^{-1} \circ g^{-1}$ is cont $\Rightarrow X \approx Z$.

Topology

Topological Properties

Topological Properties

Topological Property:
A property "P" of a topological space $(X, \mathcal{T})$ is called "Topological Property" if every space homeomorphic to X has property P.
i.e.
A property that is unchanged under homeomorphism.
Topological property is also called topological invariant.

Topological Properties

Some Topological Properties:
- Cardinality of set X.
- Cardinality of $\mathcal{T}$.
- Connectedness of space X.
- Being Discrete.

Topological Properties

$X = [0, 2]$, $Y = [0, 4]$

$L_X = 2$, $L_Y = 4$

$f: X \to Y$

$x \mapsto 2x$ Biject. + Cont + Inv. of f is also cont

$X \approx Y$

Some Non Topological Properties:
- Length
- Area
- Boundedness

Topology

Metric on a Set

Metric on a Set



A metric is a distance function i.e.
A function that defines a distance between each pair of elements of a set.

- Distance between A and B is always positive.
- Dis. bet. A and A is "0"
- "0" is the dis. only from A to A.
- Dis bet. A and B = Dis bet. B and A.
- dis. bet A and B $\leq$ dis. bet A and C + dis bet C and B.

Metric on a Set

Metric:
A metric on a non empty set X is a real valued function d defines on $X \times X$, i.e

$$d : X \times X \longrightarrow \mathbb{R}$$

such that for all $x, y, z \in X$, the following conditions are satisfied:

1. $d(x, y) \geq 0$, equality holds iff $x = y$
2. $d(x, y) = d(y, x)$ (Symmetry)
3. $d(x, z) \leq d(x, y) + d(y, z)$ (Triangle Inequality)

Metric on a Set

Metric Space:
A set X with a metric d defined on it is called a metric space.

Topology

Examples of Metric

Examples of Metric

Let us see some examples of metrics and verify all the three axioms for each metric.

Example 1:
Let $X = \mathbb{R}$ and d is defined as $d(x, y) = |x - y|$ where $x, y \in \mathbb{R}$.

Examples of Metric

- $d(x,y) = |x-y| \geq 0$

  $|x-y| = 0 \iff x - y = 0$

  $x = y$

- $d(x,y) = |x-y| = |-(y-x)|$

  $= |y-x| = d(y,x)$

- $x, y, z \in \mathbb{R}$

  $d(x,z) = |x-z| = |x-y+y-z| \leq \underbrace{|x-y|}_{d(x,y)} + \underbrace{|y-z|}_{d(y,z)}$

  $d(x,z) \leq d(x,y) + d(y,z)$

$|a| = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases}$

$|x| = |-x|$

$|a+b| \leq |a| + |b|$

Examples of Metric

Example 2:
Let $X = C[0, 1]$ (the set of all real valued continuous functions on $[0, 1]$) and d is defined as

$$d(f, g) = \int_0^1 |f(x) - g(x)|dx$$

where $f, g \in C[0, 1]$.

Examples of Metric

Example 2:
Let $X = C[0, 1]$ (the set of all real valued continuous functions on $[0, 1]$) and d is defined as

$$d(f, g) = \int_0^1 |f(x) - g(x)|dx$$

where $f, g \in C[0, 1]$.

Sol: $d(f,g) = \int_0^1 |f(x) - g(x)|dx \geq 0$

$d(f,g) = 0 \iff \int_0^1 |f(x) - g(x)|dx = 0$

$\iff f(x) - g(x) = 0 \iff f(x) = g(x) \quad \forall x \in [0,1]$

Examples of Metric

- $d(f,g) = \int_0^1 |f(x) - g(x)|dx = \int_0^1 |g(x) - f(x)|dx$

  $= d(g,f)$

- $d(f,h) = \int_0^1 |f(x) - h(x)|dx = \int_0^1 |f(x) - g(x) + g(x) - h(x)|dx$

  $\leq \int_0^1 |f(x) - g(x)|dx + \int_0^1 |g(x) - h(x)|dx$

  $= d(f,g) + d(g,h)$. □

Topology

More Examples of Metric

More Examples of Metric

Let us see some more examples of metrics.

Example 3:
Let X be any nonempty set and d is defined as

$$d(x,y) = \begin{cases} 0, & \text{if } x = y \\ 1, & \text{if } x \neq y \end{cases}$$

where $x, y \in \mathbb{R}$.

More Examples of Metric

- $d(x,y) = \begin{cases} 0 \\ 1 \end{cases} \geq 0$

  $d(x,y) = 0 \iff x = y$

- $d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases} = \begin{cases} 0 & \text{if } y = x \\ 1 & \text{if } y \neq x \end{cases}$

  $= d(y,x)$

- $x, y, z \in X \qquad d(x,z) = 1 < 1+1 = d(x,y) + d(y,z)$

More Examples of Metric

Example 4:
Let $X = C[0,1]$ (the set of all real valued continuous functions on $[0,1]$) and d' is defined as

$$d'(f,g) = sup\{|f(x) - g(x)| \ : \ x \in [0,1]\}$$

where $f, g \in C[0,1]$.

d' is the max. vertical gap between the function f and g:



Topology

Usual Metric on $\mathbb{R}^n$

Usual Metric on $\mathbb{R}^n$

Recall the metric defined on $\mathbb{R}$ in "Example 1."

$$d(x,y) = |x - y|$$

where $x, y \in \mathbb{R}$.
This is the usual metric in $\mathbb{R}$.

We can also write it in the following way

$$d(x,y) = \sqrt{(x-y)^2}$$

Usual Metric on $\mathbb{R}^n$

Usual Metric on $\mathbb{R}^2$
Consider $\mathbb{R}^2$ and d is defined as

$$d(x,y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

where $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2$.

Usual Metric on $\mathbb{R}^n$

Usual Metric on $\mathbb{R}^n$
Consider $\mathbb{R}^n$ and d is defined as

$$d(x,y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2}$$

where $x = (x_1, x_2, \cdots, x_n), y = (y_1, y_2, \cdots, y_n) \in \mathbb{R}^n$.

Topology

Unusual Metrics on $\mathbb{R}^n$

Unusual Metrics on $\mathbb{R}^n$

Let us see some unusual metrics defined on $\mathbb{R}^2$. These definitions can be generalized for $\mathbb{R}^n$ with $n > 2$.

Unusual Metrics on $\mathbb{R}^n$

Example 1:
Consider $\mathbb{R}^2$ and d_1 is defined as

$$d_1(x, y) = max(|x_1 - y_1|, |x_2 - y_2|)$$

where $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2$.

$x = (2, 3)$, $y = (7, 15)$

$d_1(x, y) = \max(|2-7|, |3-15|)$

$= \max(5, 12) = 12$



Unusual Metrics on $\mathbb{R}^n$

Example 2:
Consider $\mathbb{R}^2$ and d_2 is defined as

$$d_2(x, y) = |x_1 - y_1| + |x_2 - y_2|$$

where $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2$.

$x = (1, 3)$ $y = (7, 2)$

$d_2(x, y) = |1-7| + |3-2|$

$= 6 + 1 = 7$



Topology

Open Ball

Open Ball

Def:
Consider X with a metric d. For any $x \in X$ and any real number $r > 0$, the set of all points $y \in X$ whose distance from x is less than r is called the **open ball** of radius r centered at x.
i.e.

$$B(x, r) = \{y \in X | d(x, y) < r\}$$

Open Ball

Example:
Recall $\mathbb{R}$ with usual metric $d = |x - y|$. Consider $x = 2$ and $r = 1$.
Let us see what is $B(2, 1)$?

Open Ball

Example:
Recall $\mathbb{R}$ with usual metric $d(x,y) = |x - y|$. Consider $x = 2$ and $r = 1$.
Let us see what is $B(2, 1)$?

$B(2,1) = \{y \in \mathbb{R} \mid d(2, y) = |2 - y| < 1\} = (1, 3)$

$|2 - y| < 1$

$\Rightarrow -1 < 2 - y < 1$

$\Rightarrow 1 < y < 3$



Topology

Examples: Open Ball

Examples: Open Ball

We have already discussed one simple example of open ball. Now we will discuss some more examples of open balls with respect to different metrics.

Examples: Open Ball

Example 1:
Recall $\mathbb{R}^2$ with usual metric
$d(x,y) = \sqrt{(x_1-y_1)^2+(x_2-y_2)^2}$ where
$x=(x_1,x_2), y=(y_1,y_2) \in \mathbb{R}^2$.

Examples: Open Ball

Example 1:
Recall $\mathbb{R}^2$ with usual metric
$d(x,y) = \sqrt{(x_1-y_1)^2+(x_2-y_2)^2}$ where
$x=(x_1,x_2), y=(y_1,y_2) \in \mathbb{R}^2$.

$B(x,1) = \{y \in \mathbb{R}^2 \mid d(x,y) < 1\}$

$x=(0,0)$

$\lambda = 1$

Examples: Open Ball

Example 1:
Recall $\mathbb{R}^2$ with usual metric
$d(x,y) = \sqrt{(x_1-y_1)^2+(x_2-y_2)^2}$ where
$x=(x_1,x_2), y=(y_1,y_2) \in \mathbb{R}^2$.

$B(x,\lambda) = \{y \in \mathbb{R}^2 \mid d(x,y) < \lambda\}$

x

$\lambda > 0$

$\mathbb{R}^n$

Examples: Open Ball

Example 2:
Recall trivial metric on a set X. i.e.

$$d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

where $x, y \in X$.

Examples: Open Ball

Example 2:
Recall trivial metric on a set X. i.e.

$$d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

where $x, y \in X$.

$B(x,\lambda) = \{y \in X \mid d(x,y) < \lambda\} = \{x\}$ $\quad \lambda \leq 1$

$B(x,\lambda) = \{y \in X \mid d(x,y) < \lambda\} = X.$ $\quad (\lambda > 1)$

Examples: Open Ball

Example 3:
Recall $\mathbb{R}^2$ with $d_1(x,y) = max(|x_1-y_1|, |x_2-y_2|)$ where
$x=(x_1,x_2), y=(y_1,y_2) \in \mathbb{R}^2$.

$x=(0,0)$

$B(x,\lambda)$



Examples: Open Ball

Example 4:
Recall $C[0,1]$ with $d(f,g) = \int_0^1 |f(x) - g(x)|dx$ where $f, g \in C[0,1]$.

$B(f, \lambda)$

$\lambda = 3$

Topology

Open Ball as Open Set

Open Ball as Open Set

Lemma:
Let d be a metric defined on a nonempty set X. For $x \in X$ and a positive real number r, consider open ball $B(x,r)$. Then for every point $y \in B(x,r)$ there exists an open ball $B'(y,r')$ such that $B'(y,r') \subset B(x,r)$.

Open Ball as Open Set

Proof: Let $y \in B(x,\lambda)$

$\Rightarrow d(x,y) < \lambda$

$\Rightarrow \lambda - d(x,y) > 0$

Let $\lambda' := \lambda - d(x,y)$

Now we want to show that

$B'(y,\lambda') \subset B(x,\lambda)$

Open Ball as Open Set

Let $z \in B'(y,\lambda')$

$\Rightarrow d(z,y) < \lambda' = \lambda - d(x,y)$

$\Rightarrow |d(x,y) + d(y,z)| < \lambda$

$d(x,z) \leq d(x,y) + d(y,z) < \lambda$

$\Rightarrow d(x,z) < \lambda$

$\Rightarrow z \in B(x,\lambda)$

$B'(y,\lambda) \subset B(x,\lambda)$ □

Topology

Open Balls Centered at One Point

Open Balls Centered at One Point

Fact:
Let $B_1(x,r_1)$ and $B_2(x,r_2)$ be two open balls centered at x with radius r_1 and r_2 respectively. Then either $B_1(x,r_1) \subset B_2(x,r_2)$ or $B_2(x,r_2) \subset B_1(x,r_1)$.

Open Balls Centered at One Point

Proof: $B_1(x,\lambda_1)$, $B_2(x,\lambda_2)$

$\lambda_1, \lambda_2 \in \mathbb{R}^+$

$\Rightarrow \lambda_1 \leq \lambda_2$ or $\lambda_2 \leq \lambda_1$

$\Rightarrow B_1(x,\lambda_1) \subset B_2(x,\lambda_2)$ | $B_2(x,\lambda_2) \subset B_1(x,\lambda_1)$

$y \mid d(x,y) < \lambda_1$

Topology

Properties of
Collection of Open Balls

Properties of Collection...

Theorem:
Consider X with a metric d. Let $\mathcal{B}$ be the collection of all open balls in X. Then $\mathcal{B}$ satisfies the following properties.

1. For all $x \in X$ there exists $B(x,r) \in \mathcal{B}$
2. For all $z \in B_1(x,r_1) \cap B_2(y,r_2)$ there exists $B(z,r)$ such that $z \in B(z,r) \subset B_1(x,r_1) \cap B_2(y,r_2)$

Properties of Collection...

Proof. X, d, $\mathcal{B}$

$x \in X \quad \lambda > 0$

$x \in B(x,\lambda) \in \mathcal{B}$

② → Let $z \in B_1(x,\lambda_1) \cap B_2(y,\lambda_2)$

$\Rightarrow z \in B_1(x,\lambda_1)$ and $z \in B_2(y,\lambda_2)$

$\Rightarrow \exists\ B_3(z,\lambda_3) \subset B_1(x,\lambda_1)$ $\Rightarrow \exists\ B_4(z,\lambda_4) \subset B_2(y,\lambda_2)$

$\lambda_3, \lambda_4 \in \mathbb{R}^+$

$\lambda_3 < \lambda_4$

Properties of Collection...

$B_3(z,\lambda_3) \subset B_4(z,\lambda_4) \subset B_2(y,\lambda_2)$

$z \in B_3(z,\lambda_3) \subset B_1(x,\lambda_1) \cap B_2(y,\lambda_2)$

$B_3(z,\lambda_3) = B(z,\lambda)$

□

Topology

Metric Topology

Metric Topology

For a non empty set X we defined a metric d.
With respect to this metric we defined open balls.
And considered $\mathcal{B}$ "Collection of all open balls in X."
This collection $\mathcal{B}$ satisfies the properties of "Basis."
So $\mathcal{B}$ is able to generate a topology on X.

Dr. Hani Shakir

Metric Topology

Metric Topology:
Let X be a nonempty set with metric d. The topology $\mathcal{T}$ on X generated by the set of all open balls in X with respect to d is called the "Metric Topology."
We also say that a topology $\mathcal{T}$ on X induced by the metric d.

Metric Space:
Consider X with a metric d, this d induces a topology the "metric topology." (X,d) is called a metric space.

Topology

Examples: Metric
Topology

Examples: Metric Topology

Let us see some examples of metric topologies. In these examples we will generate the topologies induced by already defined metrics.

Examples: Metric Topology

Example 1:
Recall $\mathbb{R}$ with usual metric $d(x,y) = |x-y|$ where $x, y \in \mathbb{R}$.
Here $\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\}$ is the set of all open intervals.

And we know that the topology generated by the set of all open intervals is usual topology.
So the metric topology induced by $d(x,y) = |x-y|$ on $\mathbb{R}$ is usual topology on $\mathbb{R}$.

Examples: Metric Topology

Example 1:
Recall $\mathbb{R}$ with usual metric $d(x,y) = |x-y|$ where $x, y \in \mathbb{R}$.
Here $\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\}$ is the set of all open intervals.

$B(x,\lambda) = (x-\lambda, x+\lambda)$

And we know that the topology generated by the set of all open intervals is usual topology.
So the metric topology induced by $d(x,y) = |x-y|$ on $\mathbb{R}$ is usual topology on $\mathbb{R}$.

Examples: Metric Topology

Example 2:
Recall $\mathbb{R}^2$ with usual metric
$d(x,y) = \sqrt{(x_1-y_1)^2 + (x_2-y_2)^2}$ where
$x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2$.
Here $\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\}$ is the set of all open disks.

And we know that the topology generated by the set of all open disks is usual topology on $\mathbb{R}^2$. So the metric topology induced by $d(x,y) = \sqrt{(x_1-y_1)^2 + (x_2-y_2)^2}$ on $\mathbb{R}^2$ is the usual topology on $\mathbb{R}^2$.

Examples: Metric Topology

Example 2:
Recall $\mathbb{R}^2$ with usual metric
$d(x,y) = \sqrt{(x_1-y_1)^2 + (x_2-y_2)^2}$ where
$x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2$.
Here $\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\}$ is the set of all open disks.

And we know that the topology generated by the set of all open disks is usual topology on $\mathbb{R}^2$. So the metric topology induced by $d(x,y) = \sqrt{(x_1-y_1)^2 + (x_2-y_2)^2}$ on $\mathbb{R}^2$ is the usual topology on $\mathbb{R}^2$.

Examples: Metric Topology

Example 3:
Recall trivial metric on a set X. i.e.

$$d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

where $x, y \in X$. Here
$\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\} = \{\{x\}, X | x \in X\}$.

And we know that the topology generated by the set of all single tons is discrete topology. So the metric topology induced by the above metric is the discrete topology on X.

Examples: Metric Topology

Example 3:
Recall trivial metric on a set X. i.e.

$$d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

where $x, y \in X$. Here
$\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\} = \{\{x\}, X | x \in X\}$.

And we know that the topology generated by the set of all single tons is discrete topology. So the metric topology induced by the above metric is the discrete topology on X.

Examples: Metric Topology

Example 3:
Recall trivial metric on a set X. i.e.

$$d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

where $x, y \in X$. Here
$\mathcal{B} = \{B(x,r) | x \in X \quad r > 0\} = \{\{x\}, X | x \in X\}$.

And we know that the topology generated by the set of all single tons is discrete topology. So the metric topology induced by the above metric is the discrete topology on X.

Topology

Metrizable Spaces

Metrizable Spaces

Consider a non empty set X with a metric d defined on X. Now this metric induces a topology on X. Let's reverse this process, i.e.
We have a set X with a topology $\mathcal{T}$, now the question is
"Dose there exist a metric d such that d on X induces the topology $\mathcal{T}$?"
The answer depends on the selection of $\mathcal{T}$.

Metrizable Spaces

Metrizable Space:
A topological space $(X, \mathcal{T})$ is called metrizable space if there exists a metric d on X that induces the topology $\mathcal{T}$.
Remark:
Not all spaces are metrizable.

Topology

Examples: Metrizable Spaces

Examples: Metrizable Spaces

Let us see some examples of metrizable spaces, i.e. the topological space such that there exists a metric that induces the given topology.

Dr. Hani Shakir

Examples: Metrizable Spaces

Example 1:
Consider $\mathbb{R}$ with usual topology. Recall usual metric $d(x,y) = |x - y|$ on $\mathbb{R}$.
Now the topology induced by $d(x,y) = |x - y|$ on $\mathbb{R}$ is the usual topology on $\mathbb{R}$.
So $(\mathbb{R}, \mathcal{T}_u)$ is metrizable space.
Example 2:
Consider $\mathbb{R}^2$ with usual topology. Recall usual metric

$$d(x,y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

on $\mathbb{R}^2$. Now the topology induced by d on $\mathbb{R}^2$ is the usual topology on $\mathbb{R}^2$. So $(\mathbb{R}^2, \mathcal{T}_u)$ is metrizable space.

Examples: Metrizable Spaces

Example 3:
Consider X with discrete topology. Recall trivial metric.

$$d(x,y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

where $x, y \in X$.
The topology induced by the above metric is the discrete topology on X. So discrete space is metrizable space.

Topology

Examples:
Non-Metrizable Spaces

Examples: Non-Metrizable Spaces

Let us see some examples of non-metrizable spaces, i.e. the topological space $(X, \mathcal{T})$ such that there exists no metric which induces $\mathcal{T}$.

Examples: Non-Metrizable Spaces

Example 1:
Consider $X \neq \{x\}$ with indiscrete topology i.e. $\mathcal{T}_{ind} = \{\emptyset, X\}$. Then it is not metrizable space.

Reason:
It is because of the fact that in a metric space all the finite set are closed. But here only closed sects are $\emptyset$ and X. So $\mathcal{T}_{ind}$ can not be induced by any metric.

Caution:
It does not mean that if in some topological space all finite sets are closed then that space is metrizable.

Examples: Non-Metrizable Spaces

Example 2:
Let X be an infinite set. Consider cofinite topology on X. Then it is not metrizable space.

Reason:
There is a topological property called "Hausdorffness" which should exist in a metric space. So if there exists some metric corresponds to $(X, \mathcal{T}_{cof})$ then $(X, \mathcal{T}_{cof})$ should be Hausdorff but it is not true.
So $(X, \mathcal{T}_{cof})$ is not metrizable space.

Caution:
It does not mean that if some topological space is Hausdorff then that space is metrizable.

Examples: Non-Metrizable Spaces

Example 3:
$\mathbb{R}$ with lower limit topology is not metrizable space.

Reason:
There are topological spaces called "Second Countable Spaces" and separable metric spaces are second countable. So if there exists some metric corresponds to $(\mathbb{R}, \mathcal{T}_l)$ then $(\mathbb{R}, \mathcal{T}_l)$ should be second countable but it is not true.
So $(\mathbb{R}, \mathcal{T}_l)$ is not metrizable space.

Topology

First Countable Spaces

First Countable Spaces

Def:
A topological space $(X, \mathcal{T})$ is called a "First Countable Space" if for each $p \in X$ there exists a countable class $\mathcal{B}_p$ of open subsets of X containing p such that for every $U_p \in \mathcal{T}$ (open set containing p) there exists $B_p \in \mathcal{B}_p$ such that $p \in B_p \subset U_p$.

First Countable Spaces

Def:
A topological space $(X, \mathcal{T})$ is called a "First Countable Space" if for each $p \in X$ there exists a countable class $\mathcal{B}_p$ of open subsets of X containing p such that for every $U_p \in \mathcal{T}$ (open set containing p) there exists $B_p \in \mathcal{B}_p$ such that $p \in B_p \subset U_p$.

First Countable Spaces

Def:
A topological space $(X, \mathcal{T})$ is called a "First Countable Space" if for each $p \in X$ there exists a countable local base at p.

First Countable Spaces

Remark:
Every subspace of a first countable space is first countable.

$A \subset X$ (X is first countable)
$\Rightarrow$ A is also first countable.

Proof: Let X be a first countable space and $A \subset X$, Let $p \in A$ $\therefore X$ is first countable.
So $\exists$ a countable local base at p.
$\mathcal{B}_p = \{B_n \mid n \in \mathbb{N}^*\}$.
Now $\mathcal{B}_{p_A} = \{A \cap B_n \mid n \in \mathbb{N}^*\}$

First Countable Spaces

Now $\mathcal{B}_{p_A}$ is a local base at p. for $\mathcal{T}_A$
And $\mathcal{B}_{p_A}$ is countable So
A is first countable. □

Topology

Examples:
First Countable Spaces

Examples: First Countable Spaces

Let us discuss some examples of first countable spaces.
Example 1:
Discrete space is first countable.
Remember a local base at a point p in discrete space, i.e.

$$\mathcal{B}_p = \{\{p\}\}$$

and it is finite.

Examples: First Countable Spaces

Example 2:
Let $X = \{a, b, c, d\}$ and

$$\mathcal{T} = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}, X\}$$

be a topology defined on X.

- $\mathcal{B}_a = \{\{a\}\}$
- $\mathcal{B}_b = \{\{b\}\}$
- $\mathcal{B}_c = \{\{c, d\}\}$
- $\mathcal{B}_d = \{\{c, d\}\}$

Remark: Every finite set with any top. is first countable.

Examples: First Countable Spaces

Example 3:
$\mathbb{R}$ with usual topology is first countable.
For any real number x one may consider the following countable local base at x.

$$\mathcal{B}_x = \left\{\left(x - \frac{1}{n}, x + \frac{1}{n}\right) \mid n \in \mathbb{Z}^+\right\}$$

Examples: First Countable Spaces

Example 3:
$\mathbb{R}$ with usual topology is first countable.
For any real number x one may consider the following countable local base at x.

$$\mathcal{B}_x = \left\{\left(x - \frac{1}{n}, x + \frac{1}{n}\right) \mid n \in \mathbb{Z}^+\right\}$$

$\mathcal{B}_x$ is countable
$\Rightarrow$ $(\mathbb{R}, \mathcal{T}_u)$ is first countable

Examples: First Countable Spaces

Example 4:
Every metric space is first countable.

Sol: Let X be a metric space $(X, d, \mathcal{T})$.
Now consider the following collection of open balls for $p \in X$.

$$\mathcal{B}_p = \{ B(p, \tfrac{1}{n}) \mid n \in \mathbb{N} \}.$$

$\mathcal{B}_p$ is a local base at p. $\mathcal{B}_p$ is countable
$\Rightarrow$ X is first countable.

Topology

Second Countable Spaces

Second Countable Spaces

Def:
A topological space $(X, \mathcal{T})$ is called a "Second Countable Space" if there exists a countable class $\mathcal{B}$ of open subsets of X such that for every $U \in \mathcal{T}$ and for all $x \in U$ there exists $B \in \mathcal{B}$ such that $x \in B \subset U$.

Second Countable Spaces

Def:
A topological space $(X, \mathcal{T})$ is called a "Second Countable Space" if there exists a countable class $\mathcal{B}$ of open subsets of X such that for every $U \in \mathcal{T}$ and for all $x \in U$ there exists $B \in \mathcal{B}$ such that $x \in B \subset U$.

Dr. Hani Shakir

Second Countable Spaces

Def:
A topological space $(X, \mathcal{T})$ is called a "Second Countable Space" if there exists a countable basis for $\mathcal{T}$.

Second Countable Spaces

Remark:
Every subspace of a second countable space is second countable.

Proof: Let $A \subset X$ (X is second countable space)
So $\exists$ $\mathcal{B} = \{ B_n \mid n \in \mathbb{N} \}$ countable.
Now $\mathcal{B}_A = \{ A \cap B_n \mid n \in \mathbb{N} \}$ is a basis for $\mathcal{T}_A$. And $\mathcal{B}_A$ is countable
$\Rightarrow$ A is second countable. □

Topology

Examples: Second Countable Spaces

Examples: Second Countable

Let us see discuss some examples of second countable spaces.

Example 1:
Let X be a countable set with discrete topology. Then $(X, \mathcal{T}_{Dis})$ is second countable. Recall basis for discrete space, i.e.

$$\mathcal{B} = \{\{x\} | x \in X\}$$

Examples: Second Countable

Example 2:
Let $X = \{a, b, c, d\}$ and

$$\mathcal{T} = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}, X\}$$

be a topology defined on X. A basis for $\mathcal{T}$ is

$$\mathcal{B} = \{\{a\}, \{b\}, \{c, d\}\}$$

Examples: Second Countable

Example 2:
Let $X = \{a, b, c, d\}$ and

$$\mathcal{T} = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}, X\}$$

be a topology defined on X. A basis for $\mathcal{T}$ is

$$\mathcal{B} = \{\{a\}, \{b\}, \{c, d\}\}$$

Remark: Any finite set with any top is second countable.

Examples: Second Countable

Example 2:
Let $X = \{a, b, c, d\}$ and

$$\mathcal{T} = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}, X\}$$

be a topology defined on X. A basis for $\mathcal{T}$ is

$$\mathcal{B} = \{\{a\}, \{b\}, \{c, d\}\}$$ finite.

basis

So $(X, \mathcal{T})$ is second countable.

Examples: Second Countable

Example 3:
$\mathbb{R}$ with usual topology is second countable.
Consider the following basis for usual topology on $\mathbb{R}$.

$$\mathcal{B} = \{(a, b) \,|\, a, b \in \mathbb{Q}\}$$

Examples: Second Countable

Example 3:
$\mathbb{R}$ with usual topology is second countable.
Consider the following basis for usual topology on $\mathbb{R}$.

$$\mathcal{B} = \{(a, b) \,|\, a, b \in \mathbb{Q}\}$$

Countable basis

So $(\mathbb{R}, \mathcal{T}_u)$ is second countable.

Ex: 4. $(\mathbb{R}, \mathcal{T}_{Dis})$ is not second countable.

$\mathcal{B}_{\mathcal{T}_{Dis}} = \{\{x\} \mid x \in \mathbb{R}\}$:

Examples: Second Countable

Example 5:
$\mathbb{R}^2$ with usual topology is second countable.
Consider a basis $\mathcal{B}$ for usual topology on $\mathbb{R}^2$ consists of collection of open disks with rational radii and centers with rational coordinates. Such collection is countable.

$r, x, y \in \mathbb{Q}$

Topology

Second Countability implies First Countability

Second Countability implies...

- First Countability Axiom
- Second Countability Axiom

Theorem:
Let a topological space $(X, \mathcal{T})$ is second countable space. Then it also first countable space.

Second Countability implies...

Proof: Let X be a second countable space.
i.e $\exists$ $\mathcal{B}$ (a basis) s.t $\mathcal{B}$ is countable.
Now let $p \in X$ and consider
$\mathcal{B}_p \subset \mathcal{B}$ $\quad \mathcal{B}_p = \{B_p \mid p \in B_p\} \subset \mathcal{B}$
Now $\mathcal{B}_p$ is countable $\because$ $\mathcal{B}$ is countable
Now $\mathcal{B}_p$ is local base at p:

Second Countability implies...

$\mathcal{B}_p \subset \mathcal{B}$ (basis) $\quad \forall\ U \in \mathcal{T}$ s.t $p \in U$
$\exists\ B_p \in \mathcal{B}_p$ s.t
$p \in B_p \subset U$
$\Rightarrow$ $\mathcal{B}_p$ is a local base at p.
And this exists $\forall$ $p \in X$.
$\Rightarrow$ X is first countable $\square$

Topology

First Countability
$\not\Rightarrow$ Second Countability

First Countability $\not\Rightarrow$...

It has already been proved that every second countable space is first countable.
But what about its reveres implication, i.e.
For a first countable space what one can say about its second countability?
Before answering this question, let us first discuss some examples.

First Countability $\not\Rightarrow$...

Example 1:
Let X be an infinite set with discrete topology. Then it is first countable space. Since for any point $p \in X$.

$$\mathcal{B}_p = \{\{p\}\}$$

is a local base at p. And it is finite so it is first countable. But it is not second countable.

$$\mathcal{B} = \{\{x\} | x \in X\}$$

is the smallest basis of discrete space. Since X is infinite so $\mathcal{B}$ is infinite.

First Countability $\not\Rightarrow$...

Example 2:
Consider $\mathbb{R}$ with usual topology. It is both first countable and second countable.

Example 3:
Consider $\mathbb{R}$ with lower limit topology. It is first countable but not second countable.

Remark:
A first countable space may or may not be second countable.

Topology

Open Cover

Open Cover

Let $X = [0,1]$



$\mathcal{U} = \{[-1, 2]\}$
$[0,1] \subset [-1, 2]$ $\mathcal{U}$ is cover of $[0,1]$
$\mathcal{V} = \{[0, \frac{1}{2}), (\frac{1}{4}, 1.9], (1.8, 3)\}$

Cover:
A cover of a set X is a collection of sets

$$\mathcal{U} = \{U_i | i \in I\}$$

where I is index set, such that

$$X \subset \bigcup_{i \in I} U_i$$

Open Cover



Cover:
A cover of a set X is a collection of sets

$$\mathcal{U} = \{U_i | i \in I\}$$

where I is index set, such that

$$X \subset \bigcup_{i \in I} U_i$$

Open Cover

Cover:(In Topology)
Let $(X, \mathcal{T})$ be topological space. A cover of a X is a collection of subsets of X, i.e.

$$\mathcal{U} = \{U_i | U_i \subset X, i \in I\}$$

where I is index set, such that

$$X = \bigcup_{i \in I} U_i$$

Open Cover

Open Cover:
Let $(X, \mathcal{T})$ be topological space. An open cover of a X is a collection of open subsets of X, i.e.

$$\mathcal{U} = \{U_i | U_i \in \mathcal{T}, i \in I\}$$

where I is index set, such that

$$X = \bigcup_{i \in I} U_i$$

Open Cover

Open Cover of Subset:
Let $(X, \mathcal{T})$ be topological space and $Y \in X$. An open cover of a Y is a collection of open subsets of X, i.e.

$$\mathcal{C} = \{C_i | C_i \in \mathcal{T}, i \in I\}$$

where I is index set, such that

$$Y \subset \bigcup_{i \in I} C_i$$

Topology

Examples: Open Cover

Examples: Open Cover

Let us see some examples of open cover for the better understanding of the definitions we discussed in previous topic.

Examples: Open Cover

Example 1:
Let $X = \{a, b, c, d\}$ with topology

$$\mathcal{T} = \{\phi, \{a\}, \{a, b\}, \{a, c, d\}, X\}$$

Examples: Open Cover

Example 1:
Let $X = \{a, b, c, d\}$ with topology

$$\mathcal{T} = \{\phi, \{a\}, \{a, b\}, \{a, c, d\}, X\}$$

Examples: Open Cover

Example 1:
Let $X = \{a, b, c, d\}$ with topology

$$\mathcal{T} = \{\phi, \{a\}, \{a, b\}, \{a, c, d\}, X\}$$

$\mathcal{U} = \{\{a, b\}, \{a, c, d\}\}$ open cover of X

Examples: Open Cover

Example 2:
Let $X = \mathbb{R}$ with usual topology and consider the following collections of subset of $\mathbb{R}$

$$\mathcal{U} = \{(-n, n) | n \in \mathbb{N}\}$$

$$\mathcal{V} = \{(n, n+2) | n \in \mathbb{Z}\}$$

$$\mathcal{W} = \{(n, n+1) | n \in \mathbb{Z}\}$$

$$\mathcal{Y} = \{[n, n+1] | n \in \mathbb{Z}\}$$

Examples: Open Cover

Example 2:
Let $X = \mathbb{R}$ with usual topology and consider the following collections of subset of $\mathbb{R}$

$\mathcal{U} = \{(-n, n) | n \in \mathbb{N}\}$ open cover

$\mathcal{V} = \{(n, n+2) | n \in \mathbb{Z}\}$ open cover



$\mathcal{W} = \{(n, n+1) | n \in \mathbb{Z}\}$ Not a cover of $\mathbb{R}$



$\mathcal{Y} = \{[n, n+1] | n \in \mathbb{Z}\}$ $\mathcal{Y}$ is a cover

Examples: Open Cover

Example 3:
Let $X = \mathbb{R}$ with usual topology and consider $(0, 1) \subset \mathbb{R}$ and $[0, 1) \subset \mathbb{R}$. The following collections of subset of $\mathbb{R}$.

$$\mathcal{U} = \{(1/n, 1) | n \in \mathbb{N}^*\}$$

$$\mathcal{V} = \{(n, n+2) | n \in \mathbb{Z}\}$$

$\mathcal{V}$ is an open cover of $(0,1)$ but it is not an open cover of $[0,1)$

$\mathcal{U}$ is an open cover of $(0,1)$ and open cover of $[0,1)$.

Topology

Open Subcover

Open Subcover

Subcover:
Let $(X, \mathcal{T})$ be topological space. Consider $\mathcal{C}$ covers X, i.e.

$$\mathcal{C} = \{C_i | C_i \subset X\}$$

such that $X = \bigcup_i C_i$.
A subclass $\mathcal{D}$ of $\mathcal{C}$ that also covers X is called an open subcover of $\mathcal{C}$.

Open Subcover

Open Subcover:
Let $(X, \mathcal{T})$ be topological space. Consider $\mathcal{C}$ be an open cover of X, i.e.

$$\mathcal{C} = \{C_i | C_i \in \mathcal{T}\}$$

such that $X = \bigcup_i C_i$.
A subclass $\mathcal{D}$ of $\mathcal{C}$ that also covers X is called an open subcover of $\mathcal{C}$.

Open Subcover

Example 1:
Let $X = \{a, b, c, d\}$ with topology

$$\mathcal{T} = \{\phi, \{a\}, \{a, b\}, \{c, d\}, \{a, c, d\}, X\}$$

$\mathcal{C} = \{\{a\}, \{a, b\}, \{c, d\}\}$ is an open cover of X.

$\mathcal{D} = \{\{a, b\}, \{c, d\}\}$ it again covers X so $\mathcal{D}$ is a subcover of $\mathcal{C}$

$\mathcal{U} = \{\{a\}, \{c, d\}\}$

Examples: Open Cover

Example 2:
Let $X = \mathbb{R}$ with usual topology and consider

$$\mathcal{U} = \{(-n, n) | n \in \mathbb{N}\}$$

$\mathcal{D} \subset \mathcal{U}$

$\mathcal{D} = \{(-n, n) \mid n \in \mathbb{N} - \{1, 2, 3, 4, 5\}\}$

Examples: Open Cover

Example 2:
Let $X = \mathbb{R}$ with usual topology and consider

$$\mathcal{U} = \{(-n, n) | n \in \mathbb{N}\}$$



$\mathcal{D} \subset \mathcal{U}$

$\mathcal{D} = \{(-n, n) \mid n \in \mathbb{N} - \{1, 2, 3, 4, 5\}\}$

Topology

Lindelöf Space

Lindelöf Space

Lindelöf Space:
A topological space $(X, \mathcal{T})$ is called a Lindelöf space if and only if every open cover of X has a countable subcover.

Lindelöf Space

Example 1:
Let X be a finite set with any topology then it is Lindelöf.

Example 2:
A set X with indiscrete topology then it is Lindelöf.

Example 3:
$X = \mathbb{R}$ with usual topology is Lindelöf.

Lindelöf Space

Remark:
A subspace of a Lindelöf space needs not be Lindelöf.

Topology

Second Countable
and
Lindelof Spaces

Second Countable...

Theorem:
Every second countable space is Lindelöf space.

Second Countable...

Proof: Let $(X, \mathcal{T})$ be a second countable space. i.e $\exists$ a countable basis for $\mathcal{T}$

Say $\mathcal{B} = \{B_n\}$

Let $\mathcal{A}$ be an open cover of X.

$$X = \bigcup_i A_i \quad A_i \in \mathcal{A} \subset \mathcal{T}$$

Now for each positive integer "n" choose an element A_n of $\mathcal{A}$ s.t $B_n \subset A_n$.

Second Countable...

Now collect all such elements of $\mathcal{A}$

$$\mathcal{A}' = \{A_n \mid B_n \subset A_n\} \subset \mathcal{A}$$

Now $\mathcal{A}'$ is countable.

Second Countable...

Now collect all such elements of $\mathcal{A}$

$$\mathcal{A}' = \{A_n \mid B_n \subset A_n\} \subset \mathcal{A}$$

Now $\mathcal{A}'$ is countable.

Now we have to show that $\mathcal{A}'$ covers X.

Let $x \in X$ and consider an element A of $\mathcal{A}$ such A exists $\because \mathcal{A}$ is a open cover.

Moreover this one A is open

Second Countable...

$x \in A$, $\exists$ $B_n \in \mathcal{B}$ (Basis)

s.t $x \in B_n \subset A$

But for B_n $\exists$ $A_n \in \mathcal{A}'$

s.t $x \in B_n \subset A_n$

$\Rightarrow$ $X = \bigcup_i A_i$ $\quad A_i \in \mathcal{A}'$

$\Rightarrow$ $\mathcal{A}'$ covers X So X is Lindelöf.

Topology

Separable Spaces

Separable Spaces

Def:
A topological space $(X, \mathcal{T})$ is said to be "Separable" if there exists a countable dense subset A of X.
i.e.
$\exists$ $A \subset X$ such that
1. A is countable.
2. $\overline{A} = X$.

Separable Spaces

Def:
A topological space $(X, \mathcal{T})$ is said to be "Separable" if there exists a sequence of points

$$\{x_n\}_{n=1}^{\infty}$$

in X such that every nonempty subset of X contains at least one point of the sequence.

Separable Spaces

Remark:
Every second countable space is separable. The converse of this is not true in general.

Counter Example:
$\mathbb{R}$ with lower limit topology is separable but not second countable.

Topology

Examples: Separable Spaces

Examples: Separable Spaces

Let us discuss some examples of separable spaces.

Example 1:
Let X be a finite set. Then X with any topology is separable.

Example 2:
Consider a set X with indiscrete topology. Then X separable.

Examples: Separable Spaces

Example 3:
$\mathbb{R}$ with usual topology is separable.

Reason:
Since there exists $\mathbb{Q}$ in $\mathbb{R}$ that is countable and

$$\overline{\mathbb{Q}} = \mathbb{R}.$$

Example 4:
$\mathbb{R}$ with discrete topology is not separable.

Reason:
Since there exists no countable subset in $\mathbb{R}$ that is dense in $\mathbb{R}$.

Topology

Some More Examples: Separable Spaces

Some More Examples

Let us discuss some more examples of separable spaces. We will discuss these examples in the form of propositions.

Some More Examples

Proposition 1:
A discrete space is separable if and only if it is countable.

Proof: X, $\mathcal{T}_{dis} = P(X)$
We know that the only closed subset in X is X itself.
So the only choice for A is X s.t $\overline{A} = X$
$\Rightarrow$ X is separable iff X is countable. □

Some More Examples

Proposition 2:
(X, T_{cof}) is separable.

Proof: If X is finite $\Rightarrow \mathcal{T}_{cof} = \mathcal{T}_{dis}$
$\Rightarrow$ X is separable.
Now Let X is not finite.
Now consider $A \subset X$ (Countable (Not finite))

Recall $\mathcal{T}_{cof} = \{U \subset X \mid U^c = \text{finite on } X\}$

Some More Examples

$B \subset X$ (closed) $\Rightarrow$ B is finite

Now recall the closure of a set is the smallest closed superset of the set. So $\overline{B} = B$
Now for A the only closed superset of A is X.
$\Rightarrow$ $\overline{A} = X$ $\Rightarrow$ X is sep

Topology

Subspace of Separable Space

Subspace of Separable Space

As we proved that the subspace of a first countable space is first countable and the subspace of a second countable space is second countable. Now what about the subspace of a separable space? i.e.

Is subspace of a separable is separable?

Not in general.

Subspace of Separable Space

Example:
Consider $\mathbb{R}^2$ with topology generated by

$$\mathcal{B} = \bigcup_{a,b,c,d \in \mathbb{R}} \{(x,y) | a \leq x < b, c \leq y < d\}.$$

Now it is separable.

Subspace of Separable Space

Example:
Consider $\mathbb{R}^2$ with topology generated by

$$\mathcal{B} = \bigcup_{a,b,c,d \in \mathbb{R}} \{(x,y) | a \leq x < b, c \leq y < d\}.$$

Now it is separable. $\{(x,y) \mid a \leq x < b,\ c \leq y < d\} = [a,b) \times [c,d)$ in $\mathbb{R}^2$



Subspace of Separable Space

Consider

$$A = \{(x,y) | x = -y\} \subset \mathbb{R}^2$$

with relative topology on A relative to the said topology on $\mathbb{R}^2$.

$\mathcal{T}_A = \{ -- , \{q\}, -- \}$

" Discrete top on A



And we know that Discrete space is separable iff A is countable
But here A is not countable.
$\therefore$ $(A, \mathcal{T}_A)$ is not separable.

Topology

Metric Spaces and Separability

Metric Spaces and Separability

We have seen that every metric space is first countable space but we did not discuss about the second countability and separability of metric spaces. In fact a metric space is not second countable and not separable in general.

Example:
An infinite set X with trivial metric is not second countable and not separable.

Metric Spaces and Separability

Theorem:
A separable metric space is second countable.

Proof: Now X is separable. So $\exists\ A \subset_{\text{countable}} X$ s.t $\overline{A} = X$

Now consider a class of all open balls of the kind $B(a,r)$ where $a \in A$, $r \in \mathbb{Q}$

i.e $\mathcal{B} = \{B(a,r) \mid a \in A,\ r \in \mathbb{Q}\}$

$\because$ A and $\mathbb{Q}$ both countable So $\mathcal{B}$ is:

Metric Spaces and Separability

Now for showing X is second countable we have to prove that $\mathcal{B}$ is a basis for the metric top. on X.

Let $U \in \overline{S}$ and $x \in U$

$\because$ X is a metric space so

$\exists\ \varepsilon,\ B(x,\varepsilon)$ and $x \in B(x,\varepsilon) \subset U$ — (I)

$\because$ A is dense in X so $\exists$ a point $a \in A$ s.t $d(a,x) < \varepsilon$ Let $d(a,x) < \frac{1}{3}\varepsilon$.

Metric Spaces and Separability

Now consider a rational no δ s.t.

$$\frac{1}{3}\varepsilon < \delta < \frac{2}{3}\varepsilon$$

and consider $B(a,\delta)$

$x \in B(a,\delta)$ — (II) $\because d(a,x) < \frac{1}{3}\varepsilon < \delta$

Now Let $b \in B(a,\delta) \Rightarrow d(a,b) < \delta$

$$d(x,b) \le d(x,a) + d(a,b)$$
$$< \frac{1}{3}\varepsilon + \delta < \frac{1}{3}\varepsilon + \frac{2}{3}\varepsilon = \varepsilon$$

Metric Spaces and Separability

$\Rightarrow$ $B(a,\delta) \subset B(x,\varepsilon)$

By (I)

$x \in B(a,\delta) \subset B(x,\varepsilon) \subset U$

Countable $\mathcal{B}$ $a \in A$, $\delta \in \mathbb{Q}$

$\Rightarrow$ $\mathcal{B}$ is a basis for the metric sp X

$\Rightarrow$ X is second countable.

Topology

Separation Axioms

Separation Axioms

1. T_0 Spaces
2. T_1 Spaces
3. T_2 (Hausdorff) Spaces
4. Regular Spaces
5. T_3 Spaces
6. Normal Spaces
7. T_4 Spaces

Topology

T_0 Space

T_0 Space

Def:
A topological space (X, T) is said to be "T_0 Space" iff for each $x, y \in X$ such that $x \neq y$ there exists an open set $U_x \subset X$ containing x such that $y \notin U_x$

or

there exists an open set $V_y \subset X$ containing y such that $x \notin V_y$.

Dr. Hani Shakir

T_0 Space

Remark:
T_0 property is a topological property.

Proof: Let X be space that is T_0

Consider $f: X \longrightarrow Y$ f is bijection + Continuous f^{-1} is cont.

Now $x, y \in X$ $x \neq y$

$\exists\ U_x \ni x$ (open set)

s.t $y \notin U_x$ Now $f(U_x) \subset$ open (Y)

moreover $f(y) \notin f(U_x)$

$\Rightarrow$ Y is T_0.

Topology

Examples: T_0 Space

Examples: T_0 Space

Let us discuss some examples of T_0 spaces.

Example 1:
Every discrete space is T_0.

Example: 2:
Let $X \neq \{a\}$
Consider $\mathcal{T}_{indis} = \{\phi, X\}$
$(X, \mathcal{T}_{indis})$ is not T_0

Examples: T_0 Space

Example 3: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\phi, \{a\}, \{a, b\}, \{a, c\}, \{a, b, c\}, X\}$$

Examples: T_0 Space

Example 3: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\phi, \{a\}, \{a, b\}, \{a, c\}, \{a, b, c\}, X\}$$

$(X, \mathcal{T})$ is T_0.

Examples: T_0 Space

Example 4: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\phi, \{a, c\}, \{b, d\}, X\}$$

Examples: T_0 Space

Example 4: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\phi, \{a, c\}, \{b, d\}, X\}$$

So $(X, \mathcal{T})$ is not T_0

Topology

T_1 Space

T_1 Space

Def:
A topological space $(X, \mathcal{T})$ is said to be "T_1 Space" iff for each $x, y \in X$ such that $x \neq y$ there exist open subsets U_x, U_y of X containing x, y respectively such that $y \notin U_x$ and $x \notin U_y$.

Dr. Hani Shakir

T_1 Space



T_1 Space

Remark:
T_1 property is a topological property.

Topology

Examples: T_1 Space

Examples: T_1 Space

Let us discuss some examples of T_1 spaces.

Example 1:
Every discrete space is T_1.

Example 2:
Every metric space is T_1.



Examples: T_1 Space

Example 3: Every cofinite space is T_1.

Sol Let $(X, \mathcal{T}_{cof})$
Let X is finite.
$\mathcal{T}_{cof} = P(X)$ discrete.
If X is not finite. Let $U \in \mathcal{T}_{cof}$
iff $U = \phi$ or U^c = finite.
$x, y \in X$ $x \in U_x = X \setminus \{y\}$ $y \in U_y = X \setminus \{x\}$
but $y \notin U_x$ but $x \notin U_y$

Examples: T_1 Space

Example 4: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\phi, \{a\}, \{b\}, \{a, b\}, X\}$$

Examples: T_1 Space

Example 4: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\phi, \{a\}, \{b\}, \{a, b\}, X\}$$

$a, b \in \{c\}^c$, $\{b\}^c$ but $a \notin \{b\}^c$

c, d

$(X, \mathcal{T})$ is not T_1.

Topology

$T_0 \not\Longrightarrow T_1$

$T_0 \not\Longrightarrow T_1$

T_1 property is much stronger property than T_0 property.

i.e.

Every T_1 space is T_0 but not every T_0 space is T_1.
Let us prove first part of the claim and give an example for the second part.

$T_0 \not\Longrightarrow T_1$

$T_1 \Rightarrow T_0$
$\not\Leftarrow$

$T_0 \not\Longrightarrow T_1$

Example: A T_0 space that is not T_1.
Consider $\mathbb{R}$ with $\mathcal{T}$ generated by

$$\{(a,\infty) | a \in \mathbb{R}\}$$

$a, b \in \mathbb{R}$ $\quad a > b$

$b < c < a$

$a \in U_a = (c, \infty) \not\ni b$ So T_0-Space



Topology

Criteria for T_1 Space

Criteria for T_1 Space

We have discussed T_1 space and its examples.
Now we will state and prove a very simple criteria to check T_1 property in some space.

Criteria for T_1 Space

Theorem:
A topological space $(X, \mathcal{T})$ is T_1 iff every singleton subset $\{x\} \subset X$ is closed.

Proof: Let $(X, \mathcal{T})$ is T_1 space
Now we have to show that $\forall\, x \in X$
$\{x\} \subset X$ is closed $\Rightarrow \{x\}^c \in \mathcal{T}$
Let $y \in \{x\}^c$ so $x \neq y$
So $\exists\, U_y$ open s.t $x \notin U_y$ $\because X$ is T_1

Criteria for T_1 Space

$\Rightarrow \quad y \in U_y \subset \{x\}^c$
$\Rightarrow \{x\}^c$ is open in X
$\Rightarrow \{x\}$ is closed in X

For the other side.
Let us each singleton subset $\{x\} \subset X$ is closed.

Criteria for T_1 Space

$x \neq y \in X$
$x \in U_x = X \setminus \{y\} = \{y\}^c \in \mathcal{T}$
$y \in U_y = X \setminus \{x\} = \{x\}^c \in \mathcal{T}$

$\Rightarrow (X, \mathcal{T})$ is a T_1 space

Topology

Subspace of T_1 Space

Subspace of T_1 Space

In this topic we will discuss about the subspace of T_1 space. In fact the main aim here is to show that every subspace of a T_1 space is T_1.

Subspace of T_1 Space

Theorem:
Every subspace of a T_1 space is also a T_1 space.

Proof: Let $(X,\mathcal{T})$ be a T_1 space $Y \subset X$ and $(Y,\mathcal{T}_Y)$ be a subspace of $(X,\mathcal{T})$
We have already discussed that if in a space every singleton subset is closed then that space is T_1
Let $p \in Y \subset X$ $\because$ X is T_1.
So $\{p\} \subset X$ closed.

Subspace of T_1 Space

Ta $\{p\}^c$ is open in X
So $\rightarrow$ $Y \cap \{p\}^c$ is open in Y
But $Y \cap \{p\}^c = Y \setminus \{p\}$
$\Rightarrow$ $\{p\}$ is closed in Y
$\Rightarrow (Y, \mathcal{T}_Y)$ is T_1.

Topology

Properties of T_1 Spaces: I

Properties of T_1 Spaces: I

Let us discuss some properties of T_1 spaces. Some of these properties can also be used as a criterion of T_1 property.

Property I: Every finite subset of T_1 space is closed.

Properties of T_1 Spaces: I

Proof: Let $(X,\mathcal{T})$ be a T_1 space
$A \subset X$ finite Let $A = \{a_1, \cdots, a_n\}$
$A = \bigcup_{i=1}^{n} \{a_i\}$ $a_i \in A$ $\forall i$
$\because$ X is T_1 So every singleton subset of X is closed in X.
$\{a_i\}$ is closed $\forall$ $i = 1, 2, \cdots, n$.
$\Rightarrow$ A is a finite union of closed subset
$\Rightarrow$ A is closed in X.

Properties of T_1 Spaces: I

Property II:
Let X be a finite set. Then $(X,\mathcal{T})$ is a T_1 space if and only if $(X,\mathcal{T})$ is discrete space.

Proof: Let X is finite and $(X,\mathcal{T})$ is T_1 space. So in X every finite subset is closed.
$\Rightarrow$ Every subset of X is also open in X
$\Rightarrow \mathcal{T} = P(X)$ $\Rightarrow$ X is discrete.
Now If $(X,\mathcal{T})$ is discrete $\Rightarrow (X,\mathcal{T})$ is T_1

Topology

Properties of T_1 Spaces: II

Properties of T_1 Spaces: II

Property III:
Let A be a finite subset of a T_1 space X then A' is empty.
Proof:
Let A is finite subset of X, say A has n elements, i.e.

$$\{a_1, \ldots, a_n\}$$

Since A is finite and X is a T_1 space and we know that every finite subset of a T_1 space is closed so

$$A' \subset A$$

Now let us check one by one the points of A as limits points of A.

Properties of T_1 Spaces: II

Let's start with a_1.
Now note that

$$U_{a_1} = \{a_2, \ldots, a_n\}^c$$

is an open set containing a_1 and

$$A \cap (U_{a_1} \setminus \{a_1\}) = \emptyset$$

So a_1 is not a limit point of A.
Similarly, no other point of A is a limit point of A. So

$$A' = \emptyset$$

Topology

T_2/Hausdorff Spaces

T_2/Hausdorff Spaces

Def:
A topological space $(X, \mathcal{T})$ is said to be "T_2 Space" iff for every distinct pair of elements $x, y \in X$ there exist open subsets U_x, U_y of X containing x, y respectively such that $U_x \cap U_y = \emptyset$.

T_2 spaces are also called Hausdorff Spaces.

T_2/Hausdorff Spaces

Felix Hausdorff
(1868 - 1942)

Def:
A topological space $(X, \mathcal{T})$ is said to be "T_2 Space" iff for every distinct pair of elements $x, y \in X$ there exist open subsets U_x, U_y of X containing x, y respectively such that $U_x \cap U_y = \emptyset$.

T_2 spaces are also called Hausdorff Spaces.

T_2/Hausdorff Spaces



Remark:
T_2 property is topological Prop.

Topology

Examples: Hausdorff Spaces

Examples: Hausdorff Spaces

In this topic we will discuss some examples of Hausdorff spaces.

Example 1: Every discrete space is Hausdorff.

Example 2: A nonempty set $X \neq \{x\}$ with indiscrete topology is not Hausdorff.

Examples: Hausdorff Spaces

Example 3: $\mathbb{R}$ with lower limit topology is Hausdorff.

$[a,b) \quad (a,b)$

$x, y \in \mathbb{R} \quad x < y$

$x \in [x,c), [y,d) \ni y \quad x < c < y < d$

$U_x \cap U_y = \emptyset$

Examples: Hausdorff Spaces

Example 4: $\mathbb{R}$ with topology generated

$$\{(a,\infty) | a \in \mathbb{R}\}$$

is not Hausdorff.

$3, 6 \in \mathbb{R}$

$6 \in (5,\infty)$

$3 \in (a,\infty) \not\ni 6 \quad a < 3$

Examples: Hausdorff Spaces

Example 5: An infinite set with cofinite topology is not Hausdorff space.

Topology

Subspace of Hausdorff Space

Subspace of Hausdorff Space

In this topic we will discuss about Hausdorffness of a subspace of a Hausdorff space. In fact the main aim here is to show that the property of being a Hausdorff space is inherently.

Dr.

Subspace of Hausdorff Space

Theorem:
Let $(X, \mathcal{T})$ is a Hausdorff space and $A \subset X$ then $(A, \mathcal{T}_A)$ is Hausdorff.

Proof: Let $A \subset X$

Given $(X, \mathcal{T})$ is Hausdorff.

$\Rightarrow \forall\ x, y \in X \quad x \neq y$

$\exists\ U_x, U_y \in \mathcal{T} \Rightarrow U_x \cap U_y = \emptyset$

$x \in U_x, \ y \in U_y$

$a, b \in A \subset X \quad \exists\ a \in U_a, b \in U_b \in \mathcal{T}$ s.t

$U_a \cap U_b = \emptyset$

Subspace of Hausdorff Space

Now $U_a' = A \cap U_a$

$U_b' = A \cap U_b \in \mathcal{T}_A$

Now $a \in U_a', \quad b \in U_b'$

$U_a' \cap U_b' = (A \cap U_a) \cap (A \cap U_b)$

$= A \cap (U_a \cap U_b)$

$= A \cap (\emptyset) = \emptyset$

$\Rightarrow (A, \mathcal{T}_A)$ is Hausdorff. $\square$

Topology

Metric Spaces are Hausdorff

Metric Spaces are Hausdorff

Recall the topic where we discussed some examples of "Non Metrizable Spaces". We discussed there that an infinite set X with cofinite topology is not metrizable. It is because of non-Hausdorffness of X with respect to cofinite topology. In this topic we will discuss an important property of metric spaces called "Hausdorffness".

Metric Spaces are Hausdorff

Theorem:
Every metric space is Hausdorff.

Proof $(X,d) \longrightarrow (X,\mathcal{T}_d)$

Let $x, y \in X \quad x \neq y$

$\because$ X is metric space. $d(x,y) \neq 0$

$d(x,y) = r$ say.

Now $U_x = B(x, r/2)$, $U_y = B(y, r/2)$

My claim is $U_x \cap U_y = \emptyset$

Metric Spaces are Hausdorff

Ld it is not true. So

$\exists \; z \in U_x \cap U_y$

Now $d(x,y) \leq d(x,z) + d(z,y)$

$< r/2 + r/2 = r = d(x,y)$

$\lightning$

So $\nexists \; z \in U_x \cap U_y$

$\Rightarrow U_x \cap U_y = \emptyset \Rightarrow (X,\mathcal{T}_d)$ is Hausdorff. $\square$

Topology

KHALIL SAHIR
MC200400708

$T_1 \not\Longrightarrow T_2$

SEARCH
B
KHALIL

$T_1 \not\Longrightarrow T_2$

T_2 property is a stronger property than T_1 property.

i.e.

Every T_2 space is T_1 but not every T_1 space is T_2.

$T_1 \not\Longrightarrow T_2$

Example: An infinite set X with cofinite topology is T_1 space but not T_2 space.
Justification: Given X is infinite. Now with respect to cofinite topology on X, a subset D of X is open in X iff

$$D = \emptyset \quad \text{or} \quad D^c = \text{finite}$$

So for each pair of distinct points x, y in X one may consider

$$U_x = \{y\}^c \quad \text{and} \quad U_y = \{x\}^c$$

and it is clear that

$$x \in U_x \not\ni y \quad \text{and} \quad y \in U_y \not\ni x$$

$T_1 \not\Longrightarrow T_2$

So $(X, \mathcal{T}_{cof})$ is T_1.
Now we will show that $(X, \mathcal{T}_{cof})$ is not T_2.
Lets start contrarily i.e. with the assumption $(X, \mathcal{T}_{cof})$ is T_2. So for any distinct pair of points x, y in X, there exist U_x, U_y (open subsets of X containing x and y resp.), s.t.

$$U_x \cap U_y = \emptyset$$

Now $(U_x)^c$ and $(U_y)^c$ are finite.Now

$$(U_x \cap U_y)^c = (U_x)^c \cup (U_y)^c$$

is finite. But $(U_x \cap U_y)^c = X$ that is infinite.
Contradiction. So $(X, \mathcal{T}_{cof})$ is not T_2.

Topology

Unique Limit Point

Unique Limit Point



Recall the definition of "Convergent Sequence". Let $(X, \mathcal{T})$ be a topological space. A sequence of points $(x_i)_{i \in I}$ in X is said to be converges to a point $x \in X$ if for every open subset U_x, containing x, in X there exists $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n \in U_x$.

Unique Limit Point

Now if we consider convergent sequences in $\mathbb{R}^n$ with usual topology then our experience for such sequences is uniqueness of the limit points.
But if we critically analyze the definition of convergent sequences, we find no such condition on the uniqueness of the limit point.

Unique Limit Point

Example:
Consider $X = \{a, b, c\}$ with

$$\mathcal{T} = \{\emptyset, \{b\}, \{a, b\}, \{b, c\}, X\}$$

The constant sequence (b) converges to a, to b, and to c.

Unique Limit Point

Topological spaces in which sequences converge to unique limit points are of more importance than the topological spaces in which limit are not unique. And the reason is, very limited results can be stated/proved in spaces with multiple limit points. So one should impose some extra condition/property on the space to avoid such multiplicity. And this condition/property brings the class of spaces under consideration closer to which one's geometric intuition applies.
The property that ensure the uniqueness of limit point is the Hausdorff property.

Topology

Theorem: Unique Limit Point

Theorem: Unique Limit Point

Theorem:
Let X be a Hausdorff space then every convergent sequence of points has a unique limit point.
i.e.
for (x_i) in X if

$$a, b \in X$$

are the limit points of (x_i) then

$$a = b.$$

Theorem: Unique Limit Point

Proof:
Consider a convergent sequence (x_i) in a Hausdorff space X. Let (x_i) converges to $a, b \in X$ and

$$a \neq b$$

Since X is Hausdorff. So for $a, b \in X$ $\exists$ a pair of open subsets, U_a and V_b, of X, such that

$$U_a \cap V_b = \emptyset$$

Now if a is the limit of (x_i) then

$\exists\ N \in \mathbb{N}$ such that $\forall\ n \geq N \Rightarrow x_n \in U_a$

Theorem: Unique Limit Point

And if b is the limit of (x_i) then

$$\exists\ M \in \mathbb{N} \text{ such that } \forall\ n \geq M \ \Rightarrow\ x_n \in V_b$$

So it means that $\forall\ n \geq \max\{M, N\}$ we have

$$x_n \in U_a \cap V_b$$

Which is the contradiction of the fact

$$U_a \cap V_b = \emptyset$$

So $a = b$, i.e. in a Hausdorff space then every convergent sequence of points has a unique limit point.

Topology

Regular Spaces

KHALIL SAHIR

Regular Spaces

Def:
A topological space $(X, \mathcal{T})$ is said to be "regular" iff for every closed subset F of X and for every $x \in F^c$ there exist open subsets U_F and U_x containing F and x respectively such that

$$U_F \cap U_x = \emptyset.$$

Regular Spaces



Regular Spaces

T_3 Space:
A topological space $(X, \mathcal{T})$ is said to be "T_3 space" iff it is either
Regular+T_0
or
Regular+T_1
or
Regular+T_2

Regular Spaces

Remarks:

1. Property of being regular is a topological property.
2. Every subspace of a regular space is a regular space.
3. T_3 property is a topological property.
4. Every subspace of a T_3 space is a T_3 space.

Topology

Examples: Regular Spaces

Examples: Regular Spaces

In this topic we will discuss some examples of regular spaces.

Example 1:
Discrete space is a regular space.

Example 2:
Indiscrete space is a regular space.

Examples: Regular Spaces



Examples: Regular Spaces

Example 3: Consider $X = \{a, b, c\}$ with

$$\mathcal{T} = \{\emptyset, \{a\}, \{b, c\}, X\}.$$

$(X, \mathcal{T})$ is a regular space.



Examples: Regular Spaces

Example 3: Consider $X = \{a, b, c\}$ with

$$\mathcal{T} = \{\emptyset, \{a\}, \{b, c\}, X\}.$$

$(X, \mathcal{T})$ is a regular space.



Examples: Regular Spaces



Example 4:
$\mathbb{R}$ with usual topology is regular. Moreover this space is also T_3 space.

Example 5:
An infinite set with cofinite topology is not regular. So it is not T_3.

Examples: Regular Spaces

Example 6:
Recall the K-Topology on real line. Topology $\mathcal{T}_K$ generated by

$$\mathcal{B}_K = \{(a, b) |\ a, b \in \mathbb{R}\} \cup \{(a, b) \backslash K | a, b \in \mathbb{R}\}$$

where

$$K = \{1, 1/2, 1/3, \cdots\} \subset \mathbb{R}$$

$(\mathbb{R}, \mathcal{T}_K)$ is not a regular space.

Topology

Normal Spaces

Normal Spaces

Def:
A topological space $(X, \mathcal{T})$ is said to be "normal" iff for every pair of disjoint closed subsets, $F_1, F_2 \subset X$ there exist open subsets U_{F_1} and U_{F_2} containing F_1 and F_2 respectively such that

$$U_{F_1} \cap U_{F_2} = \emptyset.$$

Normal Spaces



Normal Spaces

T_4 Space:

A topological space $(X, \mathcal{T})$ is said to be "T_4 space" iff it is either
Normal+T_1
or
Normal+T_2

Normal Spaces

Remarks:

1. Property of being normal is a topological property.
2. Subspace of a normal space needs not be a normal space.
3. T_4 property is a topological property.

Completely Normal Space:
A topological space $(X, \mathcal{T})$ is said to be completely normal space iff every subspace of X with subspace topology is normal.

Topology

Examples: Normal Spaces

Examples: Normal Spaces

Let us see some examples for the better understanding of the definitions of the normal-axiom and related axioms.

KHALIL SAHIR

Examples: Normal Spaces

Example 1: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, \{b, c, d\}, X\}.$$

Claim: $(X, \mathcal{T})$ is a normal space.
Explanation:
The closed collection is

$$\mathcal{C}_{\mathcal{T}} = \{\emptyset, \{a\}, \{a, b\}, \{a, c\}, \{a, b, c\}, X\}$$

Examples: Normal Spaces

Example 1: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, \{b, c, d\}, X\}.$$

Claim: $(X, \mathcal{T})$ is a normal space. But $(X, \mathcal{T})$ is not T_1
Explanation:
The closed collection is

$$\mathcal{C}_{\mathcal{T}} = \{\emptyset, \{a\}, \{a, b\}, \{a, c\}, \{a, b, c\}, X\}$$

$F_1 = \emptyset$ $F_2 = X$ $U_{F_1} = \emptyset$, $U_{F_2} = X$
$U_{F_1} \cap U_{F_2} = \emptyset$

Examples: Normal Spaces

Example 2: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, \{b, c, d\}, X\}.$$

Claim: $(X, \mathcal{T})$ is not T_4 space.

Examples: Normal Spaces

Example 3: Consider $Y = \{b, c, d\}$ with

$$\mathcal{T}_Y = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, Y\}.$$

Claim: $(Y, \mathcal{T}_Y)$ is not a normal space.
Explanation:
The closed collection is

$$\mathcal{C}_{\mathcal{T}_Y} = \{\emptyset, \{b\}, \{c\}, \{b, c\}, Y\}$$

KHALIL SAHIR

Examples: Normal Spaces

Example 3: Consider $Y = \{b, c, d\}$ with

$$\mathcal{T}_Y = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, Y\}.$$

Claim: $(Y, \mathcal{T}_Y)$ is not a normal space.
Explanation:
The closed collection is

$$\mathcal{C}_{\mathcal{T}_Y} = \{\emptyset, \{b\}, \{c\}, \{b, c\}, Y\}$$

$F_1 = \{b\}$, $F_2 = \{c\}$ $\quad U_{F_1} \cap U_{F_2} \neq \emptyset$

Examples: Normal Spaces

Example 4: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, \{b, c, d\}, X\}.$$

Claim: $(Y, \mathcal{T}_Y)$ is not a completely normal space.
Explanation:
Consider $Y = \{b, c, d\}$. So the subspace topology on Y with respect to X is

$$\mathcal{T}_Y = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, Y\}.$$

Examples: Normal Spaces

Example 4: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, \{b, c, d\}, X\}.$$

Claim: $(X, \mathcal{T}_Y)$ is not a completely normal space.
Explanation:
Consider $Y = \{b, c, d\}$. So the subspace topology on Y with respect to X is

$$\mathcal{T}_Y = \{\emptyset, \{d\}, \{b, d\}, \{c, d\}, Y\}.$$

Examples: Normal Spaces

Example 5: $\mathbb{R}$ with usual topology is a T_4 space.

$(\mathbb{R}, \mathcal{T}_u)$ is Normal Space.

F_1 $\quad F_2$

$[a, b] \cap [c, d] = \emptyset$

$U_{F_1} = (a', b')$

$U_{F_2} = (b', d')$

$a' < a$

$b < b' < d$

$d' > d$

$U_{F_1} \cap U_{F_2} = \emptyset$

Topology

Metric Spaces are Normal

KHALIL SAHIR

Metric Spaces are Normal

In this topic we will discuss another important property of metric spaces, that is called "Normal Property". In fact more than this one metric spaces are T_4 space.

Theorem:
Every metric space is normal.

Metric Spaces are Normal

Definition:
Let (X, d) be a metric space and A be a nonempty subset of X. Then a continuous function

$$d_A : X \longrightarrow \mathbb{R}$$

defined as

$$d_A(x) = \inf\{d(x, a) | a \in A\},$$

provides the distance between x and A. We denote it as

$$\text{dist}(x, A) = d_A(x)$$

Metric Spaces are Normal

Proof:
Let (X, d) be a metric space and F_1 and F_2 be two disjoint closed subsets of X. Consider a subset V_{F_1} of X consists of all those elements x of X whose distance from F_1 is less than the distance from F_2, i.e.

$$V_{F_1} = \{x \in X | d_{F_1}(x) < d_{F_2}(x)\}$$

It is easy to see that $F_1 \subset V_{F_1}$.
Since $x \in F_1$ implies that $d_{F_1}(x) = 0$ and $d_{F_2}(x) > 0$, F_2 is closed.
So $d_{F_1}(x) < d_{F_2}(x)$ and we have $x \in V_{F_1}$.

Metric Spaces are Normal

Now define a map

$$\delta = d_{F_2} - d_{F_1} : X \longrightarrow \mathbb{R}$$

Under this map

$$V_{F_1} = \delta^{-1}((0, \infty))$$

So V_{F_1} is an open subset of X containing F_1, since δ is continuous. On the same lines we can show that

$$V_{F_2} = \{x \in X | d_{F_2}(x) < d_{F_1}(x)\}$$

is an open subset of X containing F_2. Moreover

$V_{F_1} \cap V_{F_2} = \emptyset \quad \Longrightarrow \quad X$ is normal.

Topology

T_i Properties: An Overview

T_i Properties: An Overview

An overview of T_i properties is given in this topic through a diagram. This diagram illustrates the relationship between the T_i spaces we studied in previous topics.

T_i Properties: An Overview



Topology

Urysohn's Lemma

Urysohn's Lemma

In this topic, one of the most classical and important results of topology will be discussed. This result is named as "Urysohn's Lemma".
Let us state this lemma.

Urysohn's Lemma

Urysohn's Lemma: Let X be a normal space; A and B be two disjoint closed subsets of X. Let $[a, b]$ be a closed interval in $\mathbb{R}$. Then there exists a continuous map

$$f : X \longrightarrow [a, b]$$

such that $f(x) = a \ \forall \ x \in A$ and $f(x) = b \ \forall \ x \in B$.

Urysohn's Lemma: (Strong Form)
X is a normal space if and only if for every two disjoint closed subsets A and B of X. There exists a continuous map

$$f : X \longrightarrow [a, b]$$

such that $f(x) = a \ \forall \ x \in A$ and $f(x) = b \ \forall \ x \in B$.

Urysohn's Lemma

One important application of Urysohn's Lemma is the partial solution of metrization problem. Here we have this solution in the form of "Urysohn's Metrization Theorem".

Urysohn's Metrization Theorem:
Every normal-T_1 space with countable basis is metrizable.

Another version of Urysohn's Metrization Theorem is

Urysohn's Metrization Theorem:
Every regular-T_1 space with countable basis is metrizable.

Topology

Compact Spaces

Compact Spaces

Recall the definitions of cover, subcover, and open cover.

Cover:
Let $(X, \mathcal{T})$ be a topological space. Let $\mathcal{C} = \{C_i\}$ be a class of subsets of X, such that

$$X = \bigcup_{i \in I} C_i,$$

then $\mathcal{C}$ is called a cover of X.

Compact Spaces

Subcover:
A subclass $\mathcal{S}$ of a cover $\mathcal{C}$ of X which is also a cover is called a subcover.

Open cover:
A cover $\mathcal{C}$ of X is said to be an open cover if and only if $\mathcal{C} \subset \mathcal{T}$.

KHALIL SAHIR

Compact Spaces

Compact Space:
A topological space $(X, \mathcal{T})$ is said to be compact if every open cover $\mathcal{C}$ of X contains a finite subcover $\mathcal{S}$ of X.

Topology

Examples: Compact Spaces

Examples: Compact Spaces

Example 1:
A set X with indiscrete topology $\mathcal{T}$ is compact.

Reason:
Since there is only one open cover of X with respect to indiscrete topology and that is

$$\{X\}$$

which is it self finite.

Dr.

Examples: Compact Spaces

Example 2:
A set X with topology $\mathcal{T}$ containing finite number subsets of X is compact.

Reason:
Since every open cover $\mathcal{C}$ of X is a subclass of $\mathcal{T}$ and $\mathcal{T}$ itself finite so $\mathcal{C}$ is finite too and any subcover $\mathcal{S}$ of $\mathcal{C}$ is also finite.

Examples: Compact Spaces

Example 3:
A finite set X with any topology $\mathcal{T}$ is compact.

Reason:
Since every topology of a finite set X is finite and from "Example 2" X is compact.

Examples: Compact Spaces

Example 4:
An infinite set X with discrete topology $\mathcal{T}$ is not compact.
Reason:

$\mathcal{T}_{dis} = \mathcal{P}(X)$

$\mathcal{C} = \{\{x\} \mid x \in X\} \cup \{\{x_1, x_2\}\}$ (fin.)

$X = \bigcup_{i \in I} C_i$ $\quad C_i = \{x_i\}$

Topology

Open Intervals in $\mathbb{R}$

Open Intervals in $\mathbb{R}$

In this topic we will discuss the compactness of open intervals in $\mathbb{R}$ with respect to the usual topology.

Fact:
An open interval in $\mathbb{R}$ with respect to usual topology is not compact.

An immediate consequence of this fact is

$(\mathbb{R}, \mathcal{T}_u)$ is not compact.

Open Intervals in $\mathbb{R}$

Proof:
Consider $(\mathbb{R}, \mathcal{T}_u)$ and (a, b) be an open interval in $\mathbb{R}$.
Consider the following covering of (a, b).

$$\mathcal{C} = \left\{\left(a + \frac{b-a}{n+2}, a + \frac{b-a}{n}\right)\right\}$$

Open Intervals in $\mathbb{R}$

Proof:
Consider $(\mathbb{R}, \mathcal{T}_u)$ and (a, b) be an open interval in $\mathbb{R}$.
Consider the following covering of (a, b).

$$\mathcal{C} = \left\{\left(a + \frac{b-a}{n+2}, a + \frac{b-a}{n}\right)\right\} \quad n \in \mathbb{Z}^+$$

$n=1 \quad (a + \frac{b-a}{3}, b)$

$n=2 \quad (a + \frac{b-a}{4}, a + \frac{b-a}{2})$



$\nexists$ subcover of $\mathcal{C}$ that is finite

So (a, b) is not compact.

Topology

Cofinite Topology and Compactness

Cofinite Topology and Compactness

In this topic we will study the compactness of cofinite space, a set X with cofinite topology $\mathcal{T}_{cof}$.

Fact:
A set X with cofinite topology $\mathcal{T}_{cof}$ is compact.

Cofinite Topology and Compactness

Proof: If X is finite. Then $\mathcal{T}_{cof} = \mathcal{P}(X)$

$\Rightarrow (X, \mathcal{T}_{cof})$ is compact.

But if X is not finite. Then

Consider an open cover $\mathcal{U}$ of X.

Now let $\emptyset \neq O \in \mathcal{U}$ $\because$ $\mathcal{U}$ is an open cover

$\Rightarrow O$ is open. $\Rightarrow O^c$ is finite

Now for every $x \in O^c$ choose an element of $\mathcal{U}$ O_x containing x.

$\{O_x \mid x \in O^c\}$ is finite

Cofinite Topology and Compactness

$\because$ D^c is finite.

Consider this subcover of $\mathcal{D}$

$\mathcal{C} = \{D\} \cup \{D_x \mid x \in D^c\}$

finite, finite

$\Rightarrow$ $\mathcal{C}$ is a finite subcover of $\mathcal{D}$

$\Rightarrow$ $(X, \mathcal{T}_{cof})$ is compact



Topology

Compactness and Open Collection

Compactness and Open Collection

The aim of this topic is to show that compactness of some space depends on the collection of open subsets of the space.

Consider $\mathbb{R}$, now

- $(\mathbb{R}, \mathcal{T}_u)$ is not compact.
- $(\mathbb{R}, \mathcal{T}_{indis})$ is compact.
- $(\mathbb{R}, \mathcal{T}_{dis})$ is not compact.
- $(\mathbb{R}, \mathcal{T}_{cof})$ is compact.

Topology

Theorem: Closed Subspace of Compact Space

Theorem: Closed Subspace...

Theorem:
Every closed subspace of a compact space is compact.

Proof:
Let $(X, \mathcal{T})$ be a compact space and A be a closed subset of X.
So $(A, \mathcal{T}_A)$ be closed subspace of X, where $\mathcal{T}_A$ is subspace topology on A.
Consider an open cover $\mathcal{D}$ of A, i.e.

Theorem: Closed Subspace...

$$A = \bigcup_{i \in I} D_i$$

where $D_i \in \mathcal{D} \subset \mathcal{T}_A$.
Now for each D_i there exists $U_i \in \mathcal{T}$ such that $D_i = A \cap U_i$.
So

$$A = \bigcup_{i \in I} D_i = \bigcup_{i \in I} (A \cap U_i)$$

But A is closed in X so A^c is an open subset of X. And we have an open covering of X.

$$X = A^c \cup (\bigcup_{i \in I} U_i)$$

Theorem: Closed Subspace...

$$A = \bigcup_{i \in I} D_i$$

where $D_i \in \mathcal{D} \subset \mathcal{T}_A$.
Now for each D_i there exists $U_i \in \mathcal{T}$ such that $D_i = A \cap U_i$.
So

$$A = \bigcup_{i \in I} D_i = \bigcup_{i \in I} (A \cap U_i) \Rightarrow A \subset \bigcup_{i \in I} U_i$$

But A is closed in X so A^c is an open subset of X. And we have an open covering of X.

$$X = A^c \cup (\bigcup_{i \in I} U_i)$$

Theorem: Closed Subspace...

But X is compact. So every open cover of X has a finite subcover, i.e. there exists $J \underset{\text{finite}}{\subset} I$ such that

$$X = A^c \cup (\bigcup_{j \in J} U_j)$$

$\Longrightarrow$

$$A \subset \bigcup_{j \in J} U_j$$

$A \cap U_j = D_j$

$\Longleftrightarrow$

$$A = A \cap (\bigcup_{j \in J} U_j) = \bigcup_{j \in J} D_j$$

Proved.

Theorem: Closed Subspace...

But X is compact. So every open cover of X has a finite subcover, i.e. there exists $J \underset{\text{finite}}{\subset} I$ such that

$$X = A^c \cup (\bigcup_{j\in J} U_j)$$

$$\Longrightarrow$$

$$A \subset \bigcup_{j\in J} U_j$$

$$\Longleftrightarrow$$

$$A = A \cap (\bigcup_{j\in J} U_j) = \bigcup_{j\in J} D_j \quad (A \cap U_j = D_j)$$

A is compact.

Proved.

Topology

Compact Subspace of Hausdorff Space

Compact Subspace of Hausdorff...

Theorem:
Let $(A, \mathcal{T}_A)$ be compact subspace of a Hausdorff space $(X, \mathcal{T})$.
Then A is closed in X.
i.e.
Every compact subspace of a Hausdorff space is closed.

Compact Subspace of Hausdorff...

Proof: Let $(X, \mathcal{T})$ is Hausdorff.
Consider $(A, \mathcal{T}_A)$ a subspace of $(X, \mathcal{T})$
Compact.
Now we have to show that A is closed in X

$\Longleftrightarrow$ $A = \bar{A}$

$\Longleftrightarrow$ $\nexists\ x \in A^c = X - A$ s.t $x \in \bar{A}$

$\longrightarrow \Longleftrightarrow$ for every $x \in A^c = X - A$ $\exists\ U_x$ (open subset of X containing x) s.t $U_x \cap A = \emptyset$

Compact Subspace of Hausdorff...

Let $x \in A^c$
Given $(X, \mathcal{T})$ is Hausdorff
So for every $y \in A \subset X$
$\exists$ U_{xy}, U_y open subsets of X containing x and y resp.
s.t $U_{xy} \cap U_y = \emptyset$
Now for every $y \in A$ such disjoint pair of subsets exists. one containing x and other containing y

Compact Subspace of Hausdorff...

So we have an open cover of A

$$A \subset \bigcup_{y\in A} U_y$$

$$\Rightarrow A = \bigcup_{y\in A} (A \cap U_y)$$

But given A is compact. Then this cover has a finite subcover.
$\exists\ y_1, y_2, \ldots, y_n \in A$ s.t

$$A = \bigcup_{i=1}^{n} (A \cap U_{y_i})$$

Compact Subspace of Hausdorff...

$$\Rightarrow A \subset \bigcup_{i=1}^{n} U_{y_i} = V \text{ (Say)}$$

Now $U_x = \bigcap_{i=1}^{n} U_{xy_i}$ an open subset of X containing x.

More than this one $U_x \cap V = \emptyset$

$$\Rightarrow U_x \cap A = \emptyset$$

$\Rightarrow$ A is closed.

Topology

KHALIL SAHIR

Compact Space under Continuous Map

Compact Space under Cont. Map

Theorem:
Let $(X, \mathcal{T})$ be a compact space and

$$f : (X, \mathcal{T}) \to (Y, \mathcal{T}')$$

be a continuous map. Then $f(X)$ is compact.
i.e.
Image of compact space under a continuous map is compact.

Compact Space under Cont. Map

Proof: Let $f : (X, \mathcal{T}) \xrightarrow[\text{cont.}]{\text{compact}} (Y, \mathcal{T}')$

Now $f(X) \subset Y$

Now consider an open cover of $f(X)$

$f(X) \subset \bigcup_{i \in I} U_i \qquad U_i \in \mathcal{T}'$

$\because$ f is cont. So $\forall\ U_i \in \mathcal{T}'$

$f^{-1}(U_i) \in \mathcal{T}$ Now $D_i = f^{-1}(U_i)$

Compact Space under Cont. Map

$X = \bigcup_{i \in I} f^{-1}(U_i) = \bigcup_{i \in I} D_i$

So we have an open cover of X

But given, X is compact.

So $\exists$ a finite subcover of this cover

i.e. $\exists\ J \subset_{\text{finite}} I$ s.t

$X = \bigcup_{j \in J} D_j = \bigcup_{j \in J} f^{-1}(U_j)$

$\Rightarrow$ $\boxed{f(X) \subset \bigcup_{j \in J} U_j}$ $\Rightarrow$ $f(X)$ is compact.

Topology

Compactness and Homeomorphism

Compactness and Homeomorphism

Corollary I:
Let $(X, \mathcal{T})$ be a compact space and $(Y, \mathcal{T}')$ be a Hausdorff space. Then a continuous map

$$f : (X, \mathcal{T}) \to (Y, \mathcal{T}')$$

is a closed map.

Compactness and Homeomorphism

Proof:
Let A be a closed subset of X. It is given, X is compact and we know that a closed subspace of a compact space is compact.

So A is compact. Now f is continuous and we know that image of a compact space under a continuous map is compact so $f(A)$ is a compact subspace of Y.

Now Y is Hausdorff and is has been proved that every compact subspace of a Hausdorff is closed, so $f(A)$ is closed.

So f is a closed map.

Compactness and Homeomorphism

Corollary II:
Let $(X, \mathcal{T})$ be a compact space and $(Y, \mathcal{T}')$ be a Hausdorff space. Then a bijective continuous map

$$f : (X, \mathcal{T}) \to (Y, \mathcal{T}')$$

is a homeomorphism.

Topology

Application: Compactness and Homeomorphism

Application: Compactness and...

Exercise I:
Consider $(\mathbb{R}, \mathcal{T}_u)$. Show that there exists no homeomorphism between an open interval of $\mathbb{R}$ and a closed interval of $\mathbb{R}$.

Exercise II:
Prove or disprove.

$$[0;1) \approx S^1$$

Application: Compactness and...

Ex(1) We know that the open intervals in $(\mathbb{R}, \mathcal{T}_u)$ are not compact.
Moreover the only compact subsets of $\mathbb{R}$ are closed and bounded subsets.
Compact $[a,b]$ (c,d) Not compact.
if $\exists$ $f: [a,b] \xrightarrow[\text{cont}]{\text{bijection}} (c,d)$
Then $f([a,b]) = (c,d)$
compact Not compact
$\not\exists$ f:

Application: Compactness and...

Ex(2) $[0,1) \not\approx S^1$
$\because$ S^1 is compact and $[0,1)$ is not compact.

$f: [0,1) \xrightarrow[\text{cont}]{\text{bij}} S^1$ ✓
$x \longmapsto (\cos 2\pi x, \sin 2\pi x)$
Can we say $\Rightarrow$ f is homeomorphism No
$\because$ $[0,1)$ is not compact.

Topology

Finite Union of Compact Spaces

Topology

Finite Union of Compact Spaces

Finite Union of Compact Spaces

Theorem:
Let $(X, \mathcal{T})$ be a topological space. And

$$\{(X_i, \mathcal{T}_i) | i \in I\}$$

be a finite family of compact subspaces of X.
Then

$$\bigcup_{i \in I} X_i$$

is compact.

Finite Union of Compact Spaces

Proof: Let $X_1, X_2 \subset_{\text{compact}} X$
Now $X_1 \cup X_2$ we want to show that this union is compact.
Let us consider an open cover of $X_1 \cup X_2$. $X_1 \cup X_2 \subset \bigcup_{i \in I} U_i$ $U_i \in \mathcal{T}$
$\Rightarrow$ $X_1 \subset \bigcup_{i \in I} U_i$ and $X_2 \subset \bigcup_{i \in I} U_i$
But X_1, X_2 both compact subspaces of X.

Finite Union of Compact Spaces

$\Rightarrow$ $X_1 \subset \bigcup_{j \in J} U_j$ and $X_2 \subset \bigcup_{j' \in J'} U_{j'}$
where $J, J' \subset I$ finite
$\Rightarrow$ $X_1 \cup X_2 \subset \left(\bigcup_{j \in J} U_j\right) \cup \left(\bigcup_{j' \in J'} U_{j'}\right)$
$\Rightarrow$ $X_1 \cup X_2$ is compact.

Topology

Compact Hausdorff Space is Normal

Compact Hausdorff Space is Normal

Theorem:
Let $(X, \mathcal{T})$ be a compact Hausdorff space. Then $(X, \mathcal{T})$ is normal.
i.e.
For every pair of disjoint closed subsets F_1, F_2 of a compact Hausdorff space X there exists a pair of disjoint open subsets V_{F_1}, V_{F_2} of X containing F_1, F_2 respectively.

Compact Hausdorff Space is Normal

Proof:
Let

$$F_1, F_2 \underset{\text{closed}}{\subset} X$$

such that

$$F_1 \cap F_2 = \emptyset$$

Consider an open covering of F_1

$$F_1 \subset \bigcup_{x \in F_1} V_x$$

Now F_1 is a closed subset of compact space X, so F_1 is compact.

Compact Hausdorff Space is Normal

So there exist $x_1, x_2, \ldots, x_n \in F_1$ such that

$$F_1 \subset \bigcup_{i=1}^{n} V_{x_i} = V_{F_1}(\text{say})$$

Given, X is Hausdorff, so for an element y of F_2 there exists $U_{y_{x_i}}$, an open set containing y such that

$$U_{y_{x_i}} \cap V_{x_i} = \emptyset \quad .$$

So $\bigcap_{i=1}^{n} U_{y_{x_i}}$ is an open set containing y and disjoint from V.

Compact Hausdorff Space is Normal

Continue with this construction one can find an open covering of F_2 by taking the union of such open subsets over $y \in F_2$, i.e.

$$F_2 \subset \bigcup_{y \in F_2} \left(\bigcap_{i=1}^{n} U_{y_{x_i}} \right)$$

Again F_2 is compact so this cover can be reduced to a finite subcover, i.e. there exist

$$y_1, y_2, \ldots, y_m \in F_2$$

Compact Hausdorff Space is Normal

such that

$$F_2 \subset \bigcup_{j=1}^{m} \left(\bigcap_{i=1}^{n} U_{y_{j_{x_i}}} \right) = U_{F_2}(\text{say})$$

And we have

$$V_{F_1} \bigcap U_{F_2} = \emptyset$$

$\Longrightarrow$
X is normal.

Topology

Separated Sets

Separated Sets

Def I:
Let A and B be two subsets of a topological space $(X, \mathcal{T})$. Then A and B are said to be separated sets if and only if

$$A \cap B = \emptyset$$
and
$$A' \cap B = \emptyset$$
and
$$A \cap B' = \emptyset$$

Separated Sets

Def II:
Let A and B be two subsets of a topological space $(X, \mathcal{T})$. Then A and B are said to be separated sets if and only if

$$\overline{A} \cap B = \emptyset$$
and
$$A \cap \overline{B} = \emptyset$$

Separated Sets

Def III:
Let A and B be two subsets of a topological space $(X, \mathcal{T})$. Then A and B are said to be separated sets if and only if there exist open subsets U_A and U_B of X containing A and B respectively such that

$$A \cap U_B = \emptyset$$
and
$$B \cap U_A = \emptyset$$

Topology

Examples: Separated Sets

Examples: Separated Sets

Example 1:
Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (0, 1)$ and $B = (4, 9]$ are separated sets.
Reason: (By Definition I)
Since

$$(0, 1) \cap (4, 9] = \emptyset$$

and

$$(0, 1)' \cap (4, 9] = \emptyset$$

and

$$(0, 1) \cap (4, 9]' = \emptyset$$

Examples: Separated Sets

Example 1:
Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (0, 1)$ and $B = (4, 9]$ are separated sets.
Reason: (By Definition I)
Since

$$(0, 1) \cap (4, 9] = \emptyset$$

and

$$(0, 1)' \cap (4, 9] = \emptyset$$

and

$$(0, 1) \cap (4, 9]' = \emptyset$$

Examples: Separated Sets

Let discuss "Example 1" with "Definition III" of separated sets.
There exist $U_A = (-0.5, 2)$ and $U_B = (3, 10)$ two open subsets of $\mathbb{R}$ containing $A = (0, 1)$ and $B = (4, 9]$ respectively such that

$$A \cap U_B = \emptyset$$
and
$$B \cap U_A = \emptyset$$

So A and B are separated sets of $(\mathbb{R}, \mathcal{T}_u)$.

Examples: Separated Sets

Example 2:
Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (-1, 3]$ and $B = [3, 5)$ are not separated sets.
Reason:
By "Definition I" both sets should be disjoint but this is not the case here, i.e.

$$(-1, 3] \cap [3, 5) = \{3\} \neq \emptyset$$

KHALIL SAHIR

Examples: Separated Sets

Example 3:
Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (-1, 3)$ and $B = [3, 5)$ are not separated sets.
Reason: (By Definition I)
Both sets are disjoint, i.e.

$$(-1, 3) \cap [3, 5) = \emptyset$$

But at the same time both sets should not contain each others limit points. But here

$$(-1, 3)' \cap [3, 5) = \{3\}$$

Examples: Separated Sets

Example 3:
Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (-1, 3)$ and $B = [3, 5)$ are not separated sets.
Reason: (By Definition I)
Both sets are disjoint, i.e.

$$(-1, 3) \cap [3, 5) = \emptyset$$

But at the same time both sets should not contain each others limit points. But here

$$(-1, 3)' \cap [3, 5) = \{3\} \neq \emptyset$$

Topology

Connected Spaces

Connected Spaces

Connected Set:
A subset A of a topological space $(X, \mathcal{T})$ is said to be connected iff there exists no pair of nonempty open subsets U and V of X such that

$$A \cap U \quad \text{and} \quad A \cap V$$

are nonempty disjoint sets and

$$A = (A \cap U) \cup (A \cap V)$$

Connected Spaces

Disconnected Set:
A subset A of a topological space $(X, \mathcal{T})$ is said to be disconnected if it is not connected, i.e.
if there exists a pair of nonempty open subsets U and V of X such that $A \cap U$ and $A \cap V$ are nonempty disjoint sets and

$$A = (A \cap U) \cup (A \cap V).$$

Connected Spaces

Connected Space:
A topological space $(X, \mathcal{T})$ is said to be connected if it can not be written as union of two nonempty open disjoint subsets U and V of X.

Connected Spaces

Disconnected Space:
A topological space $(X, \mathcal{T})$ is said to be disconnected if it can be written as union of two nonempty open disjoint subsets U and V of X.

Connected Spaces

Connected Space: (Def. II)
A topological space $(X, \mathcal{T})$ is said to be connected if

$$X = D_1 \cup D_2$$

such that D_1, D_2 both open and

$$D_1 \cap D_2 = \emptyset$$

implies that either

$$D_1 = \emptyset \quad \text{or} \quad D_2 = \emptyset.$$

Topology

Examples: Connected Spaces

Examples: Connected Spaces

Example 1: Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (0,1) \cup (3,5]$ is disconnected subset of $\mathbb{R}$.
Reason:
Since there exists a pair of nonempty open subsets

$$U = (0,1) \text{ and } V = (3,6)$$

of $\mathbb{R}$ such that
$A \cap U$ and $A \cap V$ are nonempty disjoint sets and

$$A = (A \cap U) \cup (A \cap V).$$

Examples: Connected Spaces

Example 1: Consider $(\mathbb{R}, \mathcal{T}_u)$. Then $A = (0,1) \cup (3,5]$ is disconnected subset of $\mathbb{R}$.
Reason:
Since there exists a pair of nonempty open subsets

$$U = (0,1) \text{ and } V = (3,6)$$



of $\mathbb{R}$ such that
$A \cap U$ and $A \cap V$ are nonempty disjoint sets and

$$A = (A \cap U) \cup (A \cap V).$$

$(0,1) \cup (3,5]$

Examples: Connected Spaces

Example 2: Consider $X = \{a, b, c, d, e\}$ with
$\mathcal{T} = \{\emptyset, \{a\}, \{a, b\}, \{a, b, c\}, \{a, d, e\}, \{a, b, d, e\}, X\}$.
Then $A = \{c, e\}$ is disconnected subset of X.
Reason:
Since there exists a pair of nonempty open subsets
$U = \{a, b, d, e\}$ and $V = \{a, b, c\}$ of X such that

$$A \cap U = \{e\} \quad \text{and} \quad A \cap V = \{c\}$$

are nonempty disjoint sets and

$$A = (A \cap U) \cup (A \cap V) = \{e\} \cup \{c\}.$$

Examples: Connected Spaces

Example 2: Consider $X = \{a, b, c, d, e\}$ with
$\mathcal{T} = \{\emptyset, \{a\}, \{a, b\}, \{a, b, c\}, \{a, d, e\}, \{a, b, d, e\}, X\}$.
Then $A = \{c, e\}$ is disconnected subset of X.
Reason:
Since there exists a pair of nonempty open subsets
$U = \{a, b, d, e\}$ and $V = \{a, b, c\}$ of X such that

$$A \cap U = \{e\} \quad \text{and} \quad A \cap V = \{c\}$$

$\{e\} \cap \{c\} = \emptyset$

are nonempty disjoint sets and

$$A = (A \cap U) \cup (A \cap V) = \{e\} \cup \{c\} = \{c, e\}$$

Examples: Connected Spaces

Example 3: Consider $X = \{a, b, c, d, e\}$ with
$\mathcal{T} = \{\emptyset, \{a, b\}, \{a, b, c\}, X\}$. Then X is a connected space.
Reason:
Since there exists no pair of nonempty open disjoint subsets

$$U \text{ and } V$$

of X such that

$$X = U \cup V.$$

Examples: Connected Spaces

Example 4: Consider $X = \{a, b, c, d, e\}$ with
$\mathcal{T} = \{\emptyset, \{a, b\}, \{d, e\}, \{a, b, c\}, \{a, b, d, e\}, X\}$. Then X is a disconnected space.
Reason:
Since there exist nonempty open disjoint subsets

$$U = \{a, b, c\} \text{ and } V = \{d, e\}$$

of X such that

$$X = U \cup V.$$

Topology

Connected Spaces and Open Closed Sets

Connected Spaces and Open...

Recall
Connected Space:
A topological space $(X, \mathcal{T})$ is said to be connected iff

$$X = D_1 \cup D_2$$

such that D_1, D_2 both open and disjoint implies that either

$$D_1 = \emptyset \text{ or } D_2 = \emptyset.$$

Connected Spaces and Open...

Disconnected Space:
A topological space $(X, \mathcal{T})$ is said to be disconnected iff

$$X = D_1 \cup D_2$$

such that D_1, D_2 both open and disjoint implies that neither D_1 is empty nor D_2 is empty.

KHALIL SAHIR

Connected Spaces and Open...

$X = D_1 \cup D_2$ $\quad D_1 \cap D_2 = \emptyset$

disconnected $\quad$ Both are open

$D_1 \neq \emptyset$ $\quad D_2 \neq \emptyset$

D_1 is open

$\Rightarrow D_1^c$ is closed.

$\quad\quad\quad

Connected Set and its Closure

Say $A \cap N = \emptyset$

$\Rightarrow A \subset A \cap M \Rightarrow A \subset M$

$\Rightarrow \bar{A} \subset \bar{M}$

KHALIL SAHIR

Now $A \subset B \subset \bar{A} \subset \bar{M}$

But $\bar{M} \cap N = \emptyset$

So $B \cap N = \emptyset \Rightarrow (N = \emptyset)$

$\Rightarrow$ B is connected. And this implies that $\bar{A}$ is connected.

Topology

Image of Connected Space

KHALIL SAHIR
MC200400708
03054232382

Image of Connected Space

Theorem:
Let f be a continuous function from (X, T) to (Y, T') and A be a connected subset of (X, T). Then $f(A)$ is connected.
i.e.

Image of a connected space under a continuous map is connected.

Image of Connected Space

Proof $f: (X, T) \xrightarrow[\text{Cont.}]{} (Y, T')$

$A \subset X$ Connected

Now we have to show that $f(A)$ is connected.

Let $f(A)$ is disconnected

$\Rightarrow \exists$ a separation of $f(A)$

So $f(A) = B \cup C$ B, C both Nonempty + open: Disjoint

Image of Connected Space

Now $A \subset f^{-1}(B \cup C) = f^{-1}(B) \cup f^{-1}(C)$

$\Rightarrow A = A \cap (f^{-1}(B) \cup f^{-1}(C))$

$= (A \cap f^{-1}(B)) \cup (A \cap f^{-1}(C))$

open open open open

$\because B \neq \emptyset \quad C \neq \emptyset \quad f(A) = B \cup C$

$\Rightarrow A \cap f^{-1}(B) \neq \emptyset$ and $A \cap f^{-1}(C) \neq \emptyset$

$\Rightarrow (A \cap f^{-1}(B)) \cap (A \cap f^{-1}(C)) = \emptyset \quad B \cap C = \emptyset$

Image of Connected Space

So we have a separation of A.

But A is connected.

$\exists$ no separation

$\Rightarrow f(A)$ is connected $\because \exists$ no separation of $f(A)$

Topology

Application: Image of Connected Set

Application: Image of...

Exercise I:
Consider $(\mathbb{R}, \mathcal{T}_u)$ and show that there exists no homeomorphism between $(0, 2)$ and $(0, 1] \cup [3, 4)$.

Exercise II:
Prove or disprove

$$(0, 1) \approx (0, 1]$$

Application: Image of...

Ex ① Let $f: (0,2) \longrightarrow (0,1] \cup [3,4)$

bijective + bicont.

$\because$ f is cont and w.r.t $\mathcal{T}_u$ on $\mathbb{R}$

$A=(0,2)$ is connected. $\Rightarrow$ $f(A)$ is connected.

But $(0,1] \cup [3,4)$ is not connected.

$f(A) \longrightarrow (0,1]$ or $[3,4)$

$\Rightarrow \not\exists\ f \Rightarrow (0,2) \not\simeq (0,1] \cup [3,4)$

Application: Image of...

Ex ② $(0,1) \overset{??}{\simeq} (0,1]$

Ans. $\exists$ no such f bijective + bicont.

Let $\exists\ f: (0,1] \longrightarrow (0,1)$ biject + bicont

$1 \in (0,1] \Rightarrow f(1) \in (0,1)$

$f': (0,1] \setminus \{1\} \longrightarrow (0,1) \setminus \{f(1)\} = (0, f(1)) \cup (f(1), 0)$

connected; biject + bicont (Not possible); disconnected

$\Rightarrow (0,1) \not\simeq (0,1]$

Topology

Connected Subsets of Real Line

Connected Subsets of Real Line

Theorem:
A subset A of $(\mathbb{R}, \mathcal{T}_u)$ is connected if and only if A is an interval.

Proof:
Let A be a nonempty connected subset of $\mathbb{R}$.
If A contains only one point then we have nothing to prove.
Let A contains two points, say

$$a, b \in A$$

Connected Subsets of Real Line

Consider $a < b$.
If every real number c, between a and b, belongs to A.
Then A is an interval. Proved.
If this is not the case, i.e. there exists some $c \in (a, b)$ such that

$$c \notin A$$

Then we can write A as

$$A = (A \cap (-\infty, c)) \cup (A \cap (c, \infty))$$

This provides us a separation of A, i.e. a nonempty, open, disjoint pair of sets whose union is A.

Connected Subsets of Real Line

This implies that A is disconnected and it is contradiction of the fact that A is connected.
So for all $c \in (a, b)$, $\Longrightarrow$ $c \in A \Longrightarrow$ A is an interval.
Converse:
Let A be an interval. i.e.

$$A = [a, b]$$

Now let A is disconnected. This means there exists a separation of A, i.e. there exist B, C nonempty, open, disjoint subsets such that

$$A = B \cup C$$

Connected Subsets of Real Line

Now let

$$d = \sup\{c \in A | [a, c] \subset B\} \in A$$

Now one thing is very much clear here that $d \neq b$.
Let $d \in B$, now since B is open so there exists an open set U_d containing d such that

$$U_d \subset B.$$

So we can find a real number $d_1 \in U_d$ greater than d such that

$$[a, d_1] \subset B$$

Connected Subsets of Real Line

Now let

$$d = \sup\{c \in A | [a, c] \subset B\} \in A$$

(M)

Now one thing is very much clear here that $d \neq b$.
Let $d \in B$, now since B is open so there exists an open set U_d containing d such that



$U_d \subset B$.

So we can find a real number $d_1 \in U_d$ greater than d such that

$[a, d_1] \subset B \Rightarrow d_1 \in M$

But d is sup M and $d_1 > d$

Connected Subsets of Real Line

But this is a contradiction of the fact the d is the supremum. So $d \notin B$ and it means

$$d \in C$$

Again C is open so there exists an open set V_d containing d such that $V_d \subset C$.
This implies that there exists a real number $d_2 \in V_d$, less than d. So

$$[a, d_2] \subset B \quad \Rightarrow \quad d_2 \in B \cdot$$

And we have again a contradiction of the fact $B \cap C = \emptyset$.
So A is connected.

Topology

Connectedness and Fixed Point Theorem

Connectedness and Fixed Point...

Fixed Point:
A point $x \in X$ is said to be a fixed point of a map $f : X \to X$ iff $f(x) = x$.

Example:

1. $f : \mathbb{R} \to \mathbb{R}$, defined as $f(x) = x^2$. Here 0 is a fixed point of f. 1 is also a fixed point of f.
2. $g : \mathbb{R} \to \mathbb{R}$, defined as

$$g(x) = x + 1$$

g has no fixed point.

Connectedness and Fixed Point...

Theorem:
Consider $\mathbb{R}$ with usual topology and

$$I = [0, 1]$$

be closed interval in $\mathbb{R}$. Then every continuous map

$$f : I \to I$$

has at least one fixed point.

Connectedness and Fixed Point...

Geometrical Interpretation:

$f: I \xrightarrow[\text{Cont}]{} I \qquad \exists\ x \in I$ s.t $f(x) = x$

Connectedness and Fixed Point...

Proof: $f: I \xrightarrow[\text{Cont.}]{} I$

If $f(0) = 0$ or $f(1) = 1$ then, proved.

Now suppose that $f(0) > 0$ and $f(1) < 1$

Now the graph of f is connected.

i.e $G = \{(x, f(x)) \in \mathbb{R}^2 \mid 0 \le x \le 1,\ 0 \le f(x) \le 1\} \subset \mathbb{R}^2$ Connected.

Now consider $M, N \subset G$ defined as:

Connectedness and Fixed Point...

$\to M = \{(x, f(x)) \mid x < f(x)\} \subset G$

$M \neq \emptyset \quad \because (0, f(0)) \in M$ since $f(0) > 0$

$\to N = \{(x, f(x)) \mid x > f(x)\} \subset G$

$N \neq \emptyset \quad (1, f(1)) \in N$ since $f(1) < 1$

Both are open subsets.

$\to M \cap N = \emptyset$

Connectedness and Fixed Point...

[Suppose $\exists$ no fixed point of f. i.e $\nexists\ x \in I$ s.t $x = f(x)$]

So $G = M \cup N$ Separation of G.

$\Rightarrow G$ is disconnected. ✗

But G is connected.

So $\exists\ x \in I$ s.t $f(x) = x$.

Topology

Union of Connected Subsets

KHALIL SAHIR
MC200400708
03054232382

Union of Connected Subsets

Theorem:
Let A and B be two connected subsets of a topological spaces (X, T) such that

$$x \in A \cap B$$

than

$$C = A \cup B$$

is a connected subset of X.

Union of Connected Subsets

Proof: $A, B \subset X$ connected s.t $x \in A \cap B$

Now $C = A \cup B \leftarrow$ is connected !!

Let C is not connected. i.e

$\exists$ a separation of C say $C = M \cup N$ s.t M, N both
- open
- Nonempty.
- disjoint.

Now $x \in C$ either $x \in M$ or $x \in N$

Union of Connected Subsets

Say $x \in M$ Then

$A \subset M$ also $B \subset M$

$\because x \in A$ $\because x \in B$

$\Rightarrow A \cup B \subset M$

$\Rightarrow C \subset M \Rightarrow N = \emptyset$

So $\nexists$ no such separation of C.

So C is connected

Union of Connected Subsets

Theorem:
The union of a collection of connected subspaces of (X, T), that having a point common, is connected.

Topology

Connected Component

Connected Component



Def:
Consider a topological space (X, T). Let $a \in X$ and C be a connected subset of X containing a.
Then

$$C_a = \bigcup_{a \in C} C$$

is called the connected component of X containing a.

Connected Component

(4) A space $(X, \mathfrak{I})$ is connected iff it has only one connected component.

Remarks:

1. The connected component C_a is the largest connected subset of X containing a.
2. If $b \in C_a$ then $C_b = C_a$.
3. X can be written as disjoint union of its connected components. i.e.

$$X = \coprod_{C_i} C_i$$

Topology

Examples: Connected Component

Examples: Connected Component

Example 1: Consider $X = \{a, b, c, d, e\}$ with

$$\mathcal{T} = \{\emptyset, \{a, b\}, \{d, e\}, \{a, b, c\}, \{a, b, d, e\}, X\}$$

$C_a = \bigcup_{a \in C} C$ $\{a\}, \{a,b\}, \{a,c\}, \{a,b,c\}$

$\{a,b,d\} = (B \cap \{a,b,c\}) \cup (B \cap \{d,e\})$

B

$= \{a,b,c\}$

$\Rightarrow C_a = C_b = C_c$

Now $C_d = \bigcup_{d \in C} C$ $\{d\}$ $\{d,e\}$ $\{c,d,e\}$ Not connected

$= \{d,e\} = C_e$ $X = C_a \sqcup C_d$.

Examples: Connected Component

Example 2: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{a, b\}, \{a, b, c\}, X\}$$

$C_a = \bigcup_{a \in C} C = X = C_b = C_c = C_d$

KHALIL SAHIR
MC200400708
03054232382

Topology

Connected Component is Closed

Connected Component is Closed

Theorem:
Let C be a connected component of X then

$$C = \overline{C}$$

i.e.
A Connected component is a closed subset of X.

Connected Component is Closed

Proof: Let C be a connected component of X.

$C \subset \overline{C}$ —①

But we know that C is connected and if a set is connected then its closure is connected too.

$\Rightarrow \overline{C}$ is connected.

But C is the max connected subset.

$\Rightarrow C \subset \overline{C}$ —② $\Rightarrow C = \overline{C}$ □

Topology

Connected Component is not Open

Connected Component is not Open



We have discussed about the closed nature of connected component of a topological space $(X, \mathcal{T})$.
Now one may ask the following question.
Question:
Is a connected component C of X open?
Answer:
No, not in general.

Connected Component is not Open

Example: Let $\mathbb{Q} \subset \mathbb{R}$ $(\mathbb{R}, \mathcal{T}_u)$

Now the connected components of $(\mathbb{Q}, \mathcal{T}_{\mathbb{R}_u})$ are single tons

$\mathbb{Q} = \coprod_{x \in \mathbb{Q}} \{x\}$

Now $\{x\} \subset \mathbb{Q}$ Not open

$\therefore U_x \subset_{\text{open}} \mathbb{Q}$ s.t $U_x \subset \{x\}$

Topology

Locally Connected Spaces

Locally Connected Spaces

Def:
Let $(X, \mathcal{T})$ be a topological space. X is said to be locally connected at a point $x \in X$ iff for every U_x, open subset of X containing x, there exists V_x, an open subset of X containing x, such that

- $V_x \subset U_x$
- V_x is connected.

Locally Connected Spaces



Locally Connected Spaces

Def:
A topological space $(X, \mathcal{T})$ locally connected iff it is locally connected at each of its points.
i.e.
For all $x \in X$ and for every U_x, open subset of X containing x, there exists V_x, an open subset of X containing x, such that

- $V_x \subset U_x$
- V_x is connected.

Topology

Examples: Locally Connected Spaces

Examples: Locally Connected...

Example 1: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{a, b\}, \{a, b, c\}, X\}$$

Examples: Locally Connected...

Example 1: Consider $X = \{a, b, c, d\}$ with

$$\mathcal{T} = \{\emptyset, \{a, b\}, \{a, b, c\}, X\}$$

"a" $U_a = \{a, b, c\}$ $U_a = \{a, b\}$ connected.

"b" U_b

Examples: Locally Connected...

Example 2: A discrete space is locally connected.

X $\quad \mathcal{T}_{dis} = \mathcal{P}(X)$

$x \in X \quad U_x \in \mathcal{P}(X)$

$V_x = \{x\} \subset U_x.$

Topology

Connectedness and Locally Connectedness

KHALIL SAHIR
MC200400708
03054232382

Connectedness and Locally...

Questions:

1. Does locally connectedness imply connectedness?
2. Does connectedness imply locally connectedness?

Dr. H

Connectedness and Locally...

Locally Connectedness $\not\Longrightarrow$ Connectedness

Example:

$X = B(-2;1) \cup B(2;1) \subset \mathbb{R}^2$

X is not connected.

X is Locally Connected.



Connectedness and Locally...

Connectedness $\not\Longrightarrow$ locally Connectedness

Example:

$$Y = \{(x,0) | x \in [0,1]\} \bigcup \left[\bigcup_{n \in \mathbb{Z}^+} \left\{ \left(x, \frac{x}{n}\right) | x \in [0,1] \right\} \right] \subset \mathbb{R}^2$$

$n=1 \quad \{(x,x) \mid x \in [0,1]\}$

$n=2 \quad \{(x, \frac{x}{2}) | x \in [0,1]\}$

Now Y is connected.

Connectedness and Locally...

Now Y is not Locally Connected.

$(1,0) \in Y$ s.t Y is not L.C at $(1,0)$

$(1,0)$ $U_{(1,0)}$ s.t $\nexists$ $V_{(1,0)}$ Not connected $\cup$ $(1,0)$

$U_{(1,0)}$

$\Rightarrow$ Y is not L.C



Topology

Path Connected Spaces

Path Connected Spaces

Path:
Let $(X, \mathcal{T})$ be a topological space and $x, y \in X$. A path from x to y in X is a continuous map

$$p : [0,1] \to X$$

such that

$$p(0) = x$$

and

$$p(1) = y.$$

Dr. Hani Shaki

Path Connected Spaces



Path Connected Spaces

Path Connected Space:
A topological space $(X, \mathcal{T})$ is said to be path connected iff for every $x, y \in X$ there exists a path p from x to y in X.

Examples:

- $(\mathbb{R}, \mathcal{T}_u)$
- $(\mathbb{R}^2, \mathcal{T}_u)$

Topology

Facts: Path Connected Spaces: 1

Facts: Path Connected Spaces: 1

Theorem:
Let $(X, \mathcal{T})$ be a path connected topological space then $(X, \mathcal{T})$ is connected.

i.e.

Path connectedness implies connectedness.

Dr. Hani

Facts: Path Connected Spaces: 1

Proof Let $x \in X$

$\because$ X is path connected. So $\forall\, y \in X$

$\exists$ a path from x to y.

i.e. $\gamma_y : I \to X$ s.t $\gamma_y(0) = x$, $\gamma_y(1) = y$

($I = [0,1]$)

So $X = \bigcup_{y \in X} \gamma_y(I)$

Now $\gamma_y(0) = x \quad \forall\, y \in X$

$\Rightarrow x \in \gamma_y(I) \quad \forall\, y \in X.$

Facts: Path Connected Spaces: 1

$\Rightarrow x \in \bigcap_{y \in X} \gamma_y(I)$

$\Rightarrow X = \bigcup_{y \in X} \gamma_y(I)$

$\to$ is connected.

$\gamma_y(I)$ is connected

$\because$ I is connected

And we know that The image of a connected space is connected under a cont map.

Facts: Path Connected Spaces: 1

Remarks ① Connectedness $\not\Rightarrow$ Path connectedness

② $D \subsetneq_{\text{open}} \mathbb{R}^n$

if D is connected $\Rightarrow D$ is p

Topology

Facts: Path Connected Spaces: 2

Facts: Path Connected Spaces: 2

Proposition:
Let $(X, \mathcal{T})$ be a path connected topological space and

$$f : (X, \mathcal{T}) \longrightarrow (Y, \mathcal{T}')$$

be a continuous map. Then $f(X)$ is path connected.

Facts: Path Connected Spaces: 2

Proof: $f : (X, \mathcal{T}) \xrightarrow[\text{Cont}]{} (Y, \mathcal{T}')$

$\downarrow$ Path connected.

$f(X)$ is path connected ??

Let $a, b \in f(X) \Rightarrow \exists\ x, y \in X$ s.t $f(x) = a$, $f(y) = b$

Now $\because$ X is path connected.

So $\exists$ $p : I \xrightarrow[\text{Cont}]{} X$ $\quad p(0) = x$, $p(1) = y$

Facts: Path Connected Spaces: 2

$q = f \circ p : I \longrightarrow f(X)$ ✓

$q(0) = f \circ p(0) = f(p(0)) = f(x) = a$ ✓

$q(1) = f \circ p(1) = f(p(1)) = f(y) = b$ ✓

$\Rightarrow$ $f(X)$ is path connected.

KHALIL SAHIR
MC200400708
03054232382