VU Quiz Collection

Total Quizzes: 10

Courses: MGT211 - Introduction To Business (Quiz 03), MTH603 - Numerical Analysis (Quiz No 02)

Generated: 11/25/2025, 10:59:56 AM

Powered by Firewall Bypass VU Extension
Waseem Akram | wasii.dev | 03137119351

VU Quiz Collection

Generated by Firewall Bypass VU Extension

Quiz 1 - MGT211 - Introduction To Business (Quiz 03)

Question:

How does operations management contribute to customer satisfaction?

Options:

A. By outsourcing HR functions

B. By cutting marketing costs

C. By minimizing production

D. By ensuring timely, quality delivery

Copied: 11/25/2025, 10:56:38 AM

Quiz 2 - MGT211 - Introduction To Business (Quiz 03)

Question:

Which of the following companies uses robotic assembly lines?

Options:

A. Nestlé

B. McDonald’s

C. Tesla

D. Facebook

Copied: 11/25/2025, 10:56:50 AM

Quiz 3 - MGT211 - Introduction To Business (Quiz 03)

Question:

Why is “People” important in the 7Ps of marketing?

Options:

A. It refers to product designers

B. It includes factory workers

C. It ensures automation

D. It impacts service quality

Copied: 11/25/2025, 10:57:01 AM

Quiz 4 - MGT211 - Introduction To Business (Quiz 03)

Question:

Which of the following is an example of sustainability in operations?

Options:

A. Using eco-friendly packaging

B. Expanding warehouses

C. Increasing packaging waste

D. Hiring fewer workers

Copied: 11/25/2025, 10:57:13 AM

Quiz 5 - MTH603 - Numerical Analysis (Quiz No 02)

Question:

The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric _____ definite matrices A.

Options:

A. positive

B. negative

Copied: 11/25/2025, 10:57:59 AM

Quiz 6 - MTH603 - Numerical Analysis (Quiz No 02)

Question:

While using power method, the computed vector
u (2) =#›#•#œ 1 . 8" 0 . 8 0 . 6#ž# #Ÿ u (2) = (1 . 8" 0 . 8 0 . 6) { u ^ { (2) } } = \l e f t({\b e g i n{a r r a y}{*{2 0
{1.8}\\{ - 0.8}\\{0.6}
\end{array}} \right)
will be in normalized form as

Options:

A . u (2) =" 0 . 8#›#•#œ" 1 . 8 / 0 . 8 1" 0 . 6 / 0 . 8#ž# #Ÿ u (2) =" 0 . 8 (" 1 . 8 / 0 . 8 1" 0 . 6 / 0 . 8) { u ^ { (2) } }
{ - 1.8/0.8}\\1\\{ - 0.6/0.8}
\end{array}} \right)

B . u (2) = 1 . 8#›#•#œ 1" 0 . 8 / 1 . 8 0 . 6 / 1 . 8#ž# #Ÿ u (2) = 1 . 8 (1" 0 . 8 / 1 . 8 0 . 6 / 1 . 8) { u ^ { (2) } } = 1
1\\{ - 0.8/1.8}\\{0.6/1.8}
\end{array}} \right)

C . u (2) = 0 . 6#›#•#œ 1 . 8 / 0 . 6" 0 . 8 / 0 . 6 1#ž# #Ÿ u (2) = 0 . 6 (1 . 8 / 0 . 6" 0 . 8 / 0 . 6 1) { u ^ { (2) } } = 0
{1.8/0.6}\\{ - 0.8/0.6}\\1
\end{array}} \right)

D . u (2) = 2 . 8#›#•#œ 1 . 8 / 2 . 8" 0 . 8 / 2 . 8 1#ž# #Ÿ u (2) = 2 . 8 (1 . 8 / 2 . 8" 0 . 8 / 2 . 8 1) { u ^ { (2) } } = 2
{1.8/2.8}\\{ - 0.8/2.8}\\1
\end{array}} \right)

Copied: 11/25/2025, 10:58:12 AM

Quiz 7 - MTH603 - Numerical Analysis (Quiz No 02)

Question:

If n x n matrices A and B are similar, then they have the different eigenvalues (with the same multiplicities).

Options:

A. TRUE

B. FALSE

Copied: 11/25/2025, 10:58:29 AM

Quiz 8 - MTH603 - Numerical Analysis (Quiz No 02)

Question:

While using power method, from the resultant normalize vector
u (3) = 3 . 1 7 8 9 0#›#•#œ 1 . 0 1 . 2 3 1 9 2 6 1 . 4 2 1 3 8 9#ž# #Ÿ u (3) = 3 . 1 7 8 9 0 (1 . 0 1 . 2 3 1 9 2 6 1 . 4 2 1 3 8 9
{\begin{array}{*{20}{c}}
{1.0}\\{1.231926}\\{1.421389}
\end{array}} \right)
we have the largest eigen value and the corresponding eigenvector as

Options:

A . » = 3 . 1 7 8 9 0 , (X) =#›#•#œ 1 . 0 1 . 2 3 1 9 2 6 1 . 4 2 1 3 8 9#ž# #Ÿ » = 3 . 1 7 8 9 0 , (X) = (1 . 0 1 . 2 3 1 9 2
3.17890,\,\,(X) = \left({\begin{array}{*{20}{c}}
{1.0}\\{1.231926}\\{1.421389}
\end{array}} \right)

B . » = 3 . 1 7 8 9 0 , (X) =#›#•#œ 1 . 2 3 1 9 2 6 1 . 0 1 . 4 2 1 3 8 9#ž# #Ÿ » = 3 . 1 7 8 9 0 , (X) = (1 . 2 3 1 9 2 6 1 .
3.17890,\,\,(X) = \left({\begin{array}{*{20}{c}}
{1.231926}\\{1.0}\\{1.421389}
\end{array}} \right)

C . » = 3 . 1 7 8 9 0 , (X) =#›#•#œ 1 . 2 3 1 9 2 6 1 . 4 2 1 3 8 9 1 . 0#ž# #Ÿ » = 3 . 1 7 8 9 0 , (X) = (1 . 2 3 1 9 2 6 1 .
3.17890,\,\,(X) = \left({\begin{array}{*{20}{c}}
{1.231926}\\{1.421389}\\{1.0}
\end{array}} \right)

D . » = 3 . 1 7 8 9 0 , (X) =#›#•#œ 1 . 0 1 . 4 2 1 3 8 9 1 . 2 3 1 9 2 6#ž# #Ÿ » = 3 . 1 7 8 9 0 , (X) = (1 . 0 1 . 4 2 1 3 8
3.17890,\,\,(X) = \left({\begin{array}{*{20}{c}}
{1.0}\\{1.421389}\\{1.231926}
\end{array}} \right)

Copied: 11/25/2025, 10:59:06 AM

Quiz 9 - MTH603 - Numerical Analysis (Quiz No 02)

Question:

While using power method, from the resultant normalize vector
u (2) = 9 . 8 5 4 9 0#›#•#œ 0 . 6 7 8 9 5 8 1 . 0 1 . 1 2 3 6 8 5#ž# #Ÿ u (2) = 9 . 8 5 4 9 0 (0 . 6 7 8 9 5 8 1 . 0 1 . 1 2 3 6 8 5
{\begin{array}{*{20}{c}}
{0.678958}\\{1.0}\\{1.123685}
\end{array}} \right)

we have the largest eigen value and the corresponding eigenvector as

Options:

A . » = 9 . 8 5 4 9 0 , (X) =#›#•#œ 0 . 6 7 8 9 5 8 1 . 0 1 . 1 2 3 6 8 5#ž# #Ÿ » = 9 . 8 5 4 9 0 , (X) = (0 . 6 7 8 9 5 8 1 .
9.85490,\,\,(X) = \left({\begin{array}{*{20}{c}}
{0.678958}\\{1.0}\\{1.123685}
\end{array}} \right)

B . » = 1 . 1 2 3 6 8 5 , (X) =#›#•#œ 0 . 6 7 8 9 5 8 1 . 0 9 . 8 5 4 9 0#ž# #Ÿ » = 1 . 1 2 3 6 8 5 , (X) = (0 . 6 7 8 9 5 8 1
1.123685,\,\,(X) = \left({\begin{array}{*{20}{c}}
{0.678958}\\{1.0}\\{9.85490}
\end{array}} \right)

C . » = 1 . 1 2 3 6 8 5 , (X) =#›#•#œ 9 . 8 5 4 9 0 1 . 0 0 . 6 7 8 9 5 8#ž# #Ÿ » = 1 . 1 2 3 6 8 5 , (X) = (9 . 8 5 4 9 0 1 .
1.123685,\,\,(X) = \left({\begin{array}{*{20}{c}}
{9.85490}\\{1.0}\\{0.678958}
\end{array}} \right)

D . » = 9 . 8 5 4 9 0 , (X) =#›#•#œ 1 . 1 2 3 6 8 5 1 . 0 0 . 6 7 8 9 5 8#ž# #Ÿ » = 9 . 8 5 4 9 0 , (X) = (1 . 1 2 3 6 8 5 1 .
9.85490,\,\,(X) = \left({\begin{array}{*{20}{c}}
{1.123685}\\{1.0}\\{0.678958}
\end{array}} \right)

Copied: 11/25/2025, 10:59:33 AM

Quiz 10 - MTH603 - Numerical Analysis (Quiz No 02)

Question:

If n x n matrices A and B are similar, then they have the _____ eigenvalues (with the same multiplicities).

Options:

A. different

B. same

Copied: 11/25/2025, 10:59:56 AM