VU Quiz Collection

Total Questions: 10 from 10 Quiz(es)

Courses: MTH603

Generated: 5/7/2026, 12:52:24 PM

Powered by Firewall Bypass VU Extension
Waseem Akram | wasii.dev | +1 (226) 798-2360

Quiz 1 - MTH603 (Q1/10)

Question:

The linear equation: 0x+0y =2 has -------- solution/solutions.

Options:

A. no solution

B. finite many

C. unique

D. infinite many

Copied: 5/7/2026, 12:40:08 PM

Quiz 2 - MTH603 (Q2/10)

Question:

If the order of coefficient matrix corresponding to system of linear equations is 3*3 then which of the following will be the orders of its decomposed matrices; ‘L’ and ‘U’?

Options:

A. Order of ‘L’ = 3*1, Order of ‘U’ = 1*3

B. Order of ‘L’ = 3*2, Order of ‘U’ = 2*3

C. Order of ‘L’ = 3*4, Order of ‘U’ = 4*3

D. Order of ‘L’ = 3*3, Order of ‘U’ = 3*3

Copied: 5/7/2026, 12:40:38 PM

Quiz 3 - MTH603 (Q3/10)

Question:

The linear equation: x+y=1 has --------- solution/solutions.

Options:

A. no solution

B. unique

C. finite many

D. infinite many

Copied: 5/7/2026, 12:41:26 PM

Quiz 4 - MTH603 (Q4/10)

Question:

Numerical methods for finding the solution of the system of equations are classified as direct and ……. methods

Options:

A. Jacobi

B. Indirect

C. Iterative

D. None of the given choices

Copied: 5/7/2026, 12:41:57 PM

Quiz 5 - MTH603 (Q5/10)

Question:

Full pivoting, in fact, is more …….than the partial pivoting.

Options:

A. Complicated

B. Easiest

Copied: 5/7/2026, 12:42:46 PM

Quiz 6 - MTH603 (Q6/10)

Question:

Which of the following rearrangement make strictly diagonal dominant, the system of linear equations; x-3y+z= –2, –6x+4y+11z=1, 5x–2y–2z=9?

Options:

A. 5x–2y–2z=9, –6x+4y+11z=1, x–3y+z= –2

B. –6x+4y+11z=1, x–3y+z= –2, 5x–2y–2z=9

C. No need to rearrange as system is already in diagonal dominant form.

D. 5x–2y–2z=9, x–3y+z= –2, –6x+4y+11z=1

Copied: 5/7/2026, 12:43:24 PM

Quiz 7 - MTH603 (Q7/10)

Question:

While using the Gauss-Seidel Method for finding the solution of the system of equation, the following system
x+2y+2z=3x+3y+3z=2x+y+5z=2x+2y+2z=3x+3y+3z=2x+y+5z=2\begin{array}{l}
x + 2y + 2z = 3\\x + 3y + 3z = 2\\x + y + 5z = 2
\end{array}
can be rewritten as

Options:

A. x = 3" 2 y" 2 z y = 2" x" 3 z z = 2" y" x x = 3" 2 y" 2 z y = 2" x" 3 z z = 2" y" x \ b e g i n { a r r a y } { l }
x = 3 - 2y - 2z\\y = 2 - x - 3z\\z = 2 - y - x
\end{array}

B. x = 1" 2 3 y" 2 3 z y = 1" 3 2 x" 3 2 z z = 1" y 2" x 2 x = 1" 2 3 y" 2 3 z y = 1" 3 2 x" 3 2 z z = 1" y 2" x 2 \ b e g
x = 1 - \frac{2}{3}y - \frac{2}{3}z\\y = 1 - \frac{3}{2}x - \frac{3}{2}z\\z = 1 - \frac{y}{2} - \frac{x}{2}
\end{array}

C. x = 3" 2 y" 2 z y = 2 3" x 3" z z = 2 5" x 5" y 5 x = 3" 2 y" 2 z y = 2 3" x 3" z z = 2 5" x 5" y 5 \ b e g i n { a r r a y
x = 3 - 2y - 2z\\y = \frac{2}{3} - \frac{x}{3} - z\\z = \frac{2}{5} - \frac{x}{5} - \frac{y}{5}
\end{array}

D. x = 3 2" y" z y = 2" x 3" z z = 2 5" x 5" y 5 x = 3 2" y" z y = 2" x 3" z z = 2 5" x 5" y 5 \ b e g i n { a r r a y } { l }
x = \frac{3}{2} - y - z\\y = 2 - \frac{x}{3} - z\\z = \frac{2}{5} - \frac{x}{5} - \frac{y}{5}
\end{array}

Copied: 5/7/2026, 12:44:57 PM

Quiz 8 - MTH603 (Q8/10)

Question:

In Gauss-Jacobi’s method, the corresponding elements of x(r+1)ixi(r+1)x_i^{(r + 1)} replaces those of x(r)ixi(r)x_i^{(r)} as soon as they become available.

Options:

A. False

B. True

Copied: 5/7/2026, 12:48:59 PM

Quiz 9 - MTH603 (Q9/10)

Question:

In ……. method, matrix [A] of the system of equations is decomposed into the product of two matrices [L] and [U], where [L] is a lower-triangular matrix and [U] is an upper-triangular matrix with 1’s on its main diagonal.

Options:

A. Gaussian elimination

B. Crout’s Reduction

C. None of the given choices

D. Gauss-Seidel

Copied: 5/7/2026, 12:50:19 PM

Quiz 10 - MTH603 (Q10/10)

Question:

Which of the following system of equation is diagonally dominant?

Options:

A . x + 2 y + 9 z = 3 8 x + y" z = 5 2 x + 1 3 y + 5 z = 7 x + 2 y + 9 z = 3 8 x + y" z = 5 2 x + 1 3 y + 5 z = 7 \ b e g i {x + 2y + 9z = 3} \\\

{8x + y - z = 5} \\\
{2x + 13y + 5z = 7}
\end{array}

B. 2x+13y+5z=7 8x+y" z=5 x+2y+9z=3 2x+13y+5z=7 8x+y" z=5 x+2y+9z=3 \begin{array}{*{20}{l}}
{2x + 13y + 5z = 7} \\\
{8x + y - z = 5} \\\
{x + 2y + 9z = 3}
\end{array}

C. 8x+y" z=5 x+2y+9z=3 2x+13y+5z=7 8x+y" z=5 x+2y+9z=3 2x+13y+5z=7 \begin{array}{*{20}{l}}
{8x + y - z = 5} \\\
{x + 2y + 9z = 3} \\\
{2x + 13y + 5z = 7}
\end{array}

D. 8x+y" z=5 2x+13y+5z=7 x+2y+9z=3 8x+y" z=5 2x+13y+5z=7 x+2y+9z=3 \begin{array}{*{20}{l}}
{8x + y - z = 5} \\\
{2x + 13y + 5z = 7} \\\
{x + 2y + 9z = 3}
\end{array}

Copied: 5/7/2026, 12:51:22 PM