ZOO 405
ECOLOGY
MID-PPT
Module(001-056)

- MERGED BY BAQIR HUSSAIN BUZDAR
- BS ZOOLOGY
- CELL NO. 03336361244

Module# 01

What is Ecology

- Ecology is a branch of science, including human science, population, community, ecosystem and biosphere.
- Ecology is the study of organisms, the environment and how the organisms interact with each other and their environment.
- It is studied at various levels, such as organism, population, community, biosphere and ecosystem.

What is Ecology

- An ecologist's primary goal is to improve their understanding of life processes, adaptations and habitats, interactions and biodiversity of organisms.

- Merged by
- BAQIR HUSSAIN BS ZOOLOGY
- 03336361244

MODULE# 02

Biotic and Abiotic Factors

Biotic and Abiotic Factors

- Biotic and abiotic factors include the living and non-living factors and their interaction with the environment.

Biotic components

Biotic components are living factors of an ecosystem. A few examples of biotic components include bacteria, animals, birds, fungi, plants, etc.




Abiotic components

- Abiotic components are non-living chemical and physical factors of an ecosystem. These components could be acquired from the atmosphere, lithosphere and hydrosphere. A few examples of abiotic components include sunlight, soil, air, moisture minerals and more.
- Living organisms are grouped into biotic components, whereas non-living components like sunlight, water, topography are listed under abiotic components.


ABIOTIC COMPONENTS
1
2
3

MODULE# 03 Types of Ecology



Ecology can be classified into different types. The different types of ecology are given below:


Global Ecology

It deals with interactions among earth's ecosystems, land, atmosphere and oceans. It helps to understand the large-scale interactions and their influence on the planet.

Landscape Ecology

It deals with the exchange of energy, materials, organisms and other products of ecosystems. Landscape ecology throws light on the role of human impacts on the landscape structures and functions.

MODULE# 04 Types of Ecology



Ecosystem Ecology

It deals with the entire ecosystem, including the study of living and non-living components and their relationship with the environment. This science researches how ecosystems work, their interactions, etc.

Community Ecology

It deals with how community structure is modified by interactions among living organisms. Ecology community is made up of two or more populations of different species living in a particular geographic area.


- **Population Ecology**

- It deals with factors that alter and impact the genetic composition and the size of the population of organisms. Ecologists are interested in fluctuations in the size of a population, the growth of a population and any other interactions with the population.

- In biology, a population can be defined as a set of individuals of the same species living in a given place at a given time. Births and immigration are the main factors that increase the population and death and emigration are the main factors that decrease the population.

- Population ecology examines the population distribution and density. Population density is the number of individuals in a given volume or area. This helps in determining whether a particular species is in endanger or its number is to be controlled and resources to be replenished.

MODULE# 05 Types of Ecology



Organismal Ecology

Organismal ecology is the study of an individual organism's behaviour, morphology, physiology, etc. in response to environmental challenges. It looks at how individual organisms interact with biotic and abiotic components. Ecologists research how organisms are adapted to these non-living and living components of their surroundings.

Individual species are related to various adaptations like physiological adaptation, morphological adaptation, and behavioural adaptation.

Molecular Ecology

The study of ecology focuses on the production of proteins and how these proteins affect the organisms and their environment. This happens at the molecular level.

DNA forms the proteins that interact with each other and the environment. These interactions give rise to some complex organisms

MODULE# 06

Importance of Ecology

Conservation of Environment

- Ecology helps us to understand how our actions affect the environment. It shows the individuals the extent of damage we cause to the environment.

- Lack of understanding of ecology has led to the degradation of land and the environment. It has also led to the extinction and endangerment of certain species. For eg., dinosaurs, white shark, mammoths, etc. Thus, the study of the environment and organisms helps us to protect them from any damage and danger.

Resource Allocation

With the knowledge of ecology, we are able to know which resources are necessary for the survival of different organisms. Lack of ecological knowledge has led to scarcity and deprivation of these resources, leading to competition.

MERGED BY BAQIR HUSSAIN BS ZOOLOGY 03336361244

Energy Conservation

All organisms require energy for their growth and development. Lack of ecological understanding leads to the over-exploitation of energy resources such as light, nutrition and radiation, leading to its depletion.Proper knowledge of ecological requirements prevents the unnecessary wastage of energy resources, thereby, conserving energy for future purposes.

Eco-Friendliness


Ecology encourages harmonious living within the species and the adoption of a lifestyle that protects the ecology of life.

MODULE# 07

Laws of thermodynamics

Definition:

- The laws of thermodynamics are important unifying principles of biology.
- These principles govern the chemical processes (metabolism) in all biological organisms.


Energy and human life
Chemical energy
- Carbohydrates
- Fats
- Others
Chemical waste
- Carbon dioxide
- Water
Heat
ATP
- body's "energy currency"
metabolism
Heat

- The First Law of Thermodynamics, also known as the law of conservation of energy, states that energy can neither be created nor destroyed. It may change from one form to another, but the energy in a closed system remains constant.

- The Second Law of Thermodynamics states that when energy is transferred, there will be less energy available at the end of the transfer process than at the beginning. Due to entropy, which is the measure of disorder in a closed system, all of the available energy will not be useful to the organism. Entropy increases as energy is transferred.

MODULE# 08

First Law of Thermodynamics

- All biological organisms require energy to survive. In a closed system, such as the universe, this energy is not consumed but transformed from one form to another.
- Cells, for example, perform a number of important processes. These processes require energy. In photosynthesis, the energy is supplied by the sun.
- Light energy is absorbed by cells in plant leaves and converted to chemical energy.

- The chemical energy is stored in the form of glucose, which is used to form complex carbohydrates necessary to build plant mass.

- The energy stored in glucose can also be released through cellular respiration.

- This process allows plant and animal organisms to access the energy stored in carbohydrates, lipids, and other macromolecules through the production of ATP.

- This energy is needed to perform cell functions such as DNA replication, mitosis, meiosis, cell movement, endocytosis, exocytosis, and apoptosis.

MODULE# 09

Second Law of Thermodynamics

As with other biological processes, the transfer of energy is not 100% efficient. In photosynthesis,

for example, not all of the light energy is absorbed by the plant. Some energy is reflected and some is lost as heat.

The loss of energy to the surrounding environment results in an increase of disorder or entropy.

Unlike plants and other photosynthetic organisms, animals cannot generate energy directly from the sunlight. They must consume plants or other animal organisms for energy.

The higher up an organism is on the food chain, the less available energy it receives from its food sources.

Much of this energy is lost during metabolic processes performed by the producers and primary consumers that are eaten.

Therefore, much less energy is available for organisms in higher trophic levels. The lower the available energy, the less number of organisms can be supported.

This is why there are more producers than consumers in an ecosystem.

Living systems require constant energy input to maintain their highly ordered state.

Cells, for example, are highly ordered and have low entropy. In the process of maintaining this order, some energy is lost to the surroundings or transformed.

So while cells are ordered, the processes performed to maintain that order result in an increase in entropy in the cell's/organism's surroundings.

The transfer of energy causes entropy in the universe to increase

MODULE# 10

Food chains and food webs Key points

Producers, or autotrophs, make their own organic molecules.

Consumers, or heterotrophs, get organic molecules by eating other organisms.

Food chain is a linear sequence of organisms through which nutrients and energy pass as one organism eats another.

- In a **food chain**, each organism occupies a different trophic level, defined by how many energy transfers separate it from the basic input of the chain.

- **Food webs** consist of many interconnected food chains and are more realistic representation of consumption relationships in ecosystems.

- Energy transfer between trophic levels is inefficient (with a typical efficiency around 10 %). This inefficiency limits the length of food chains.

- Merged by BAQIR HUSSAIN
- BS ZOOLOGY
- 03336361244

MODULE# 11 Autotrophs

- Autotrophs ("self-feeders"), can make their own food – that is, their own organic compounds.

Two basic types of autotrophs:

1. Photoautotrophs
2. Chemoautotrophs

Photoautotrophs

- Such as plants, use energy from sunlight to make organic compounds (sugars) out of carbon dioxide in photosynthesis.

- Other examples of photoautotrophs include algae and cyanobacteria.

Chemoautotrophs

- Chemoautotrophs use energy from chemicals to build organic compounds out of carbon dioxide (or similar molecules). This is called chemosynthesis.


- For instance, there are hydrogen sulfide-oxidizing chemoautotrophic bacteria found in undersea vent communities (where no light can reach).


MODULE# 12

Heterotrophs

- Heterotrophs (“other-feeders”) such as humans can't capture light or chemical energy to make their own food out of carbon dioxide.

- Instead, they get organic molecules by eating other organisms or their by-products. Animals, fungi, and many bacteria are heterotrophs.


Autotroph vs Heterotroph
AUTOTROPH
Producers
An autotroph is an organism that produces its own food.
Plants
Algae
Some Bacteria
Phytoplankton
HETEROTROPH
Consumers
A heterotroph is an organism that does not make its own food.
Animals
Fungi
Most Protozoa
Most Bacteria
sciencenotes.org


Heterotrophs
Autotrophs
Light Energy
Oxygen
Carbon Dioxide
Water and Nutrients from Soil

MODULE# 13

Food chains

- Food chain is a linear sequence of organisms through which nutrients and energy pass as one organism eats another.
- At the base of the food chain lie the **primary producers**. The primary producers are autotrophs, and are most often photosynthetic organisms (such as plants, algae, or cyanobacteria).

- The organisms that eat the primary producers are called **primary consumers**. Primary consumers are usually herbivores (plant-eaters), though they may be algae or bacteria eaters.


- The organisms that eat the primary consumers are called **secondary consumers**. Secondary consumers are generally meat-eaters (carnivores).


- The organisms that eat the secondary consumers are called **tertiary consumers**. These are carnivore-eating carnivores, like eagles or big fish.

- Some food chains have additional levels, such as **quaternary consumers** (carnivores that eat tertiary consumers).

- Organisms at the very top of a food chain are called the **apex consumers**.

Food Chains



grass
(producer)
grasshopper
(primary consumer)
bluebird
(secondary consumer)
snake
(tertiary consumer)
owl
(apex predator)
(decomposers)



teachoo.com
Food Chains
Trees
Deer
Lion
Flower
Flies
Frog
Snake
Eagle
Flower
Small Fishes
Larger Fish
Birds

MODULE# 14

Trophic level

The trophic level of an organism is the number of steps it is from the start of the chain. A food web starts at trophic level 1 with primary producers such as plants, can move to herbivores at level 2, carnivores at level 3 or higher, and typically finish with apex predators at level 4 or


Energy flow and trophic levels
eagle
tertiary consumers
10 kcal
heat loss
snake
secondary consumers
100 kcal
heat loss
rabbit
primary consumers
1,000 kcal
energy
from
Sun
heat loss
producers
10,000 kcal
© Encyclopædia Britannica, Inc.


carnivore
omnivore
herbivore
producers
decomposer
(ex. detritivore)
trophic levels

Trophic levels divide up energy levels within an ecosystem

An organism is placed into a trophic level based on how it receives its energy. The levels are classified as the following:

Primary Producers:


Primary producers receive their energy from sunlight. So, all primary producers are autotrophs. Without primary producers, the sun's energy would not be converted into usable energy for the rest of the ecosystem.

Primary Consumers:


Primary consumers are the first level of consumers. They feed exclusively on producers. So, all primary consumers are heterotrophs.

Secondary Consumers:

Secondary consumers feed off of primary consumers secondary consumers are carnivores.



Trophic levels are helpful because they allow us to group organisms based on their eating habits and define each step of a food chain.


TROPHIC LEVELS
apex predator
tertiary consumer
secondary consumers
primary consumers
producers


MODULE# 15 Nitrogen Cycle


- Nitrogen Cycle is a biogeochemical process through which nitrogen is converted into many forms, consecutively passing from the atmosphere to the soil to organism and back into the atmosphere.

- It involves several processes such as nitrogen fixation, nitrification, denitrification, decay and putrefaction.

Process of the Nitrogen Cycle consists of the following steps

- Nitrogen fixation
- Nitrification
- Assimilation
- Ammonification
- Denitrification


NITROGEN CYCLE
BYJU'S
The Learning App
N_2
NITROGEN FIXATION
DENITRIFICATION
AMMONIFICATION
NH_4
NITROGEN FIXING BACTERIA
NO_2
ASSIMILATION
DENITRIFICATION BACTERIA
NH_3
NITRIFICATION
NO_3
NITRIFYING BACTERIA


The Nitrogen Cycle
Nitrous oxide from fossil fuels falls as nitric acid in rainwater
Lighting creates soil nitrates
The air is 78% nitrogen gas
Nitrogen compounds eventually break down into gas and return to the air
Runoff from nitrates in farming enters water
Animals eat plants
Plants obtain nitrate ions from soil and water bacteria
Bacteria in soil and water 'fix' nitrogen from the air
Decomposer bacteria break down animal and plant matter into nitrogen compounds

Nitrogen Fixation

- It is the initial step of the nitrogen cycle. Here, Atmospheric nitrogen (N_2) which is primarily available in an inert form, is converted into the usable form -ammonia (NH_3).
- During the process of Nitrogen fixation, the inert form of nitrogen gas is deposited into soils from the atmosphere and surface waters, mainly through precipitation.
- The entire process of Nitrogen fixation is completed by symbiotic bacteria, which are known as Diazotrophs. Azotobacter and Rhizobium also have a major role in this process

Nitrification

- In this process, the ammonia is converted into nitrate by the presence of bacteria in the soil. Nitrites are formed by the oxidation of ammonia with the help of Nitrosomonas bacteria species. Later, the produced nitrites are converted into nitrates by Nitrobacter. This conversion is very important as ammonia gas is toxic for plants.

- The reaction involved in the process of Nitrification is as follows:

- $2NH_3 + 3O_2 \rightarrow 2NO_2^- + 2H^+ + 2H_2O$

- $2NO_2^- + O_2 \rightarrow 2NO_3^-$

Denitrification

- Denitrification is the process in which the nitrogen compounds make their way back into the atmosphere by converting nitrate (NO_3^-) into gaseous nitrogen (N). This process of the nitrogen cycle is the final stage and occurs in the absence of oxygen. Denitrification is carried out by the denitrifying bacterial species- Clostridium and Pseudomonas, which will process nitrate to gain oxygen and gives out free nitrogen gas as a byproduct.

MODULE# 16

Assimilation


- Primary producers – plants take in the nitrogen compounds from the soil with the help of their roots, which are available in the form of ammonia, nitrite ions, nitrate ions or ammonium ions and are used in the formation of the plant and animal proteins. This way, it enters the food web when the primary consumers eat the plants.

Ammonification

- When plants or animals die, the nitrogen present in the organic matter is released back into the soil. The decomposers, namely bacteria or fungi present in the soil, convert the organic matter back into ammonium. This process of decomposition produces ammonia, which is further used for other biological processes.

Denitrification

- Denitrification is the process in which the nitrogen compounds make their way back into the atmosphere by converting nitrate (NO_3^-) into gaseous nitrogen (N). This process of the nitrogen cycle is the final stage and occurs in the absence of oxygen. Denitrification is carried out by the denitrifying bacterial species- Clostridium and Pseudomonas, which will process nitrate to gain oxygen and gives out free nitrogen gas as a byproduct.


Nitrogen Gas
N_2
Nitrogen fixation
Organic Nitrogen
Ammonification
Ammonium
NH_4^+
Nitrification
Nitrite, Nitrate
NO_2^- NO_3^-
Denitrification

Importance of Nitrogen Cycle

- Helps plants to synthesise chlorophyll from the nitrogen compounds.
- Helps in converting inert nitrogen gas into a usable form for the plants through the biochemical process.
- In the process of ammonification, the bacteria help in decomposing the animal and plant matter, which indirectly helps to clean up the environment.

- Nitrates and nitrites are released into the soil, which helps in enriching the soil with the necessary nutrients required for cultivation.

- Nitrogen is an integral component of the cell and it forms many crucial compounds and important biomolecules.

MODULE#17

Phosphorus Cycle


- Phosphorus is an important element for all living organisms. It forms a significant part of the structural framework of DNA and RNA. They are also an important component of ATP. Humans contain 80% of phosphorus in teeth and bones.
- Phosphorus cycle is a very slow process. Various weather processes help to wash the phosphorus present in the rocks into the soil. Phosphorus is absorbed by the organic matter in the soil which is used for various biological processes.


Steps of Phosphorus Cycle

Following are the important steps of phosphorus cycle

- Weathering
- Absorption by Plants
- Absorption by Animals
- Return to the Environment through Decomposition

- MERGED BY BAQIR HUSSAIN BS ZOOLOGY
- CELL NUMBER: 03336361244


Phosphorus Cycle
Plants
Animals
fertilizer runoff
Soils
Weathering
and
erosion
Marine animals
Marine plants
Dissolved pohsphates
Phosphate
rock formation
Sediments


PHOSPHORUS CYCLE
Weathering and Erosion
Rivers
Fertilizer Runoff
Phosphate Rock Formation
Plants
Animals
Marine Plants
Marine Animals
Dissolved Phosphates
SEDIMENTS
SOIL
Loss in Drainage
shutterstock
IMAGE ID: 1549733282
www.shutterstock.com

Weathering

- Phosphorus is found in the rocks in abundance. That is why the phosphorus cycle starts in the earth's crust. The phosphate salts are broken down from the rocks. These salts are washed away into the ground where they mix in the soil.

Absorption by Plants

- The phosphate salts dissolved in water are absorbed by the plants. However, the amount of phosphorus present in the soil is very less. That is why the farmers apply phosphate fertilizers on agricultural land.

- The aquatic plants absorb inorganic phosphorus from lower layers of water bodies. Since phosphate salts do not dissolve in water properly, they affect plant growth in aquatic ecosystems.

Absorption by Animals

- The animals absorb phosphorus from the plants or by consuming plant-eating animals. The rate of the phosphorus cycle is faster in plants and animals when compared to rocks.

Return of Phosphorus Back to the Ecosystem

- When the plants and animals die they are decomposed by microorganisms During this process, the organic form of phosphorus is converted into the inorganic form, which is recycled to soil and water.

- Soil and water will end up in sediments and rocks, which will again release phosphorus by weathering. Thus, the phosphorus cycle starts over.

MODULE# 18


Sulphur Cycle

- Sulphur is one of the most abundant elements on the earth. It is a yellow, brittle, tasteless, odourless non-metal. Sulphur is present in all kinds of proteins. Plants directly absorb sulphur-containing amino acids such as methionine, cystine, and cysteine.
- Sulphur is released into the atmosphere by the burning of fossil fuels, volcanic activities, and decomposition of organic molecules.

Steps of Sulphur Cycle

Following are the important steps of the sulphur cycle:

- Decomposition of Organic Compounds
- Oxidation of Hydrogen Sulphide to Elemental Sulphur
- Oxidation of Elemental Sulphur
- Reduction of Sulphates


The Sulfur Cycle
Sulfate in the atmosphere
SO_2
SO_2
Wet deposition
Sulfate in water
Leaching
Plant residues
Decomposition and wast products
Plant uptake
Bacteria
SO_4
Sedimentation

Decomposition of Organic Compounds

- Protein degradation releases amino acids that contain sulphur. Sulphates are reduced to H_2S by the action of Desulfotomaculum bacteria.

Oxidation of Hydrogen Sulphide to Elemental Sulphur

- Hydrogen sulphide oxidises to produce elemental sulphur. Certain photosynthetic bacteria from the families Chlorobiaceae and Chromatiaceae initiate the oxidation process.

Oxidation of Elemental Sulphur

- Elemental sulphur present in the soil cannot be utilized directly by the plants. Therefore, it is converted into sulphates by chemolithotrophic bacteria.

Reduction of Sulphates


Sulphates are reduced to hydrogen sulphide by Desulfovibrio desulfuricans. This occurs in two steps:

- Firstly, the sulphates are converted to sulphites utilizing ATP.
- Secondly, the reduction of sulphite to hydrogen sulphide.

MODULE# 19


Water Cycle


The water cycle describes how water evaporates from the surface of the earth, rises into the atmosphere, cools and condenses into rain or snow in clouds, and falls again to the surface as precipitation.


Water Cycle Diagram
Sun
Condensation
Precipitation
Transpiration
Evaporation
River
Lake
Ocean
Underground Water

Steps of Water Cycle

- Evaporation
- Sublimation
- Condensation
- Precipitation
- Infiltration
- Runoff


The Hydrologic Cycle (Water Cycle)
Condensation
Transpiration
Precipitation
Evaporation
Evaporation
Snowmealt
Lake
Ocean
Percolation
Groundwater Flow


THE WATER CYCLE
CONDENSATION
PRECIPITATION
EVAPORATION
COLLECTION

Module# 20

Steps of Water Cycle

Evaporation

- The sun is the ultimate source of energy, and it powers most of the evaporation that occurs on earth. Evaporation generally happens when water molecules at the surface of water bodies become excited and rise into the air. These molecules with the highest kinetic energy accumulate into water vapour clouds. Evaporation usually takes place below the boiling point of water. Another process called evapotranspiration occurs when evaporation occurs through the leaves of plants. This process contributes to a large percentage of water in the atmosphere.

Sublimation

Sublimation occurs when snow or ice changes directly into water vapour without becoming water. It usually occurs as a result of dry winds and low humidity. Sublimation can be observed on mountain peaks, where the air pressure is quite low. The low air pressure helps to sublimate the snow into water vapour as less energy is utilised in the process. Another example of sublimation is the phase where fog bellows from dry ice. On earth, the primary source of sublimation is from the ice sheets covering the poles of the earth.

Condensation

- The water vapour that accumulated in the atmosphere eventually cools down due to the low temperatures found at high altitudes. These vapours become tiny droplets of water and ice, eventually coming together to form clouds.

Condensation

- The water vapour that accumulated in the atmosphere eventually cools down due to the low temperatures found at high altitudes. These vapours become tiny droplets of water and ice, eventually coming together to form clouds.

Precipitation

- Above o degrees centigrade, the vapours will condense into water droplets. However, it cannot condense without dust or other impurities. Hence, water vapours attach itself on to the particle's surface. When enough droplets merge, it falls out of the clouds and on to the ground below. This process is called precipitation (or rainfall). In particularly cold weather or extremely low air pressure, the water droplets freeze and fall as snow or hail.

Infiltration

- Rainwater gets absorbed into the ground through the process of infiltration. The level of absorption varies based on the material the water has seeped into. For instance, rocks will retain comparatively less water than soil. Groundwater can either follows streams or rivers. But sometimes, it might just sink deeper, forming aquifers.


Runoff

- If the water from rainfall does not form aquifers, it follows gravity, often flowing down the sides of mountains and hills; eventually forming rivers. This process is called runoff. In colder regions, icecaps form when the amount of snowfall is faster than the rate of evaporation or sublimation. The biggest icecaps on earth are found at the poles.

Module# 21

Carbon Cycle

Carbon cycle is the process where carbon compounds are interchanged among the biosphere, geosphere, pedosphere, hydrosphere, and atmosphere of the earth.


Carbon Cycle
Sunlight
CO_2 in the atmosphere
Photosynthesis by producers
Plant respiration
Fossil fuel combustion
Animal respiration
Carbon fixation by consumers
Decomposition
Fossils and fossil fuels
ScienceFacts.net


Steps of Carbon Cycle


Following are the major steps involved in the process of the carbon cycle:

- Carbon present in the atmosphere is absorbed by plants for photosynthesis.
- These plants are then consumed by animals and carbon gets bioaccumulated into their bodies.
- These animals and plants eventually die, and upon decomposing, carbon is released back into the atmosphere.

- Some of the carbon that is not released back into the atmosphere eventually become fossil fuels.

- These fossil fuels are then used for man-made activities, which pumps more carbon back into the atmosphere.


CARBON CYCLE
BYJU'S
The Learning App
Sunlight
CO_2
Photosynthesis
Auto and
factory
emissions
Animal
respiration
Organic
Carbon
Decay
organisms
Root
respiration
Dead organisms and
waste products
Fossils and fossil fuel


Carbon Cycle
CO_2
Photosynthesis
Plant respiration
Combustion
Animal respiration
Buzzle.com
Decay organisms
Mineral carbon
Root respiration


Module# 22

Limiting Factors

Limiting Factors are


- Conditions of the environment that limit the growth of a species.
- Biotic and abiotic factors that prevent the continuous growth of a population.

- Populations would continue to increase if they had all of the resources they require in unlimited amounts, but there are always factors that limit their increase.



Biology Online

Limiting factor



A limiting factor is an environmental factor or variable that has the capacity to restrict the growth, abundance, or distribution of a population in an ecosystem.

FOR EXAMPLE:
LIMITING FACTORS OF A PLANT GROWTH ARE THE AMOUNT OF AVAILABLE SUNLIGHT, WATER, AND NUTRIENTS THAT ARE ESSENTIAL IN PHOTOSYNTHESIS.

Carrying capacity


- Because of these limiting factors, each ecosystem has a finite capacity for growth connected to its carrying capacity.
- Carrying capacity is the number of individuals of a species that an ecosystem can support.


Module# 23

Biotic And Abiotic Factors

- **Biotic components** are the living things that have a direct or indirect influence on other organisms in an environment. For example plants, animals, and microorganisms and their waste materials.

- **Abiotic components** of an ecosystem include all chemical and physical elements i.e. non-living components.
- Abiotic components can vary from region to region, from one ecosystem to another. They mainly take up the role of life supporter.
- They determine and restrict the population growth, number, and diversity of biotic factors in an ecosystem. Hence, they are called limiting factors.



BIOTIC
FACTORS
BACTERIA
FUNGI
PLANTS
ARCHEA
ANIMALS
PROTISTS
ABIOTIC
FACTORS
AIR
SALINITY
SOIL
TEMPERATURE
LIGHT
WATER
MINERALS
pH
HUMIDITY
dreamstime.com
ID 217284856 © VectorMine


Biotic and Abiotic Factors
Biotic factors are living, while abiotic factors are non-living.
abiotic
biotic
abiotic
abiotic
biotic
biotic
biotic
abiotic
abiotic
biotic
sciencenotes.org

Biotic vs Abiotic Factors



Biotic Factors




Abiotic Factors

Module# 24


Producers

Producers:

- These include all the autotrophs. They use light energy and synthesize food on their own, e.g. plants, green algae, etc.
- Producers are those organisms that use photosynthesis to capture energy by using sunlight, water and carbon dioxide to create carbohydrates, and then use that energy to create more complex molecules like proteins, lipids and starches that are crucial to life processes.


Sunlight
Carbon Dioxide from the atmosphere
Glucose produced used as energy source by whole plant
Oxygen Released
Water from the soil


Producers



Consumers



Decomposers



- Producers funnel into the ecosystem the energy needed for its biological processes.

- The carbohydrates and other organic chemicals formed by the producers are utilized by the heterotrophs, or consumers.

- First by the herbivores who eat the plants--the primary consumers--then by the predators who eat the herbivores--the secondary, tertiary, and so on consumers.

- But at each step, much energy is lost. Less than 10 percent of the energy stored in plants is converted to herbivore mass. The loss from herbivore to predator is similar. Thus energy needs to be added to the ecosystem continuously.


Module# 25

Consumers

- Consumers are organisms (including humans) that get their energy from producers, regarding the flow of energy through an ecosystem.

- For example, producers, (such as plants), make their own food by the process of photosynthesis. An organism ate this plant, than it would be a primary consumer.

- The animal that eats this animal is known as the second order consumer.
- Scientifically, all consumers are either herbivores, carnivores, omnivores or detrivores (decomposers and other organism that break down organic matter).


EXAMPLES OF PRIMARY CONSUMERS
tertiary consumer
secondary consumer
producer
primary consumer
MAIL
Study.com


PRODUCERS
PRODUCE THEIR OWN FOOD


CONSUMERS
EAT OTHER THINGS

Module# 26 Decomposers

- **Decomposers**
  - *An organism that gets energy by breaking down the remains of dead organisms and the wastes of living things.*
  - Another kind of heterotroph
  - Most fungi and bacteria fall into this category

- Earthworms and insects also
- They are often called nature's recyclers
- They return carbon and nitrogen to the air and soil.

Decomposers



Worm Mushroom Insects Bacteria

Decomposer

Decomposers feed on the remains of other animals. Decomposers play an important role in the ecosystem. By digesting dead matter they put nutrients back into the soil, making them available to producers.




Decomposers

1. Decomposers are organisms that breakdown dead or decaying organisms.
2. Micro-organisms like the bacteria and fungi that carry out decomposition of the dead organic material.



Worm Mushroom Insects Bacteria



Module# 27

Abiotic Factors

- Abiotic components are the physical and/or the chemical factors that act on the living organisms at any part of their life.

- The physical and chemical factors are characteristic of the environment. Light, air, soil, and nutrients, etc. form the abiotic components of an ecosystem.


- The abiotic factors vary from ecosystem to ecosystem.

- In an aquatic ecosystem, the abiotic factors may include water pH, sunlight, turbidity, water depth, salinity, available nutrients and dissolved oxygen.

- Similarly, abiotic factors in terrestrial ecosystems can include soil, soil types, temperature, rain, altitude, wind, nutrients, sunlight etc.

ABIOTIC FACTORS

these are the non-living components of the ecosystem



- Sun is the energy source. Producers/plants use this energy to synthesize food in the presence of carbon dioxide and chlorophyll. The energy from the sun, through several chemical reactions, turns into chemical energy.

Biotic vs. Abiotic Factors

- Living
- Examples
  - Plants
  - Animals
  - Fungi
  - Bacteria






- Non-Living
- Examples
  - Water
  - Sunlight
  - Soil
  - Air
  - Temperature









Module# 28 Abiotic Factors


Climate

- The climate refers to the weather conditions and general temperature of an ecosystem. This plays a huge role in the maintenance of the ecosystem through the regulation of the temperature and weather conditions.

- Any organisms living in a particular ecosystem evolve to suit that ecosystem, so even minute changes in climate can seriously impact the lives of the organisms living there.

pH Balance:

- The pH balance of an ecosystem refers to the general level of acidity or alkalinity present in the environment.
- It is a scientific method of measuring whether a particular mixture or solution is acidic, neutral, or alkaline.
- Pure water has a pH of 7, meaning it is neutral. Acidic mixtures have a pH balance of less than 7 while alkaline mixtures have a pH balance of more than 7.
- This also affects the organisms in an environment, because many creatures or plants, or microorganisms cannot survive in certain pH ranges.


AIR
SUNLIGHT
WATER
SOIL

Light:


- It may be interesting to note that even something as simple as the level of light present in an environment or ecosystem can greatly affect the organisms in that ecosystem.
- Photosynthesis is the main form of energy consumption for plants, which means that all plants need a certain level of light to create their own food.
- Some plants in dark ecosystems have evolved to a point where they make do with minimal amounts of light, such as the plants found deep in the oceans where light doesn't reach.

Water:

- Currents in an underwater or ocean ecosystem, water currents play a huge role in the regulation of the ecosystem.
- Water currents are some of the most important and useful parts of an oceanic ecosystem and at the same time also one of the most destructive.
- They are important because they redistribute minerals, nutrients, heat, oxygen, etc. However, strong ocean currents can also uproot and carry away plants and other organisms, which simultaneously destroys part of the ecosystem and also delivers a source of food to another part of the ecosystem.

Biotic vs. Abiotic Factors

- Living
- Examples
  - Plants
  - Animals
  - Fungi
  - Bacteria






- Non-Living
- Examples
  - Water
  - Sunlight
  - Soil
  - Air
  - Temperature










Module# 29 Atmosphere

The Earth is a system consisting of four major interacting components:

- Atmosphere
- Hydrosphere
- Lithosphere
- Biosphere


- Atmosphere - atmo referring to steam and vapor
- The Earth is surrounded by a blanket of air, which we call the atmosphere.
- Earth's atmosphere is composed of about 78% nitrogen, 21% oxygen, and one percent other gases.


Other Gases 1%
Oxygen 21%
Nitrogen 78%


The atmosphere is also responsible for the weather as the weather occurs within the lower atmosphere.

The atmosphere consists of four unique layers

- The troposphere
- The stratosphere
- The mesosphere
- The thermosphere


ATMOSPHERE
LAYERS DIAGRAM
ATMOSPHERE
is considered to be the area around the Earth in which the gaseous medium rotates together with the Earth as a whole.
EXOSPHERE
EXOBASE
THERMOSPHERE
KARMAN LINE
MESOSPHERE
STARTOSPHERE
OZON LAYER
TROPOSPHERE
SUBTITLE TEXT
The atmosphere of the Earth arose the evaporation of the substance of cosmic bodies when they fall to the Earth
POWERSLIDES
1
WWW.POWERLIDES.COM

LAYERS OF THE ATMOSPHERE



Module# 30


ATMOSPHERE LAYERS

Troposphere:

- The bottom layer of the atmosphere is known as the troposphere.
- The troposphere is where the weather happens.
- It is the warmest near the Earth because of the heat rising from the earth's surface but it becomes colder with altitude

- This layer is separated from the next by what is known as tropopause.


- The tropopause is the point in which temperatures will begin to change due to the increase of altitude.


ATMOSPHERE OF EARTH
EXOSPHERE
EXOBASE
THERMOSPHERE
KARMAN LINE
MESOSPHERE
STRATOSPHERE
OZONE LAYER
TROPOSPHERE


Troposphere

Stratosphere:

- The stratosphere layer is above the tropopause.
- The stratosphere is where there large concentration of ozone gas is found.
- The ozone gasses are essential as they absorb a large percent of radiant solar energy, protecting the earth from harmful ultra violet rays also known as UV.


MESOSPHERE
STRATOPAUSE
-15° CELSIUS
(5° FAHRENHEIT)
50 KM
(30 MILES)
OZONE LAYER
STRATOSPHERE
OZONE LAYER
-51° CELSIUS
(-60° FAHRENHEIT)
10 KM
(6 MILES)
TROPOPAUSE
TROPOSPHERE

Mesosphere:

- The coldest of spheres is known as the mesosphere this is where the water vapor often freezes to create clouds that are purely made of ice.

- The mesosphere is separated from the thermosphere by the mesopause.

Thermosphere:


- The topmost layer is known as the thermosphere, this is where many satellites circle the earth.


- Due to the thin air and proximity of the sun, the temperatures in the thermosphere tend to rapidly increase and decrease.

Module#31 HYDROSPHERE

THE HYDROSPHERE:

- The hydrosphere refers to the most important resource.
- The hydrosphere includes all forms of water in the Earth's environment.
- The forms of water include things such as the ocean, lakes, rivers, snow and glaciers, water underneath the earth's surface and even the water vapour that is found in the atmosphere.


Atmosphere (air)
Biosphere
Hydrosphere (water)
Lithosphere (earth)



The Hydrosphere

- The hydrosphere is always in motion as seen through the movement and flow of water in rivers, streams and the ocean (beach).

- Plant and animal organisms rely on the hydrosphere for their

Module# 32 Lithosphere

- **Lithosphere** - litho referring to rocks and minerals
- It is believed the lithosphere evolved about 4.6 billion years ago.
- The lithosphere refers to the solid, rocky crust that covers the entire planet.
- This solid, rocky crust is composed of a number of different rocks that have been grouped into three categories based on how they are formed.



Earth's Layers
Exosphere
Thermosphere
Mesosphere
Stratosphere
Troposphere
Lithosphere
Asthenosphere
Crust 0–100km thick
Crust
Mantle
Outer Core
Inner Core
Liquid
Solid
Core

- **Metamorphic rocks** – Metamorphic rocks are formed by heat and / or pressure from pre-existing rocks.

- **Igneous rocks** – igneous rocks are formed by the cooling of hot molten rock also known as magma. When the hot magma cools it begins to harden meaning once it had fully cooled it create what is known to be an igneous rock.


- **Sedimentary rocks** – sedimentary rocks are formed from pre-existing rocks. When rocks erode and mix with other dirt, clay and particles then settle together the mix together to form a sedimentary rock.

- Merged by Baqir Hussain
- BS Zoology
- 03336361244



The Rock Cycle
Erosion
Deposition
Igneous rock
Cooling
Sedimentary rock
Metamorphic rock
Melting


Module# 33 BIOSPHERE

- The ecosphere (also sometimes called the 'biosphere)
- The biosphere is composed of all living organisms, including; plants and animals.
- It is believed that all life exists in the biosphere.
- Most of the living organisms are found from up to three meters below ground to thirty meters above it and also in the to 200 meters of the ocean and seas.


BIOSPHERE
BYJU'S
The Learning App
Birds
Vegetation
Animals in grounds
Animals
Sea creatures
Marine living oraganisms
Dead animals / plants

- The biosphere could not survive if it wasn't for the other spheres as all organisms need water from the hydrosphere, minerals for the lithosphere and gases from the atmosphere.
- Energy flow is essential to maintain the structure of organisms by the splitting of phosphate bonds.


Atmosphere - Air
BIOSPHERE
Hydrosphere - Water
Lithosphere - Earth


ATMOSPHERE - AIR
BIOSPHERE
LITHOSPHERE - EARTH
HYDROSPHERE - WATER

Module# 34 Ecological niche

- **Ecology** is the study of the interactions between organisms and their environments, which comprise an ecosystem. The places organisms live in are called habitats.

- An **ecological niche**, in contrast, is the ecological role an organism plays within its habitat.

Ecology Definitions

Habitat – The place where an organism lives

Population – A group of organisms belonging to the same species

Community – All the populations of different organisms living and interacting in the same space at the same time

Ecosystem – A community of living organisms and the abiotic factors which affect them

Abiotic – The physical and chemical features of the environment


Biotic – The biological features of the environment (living)

Niche – A species role within it's habitat

Adaptation – A feature that members of a species have to increase their chance of survival

Niche

- Each species occupies a niche in the community. A niche is the role the species plays, and includes the type of food it eats, where it lives, where it reproduces, and its relationships with other species.



Ecological niche

- The ecological niche describes the role an organism plays in its environment. It consists of:
- The species habitat.
- The organisms activity, the period of time it is active.
- The resources it obtains from the environment.
- Eg. A pukeko lives in marsh areas. It feeds of small insects and grasses.



- Several branches of ecology have adopted the concept of the **ecological niche**.

- The ecological niche describes how a species interacts within an ecosystem. The niche of a species depends on both biotic and abiotic factors, which affect the ability of a species to survive and endure.

- **Biotic factors** affecting a species' niche include food availability and predators. **Abiotic factors** affecting ecological niche include temperature, landscape characteristics, soil nutrients, light and other non-living factors.

- An example of an ecological niche is that of the dung beetle. The dung beetle, as its name suggests, consumes dung both in larval and adult form. Dung beetles store dung balls in burrows, and females lay eggs within them.

- This allows hatched larvae immediate access to food. The dung beetle in turn influences the surrounding environment by aerating soil and rereleasing beneficial nutrients. Therefore, the dung beetle performs a unique role in its environment.

- The definition of a niche has changed since it was first introduced. A field biologist named Joseph Grinnell took the basic concept of the niche and further developed it, claiming that a niche distinguished between different species that occupied the same space. In other words, only one species could have a particular niche. He was influenced by species distribution.

Module# 35

Types of Ecological Niches


- Ecologist Charles Elton's definition of niche focused on the role of a species, such as its trophic role. His tenets emphasized more on community similarity and less on competition.
- In 1957, Zoologist G. Evelyn Hutchinson provided a sort of compromise of these trains of thought. Hutchinson described two forms of niche.

- **Fundamental niche**
- **Realized niche**

- The **fundamental niche** focused on the conditions in which a species could exist with no ecological interactions. The **realized niche**, in contrast, considered the population's existence in the presence of interactions, or competition.
- The adoption of the ecological niche concept has allowed ecologists to understand the roles of species in ecosystems

Fundamental niche and realised niche

- The **fundamental niche** of a species includes the total range of environmental conditions that are suitable for existence without the influence of **interspecific competition** or **predation** from other species. The **realized niche** describes that part of the fundamental niche actually occupied by the species.




Importance of Ecological Niches

- Ecologists use the concept of the ecological niche to help understand how communities relate to environmental conditions, fitness, trait evolution and predator-prey interactions in communities.

- This becomes ever more important as climate change affects community ecology.

- Ecological niches allow species to exist in their environment. Under the right conditions, the species will thrive and play a unique role.

- Without the ecological niches, there would be less biodiversity, and the ecosystem would not be in balance.


Yellow bird inhabits upper tree regions
Red bird inhabits middle boughs of tree
Blue bird inhabits trunk and lower branches
Fundamental Niche = Whole Tree
Realised Niche = Specific Elevations

Module# 36

Forest Ecosystem

Forest Ecosystem: A Forest Ecosystem refers to the terrestrial system in which living organisms such as trees, insects, animals and human beings interact with each other.

It is the smaller classification of the ecosystem as a whole, which is the biggest functional unit comprising all the geographical features and living organisms on Earth.




- There are many types of forest ecosystems and the classifications are based on the temperature and rainfall in the area of the particular forest ecosystem under observation.


- A forest ecosystem is the most robust of all as it does not undergo major changes by the effect of weather, forces of nature or human intervention

Types of Forest Ecosystem

Types of Forest Ecosystem

1. Temperate Forest Ecosystem
2. The Tropical Rainforest Ecosystem
3. Boreal or Taiga Forests


Components of a Forest
Wildlife
Microbes
Fungi
Air
Upper-Canopy
Mid-Canopy
Lower-Canopy
Understory
Soil-surface
Upper-soil
Deep-soil
Nutrients
Water
Geologic substrate


Forest
Ecosystem

- There are **Producers** who prepare food for the entire forest ecosystem. Trees and plants are thus called the **primary producers**.

- **Consumers** are the one who cannot produce their own food and thus depend on producers for their food and energy sources.

- Organisms that only eat plants are referred to as primary consumers. For example, herbivores such as deer and rabbits are **primary consumers.**

- **Secondary consumers** feed on herbivores and are called as **Carnivores**. Omnivores are the consumers that feed both on plant and animals.

- Organisms like worms, microbes, fungi, ants, and other bugs are called **Decomposers** as they break down the plant and animal wastes into small particles which ultimately blends with the environment. Human beings are omnivores as they feed on both flora and fauna and thus are a part of this forest ecosystem.

Module# 37 Forest Biome

Forests can be divided into 5 main categories, depending on the types of trees found in them.


1) Coniferous Forest


As the name suggests, the trees found in these forests mainly consists of cone-bearing trees like the spruce, pine, fir, and hemlock. These are found mostly in the northern parts of North America, Asia, and Europe.


2) Deciduous Forest

The trees of these forests are broadleaved and are shed between late summer to early autumn. The leaves are usually green but later acquires yellow, red and orange colours gradually. These forests are found in America, Western and Central Europe, and Northeastern Asia where the winters are cold and summers are warm.

Coniferous Trees




Deciduous Forest



The Four Seasons of Deciduous Forests
Spring
Summer
Fall
Winter
tree layer
(tree stratum)
sapling or small
tree layer
shrub layer
herb layer
ground layer

3) Mixed Forests

- Deciduous and Coniferous trees as both types constitute mixed forests, which are mostly found in mountainous areas. These are found almost every part of the world.

4) Mediterranean Forests

- They are also called scrublands as its temperature is suitable for short oaks and pines to grow. The Mediterranean forest contains a wide variety of wildflowers and insect-eating birds. This forest is also termed as “maquis”.


TEMPERATE
BROADLEAF AND MIXED
FOREST


Common Oak Species of North America
Black oak
(Quercus velutina)
Bur oak
(Quercus macrocarpa)
Laurel oak
(Quercus laurifolia)
Live oak
(Quercus virginiana)
Pin oak
(Quercus palustis)
Water oak
(Quercus nigra)

5) Tropical Rainforest

- These forests are situated in the areas with hot temperatures like South America, Africa, Asia and Australia. Thus they are called Tropical and due to continuous rainfall throughout the year, they are called Rainforests. It is a habitat for various insects, colourful birds and mammals.



Module# 38

Grassland Ecosystem

- Grasslands are areas dominated by grasses. They occupy about 20% of the land on the earth surface.
- Grasslands occur in both in tropical and temperate regions where rainfall is not enough to support the growth of trees.
- Grasslands are found in areas having well-defined hot and dry, warm and rainy seasons.

Components of Grassland Ecosystem

- The different structural components of the grassland ecosystem can be classified as abiotic and biotic components.

Biotic Components

- **Producers –** In grassland, producers are mainly grasses; though, a few herbs & shrubs also contribute to the primary production of biomass.
- **Consumers –** In a grassland, consumers are of three main types:
- **Primary Consumers –** The primary consumers are herbivores feeding directly on grasses. Herbivores such as grazing mammals (e.g., cows, sheep, deer, rabbit, buffaloes, etc), insects (e.g., Dysdercus, Coccinella, Leptocorisa, etc), some termites and millipedes are the primary consumers.


Secondary Consumers – These are carnivores that feed on primary consumers (Herbivores). The animals like foxes, jackals, snakes, frogs, lizards, birds etc., are the carnivores feeding on the herbivores. These are the secondary consumers of the grassland ecosystem.

Tertiary Consumers – These include hawks etc. which feed on secondary consumers.


Decomposers – These include bacteria of death and decay, moulds and fungi (e.g., Mucor, Penicillium, Aspergillus, Rhizopus, etc). These bring the minerals back to the soil to be available to the producers again.

Abiotic Components

- These include the nutrients present in the soil and the aerial environment.
- The elements required by plants are hydrogen, oxygen, nitrogen, phosphorus and sulphur.
- These are supplied by the soil and air in the form of CO_2, water, nitrates, phosphates and sulphates.
- In addition to these, some trace elements are also present in the soil.


Grassland Ecosystem
Grass hopper
Lizard
Plants
Rabbit
Hawk
Mouse
Snake
Atmosphere
Inner Mongolia
Cattle
Sheep
Plant

Grassland Food Web



Flora and Fauna of Grassland Ecosystem

- Grasses are the dominating plants, with scattered drought resistant thorny trees in the tropical grasslands.
- Badgers, fox, ass, zebra, antelope are found grazing on grasslands that support the dairy and leather industries.
- Grasslands also support the large population of rodents, reptiles and insects.

Module# 39

Functions of Grassland Ecosystem

They perform certain functions. These are:

- Energy flow through the food chain
- Nutrient cycling (biogeochemical cycles)
- Ecological succession or ecosystem development
- Homeostasis (or cybernetic) or feedback control mechanisms.
- To increase the fertility of the soil and to regulate the productivity of the ecosystem.
- To reduce the leaching of minerals due to low rainfall.

Economic Importance of Grasslands

- Grasslands are the grazing areas of many rural communities.
- Farmers who keep cattle or goats, as well as shepherds who keep sheep, are highly dependent on grasslands.
- Domestic animals are grazed in the 'common' land of the village.
- Fodder is collected and stored to feed cattle when there is no grass left for them to graze in summer.
- The grass is also used to thatch houses and farm sheds.

- The thorny bushes and branches of the few trees that are seen in grasslands are used as a major source of fuelwood.
- Overgrazing by huge herds of domestic livestock has degraded many grasslands.
- Grasslands have diverse species of insects that pollinate crops.
- There are also predators of these insects such as small mammals like shrews, reptiles like lizards, birds of prey, and amphibia such as frogs and toads.
- All these carnivorous animals help to control insect pests in adjoining agricultural lands.

Classification of Grasslands

- As climate plays an important role in the formation of grasslands, it is generally used as a basis to divide the world's grasslands into two broad categories:

1. Tropical regions

2. Temperate region

Tropical Grasslands

- These occur on either side of the equator and extend to the tropics.
- This vegetation grows in areas of moderate to a low amount of rainfall.
- The grass can grow very tall, about 3 to 4 metres in height.
- Savannah grasslands of Africa are of this type.
- Elephants, zebras, giraffes, deer, leopards are common in tropical grasslands

Savannah grasslands of Africa



Temperate Grasslands

- These are found in the mid latitudinal zones and in the interior part of the continents.
- Usually, the grass here is short and nutritious.
- Wild buffaloes, bison, antelopes are common in the temperate region.




Module# 040 Desert ecosystem


- The desert ecosystem is the driest terrestrial ecosystem in the world, therefore, it has less biodiversity.
- The plants in this desert ecosystem are usually short-lived in nature. They grow, germinate, and perish in a very short time.
- In this type of ecosystem, thorny shrubs are scattered around. The roots of these plants extend far below the ground. Their roots are located at a depth of 1 meter from the surface. As a result, they can easily collect rainwater.
- In many cases, their leaves are transformed into thorns to reduce the cost of water through evaporation.


Component of a desert ecosystem

- The desert ecosystem has less density of plants due to low rainfall. These regions are characterized by less than 255 mm of rainfall and high or low temperatures.
- The evaporation rate of this ecosystem is very high. Dates, cotton, millet, etc. are cultivated in different places depending on the availability of water.
- Like other ecosystems, desert ecosystems are made up of some biotic and abiotic elements


Around One-Third of the
Earth's surface is covered
in Deserts.
Antarctica is the only continent in the
world which is entirely covered by a
Antarctic desert.




Food Chain in Desert Ecosystem
Abiotic Factors
Light
Heat
Water
Minerals
Biotic Factors
Producers
Consumers
Primary
Secondary
Tertiary
Decomposers & Detritivores
Producers
Primary Consumers
Secondary Consumers
Tertiary Consumers
Food Web in Desert Ecosystem
Owl
Golden Jackal
Vulture
Carrion
Desert Plant
Desert Monitor Lizard
Desert Cat
Small Lizard
Desert Rat
Ant
Centipede
Scorpion
Bacteria & other Microbes


Negev Desert
Lut Desert
Chihuahuan Desert
Sahara Desert
Arabian Peninsul
Kalahari Desert
Gibson Deser
Namib Desert

Component of a desert ecosystem

- The abiotic components of desert ecosystems are soil, high temperature, low rainfall, dry climate. These severely affect the desert ecosystem.

1. High temperature or low temperature

- Desert ecosystems usually have either high or have low temperatures. Along with this hot deserts has a difference in day and night temperature. During the day, the temperature reaches a high point due to the heat of the sun. And very cold weather prevails at night. The temperature drops below zero degrees. Organisms that cannot adapt to these temperatures have difficulty surviving in the desert. T

2. Soil

- The nature of the vegetation in that region depends on the type of soil in all ecosystems. Plants grow according to the type of soil. Desert soils do not have any organic matter like nitrogen, phosphorus, etc. which are essential for plant growth. In the desert ecosystem, the soil is dry and rocky, sandy. The soil is not ideal for plant growth therefore, there are no big trees. Cactus is the most common plant in the desert ecosystem.

3. Low rainfall

- Low rainfall is a major feature of desert ecosystems. Rainfall is usually less than 500 mm per year. The plants and animals that live in this desert ecosystem must be able to survive with little water. Cactus develop by storing water in their stems during the rainy season .

4. Sunlight

- Sunlight is an important element of the desert ecosystem. This deeply affects the flora and fauna of the ecosystem.

Biotic components of the desert ecosystem

- The biotic components of desert ecosystems are producers, consumers, and decomposers.

1. Producers

- Grasses, thorny bushes, some species of trees like zizyphus, acacia, etc. are the producers of the desert ecosystem. And many species of succulents like euphorbia, agave, asparagus, etc. also grows in this ecosystem known as xerophytes.

2. Consumers

- Rats, goats, rabbits, camels, and many species of birds are the primary consumers of the desert ecosystem. The secondary consumer in the desert ecosystems are many species of reptiles, insects, owls, etc. And other prey animals are tertiary consumers of the desert ecosystems.

The primary consumers survive by taking the producers as food. And the secondary consumer survives by eating the primary consumer. The tertiary consumer takes primary and secondary consumers as food. In this way, the food chain is maintained in the desert ecosystem .

3. Decomposers

- The number of decomposers in the desert ecosystems is very low. Some fungi, bacteria, etc. are the decomposers of the ecosystem. They decompose dead plants and consumers in desert ecosystems into simple chemicals.

Module# 41

Type of desert ecosystem

Desert ecosystems are mainly divided into four types

1. Hot and dry
2. semi-arid
3. coastal
4. cold desert

Hot and Dry Desert Ecosystem

These kinds of the desert ecosystem have hot and dry climatic conditions through the air and have very low annual rainfall.

The hot desert ecosystem is basically found in Central America, South Asia, North America, Africa, Australia etc. There are extreme variations in temperature and soil is rough and harsh.

Semi-arid desert ecosystem

- This desert ecosystem is quite similar to the Hot and Dry desert ecosystem. This kind of ecosystem has hard rocks, stable ground, less sand dunes.
- Temperature is not as extreme as a hot and dry desert ecosystem. Great Basin is an example of Semi-arid desert ecosystem.
- It receives a lot of rain as compared to the normal desert's ecosystem.

Coastal desert ecosystem


- The Atacama Desert in Chile and Namib in Africa are a good example of Coastal desert ecosystem.
- Such desert ecosystems are found near the coastal lines of big water bodies like oceans and seas and are generally affected by the ocean currents.
- Winter fogs are common here. They are more hospitable than other desert ecosystem and therefore they have a more flora and fauna than others.


Cold desert ecosystem


- This desert ecosystem comprises of abundant rainfall throughout the winters and less in summers and generally has chilling winters with snowfall.
- The summers are short, moderately hot and moist here. These are usually covered with snow dunes. Such desert ecosystem can be found in Greenland, Antarctica, and Nearctic realm.

Types of deserts

Desert biomes can be classified according to several characteristics. There are four major types of deserts:










Hot, Dry Desert


Semi Arid Desert


Coastal Desert


Cold Desert

Importance of the desert ecosystem

- Like other ecosystems, desert ecosystems occupy a significant place in the world. This ecosystem plays an important role in maintaining the balance of the earth despite the low rainfall and dry climate.

- This ecosystem is a habitat for many species of plants and animals. These animals and plants balance by establishing relationships with each other.

- Different types of bacteria are present in the sand of the desert ecosystem. These bacteria can store carbon dioxide. And acts as a carbon sink, preventing carbon dioxide from entering the atmosphere.
- These ecosystems are often the source of mineral resources, natural gas.

- Merged BY BAQIR HUSSAIN
- BS ZOOLOGY 03336361244

Module# 42

Tundra Ecosystem

- In this type of ecosystem, they are free from trees as it is covered by snow for most of the year.
- These ecosystems are mainly present in those areas where there is a cold climate and very limited or no rainfall.
- One of the common regions where tundra ecosystems are found is the polar regions.





There are two types of tundra ecosystems, they are discussed below

1.Arctic:

- This tundra is mainly located in the northern hemisphere, encircling the north pole and extending towards the south.
- This tundra is known for its cold and desert-like conditions. In this region temperatures during winters are about 3-120 C during day time and -340 C during night time.
- This temperature enables biomes to sustain in this condition. Approx 1,700 kinds of plants are found in the arctic region with good adapting capability.
- Some of the common plants are low shrubs, sedges, reindeer mosses, liverworts, and grasses; 400 varieties of flowers example: crustose and foliose lichen.

2. Alpine:

1. They are mainly located in mountainous areas throughout the world and in this area trees can not be grown. In this area, the temperature is below the freezing point during nighttime.
2. The soil of this region is well-drained. Some of the plants which are present in this tundra region are tussock, dwarf trees, small-leafed shrubs, heaths, etc. Animals in this region have high adapting capabilities.
3. Examples of mammals that are mainly found for this reason are pikas, marmots, mountain goats, sheep, elk : Birds: grouse like birds.

Distribution of Tundra Ecosystem

- 10 percent of the total earth surface is covered by tundra ecosystems. From the southern boundary, there is a start of the arctic tundra which goes through the northern edge of the coniferous forest belt, whereas the alpine tundra is mainly found in the Northern Hemisphere.

Module# 43 Features of Tundra Ecosystem

Some of the common and well-known features of the tundra ecosystem are discussed below:

- The Tundra ecosystem mainly has a long cold winter and a very short hot summer, that is why they are known for winters.

- This ecosystem mainly has winds with high humidity and less amount of rainfall.

- Some of the common nutrients present in this reason are nitrogen and phosphorus. Production of phosphorus is mainly by a precipitation process, whereas a biochemical process produces nitrogen.

- At night time temperature in this region is below the freezing point whereas in day time it is a little bit normal which provides favourable conditions for plant growth.

Plants Found in Tundra Region


Some of the known plants which are found in this region are:

- Cottongrass
- Dwarf heath
- Sedge
- Rushes
- Dryland sedges
- Sphagnum moss
- Birches

Cottongrass



Dwarf heath



Animals Found in Tundra Region

- In this region, all kinds of animals are found like omnivores, carnivores, herbivores.

- Some of the common examples of animals found in this region are:

Wolf, foxes, hare, collembolans, muskox, reindeer, marmots, mountain goats, sheep, elk and other different types of worms, flies, butterflies, wild insects.

Wolf



Collembolans



Module# 44

Agricultural Ecosystems

- Agricultural ecosystems comprise 30 % of the Earth's surface.
- They provide ecosystem services including food and non-food biomass production, carbon sequestration, nutrient cycling, and climate regulation.

Concept of Agroecosystem

Agriculture + Ecosystem = Agroecosystem





- Agricultural ecosystems serve as important habitats for many wild plant and animal species.

- Management practices used in agricultural ecosystems determine the state of the global environment.

- MERGED BY BAQIR HUSSAIN
- BS ZOOLOGY 03336361244





What is an agricultural ecosystem

- Made up of largely domesticated animals & plants used to produce food for humans.
- Considerable energy losses occur at each trophic level of a food chain.
- Humans are usually at the 3rd, or even 4th, trophic level of a food chain.
- Agriculture tries to ensure that as much as possible of the Sun's energy is transferred to humans.
- Agriculture channels the energy flowing through a food web into the human food chain and away from other food chains. This increases the productivity of the human food chain.



Components of Agro ecosystem

- **Primary producer:** Crops and weeds of the field are the primary producer of agro ecosystem. e.g. In a Rice field, there are many producer like durba, mutha, syma etc also present with rice.

- **Consumer:** Among consumer grasshoppers, aphids, bugs, ants, rats, birds, man etc are macro consumer and frog, snake, hack are micro consumer.

Properties of Agro ecosystem

1. **Productivity-**

It is net increment of values products per unit resources (land, labour, energy, capital) and is commonly measured as annual yield /hectare

2. Stability-

It is the degree to which, productivity remain const ant, inspite of normal small scale fluctuation in environmental variables such as climate or in the economic condition in market.

3. Sustainability- It is defined as the ability of the system to maintain its productivity when subject to stress or perturbation. A stress is defined as regular, sometimes continues, relatively small and predictable disturbance. e.g. Affect of growing soil salinity. A perturbation by contrast is an irregular, in frequent relatively long and unpredictable disturbance such as drought or flood or a new pest.

4. Equitability- It is a measure of how evenly the produce of Agro ecosystem is distributed among its human beneficial. The more equitable the system, the more evenly are the products to fed shared among the population of the farm, village, regions or nation.

Module# 45

Marine Ecosystem

- The marine ecosystem covers more than 70% of the Earth's surface.
- The marine ecosystem provides a wide range of habitat for flora and fauna of the marine aquatic system that ranges from seashores to the deep seabed.
- It is a home for millions of species.



Marine Ecosystem

- The marine ecosystem plays an important role in the protection of the environment.
- For example – like land plants, the water plants also help in minimizing the carbon level in the atmosphere.
- Aquatic plants absorb carbon dioxide from the air, and it releases oxygen back in the atmosphere.

- The marine ecosystem is a wonderful combination of biotic and abiotic components supporting each other for survival.

- It is an exciting part of the ecology that will grab your attention once you start knowing it.

- You can go through this article to get a close-up experience of the marine ecosystem.


MARINE ECOSYSTEM
Sun
Ocean acidification
Seabirds
Estuaries
Farm land run off
Crabs
Red cod
Dolphins
Commercial fishing
Phytoplankton
Fish
Bivalve molluscs
Arrow squid
Cockles
Zooplankton
Bryozoans
Bryozoan thickets
Seaweed
Sea stars
Bacteria
© Copyright. The University of Waikato. All rights reserved. I www.sciencelearn.org.nz

Module# 46 Characteristics of Marine Ecosystem

The temperature of the marine ecosystem

- Temperature is one of the important characteristics of an ecosystem.
- The average temperature of oceanic water is 30 degrees Fahrenheit.
- However, this temperature may vary (maybe colder or warmer) depending on its location on the Earth's surface.

- The marine ecosystem near the equator experience comparative warm water temperature, whereas the one which is close to the poles, have cold water temperature.

- The temperature of the marine ecosystem plays an essential role in the lives of living creatures underwater.

Water in the marine ecosystem

- We know the earth as a blue planet. Well, the most credit goes to the water bodies present on the Earth's surface.

- Almost ¾ part of Earth's surface is occupied by water, out of which approximately 90% is covered by marine water bodies like seas and oceans.

- We call it marine water due to its saline nature i.e., the saltiness of the water.

- Marine water contains approximately 96.5% pure water, and the rest 3.5% are the dissolved compounds such as chlorine, sodium, magnesium, sulfur, calcium, potassium, etc.

- However, the salinity of the marine ecosystem not only depends on these dissolved compounds but on other factors as well, like the latitude, erosion, atmospheric activity, depth, volcanic activity, biological activity, etc.

Importance of sunlight

- Sunlight has always been an important factor for an ecosystem. All the living creatures depend on sunlight directly or indirectly to fulfill their need for the food.

- The plants of the marine ecosystem are known as the primary producer as they make food using the sunlight. The food chain starts from these primary producers and goes successful on top levels predators. Sunlight acts as a primary component to maintain proper balance in the marine ecosystem food chain.

Biodiversity in the marine ecosystem


- The marine ecosystem supports great biodiversity as compared to other ecosystems of the planet. The species of the marine ecosystem have adapted the aquatic culture for its survival. For example – the fishes have gills that help them to breathe in water.


- Many plant and animal species live together in the marine ecosystem. Some of the common aquatic species are-

- Plants like kelp, phytoplankton, seaweeds, seagrasses, mangroves, etc.

- Fishes like sharks, tuna, grouper, eels, seahorse, gars, swordfish, clownfish, stingray, flatfish, rockfish, sunfish mola, etc.

- Mammals like seals, dolphins, blue whales, walruses, manatees, otters, etc.




Oceanic Ecosystem
Sunlight
Surface
(euphotic zone)
phytoplankton
Nutrient
zooplankton
eat
eat
nekton
dead body/excrement
Upwelling
(only at specific
upwelling sites)
SINK
dead body/excrement
Deep Sea
digest
eat
eat
Nutrient
bacteria
benthos

Module# 47

Types of Marine Ecosystems

Open Marine Ecosystem

- Whenever you think of the marine ecosystem, the first picture that comes to your mind is the open water surface. This open water surface is known as the marine ecosystem; This is the upper layer of the ocean where sun rays reach quite easily.

- The open marine ecosystem extends up to 150 meters from the ocean surface. The open marine ecosystem provides habitat to various sea creatures such as plankton, algae, whales, jellyfish, etc.



Deep-Sea Marine Ecosystem

- The ecosystem deep inside the oceans at its floor is known as the deep sea marine ecosystem. Deep-sea marine ecosystem inhabiting various animal species in the seabed up to 1000 meters depth underwater.
- Sunlight is one of the big challenges to penetrate at the seafloor; however, the species have adapted the marine environment of this region. A huge range of animal species is found in deep-sea marine ecosystems such as squids, fishes, elephant seals, sperm whales, crabs, worms, some sharks, etc.



Coral Reef Marine Ecosystem

- Coral reefs are special type of marine ecosystem found in the seafloor. This ecosystem is mostly found in tropical waters and is quite a productive ecosystem found on the Earth. The coral reefs are the skeleton made up of limestone i.e., calcium carbonate.

- The coral reef ecosystem provides food and shelter to almost ¼ of marine water species. Coral reefs attract exotic color fishes like sponges, snails, seahorses, sometimes large animals like sharks & dolphins, etc.


CORAL REEFS

Saltwater Wetland Marine Ecosystem

- The coastal areas of oceans and seas are known as the saltwater wetland ecosystem. The saltwater wetland marine ecosystem is again classified into two types – saltwater swamps and salt marshes.
- Saltwater swamps are regions dominated with trees, whereas salt marshes are covered with grasses. Commonly found water species of saltwater wetland ecosystem are amphibians, reptiles, some migratory birds, shellfishes, few fishes, etc.



Module# 48

Estuary Marine Ecosystem

- The area around the river mouth where it merges with marine water is usually termed as an estuary marine ecosystem. The salinity in this ecosystem varies with the tides. The estuary ecosystem does not support a variety of species.
- Mostly the species of the neighbouring ecosystem are found in the Estuary marine ecosystem. This ecosystem also plays an essential role in nurseries for various kinds of fishes, shrimps, etc.

Sandy Beach Ecosystem

- Sandy's ecosystem is quite poor in terms of biodiversity as compared to different marine ecosystems. The environment in the sandy beach ecosystem varies a lot. However, the species that are found in the sandy beach ecosystem are very much adapted to a constantly variable environment.

- The species that found a home in the sandy beach ecosystem mostly includes sea turtle, sea lion, seal, algae, plankton, invertebrates like snails, crabs, clams, etc. birds like plovers, willets, gulls, terns, ruddy turnstone, curlews, etc.


Sandy Beach
Ecosystem

Polar Marine Ecosystem

- As the climate of Polar Regions is extremely cold; thus, the temperature of this type of marine ecosystem is also too cold. The species found in the polar marine ecosystem have adapted to the adverse climatic conditions of this region.
- The most common species found in the polar marine ecosystem includes planktons, algae, birds like penguins, polar bears, seals, walruses, etc.



Rocky Marine Ecosystem

- The rock shores, rock cliffs, boulders, tide pools, etc. combine to make rocky marine ecosystems. The biodiversity of rocky marine ecosystem generally have species like lichens, birds, invertebrate (lobsters, urchins, barnacles, sea stars, sea squirts, seal, etc.

- The marine ecosystem is a unique ecosystem that supports various plant and animal lives. All these species set an excellent example of a great food chain as compared to other ecosystems.


- The sad part is that we are spoiling such an amazing ecosystem by polluting it through disposing of unwanted wastes. We are disposing of our wastes into the oceans and seas that are affecting the marine ecology big time.



Module# 49

Estuarine Ecosystem

- An estuary is a partially enclosed body of coastal water where freshwater from rivers and streams mixes with salt water from the oceans.
- Estuaries and their surrounding lands are thought about locations of transition from land to sea.
- Although influenced by tides, they are likewise secured from the full hits of ocean waves, winds, and storms by landforms such as barrier islands or peninsulas.


Large island
High-elevation lakes
Montane reaches
Slopes reaches
Floodplain habitats
Low-elevation lakes
Lowland reaches
Tidal fresh waters
Estuaries


Small, steep island
Montane reaches
Slopes reaches
Lowland reaches

- Estuarine ecosystems are among the most efficient in the world, developing more organic matter each year than similar-sized locations of the forest, grassland, or farming land.

- Protected estuary waters also home to unique communities of plants and animals that have actually adapted to life on the edge of the sea.



- It is the part in which the watercourses mix into the ocean circulation. They are typically located in areas where the tides are large with beaches to the sides, that when they disappear, display their flora.

Module# 050

Freshwater and Wetlands ecosystems


Freshwater ecosystems


Freshwater ecosystems are interactive systems within which biotic species and their growth and adaptation, and associated biological productivity, nutrient cycling, and energy flows among inland aquatic microbial, plant, and animal communities, are integrated with their environment.

These inland waters include lakes, reservoirs, rivers, streams, and wetlands. The subject discipline, termed limnology, evaluates how these interactive relationships are regulated by natural and modified variations in their physical, chemical, and biotic environments.

- The standing water (lentic) lake or reservoir ecosystem is a system that is intimately coupled with the land surrounding it in its drainage area via chemical components transported to the lake by groundwater or running surface waters (lotic) of streams and rivers.

Classification of Ecosystem




Freshwater Ecosystem
Lotic
Lentic
Wet Land
Rivers
Streams
Springs etc.
Ponds
Lake
Pools etc.
Marshes
Swamps

Module# 51


Freshwater Habitat

- Any waterbody having a low salt concentration usually less than 1% is called a freshwater body. Animals as well as plants in these freshwaters are adapted to the low salt concentrations and are unable to survive high concentration of salt present in oceans.

- The study of relationship between living organisms and the fresh water environment constitutes the freshwater ecology. The study of physical, chemical, geological and biological aspects of freshwaters is called **limnology**.

Types of Freshwater Habitat

- Freshwater habitats are studied under three major categories for convenience:
  - Standing water or **Lentic** (*lenis,* calm) Habitat: e.g., Lake, pond, swamps etc.
  - Running water or **Lotic** (*lotus,* washed) Habitat: e.g., spring, streams, rivers etc.
  - **Wet Lands**: Marshes and swamps.


Freshwater Ecosystem
Lotic
Lentic
Wet Land
Rivers
Streams
Springs etc.
Ponds
Lake
Pools etc.
Marshes
Swamps

Fresh Water Lake (Lentic Habitat)



River (Lotic Habitat)



- Freshwater habitats occupy a relatively small portion of earth‘s surface as compared to marine and terrestrial habitats. However, they play a far more important role for Humans because of the following reasons:

1. They are the cheapest and the most convenient source of water for industrial as well as domestic needs.

2. They are the —bottle-neck of the hydrological cycle.

3. They are a convenient and cheapest sinks for waste disposal systems.

Module# 052

Limiting Factors of Freshwater Ecosystems

Temperature

The unique properties of water maintain the temperature fluctuation in the water bodies. The variation is small, decreasing gradually from the surface to the bottom of the water body. Some of these thermal properties are:

- High specific heat: To change the temperature of water, a relatively very large amount of heat is required compared to other substance of same mass. One gram-calorie (gcal) of heat is required to raise one milliliter (ml) of water by one degree centigrade (between 150C to 160C).
- High latent heat of fusion: 80 calories are required to change one gram of ice into water without changing its temperature.

- **Highest latent heat of evaporation: 536** calories per gram are absorbed during evaporation which occurs more or less continually from vegetation, water, and ice surfaces. Major portion of the solar energy dissipates in the evaporation of water from the ecosystems of the world. This flow of energy moderates the climate which makes life possible on earth.
- Highest density at 40 C helps in expanding water from both above and below the surface and making it lighter. This property prevents the water bodies like lake from freezing.

Transparency

- Transparency is an important parameter to measure the penetration of light. Penetration of light is limited by suspended materials, restricting the photosynthetic zone wherever aquatic habitats have appreciable depth.
- Turbidity is caused either by clay and split particles or by the excretory products of the living organisms. Turbidity caused due to suspended clay and split particles is considered as an important limiting factor.
- On the other hand turbidity caused by living organism is the measurement of productivity of the water body.

- Turbidity has a direct impact on the penetration of light. Turbid water restricts the amount of light reaching the bed.
- Transparency can be measured with an instrument called **Secchi disk**.
- In the most widely studied Wisconsin lakes, Secchi disk transparency represents the zone of light penetration down to about 5% of the solar radiation reaching the surface. The 5% level marks the lowest limit of major photosynthetic zone.

Turbid and Clear freshwater Lake in Maryland



Use of Secchi Disk to measure the transparency of lake



Module# 53

Limiting Factors of Freshwater Ecosystems

Current

It plays important role in estimating the distribution of salts, vital gases and small organisms. It serves as an important limiting factor for the lotic systems especially for streams.

Concentration of respiratory gases

- Oxygen and the carbon dioxide concentration are often limiting in freshwater environment. Dissolved oxygen content (DO) and Biological oxygen demand (BOD) are important determinants for measuring the biological and physical health of a waterbody.

- These are one of the most intensively studied physical factors. Dissolved oxygen reflects the productivity. Biochemical Oxygen Demand or Biological Oxygen Demand (BOD), measures the amount of dissolved oxygen (DO) which is utilized by the aerobic microorganisms while decomposing the organic matter in waterbody, at certain temperature and over a specific period of time.

- BOD affects the amount of dissolved oxygen (DO) in streams and rivers. The oxygen consumption rate is affected by many variables: viz. temperature, pH, the presence of certain kinds of microorganisms, and the type of organic and inorganic material in the water.
- The greater the BOD, the more rapidly oxygen is depleted in the stream. Less BOD indicates less availability of oxygen to higher forms of aquatic life. A high BOD mean low dissolved oxygen (DO). As a result of such conditions aquatic organisms become stressed, suffocate, and die.

Isolation due to land barriers

- Distribution and number of organisms in freshwater habitats is also influenced by the ability of the organisms to disperse in their surroundings. Isolation due to the presence of land barriers does not allow free movement of the organism to become established in places otherwise favorable.
- Example: Fishes of the streams, which are separated only by a few miles from land but isolated by water, may have their niches occupied by different species. On the other hand, most small organisms-algae, crustacea, protozoa and bacteria, etc. have a wide range of dispersal. Thus, same kinds of water flea are observed (Daphnia, for example) in a pond in the United States as in England.


Freshwater
Ecosystems

Module# 54

Lotic Ecosystems

Merged by BAQIR HUSSAIN BS ZOOLOGY 03336361244

The Lotic Aquatic System

- Lotic aquatic systems are the fresh water bodies with flowing waters.
- Water in lotic system is in a state of constant motion.
- Rivers and streams are the most common examples of such systems.
- The basic function of lotic water bodies is assumed to carry the excess of rain water back to the sea.
- It is estimated that the amount water that these lotic bodies carry to the oceans amounts to approximately 25 cm of rains that can be distributed evenly on the land surface.

Lotic Systems (Running Water)







- **Rivers, streams, creeks**

Lotic Ecosystem

- The ecosystem of a river, stream, or spring.
- Lotic refers to flowing water.
- Included in the lotic environment are the biotic interactions (plants, animals and micro-organisms) as well as the abiotic interactions (physical and chemical).



Lotic Ecosystem

- A natural system provides a variety of habitats for many species of aquatic plants and animals.
- The instream habitats provide areas for feeding, resting, and reproduction and generally support a great diversity of organisms.



Examples include:

- Rivers
- Streams
- Brooks
- Creeks
- Human made channels



Typically, the lotic aquatic systems have the following properties:

- The flow of water in continuous and unidirectional in a lotic system. Water mostly flows in narrow channels that are less deep than the lakes. Water from diverse sources can pool into a lotic body. The bottoms of such water bodies continuously receive materials from the biomass of surface water.
- The volume of water changes rapidly which results in change in the velocity of water currents. There is also a wide range of fluctuation in the water level in a lotic system.
- Water in these systems act as effective agent of transportation, transfer, and dilution. They continuously erode materials along their channels and deposit them to other places.

- There is no vertical stratification and is a thorough mixing of contents.
- The physico-chemical statistics of water quality is in a state of perpetual change. The environment of a lotic aquatic system is relatively unstable as compared to the lentic system.
- There is a rich supply of oxygen which is derived from air above and also the photosynthetic outputs of the autotrophs. Oxygen is evenly distributed across the water body and oxygen depletion is very rare in unpolluted lotic systems.
- Turbidity limits the light penetration to the deeper zones. Very less amount of carbon dioxide present in these waters. Such conditions limit productivity. Therefore, algae bloom and growth of other organisms is rarely observed in moving waters.


Module# 55

lentic ecosystem

lentic ecosystem

- A **lentic ecosystem** entails a body of standing water, ranging from ditches, seeps, ponds, seasonal pools and lakes.
- Deeper waters, such as lakes, may have layers of ecosystems, influenced by light.
- Ponds, due to their having more light penetration, are able to support a diverse range of water plants.

- Animals and plants in a lentic system behave and adapt differently because there are various zones of every lentic system with different abiotic conditions.
- The diagram below illustrates the zones of a lentic system.



Littoral Zone:

- This zone forms the banks or boundary of the water. There is usually warmer and shallow water, with lots of plants such as reeds and cattails. Many of the plants are rooted in the muddy floor and their leaves form a good hideout for insects and fish. There is a lot of fish, frogs, reptiles, and insects in this zone.

Limnetic Zone:

- It is the zone after the Littoral zone, extending a few meters into the water body. Fish swim freely here, with some plants floating on the water. There is enough sunlight here, and algae and plankton usually thrive here.

Profundal Zone:

- This zone extends beyond the Limnetic zone and is a deeper region of the water. Here, there is usually very little or no sunlight. It is cooler. Living things in this region depend on drifting organic matter for food.

Benthic Zone:

- This is the deepest region of the water body and farthest from the banks. There is no sunlight in this region, and therefore very little or no photosynthesis occurs here. Organisms that live here include worms, bacteria, decomposers, and other scavengers.


Lotic
Lentic

Lentic Ecosystem

- Includes lakes, ponds, dammed rivers, and other static freshwater habitats






Aquatic Ecosystems - Lentic

Lake





Pond



Bog

Swamp



Marsh



Module# 56

Wetlands Ecosystem

Wetlands Ecosystem

- Wetlands are some of the most valuable ecosystems on Earth. They act like giant sponges or reservoirs.
- During heavy rains, wetlands absorb excess water, limiting the effects of flooding. Wetlands also protect coastal areas from storm surges that can wash away fragile beaches and coastal communities.
- Saltwater swamps and tidal salt marshes help secure coastal soil and sand.





Wetlands
tamarack tree
nesting osprey
cedar tree
black-toed woodpecker
yellow spatterdock
viceroy butterfly
dragonfly
red-winged blackbird
jack-in-the-pulpit
smoky shrew
pickerelweed
sundew
blueberries
blue-winged teal
mink
spring peeper
cattails
pitcher plant
©Sheri Amsel
www.exploringnature.org



- Wetland ecosystems also act as water-treatment facilities.
- The plants, fungi, and algae of a wetland filter wastes and purify water.
- Nitrates and other runoff chemicals often wash into wetlands from urban areas and farms.
- Organisms there absorb the harmful chemicals. Pollutants not absorbed by plants slowly sink to the bottom, where they are buried in sand and other sediment.

- Wetlands, especially marshes and swamps, are home to a wide variety of plant and animal life.
- Some animals, such as shrimp, live in tidal marshes. Many marine fishes, such as striped bass, enter coastal wetlands to spawn.
- Bass swim from the ocean and into salt marshes to lay their eggs. When the eggs hatch, the young bass find plenty of food and some protection in the grasses or tree roots.
- Oysters live in huge reefs in salt marshes. All of these wetlands are home to economically valuable fisheries.


Benefits of Wetlands
Wetlands create tremendous recreational, research, and tourism opportunities.
Wetlands act as sponges, temporarily storing flood waters and releasing them slowly, thus reducing flood damage.
Wetlands are considered "nature's nurseries" by providing critical habitat for fish, wildlife, and waterfowl.
Wetlands act as "nature's kidneys" by removing pollutants, such as nutrients and sediments, from water flowing through them.
Wetlands provide protection from storms and ice by absorbing wave energy and buffering shorelines against erosion.
Illustration: Tip of the Mitt Watershed Council

Benefits of wetland ecosystems



Controls Flood



Provides Clean Source Of Water



Improves Air Quality



Helps sustain healthy livelihood



Filters out toxins & adds in nutrition



Habitats to hundreds of species, including the endangered ones


What makes a freshwater wetland?
Rivers and Streams
Bogs
Ponds
Vernal Pools
Wet Meadows
Marshes
Swamps

Module# 57 Population Ecology

- Population ecology is the study of populations in relation to the environment
- Includes environmental influences on population density and distribution, age structure, and variations in population size

Definition of a Population

- A population is the total number of individuals of a given biological species found in one place at one time.

- In practice, ecologists often deal with density—numbers per unit area for land organisms and numbers per unit volume in aquatic systems—or even weight rather than raw numbers.



Population Ecology





- Population ecology is the study of populations in relation to environment
  - Including environmental influences on population density and distribution, age structure, and variations in population size





Module# 58


Characteristics of Population Ecology

- All species occur in groups called populations. There are certain characteristics that all populations have, such as population density, spatial distribution, and growth rate. These characteristics are used to classify all populations of organisms, including bacteria, animals, and plants.

Characteristics of a Population

- **Population** - individuals inhabiting the same area at the same time

- **Population Dynamics:** Population change due to
  - **Population Size** - number of individuals
  - **Population Density** - population size in a certain space at a given time
  - **Population Dispersion** - spatial pattern in habitat
  - **Age Structure** - proportion of individuals in each age group in population


Characteristics
That Determine a Population
Distribution
Population
Age
Density
Sex Ration

Population Ecology: Population Characteristics

Population Characteristics

1. **Population Density:**
   - The number of organisms per unit area
2. **Spatial Distribution:**
   - Dispersion: The pattern of spacing a population within an area
     - 3 main types of dispersion
       - Clumped
       - Uniform
       - Random
   - The primary cause of dispersion is resource availability

Module#59 Population density

Population density one characteristic of a population is its population density, which is the number of organisms per unit area.

For example, the population density of cattle egrets, shown with the water buffalo in Figure, is greater near the buffalo than farther away. Near the water buffalo, there might be three birds per square meter. Fifty meters from the water buffalo, the density of birds might be zero



Spatial distribution

- Another characteristic of a population is called dispersion—the pattern of spacing of a population within an area. Figure shows the three main types of dispersion—uniform, clumped groups, and random. Black bears are typically dispersed in a uniform arrangement. American bison are dispersed in clumped groups or herds. White-tailed deer are dispersed randomly with unpredictable spacing. One of the primary factors in the pattern of dispersion for all organisms is the availability of resources such as foo

Population density describes how many individual organisms live in a given area. Dispersion describes how the individuals are spaced within that area. Population range describes a species' distribution.


Black Bear



Density: one bear per several hundred square kilometers

Dispersion: American black bear males usually are dispersed uniformly within territories as large as several hundred square kilometers. Females have smaller territories that overlap those of males.

Black Bear Distribution (in purple)




American Bison



Density: four bison/km^2 in Northern Yellowstone in 2000

Dispersion: American bison are found in clumped groups called herds.

Bison Distribution (historic range prior to 1865 in orange)




White-tailed Deer



Density: 10 deer/km^2 in some areas of the northeastern United States

Dispersion: White-tailed deer are dispersed randomly throughout appropriate habitats.

White-tailed Deer Distribution (in blue)



- Uniform dispersion refers to organisms that live in a specific territory. One example would be penguins. Penguins live in territories, and within those territories the birds space themselves out relatively uniformly.
- Random dispersion refers to the spread of individuals such as wind-dispersed seeds, which fall randomly after traveling.

- Clustered or clumped dispersion refers to a straight drop of seeds to the ground, rather than being carried, or to groups of animals living together, such as herds or schools. Schools of fish exhibit this manner of dispersion.

Module# 60 Population ranges

Population ranges No population, not even the human population, occupies all habitats in the biosphere. Some species, such as the iiwi (EE ee wee) shown in Figure, have a very limited population range. This songbird is found only on some of the islands of Hawaii.



The iiwi lives only on some of the Hawaiian islands. The peregrine falcon is found worldwide.

- Other species, such as the peregrine falcon shown in Figure, have a vast distribution. Peregrine falcons are found on all continents except Antarctica.

- Merged by BAQIR HUSSAIN
- BS ZOOLOGY 03336361244



Peregrine falcon

- A species might not be able to expand its population range because it cannot survive the abiotic conditions found in the expanded region.
- A change in temperature range, humidity level, annual rainfall, or sunlight might make a new geographic area uninhabitable for the species.
- In addition, biotic factors, such as predators, competitors, and parasites, present threats that might make the new location difficult for survival