Virtual University

V OXFORD U BOOK SHOP

VU Book Shop

کتابوں کی فکر ختم، پڑھائی پر توجہ بڑھائیں!



- Verified Printed Handouts
- Fast Delivery All Over Pakistan
- Updated for Final Term
- One Click WhatsApp Order

Order Now: 0344-2023473

Visit: www.vubookshop.net

Your Study Partner for Every Semester!

ورچوئل یونیورسٹی کی کتابیں گھر بیٹھے آرڈر کریں

ONE CLICK WHATSAPP ORDER

03442023473

PSY516 - Statistics in Psychology (Quiz 4)

Quiz Start Time: 10:57 PM, 18 January 2024

Question # 1 of 10 (Start time: 10:57:37 PM, 18 January 2024)

Total Marks: 1

Which of the following statistical tests will be considered, If a researcher wants to know whether exam performance differs based on test anxiety levels among students, dividing students into three independent groups (e.g., low, medium and high-stressed students)?

Select the correct option

Reload Math Equations

- Z test
- F test
- Anova
- T test

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 10:59:21 PM, 18 January 2024)

If two variables are moving in the same direction, it is called as_____ correlation.

Select the correct option

○	Negative
○	Non-linear
○	Linear
○	Positive ✓

PSY516 - Statistics in Psychology (Quiz 4)

Question # 3 of 10 (Start time: 11:00:35 PM, 18 January 2024)

Pearson's Correlation Analysis is also known as:

Select the correct option

○	Spearman's Correlation Analysis
○	Kruskal-Wallis Correlational Analysis
○	Pearson's Product-Moment ✓
○	Chi-Squared Product-Moment

PSY516 - Statistics in Psychology (Quiz 4)

Question # 5 of 10 (Start time: 11:03:15 PM, 18 January 2024)

Two-way ANOVA means groups are defined by two _____ variables.

Select the correct option

○	Dependent
○	Partial
○	Extraneous
○	Independent



PSY516 - Statistics in Psychology (Quiz 4)

Question # 7 of 10 (Start time: 11:05:50 PM, 18 January 2024)

Which of the following is an advantage of ANOVA?

Select the correct option

○	It cannot tolerate data that is non-normal
○	It performs all three comparisons simultaneously in one hypothesis test
○	It decides if the population mean is equal to a specific value or not
○	All of the above

PSY516 - Statistics in Psychology (Quiz 4)

Question # 10 of 10 (Start time: 11:08:56 PM, 18 January 2024)

What does ANOVA calculate?

Select the correct option

○	Z-scores
○	T-scores
○	F ratios ✓
○	R ratios

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 11:31 PM, 18

Question # 1 of 10 (Start time: 11:31:13 PM, 18 January 2024)

In ANOVA, which of the following tests is preferably used when the assumption of the homogeneity of variance is violated?

Select the correct option

Reload Ma

- Shapiro-Wilk
- Welch and Brown-Forsythe
- Kolmogorov-Smirnov tests
- Levene's test

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 11:32:53 PM, 18 January 2024)

What would be the best interpretation of Ho: ρ = 0?

Select the correct option

○	There is negative correlation
○	There is no population correlation ✓
○	There is real correlation
○	There is covariance

PSY516 - Statistics in Psychology (Quiz 4)

Question # 3 of 10 (Start time: 11:33:59 PM, 18 January 2024)

What would be accurate interpretation of the following hypothesis H_1: $\rho \neq 0$?

Select the correct option

○	There is no population correlation
○	There is negative correlation
○	There is covariance
○	There is real correlation ✓

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start

Question # 4 of 10 (Start time: 11:35:09 PM, 18 January 2024)

In one-way ANOVA, _____ is used to determine which of given groups differ from each other.

Select the correct option

- F test
- Post hoc test
- Levene's test
- T test

PSY516 - Statistics in Psychology (Quiz 4)

Question # 8 of 10 (Start time: 11:41:21 PM, 18 January 2024)

In ANOVA, difference exists between the treatment conditions is termed as:

Select the correct option

○	Treatment variance
○	No treatment variance
○	Between-treatments variance ✓
○	Within treatments variance

PSY516 - Statistics in Psychology (Quiz 4)

Quiz Start Time: 11:52 PM, 18 January 2024

Question # 1 of 10 (Start time: 11:52:27 PM, 18 January 2024)

Total Marks: 1

ANOVA or F-test is also robust to moderate departures from normality when sample sizes are reasonably _____ and are equal.

Select the correct option

Reload Math Equations

- Large
- Small
- Skewed
- Moderate

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 11

Question # 9 of 10 (Start time: 11:42:34 PM, 18 January 2024)

Which of the following statement is true for the Pearson correlation?

Select the correct option

○	The relationship between two variables while controlling the influence of a third variable by holding it constant.
○	All of the above ✓
○	It measures the degree and the direction of the linear relationship between two variables.
○	It measures the degree and the direction of the non-linear relationship between two or more variables.

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 11:53:59 PM, 18 January 2024)

What are the two types of variance which can occur in your data?

Select the correct option

○	Repeated and extraneous groups
○	Independent and confounding groups
○	Experimenter and participant groups
○	Between or within groups



PSY516 - Statistics in Psychology (Quiz 4)

Question # 4 of 10 (Start time: 11:56:39 PM, 18 January 2024)

In which of the following cases could you use a paired-samples t-test?

Select the correct option

○	When comparing two separate groups
○	When comparing men and women's scores
○	When comparing the same participants performance before and after training ✓
○	When assessing three groups or more

PSY516 - Statistics in Psychology (Quiz 4)

Question # 5 of 10 (Start time: 11:57:58 PM, 18 January 2024)

For ANOVA, independent variable is always_____ in nature.

Select the correct option

○	Interval
○	Ratio
○	Categorical ✓
○	Continuous

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 11:52 PM, 1

Question # 6 of 10 (Start time: 11:59:46 PM, 18 January 2024)

The population correlation is positive in a directional hypothesis. This type of hypothesis can be written statistically as:.

Select the correct option

Reload M

- $H_0: \rho \leq 0$
- $H_0: \rho = 0$
- $H_1: \rho < 0$ ✓
- $H_1: \rho \neq 0$

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 11:52 PM, 18 Ja

Question # 8 of 10 (Start time: 12:03:30 AM, 19 January 2024) To

Which of the following refers to the proportion of variability in the data that is explained by the relationship between X and Y?

Select the correct option

Reload Math

- Coefficient of determination
- Coefficient of variance
- Regression analysis
- Coefficient of correlation

PSY516 - Statistics in Psychology (Quiz 4)

Question # 7 of 10 (Start time: 12:01:52 AM, 19 January 2024)

Coefficient of determination is denoted by:

Select the correct option

○	F
○	r^2 ✓
○	r
○	T

PSY516 - Statistics in Psychology (Quiz 4)

Quiz Start Time: 11:52 PM, 18 January 2024

Question # 9 of 10 (Start time: 12:05:05 AM, 19 January 2024)

Total Marks: 1

The one-way analysis of variance (ANOVA) is used to determine whether there are any statistically significant differences between the means of _____independent (unrelated) groups.

Select the correct option

Reload Math Equations

- Two
- Less than two
- One
- Two or more 

PSY516 - Statistics in Psychology (Quiz 4)

Question # 10 of 10 (Start time: 12:06:49 AM, 19 January 2024)

What would be accurate interpretation of the following hypothesis H_1: $\rho \neq 0$?

Select the correct option

○	There is real correlation
○	There is no population correlation ✓
○	There is negative correlation
○	There is covariance

PSY516 - Statistics in Psychology (Quiz 4)

Quiz Start Time: 12:27 AM, 19 January 2024

Question # 1 of 10 (Start time: 12:27:24 AM, 19 January 2024)

Total Marks: 1

Which of the following statistical tests will be considered, If a researcher wants to know whether exam performance differs based on test anxiety levels among students, dividing students into three independent groups (e.g., low, medium and high-stressed students)?

Select the correct option

Reload Math Equations

- F test
- T test
- Anova
- Z test

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 12:27 AM, 19 Jar

Question # 2 of 10 (Start time: 12:28:58 AM, 19 January 2024) To

Which of the following is a statistical technique that is used to measure and describe the relationship between two variables?

Select the correct option

Reload Math E

- Standard deviation
- Correlation
- T-test
- Mean

PSY516 - Statistics in Psychology (Quiz 4)

Question # 3 of 10 (Start time: 12:30:06 AM, 19 January 2024)

What does ANOVA calculate?

Select the correct option

○	F ratios
○	T-scores
○	R ratios
○	Z-scores

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 12:27 AM,

Question # 4 of 10 (Start time: 12:30:48 AM, 19 January 2024)

The population correlation is positive in a directional hypothesis. This type of hypothesis can be written statistically as:.

Select the correct option

Reload

- $H_0: \rho = 0$
- $H_0: \rho \leq 0$
- $H_1: \rho \neq 0$
- $H_1: \rho < 0$

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 12:27 AM, 19

Question # 7 of 10 (Start time: 12:35:54 AM, 19 January 2024)

In ANOVA, which of the following tests is preferably used when the assumption of the homogeneity of variance is violated?

Select the correct option

Reload Mo

- Kolmogorov-Smirnov tests
- Welch and Brown-Forsythe
- Shapiro-Wilk
- Levene's test

PSY516 - Statistics in Psychology (Quiz 4)

Question # 8 of 10 (Start time: 12:37:04 AM, 19 January 2024)

What would be the best interpretation of Ho: ρ = 0?

Select the correct option

○	There is real correlation
○	There is covariance
○	There is no population correlation ✓
○	There is negative correlation

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start T

Question # 10 of 10 (Start time: 12:40:57 AM, 19 January 2024)

In one-way ANOVA, _____ is used to determine which of given groups differ from each other.

Select the correct option

- ○ Levene's test
- ○ T test
- ○ Post hoc test
- ○ F test

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 10:00 PM, 20 Janu

Question # 1 of 10 (Start time: 10:00:14 PM, 20 January 2024) Tota

Which of the following refers to a hypothesis-testing procedure that is used to evaluate mean differences between two or more treatments (or populations)?

Select the correct option

Reload Math Equ

- Analysis of variance (ANOVA)
- Chi-square
- Correlation
- Analysis of Regression

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 10:01:38 PM, 20 January 2024)

How many dependent variables you must have for an ANOVA to be conducted?

Select the correct option

○	1 continuous variable
○	3 ratio variables
○	3 interval variables
○	2 nominal variables

PSY516 - Statistics in Psychology (Quiz 4)

Question # 3 of 10 (Start time: 10:03:14 PM, 20 January 2024)

What does ANOVA calculate?

Select the correct option

○	Mann Whitney
○	F ratios ✓
○	Z score
○	T score

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 10:00 PM, 20 Ja

Question # 4 of 10 (Start time: 10:04:14 PM, 20 January 2024)

Which of the following is the total probability of a Type I error accumulated from all of the separate tests in the experiment?

Select the correct option

Reload Math

- Group wise alpha level
- Individual alpha level
- Test wise alpha level
- Experiment wise alpha level

PSY516 - Statistics in Psychology (Quiz 4)

Quiz Start Time: 10:00 PM, 20 January 2024

Question # 5 of 10 (Start time: 10:05:27 PM, 20 January 2024)

Total Marks: 1

The one-way analysis of variance (ANOVA) is used to determine whether there are any statistically significant differences between the means of _____independent (unrelated) groups.

Select the correct option

Reload Math Equations

- Two
- Less than two
- One
- Two or more

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 10:00 PM, 2

Question # 6 of 10 (Start time: 10:06:35 PM, 20 January 2024)

In within-subject design, dependent variable is measured _____ times for each individual in a single sample.

Select the correct option

Reload M

- One
- Two
- Four
- Three

PSY516 - Statistics in Psychology (Quiz 4)

Question # 7 of 10 (Start time: 10:08:18 PM, 20 January 2024)

For ANOVA, dependent variable is always_____ in nature.

Select the correct option

○	Interval
○	Ratio
○	Categorical
○	Continuous

Question # 9 of 10 (Start time: 10:11:22 PM, 20 January 2024)

A repeated measures t-test can be used to assess:

Select the correct option

○	Dfferences between scores obtained on two separate occasions from the same participants ✓
○	Differences between two groups of participants
○	Relationships between two interval data sets
○	How many factors there are in questionnaire data

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 10:22 PM, 20

Question # 1 of 10 (Start time: 10:22:20 PM, 20 January 2024)

In ANOVA, which of the following tests is preferably used when the assumption of the homogeneity of variance is violated?

Select the correct option

Reload Mat

- Shapiro-Wilk
- Kolmogorov-Smirnov tests
- Welch and Brown-Forsythe
- Levene's test

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 10:22:36 PM, 20 January 2024)

How many dependent variables you must have for an ANOVA to be conducted?

Select the correct option

○	1 continuous variable ✓
○	3 ratio variables
○	3 interval variables
○	2 nominal variables

PSY516 - Statistics in Psychology (Quiz 4)

Question # 3 of 10 (Start time: 10:22:52 PM, 20 January 2024)

In ANOVA, difference exists between the treatment conditions is termed as:

Select the correct option

○	Treatment variance
○	Within treatments variance
○	No treatment variance
○	Between-treatments variance ✓

PSY516 - Statistics in Psychology (Quiz 4)

Question # 4 of 10 (Start time: 10:24:00 PM, 20 January 2024)

How many levels must there be in one independent variable for an ANOVA to be used?

Select the correct option

○	1
○	2
○	4
○	3

PSY516 - Statistics in Psychology (Quiz 4)

Question # 5 of 10 (Start time: 10:25:19 PM, 20 January 2024)

In ANOVA, the homogeneity of variance assumption will be tested by:

Select the correct option

○	Levene's test
○	Kolmogorov-Smirnov tests
○	Welch and Brown-Forsythe
○	Shapiro-Wilk

PSY516 - Statistics in Psychology (Quiz 4)

Question # 6 of 10 (Start time: 10:26:35 PM, 20 January 2024)

The population correlation is traditionally represented by the Greek letter _____.

Select the correct option

○	t
○	r^2
○	ρ
○	H_0

Question # 7 of 10 (Start time: 10:28:41 PM, 20 January 2024)

Which of the following statement is true for the Pearson correlation?

Select the correct option

○	It measures the degree and the direction of the non-linear relationship between two or more variables.
○	All of the above ✓
○	It measures the degree and the direction of the linear relationship between two variables.
○	The relationship between two variables while controlling the influence of a third variable by holding it constant.

PSY516 - Statistics in Psychology (Quiz 4)

Quiz Start Time: 10:22 PM, 20 Ja

Question # 9 of 10 (Start time: 10:29:07 PM, 20 January 2024)

When there is a _____ linear relationship in Pearson correlation, every change in the X variable is accompanied by a corresponding change in the Y variable.

Select the correct option

Reload Math

- Perfect
- Positive
- Zero
- Negative

PSY516 - Statistics in Psychology (Quiz 4)

Question # 8 of 10 (Start time: 10:28:54 PM, 20 January 2024)

Which of the following is an advantage of ANOVA?

Select the correct option

○	All of the above
○	It cannot tolerate data that is non-normal
○	It performs all three comparisons simultaneously in one hypothesis test
○	It decides if the population mean is equal to a specific value or not

PSY516 - Statistics in Psychology (Quiz 4)

Question # 10 of 10 (Start time: 10:29:18 PM, 20 January 2024)

For two-way ANOVA, we calculate _____ kinds of degrees of freedom.

Select the correct option

○	Two
○	Three ✓
○	One
○	Four

PSY516 - Statistics in Psychology (Quiz 4)

Question # 1 of 10 (Start time: 11:02:38 PM, 20 January 2024)

Two-way ANOVA means groups are defined by two _____ variables.

Select the correct option

○	Partial
○	Independent
○	Extraneous
○	Dependent

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 11:04:26 PM, 20 January 2024)

Pearson's Correlation Analysis is also known as:

Select the correct option

○	Kruskal-Wallis Correlational Analysis
○	Pearson's Product-Moment ✓
○	Spearman's Correlation Analysis
○	Chi-Squared Product-Moment

PSY516 - Statistics in Psychology (Quiz 4)

Question # 3 of 10 (Start time: 11:04:41 PM, 20 January 2024)

Coefficient of determination is denoted by:

Select the correct option

○	F
○	r
○	r^2 ✓
○	T

PSY516 - Statistics in Psychology (Quiz 4)

Question # 5 of 10 (Start time: 11:05:13 PM, 20 January 2024)

In which of the following cases could you use a paired-samples t-test?

Select the correct option

- ○ When assessing three groups or more
- ○ When comparing two separate groups
- ○ When comparing the same participants performance before and after training ✓
- ○ When comparing men and women's scores

PSY516 - Statistics in Psychology (Quiz 4)

Question # 6 of 10 (Start time: 11:06:42 PM, 20 January 2024)

Which of the following is NOT an application of correlation coefficient?

Select the correct option

○	Prediction
○	Validity
○	Reliability
○	Variation

PSY516 - Statistics in Psychology (Quiz 4)

Question # 4 of 10 (Start time: 11:05:02 PM, 20 January 2024)

How many dependent variables you must have for an ANOVA to be conducted?

Select the correct option

○	2 nominal variables
○	1 continuous variable 
○	3 ratio variables
○	3 interval variables

PSY516 - Statistics in Psychology (Quiz 4)

Question # 8 of 10 (Start time: 11:08:12 PM, 20 January 2024)

How should a paired-samples t-test be reported in APA format?

Select the correct option

○	$F\ (12, 29) = 9.67, p < 0.001$
○	$Z = 4.87, t\ (29) = 5.93, p = 0.000$
○	$r\ (18) = +8.97, p > 0.005$
○	$t(37) = +.3.67, p < 0.05$



PSY516 - Statistics in Psychology (Quiz 4)

Question # 9 of 10 (Start time: 11:10:01 PM, 20 January 2024)

Which of the following statement is true for the Pearson correlation?

Select the correct option

○	Generally computed for dispersed data
○	Generally computed for sample data ✓
○	Generally computed for population data
○	Generally computed for scattered data

PSY516 - Statistics in Psychology (Quiz 4)

Question # 10 of 10 (Start time: 11:11:19 PM, 20 January 2024)

What does ANOVA calculate?

Select the correct option

○	F ratios
○	T-scores
○	Z-scores
○	R ratios

sec(s)

PSY516 - Statistics in Psychology (Quiz 4) Quiz Start Time: 07:29 PM, 20 January 2024

Question # 2 of 10 (Start time: 07:31:02 PM, 20 January 2024) Total Marks: 1

Which of the following is the total probability of a Type I error accumulated from all of the separate tests in the experiment?

Select the correct option

Reload Math Equations

- Group wise alpha level
- Experiment wise alpha level
- Test wise alpha level 
- Individual alpha level

PSY516 - Statistics in Psychology (Quiz 4)

Question # 2 of 10 (Start time: 10:59:21 PM, 18 January 2024)

If two variables are moving in the same direction, it is called as_____ correlation.

Select the correct option

○	Negative
○	Non-linear
○	Linear
○	Positive ✓

Question # 5 of 10 (Start time: 12:36:54 AM, 18 January 2024)

What is the alternative name for a repeated-measures t-test?

Select the correct option

○	A Wilcoxon test
○	Spearman's Rho
○	Pearson's Product-Moment
◉	A paired-samples t-test