BIO5101 Introduction to Biotechnology (Topic 141-170)

Topic 141 – Biotechnology Role in Food

1. Biofortification (Nutritional Enhancement)

Biofortification is the process of increasing the density of vitamins and minerals in a crop through genetic engineering.

- **Golden Rice:** The most famous example. Rice is engineered with genes from daffodils and soil bacteria (*Erwinia*) to produce **Beta-carotene** (Pro-vitamin A). This helps combat Vitamin A deficiency, which causes blindness.
- **Iron/Zinc enrichment:** Developing cereal crops with higher mineral content to address "hidden hunger" in developing nations.

2. Delayed Ripening (Shelf-Life Extension)

Fruits like tomatoes produce **Ethylene gas**, which signals them to ripen and eventually rot.

- **The Flavr Savr Tomato:** The first GM food sold. Scientists used **Antisense RNA technology** to block the enzyme **Polygalacturonase (PG)**. This enzyme breaks down pectin in cell walls; by blocking it, the tomato stays firm for longer, allowing it to ripen on the vine for better flavor without spoiling during transport.

3. Modification of Food Composition

Biotechnology allows us to change the "ingredients" within a plant to make them healthier:

- **Healthier Fats:** Engineering oilseed crops (like canola or soybean) to produce high levels of **Omega-3 fatty acids** or reduced saturated fats.
- **Reduced Allergens:** Using gene silencing to remove proteins that cause allergies in peanuts or wheat.
- **Sweetener Production:** Producing low-calorie sweeteners like **Stevia** or **Thaumatin** in transgenic plants or microbes.

4. Food Processing and Enzymes

Biotechnology isn't just about the crops; it's about the tools used to make food:

- **Chymosin (Rennet):** Traditionally taken from the stomachs of calves to make cheese. Today, most cheese is made using **recombinant Chymosin** produced by engineered yeast or fungi.

5. Safety and Detection

Biotechnology provides tools to ensure food safety:

- **Biosensors:** Using engineered proteins to detect toxins or bacteria (*E. coli*, *Salmonella*) in food products instantly.
- **GMO Detection:** Using PCR (Polymerase Chain Reaction) to verify if food contains genetically modified ingredients for labeling purposes.

---

10 Short Questions & Answers

1. **Q: What is Golden Rice? A:** A variety of rice engineered to produce beta-carotene, a precursor of Vitamin A.
2. **Q: What does the enzyme Polygalacturonase (PG) do? A:** It breaks down pectin in the cell walls of fruit, causing softening and rotting.
3. **Q: Define "Biofortification." A:** Improving the nutritional value of food crops through biotechnology or breeding.
4. **Q: How was the Flavr Savr tomato created? A:** Using Antisense RNA technology to silence the PG gene.
5. **Q: What is the benefit of high-oleic acid soybean oil? A:** It is more heat-stable for frying and contains less saturated fat, making it heart-healthy.
6. **Q: Name the "Stress Hormone" in fruits responsible for ripening. A:** Ethylene.
7. **Q: What is Chymosin used for? A:** It is an enzyme used to curdle milk in the cheese-making process.
8. **Q: Why is Golden Rice yellow/orange? A:** Because of the presence of beta-carotene (the same pigment found in carrots).
9. **Q: What is "Hidden Hunger"? A:** A form of malnutrition where people get enough calories but lack essential vitamins and minerals.
10. **Q: How does biotechnology help people with peanut allergies? A:** By using gene silencing to "knock out" the specific proteins (allergens) that trigger the allergic reaction.

---

5 Long Questions & Answers

1. **Q: Explain the molecular strategy behind the development of Golden Rice.**
   A: Rice naturally produces the precursor for beta-carotene (GGPP) in its leaves, but the pathway is "turned off" in the edible grain (endosperm). To fix this, scientists inserted two genes: psy (phytoene synthase) from Daffodil and crtI from the soil bacterium Erwinia uredovora. These genes provide the enzymes necessary to complete the biochemical pathway, allowing the rice grain to synthesize and accumulate beta-carotene. This biofortified rice can provide up to 50% of the daily Vitamin A requirement in a standard diet.
2. **Q: Describe the mechanism of Antisense RNA technology in the Flavr Savr tomato.**
   A: In a normal tomato, the PG gene produces mRNA, which is translated into the PG enzyme that softens the fruit. In the Flavr Savr tomato, scientists inserted a "reverse" version of the PG gene. This produces an Antisense RNA strand that is complementary to the normal "Sense" mRNA. The two strands bind together to form double-stranded RNA. The cell's machinery cannot

translate double-stranded RNA into a protein, effectively "silencing" the PG enzyme and keeping the tomato firm.

3. **Q: Discuss the advantages of Recombinant Chymosin over traditional Rennet.**
   A: Traditional rennet is extracted from the stomach lining of slaughtered calves, which is expensive, inconsistent in quality, and ethically problematic for vegetarians. Recombinant Chymosin (FPC) is produced by inserting the bovine chymosin gene into microbes like Kluyveromyces lactis. This produces a pure, high-quality enzyme that is identical to the animal version but is cheaper, "vegetarian-friendly," and avoids the risks of animal-borne diseases.
4. **Q: How can biotechnology be used to improve the fatty acid profile of cooking oils?**
   A: Scientists target the enzymes involved in the lipid biosynthetic pathway, such as desaturases. For example, by silencing the gene that converts oleic acid (healthy) to linoleic acid (less stable), they can create "High-Oleic" oils. These oils have a longer shelf-life, produce fewer trans-fats when heated, and offer cardiovascular benefits similar to olive oil, but in high-yield crops like soybean or canola.
5. **Q: Evaluate the role of biotechnology in ensuring food safety and quality control.**
   A: Biotechnology provides high-precision tools for monitoring. PCR-based assays can detect the DNA of pathogens like Listeria or Salmonella much faster than traditional culturing. ELISA tests use antibodies to detect toxins or allergens. Additionally, Biosensors (using bioluminescent bacteria or enzymes) can be integrated into food packaging to change color if the food begins to spoil or if harmful gases are produced, providing a real-time safety indicator for consumers.

---

15 Multiple Choice Questions (MCQs)

1. Golden Rice was developed to address deficiency in:

   a) Vitamin C

   b) Vitamin A ✓

   c) Iron

   d) Iodine

2. The first genetically modified food to be commercially sold was:

   a) Flavr Savr Tomato ✓

   b) Bt Corn

   c) Golden Rice

   d) Roundup Ready Soybean

3. The Flavr Savr tomato used _____ technology.

   a) Gene Gun

   b) Antisense RNA ✓

c) Electroporation

d) Protoplast fusion

4. Beta-carotene is a precursor for:

a) Vitamin D

b) Protein

c) Vitamin A ✓

d) Vitamin K

5. Which enzyme is responsible for the softening of tomatoes?

a) Polygalacturonase ✓

b) Amylase

c) Cellulase

d) Lipase

6. The gene for beta-carotene in Golden Rice was taken from Daffodil and:

a) Human

b) Yeast

c) Erwinia (Bacterium) ✓

d) Agrobacterium

7. "Hidden Hunger" refers to a lack of:

a) Calories

b) Micronutrients ✓

c) Carbohydrates

d) Water

8. Most cheese today is made using recombinant:

a) Insulin

b) Lipase

c) Chymosin ✓

d) Protease

9. Ethylene is a plant hormone that functions as a/an:

a) Ripening signal ✓

b) Growth inhibitor

c) Antifreeze

d) Pigment

10. Gene silencing in peanuts is used to remove:

a) Oil

b) Allergens ✓

c) Shells

d) Roots

11. PCR is used in the food industry to:

a) Detect GMOs and pathogens ✓

b) Fry food

c) Increase sugar content

d) Change food color

12. The pigment Beta-carotene belongs to the _____ family.

a) Protein

b) Carotenoid ✓

c) Alkaloid

d) Lipid

13. Antisense RNA works by preventing:

a) DNA replication

b) Translation of mRNA ✓

c) Root growth

d) Water absorption

14. Which of these is a low-calorie sweetener produced by biotechnology?

a) Glucose

b) Fructose

c) Thaumatin ✓

d) Sucrose

15. Biotechnology can increase the "Shelf-life" of a product, meaning it:

    a) Stays fresh for a longer time ✓

    b) Grows faster in the field

    c) Needs less fertilizer

    d) Is cheaper to plant

Topic 142 – Ethical Issues of GM Food

1. Human Health and Safety Concerns

The primary ethical debate revolves around the "Right to Safety."

- **Allergenicity:** There is a fear that moving genes from an allergenic source (like Brazil nuts) into a non-allergenic crop (like soy) could cause unexpected allergic reactions in consumers.
- **Antibiotic Resistance:** Many GM crops use "Antibiotic Resistance Marker Genes" during lab development. Critics worry these genes could transfer to bacteria in the human gut, making medical antibiotics less effective.

2. Environmental Ethics

This involves our responsibility toward the "Balance of Nature."

- **Gene Flow (Superweeds):** The risk that herbicide-resistance genes will escape via pollen to wild relatives, creating "superweeds" that cannot be controlled.
- **Non-Target Organisms:** The ethical concern of harming "innocent" species. A famous example is the debate over whether **Bt Corn** pollen could harm **Monarch Butterfly** larvae.

3. Socio-Economic and Corporate Ethics

- **Corporate Monopoly:** A few large companies own the patents to most GM seeds. This raises the ethical question of "Food Sovereignty"—should a private company control the global food supply?
- **Terminator Technology:** Theoretically, seeds could be engineered to be sterile in the second generation. This would force poor farmers to buy new seeds every year instead of saving them, which is a traditional practice.

4. Religious and Cultural Concerns

- **"Playing God":** Some argue that crossing species boundaries (e.g., putting a fish gene into a tomato) is an unnatural violation of the "integrity of species."

- **Dietary Restrictions:** For individuals with religious dietary laws (Halal, Kosher, or Vegetarianism), a crop containing an animal gene may be considered "unclean" or unacceptable.

5. The Solution: Labeling and Regulation

The ethical principle of **Autonomy** (Informed Consent) suggests that consumers have the right to know what is in their food. This has led to mandatory **GMO Labeling** in many countries, allowing individuals to make choices based on their own ethical beliefs.

---

10 Short Questions & Answers

1. **Q: What is the main concern regarding allergens in GM food? A:** That a gene from a known allergen (like nuts) could be transferred to a common food (like corn) without the consumer knowing.
2. **Q: Define "Horizontal Gene Transfer." A:** The movement of genetic material between different species, such as from a GM plant to a gut bacterium.
3. **Q: What is a "Terminator Seed"? A:** A seed engineered to be sterile, preventing farmers from saving and replanting them.
4. **Q: Why is labeling of GM food an ethical issue? A:** It relates to the consumer's "Right to Know" and the ability to make an informed choice.
5. **Q: What does "Playing God" mean in biotechnology? A:** An ethical objection that humans should not interfere with the fundamental genetic makeup of living organisms.
6. **Q: How can GM crops affect biodiversity? A:** If one GM variety dominates the market, local traditional varieties (landraces) may be lost forever.
7. **Q: What are "Antibiotic Resistance Markers"? A:** Genes used in the lab to identify successfully transformed cells; they raise concerns about the spread of antibiotic resistance.
8. **Q: Who regulates the safety of GM foods globally? A:** Organizations like the FAO, WHO, and national agencies like the FDA or EPA.
9. **Q: What is the "Precautionary Principle"? A:** The idea that if an action has a risk of causing harm, one should avoid it even if there is no full scientific certainty yet.
10. **Q: Can GM foods affect non-target insects? A:** Yes, for example, a toxin meant for a pest might accidentally harm beneficial insects like bees or butterflies.

---

5 Long Questions & Answers

1. **Q: Discuss the ethical implications of corporate ownership and patenting of life forms.**
   A: When a company patents a GM seed, they own the intellectual property of that living organism. Ethically, this creates a power imbalance. Farmers become dependent on corporations for seeds and chemicals, often leading to a loss of traditional farming knowledge. There is also the concern that profit motives might overshadow safety or environmental concerns, particularly in developing nations where regulations may be weaker.

2. **Q: Explain the conflict between "Informed Consent" and the "Substantial Equivalence" doctrine.**
   A: Substantial Equivalence is the idea that if a GM food is chemically similar to its natural counterpart, it doesn't need special labeling or testing. However, the ethical principle of Informed Consent argues that because the technology is new, consumers must be told if a food is GM so they can decide for themselves whether to eat it, regardless of chemical similarity. This is why many countries now require mandatory GMO labels.
3. **Q: Analyze the environmental ethics regarding the impact of Bt crops on non-target species**.
   A: Bt crops produce a toxin that targets specific pests. However, an ethical dilemma arises when that toxin (through pollen or leaf litter) enters the wider ecosystem. If the pollen lands on weeds eaten by the Monarch butterfly, it may kill them. Ethically, we must weigh the "Good" (reduced chemical pesticide use) against the "Harm" (unintended death of non-target species). Scientists use "Risk-Benefit Analysis" to decide if the technology should be used.
4. **Q: How do GM foods intersect with religious and cultural values?**
   A: Many cultures have a deep, spiritual connection to their food. For instance, corn is sacred in some Indigenous cultures. Altering its DNA is seen as a violation of that sacredness. Similarly, Muslims or Jews may have concerns if genes from "prohibited" animals are used in plants. Ethically, biotechnology must respect "Cultural Integrity" and ensure that "Transgenic" borders do not offend religious beliefs.
5. **Q: What is "Biopiracy" and how does it relate to biotechnology ethics?**
   A: Biopiracy occurs when researchers or companies take genetic resources from developing countries (like a traditional medicinal plant) and patent them without permission or sharing the profits with the local community. Ethically, this is seen as a form of "Genetic Colonialism." International treaties, like the Nagoya Protocol, aim to ensure "Access and Benefit Sharing" to prevent such exploitation.

---

15 Multiple Choice Questions (MCQs)

1. The concern that GM crops could harm Monarch butterflies is an example of:

   a) Human health risk

   b) Impact on non-target organisms ✓

   c) Corporate monopoly

   d) Antibiotic resistance

2. "Substantial Equivalence" means a GM food is:

   a) Chemically similar to a traditional food ✓

   b) More expensive

   c) Always toxic

   d) Produced only in labs

3. Labeling of GM products supports the ethical principle of:
    a) Profit
    b) Secrecy
    c) Consumer Autonomy (Right to Know) ✓
    d) Complexity
4. A "Superweed" is a weed that has:
    a) Grown very tall
    b) Become edible
    c) Acquired resistance to herbicides via gene flow ✓
    d) Died out
5. Antibiotic resistance markers are used during:
    a) The transformation process in the lab ✓
    b) Harvesting in the field
    c) Cooking the food
    d) Selling the seeds
6. The ethical objection that humans should not "Play God" is a/an:
    a) Economic objection
    b) Moral/Philosophical objection ✓
    c) Scientific objection
    d) Legal objection
7. The movement of a transgene from a crop to a wild relative is called:
    a) Transpiration
    b) Gene Flow ✓
    c) Mutation
    d) Respiration
8. Terminator technology refers to:
    a) Making seeds sterile so they cannot be replanted ✓
    b) Using robots to farm

c) Killing all insects

d) Destroying the lab

9. The fear that a fish gene in a tomato might trigger an allergy is a concern of:

a) Allergenicity ✓

b) Nutrition

c) Soil quality

d) Climate change

10. Which group is most concerned about "Access and Benefit Sharing"?

a) Seed companies

b) Indigenous/Local communities ✓

c) Software engineers

d) Urban consumers

11. The "Precautionary Principle" suggests that we should:

a) Proceed with caution when risks are unknown ✓

b) Ignore risks if profits are high

c) Stop all science

d) Only use old technology

12. Corporate control of the seed market raises the issue of:

a) All-natural farming

b) Over-watering

c) Food Security and Sovereignty ✓

d) Pest control

13. Biopiracy involves the unauthorized use of:

a) Computers

b) Genetic resources from other countries ✓

c) Tractors

d) Fertilizers

14. GMO labeling is mandatory in:

a) No countries

b) Only the USA

c) The European Union and many other nations ✓

d) Only in space

15. A "Landrace" is a:

a) Traditional, locally adapted variety of a crop ✓

b) Type of fast-growing GM corn

c) Type of tractor

d) Chemical fertilizer

Topic 143 – Risk Factors of GM Food

1. Human Health Risks

The scientific community evaluates three primary areas of concern for human health:

- **Toxicity:** There is a possibility that the inserted gene might produce a protein that is toxic to humans or that the insertion process might disrupt a native plant gene, causing the plant to produce its own natural toxins at higher levels.
- **Allergenicity:** If a gene is taken from a known allergenic food (like Brazil nuts or wheat) and put into another crop, the new crop might trigger allergic reactions in sensitive individuals.
- **Antibiotic Resistance:** During the lab process, **Antibiotic Resistance Marker Genes (ARMGs)** are often used. There is a theoretical risk that these genes could transfer to bacteria in the human gut, contributing to the global problem of antibiotic-resistant "superbugs."

2. Environmental Risks

The release of GMOs into the open environment presents several "Ecological Risks":


- **Gene Flow (Genetic Pollution):** Transgenes can spread via pollen to wild relatives of the crop. This could create **"Superweeds"** that are resistant to herbicides and can outcompete natural vegetation.
- **Impact on Non-Target Organisms:** Insects that are not pests (like bees, butterflies, or lacewings) might be harmed by eating the pollen or tissues of insect-resistant crops (like Bt corn).
- **Loss of Biodiversity:** If farmers only grow a few "high-performing" GM varieties, traditional local varieties (**landraces**) may disappear, reducing the genetic diversity available for future breeding.

3. Evolution of Resistance

- **Insect Resistance:** Constant exposure to Bt toxins in crops can lead to the evolution of resistant insect populations.
- **Weed Resistance:** Overuse of a single herbicide (like Glyphosate) on resistant crops leads to the emergence of weeds that the chemical can no longer kill.

4. Horizontal Gene Transfer (HGT)

HGT is the movement of genetic material between unicellular and/or multicellular organisms other than via traditional reproduction. The risk is that DNA from GM food could be taken up by microbes in the soil or the human digestive tract, leading to unpredictable ecological or health consequences.



10 Short Questions & Answers

1. **Q: What is a "Superweed"? A:** A wild plant or weed that has acquired herbicide-resistance genes from a GM crop through cross-pollination.
2. **Q: Name one health risk associated with GM food. A:** Potential for increased allergenicity or toxicity.
3. **Q: What are Non-Target Organisms? A:** Beneficial or neutral insects and animals that are not the intended "target" of a GM trait (e.g., honeybees).
4. **Q: Define Gene Flow. A:** The transfer of genetic material from one population to another, often via pollen or seeds.
5. **Q: Why are Antibiotic Resistance Marker Genes used? A:** To identify and select only those cells that have successfully taken up the foreign DNA in the laboratory.
6. **Q: What is "Genetic Pollution"? A:** An informal term for the unintended spread of transgenes into the wild or organic gene pools.
7. **Q: How does the "Refuge Strategy" help? A:** It slows down the evolution of insect resistance by providing a place for susceptible insects to breed.
8. **Q: What is the risk of "Gene Silencing"? A:** The risk that the plant might eventually turn off the transgene, making the crop lose its beneficial trait.
9. **Q: Can GM DNA survive human digestion? A:** Most DNA is broken down, but there is a theoretical concern that small fragments could remain functional.
10. **Q: What is a "Secondary Pest"? A:** A pest that was previously minor but becomes a major problem after the primary pest is controlled by a GM crop.

5 Long Questions & Answers

1. **Q: Explain the potential risk of allergenicity in GM crops with a real-world example.**
   A: Allergenicity is a major safety concern. A famous case involved the Pioneer Hi-Bred company, which tried to improve the nutritional quality of soybeans by adding a gene from Brazil nuts. Tests later showed that the transgenic soy triggered reactions in people allergic to nuts. Because of these safety protocols, the project was cancelled before the product reached the market. This demonstrates that while the risk exists, rigorous testing can prevent dangerous products from being sold.
2. **Q: Describe the environmental consequences of "Superweeds" and how they form.**
   A: Superweeds form through pollen-mediated gene flow. If a herbicide-resistant GM crop (like Roundup Ready Soy) is planted near a wild relative (like wild mustard), the pollen can fertilize the wild plant. The offspring may inherit the resistance gene. If farmers continue to use only one herbicide, these resistant wild plants will survive and multiply, eventually requiring more toxic chemicals or mechanical plowing to control, which harms soil health.
3. **Q: What is the significance of "Horizontal Gene Transfer" in the context of GM risk?**
   A: Horizontal Gene Transfer (HGT) involves the movement of DNA between different species

without mating. The primary risk factor is that Antibiotic Resistance Marker Genes (like kanR or ampR) used in GM plants could be absorbed by bacteria in the human gut or the soil. If these bacteria become resistant to antibiotics used in human medicine, it could make treating common infections much more difficult. Although the probability is considered low, it remains a key point of biosafety assessment.

4. **Q: Discuss the impact of Bt crops on non-target insects like the Monarch Butterfly.**
   A: Bt crops produce a toxin specifically for pests like the Corn Borer. However, the toxin is also present in the pollen. If wind carries this pollen onto Milkweed plants (the primary food for Monarch larvae) located near the field, the larvae might consume it and die. This raises an ethical and ecological risk factor regarding the loss of beneficial insects and the disruption of the local food web.
5. **Q: How does the "Loss of Biodiversity" occur due to the dominance of GM crops?**
   A: As a few patented, high-yielding GM varieties become popular, farmers stop planting traditional landraces (local varieties). These local varieties contain unique genes for drought tolerance, flavor, or disease resistance that have evolved over centuries. If they are no longer planted, those genes may be lost forever. This "Genetic Erosion" makes the global food supply more vulnerable to new diseases or climate changes that the dominant GM varieties might not be able to handle.

---

15 Multiple Choice Questions (MCQs)

1. Which of these is a potential health risk of GM food?

   a) Faster cooking time

   b) Allergenicity ✓

   c) Improved flavor

   d) Higher yield

2. The movement of genes from GM crops to wild relatives is called:

   a) Respiration

   b) Gene Flow ✓

   c) Photosynthesis

   d) Transpiration

3. Marker genes used in GM labs often provide resistance to:

   a) Antibiotics ✓

   b) Cold

c) Sunlight

d) Water

4. A "Superweed" is a weed that has acquired:

   a) Better seeds

   b) Herbicide resistance ✓

   c) More leaves

   d) Fruit-bearing ability

5. Monarch butterflies are considered _____ in the context of Bt corn.

   a) Non-target organisms ✓

   b) Target pests

   c) Soil microbes

   d) Pollinators

6. "Genetic Erosion" refers to the loss of:

   a) Soil

   b) Biodiversity/Landraces ✓

   c) Water

   d) Fertilizer

7. Which mechanism allows DNA to move from a plant to a bacterium?

   a) Vertical gene transfer

   b) Horizontal gene transfer ✓

   c) Cross-pollination

   d) Grafting

8. The "Refuge Strategy" is designed to manage:

   a) Insect resistance ✓

   b) Soil pH

   c) Weed growth

   d) Market prices

9. The concern about Brazil nut genes in soybeans was related to:

   a) Color

b) Allergenicity ✓

c) Shape

d) Growth speed

10. Bt toxin is primarily toxic to the larvae of:

a) Lepidoptera (Butterflies/Moths) ✓

b) Mammals

c) Fish

d) Birds

11. Herbicide resistance in weeds is a result of:

a) Lack of sun

b) Continuous selection pressure ✓

c) Too much water

d) Low temperature

12. The use of markers helps scientists to:

a) Select transformed cells ✓

b) Kill the plant

c) Increase vitamin content

d) Change the flower color

13. Which organization evaluates the safety of GM foods?

a) NASA

b) WHO / FAO ✓

c) Olympic Committee

d) Interpol

14. Secondary pests emerge when:

a) The primary pest is eliminated by GM crops ✓

b) The plant dies

c) There is a flood

d) Too much fertilizer is used

15. A major environmental risk of GM crops is the disruption of:

    a) Gravity

    b) Local ecosystems/Food webs ✓

    c) Moon cycles

    d) Magnetic fields

Topic 144 – Genetic Modification of Animals

1. Methods of Gene Delivery in Animals

Unlike plants, animal cells do not have a cell wall, but delivering DNA to a developing embryo is a precise task:

- **Microinjection:** The most common method. A microscopic glass needle is used to inject DNA directly into the male pronucleus of a fertilized egg (zygote).
- **Retroviral Vectors:** Using modified viruses as "shuttles" to carry genes into animal cells.
- **Embryonic Stem (ES) Cell-Mediated Transfer:** DNA is introduced into stem cells in a lab. These cells are then inserted into a developing embryo (blastocyst), creating a "chimera."
- **Somatic Cell Nuclear Transfer (SCNT):** Known as "Cloning." A donor nucleus is placed into an enucleated egg (e.g., Dolly the Sheep).

2. Applications of Transgenic Animals

- **Medical Research (Disease Models):** "Knockout Mice" are engineered to lack specific genes (like those for cancer or Alzheimer's) so scientists can study how the disease works.
- **Biopharming:** Animals act as bioreactors to produce human proteins in their milk. For example, **ATryn** (an anticoagulant) is produced in the milk of transgenic goats.
- **Enhanced Agriculture:** Animals engineered for faster growth (AquaAdvantage Salmon) or better wool production.
- **Xenotransplantation:** Modifying animals (like pigs) so their organs are not rejected by the human immune system when transplanted.

3. Key Examples

- **Dolly the Sheep:** The first mammal cloned from an adult somatic cell (1996).
- **AquaAdvantage Salmon:** The first GM animal approved for human consumption; it grows to market size in half the time of regular salmon.

4. Challenges and Ethics

- **Animal Welfare:** Does the modification cause pain or suffering to the animal?

- **Efficiency:** The success rate of creating a healthy transgenic animal is very low (often <5%).
- **Complexity:** Animal genomes are highly regulated and complex compared to microbes.

---

10 Short Questions & Answers

1. **Q: What is a Transgenic Animal? A:** An animal whose genome has been altered by the introduction of one or more foreign DNA sequences.
2. **Q: Define "Microinjection." A:** A technique using a fine glass pipette to inject DNA directly into the nucleus of a cell or embryo.
3. **Q: What is a "Knockout Mouse"? A:** A mouse in which a specific gene has been made inoperative (turned off) to study that gene's function.
4. **Q: What is "Biopharming" in animals? A:** Using transgenic animals to produce pharmaceutical proteins in their milk, blood, or eggs.
5. **Q: Why is milk a preferred site for recombinant protein production? A:** It is easy to harvest, produced in large quantities, and does not harm the animal.
6. **Q: What was unique about Dolly the Sheep? A:** She was the first mammal cloned from an adult (somatic) cell, not an embryo.
7. **Q: Define "Xenotransplantation." A:** The transplantation of living cells, tissues, or organs from one species to another (e.g., pig to human).
8. **Q: What is a "Chimera"? A:** An animal composed of two or more genetically distinct cell populations.
9. **Q: Which animal is most commonly used in transgenics? A:** The mouse (due to fast reproduction and well-mapped genome).
10. **Q: What is the male pronucleus? A:** The nucleus of the sperm cell after it enters the egg but before it fuses with the egg nucleus; it is the target for microinjection.

---

5 Long Questions & Answers

1. **Q: Describe the process of Microinjection in creating transgenic mice.**
   A: The process begins with superovulation of a female mouse to obtain many eggs. These eggs are fertilized in vitro. Under a specialized microscope, a very fine glass needle is used to inject the transgene into the male pronucleus (which is larger than the female one). The injected eggs are then cultured and surgically transferred into the oviduct of a "pseudo-pregnant" surrogate mother. The resulting offspring are screened via PCR to see which ones have successfully integrated the gene into their DNA.
2. **Q: Explain the role of "Biopharming" and provide an example of a product.**
   A: Biopharming uses animals as "living factories." Scientists link a human gene (like a blood-clotting factor) to a promoter that only functions in the mammary glands (like the casein promoter). Consequently, the protein is only secreted in the animal's milk. A famous example is ATryn, a human antithrombin protein produced in the milk of transgenic goats. It is used to prevent blood clots in patients with a hereditary deficiency during surgery or childbirth.

3. **Q: Discuss the scientific and ethical hurdles of Xenotransplantation.**
   A: Scientifically, the biggest hurdle is Hyperacute Rejection, where the human immune system immediately attacks the animal organ because of specific sugars (Gal) on the animal's cells. Biotechnology is used to "knock out" these sugar-producing genes. Ethically, concerns include the risk of "Zoonosis" (transfer of animal viruses to humans) and the moral status of raising animals solely as organ donors.
4. **Q: How does Somatic Cell Nuclear Transfer (SCNT) differ from traditional breeding?**
   A: Traditional breeding involves the recombination of two parents' DNA, resulting in a unique offspring. SCNT (Cloning) bypasses fertilization. A nucleus from an adult body cell (somatic cell) is inserted into an egg that has had its own nucleus removed. This creates a genetic "twin" of the donor. It allows for the exact replication of an animal with high-value traits (like high milk yield or a specific genetic modification).
5. **Q: Evaluate the benefits and risks of AquaAdvantage Salmon.**
   A: The benefit is Efficiency: the fish carry a growth hormone gene from Chinook salmon and a promoter from an ocean pout, allowing them to grow year-round and reach market size in 18 months instead of 3 years. This reduces the pressure on wild fish stocks. The risk is Environmental: if these GM salmon escape into the wild, they could outcompete wild salmon, leading to ecological disruption, though they are engineered to be sterile to prevent this.

---

15 Multiple Choice Questions (MCQs)

1. The most common method for creating transgenic animals is:

   a) Viral infection

   b) Microinjection ✓

   c) Electroporation

   d) Chemicals

2. The first cloned mammal, Dolly the Sheep, was created using:

   a) Microinjection

   b) Sperm-mediated transfer

   c) Somatic Cell Nuclear Transfer (SCNT) ✓

   d) Gene gun

3. In microinjection, the DNA is injected into the:

   a) Cytoplasm

   b) Male Pronucleus ✓

c) Vacuole

d) Mitochondria

4. Knockout mice are primarily used to study:

a) How to kill mice

b) Gene function and human diseases ✓

c) Agricultural yield

d) Faster growth

5. Recombinant proteins in biopharming are usually harvested from:

a) Milk ✓

b) Hair

c) Hooves

d) Teeth

6. AquaAdvantage Salmon contains a growth hormone gene from:

a) Chinook Salmon ✓

b) Pig

c) Human

d) Bacteria

7. A "Chimera" is an animal that:

a) Is a clone

b) Contains cells from two different sources ✓

c) Has no DNA

d) Cannot reproduce

8. The enzyme/promoter used to target milk production is usually:

a) Insulin

b) PEP

c) Casein or \beta-lactoglobulin ✓

d) Amylase

9. Xenotransplantation involves transferring organs from:

a) Humans to pigs

b) Animals to humans ✓

c) Plants to animals

d) Lab to lab

10. Embryonic stem cells are taken from which stage of the embryo?

a) Zygote

b) Blastocyst ✓

c) Adult

d) Sperm

11. Dolly the sheep was a clone of her _____ donor.

a) Egg

b) Nuclear (Somatic) ✓

c) Surrogate mother

d) Father

12. Which animal is a preferred "Bioreactor" for large-scale protein production?

a) Goat or Cow ✓

b) Mouse

c) Insect

d) Fish

13. "Zoonosis" refers to:

a) Animal growth

b) Sleep patterns

c) Disease transfer from animals to humans ✓

d) Genetic mutation

14. The success rate of microinjection is generally:

a) 100%

b) 50%

c) Very low (<5%) ✓

d) Increasing to 90%

15. Transgenic animals used for organ donation are usually:

    a) Pigs ✓

    b) Monkeys

    c) Cats

    d) Mice

Topic 145 – Strategies to Transform Animal Cells

1. Introduction to Animal Cell Transformation

Transformation (often called **transfection** in animal biology) is the process of intentionally introducing nucleic acids into cells. In animal cells, this can be "Transient" (temporary) or "Stable" (integrated into the genome).

2. Physical Methods

These methods use physical force to penetrate the plasma membrane:

- **Microinjection:** DNA is injected directly into the nucleus using a glass capillary. It is the most efficient but most labor-intensive method.
- **Electroporation:** Short, high-voltage pulses create temporary "nanopores" in the cell membrane, allowing DNA to enter via diffusion.
- **Biolistics (Gene Gun):** Heavy metal particles (gold/tungsten) coated with DNA are "fired" into tissues. This is often used for skin or muscle cells *in vivo*.

3. Chemical Methods

Chemicals are used to mask the negative charge of DNA and facilitate its entry:

- **Calcium Phosphate Precipitation:** DNA is mixed with calcium chloride and phosphate buffer. The resulting DNA-calcium phosphate precipitate is taken up by cells via **endocytosis**.
- **Lipofection (Liposome-mediated):** DNA is encapsulated in artificial lipid vesicles called **liposomes**. These fuse with the plasma membrane, releasing the DNA into the cytoplasm.
- **DEAE-Dextran:** A positively charged polymer that binds to negatively charged DNA, facilitating its attachment to the cell surface and subsequent uptake.

4. Biological Methods (Transduction)

This involves using **Viral Vectors**. Viruses are natural "gene delivery machines."

- **Retroviruses:** Efficiently integrate DNA into the host genome.
- **Adenoviruses:** Often used for transient expression; they do not integrate into the genome.
- **Lentiviruses:** A subclass of retroviruses that can infect both dividing and non-dividing cells.

---

10 Short Questions & Answers

1. **Q: What is "Transfection"? A:** The process of introducing foreign DNA into animal cells using non-viral methods.
2. **Q: Why is "Lipofection" popular? A:** It is highly efficient, easy to use, and works on a wide variety of cell types.
3. **Q: Define "Transient Expression." A:** When the introduced DNA remains in the nucleus but does not integrate into the host chromosome, eventually being lost as cells divide.
4. **Q: What is the role of Calcium Phosphate in transformation? A:** It forms a precipitate with DNA that cells ingest through endocytosis.
5. **Q: Name one drawback of Electroporation. A:** High cell mortality; many cells die due to the intensity of the electric pulse.
6. **Q: What are "Liposomes"? A:** Small, spherical vesicles made of a lipid bilayer used to carry DNA into cells.
7. **Q: Which method is best for transforming large numbers of cells simultaneously? A:** Electroporation or Chemical methods (Calcium Phosphate).
8. **Q: What is a "Viral Vector"? A:** A virus that has been modified to carry a specific gene of interest instead of its own disease-causing genes.
9. **Q: Define "Endocytosis." A:** The process by which a cell takes in materials by engulfing them in a section of its membrane.
10. **Q: How does DEAE-Dextran facilitate DNA entry? A:** It provides a positive charge that allows negatively charged DNA to stick to the negatively charged cell membrane.

---

5 Long Questions & Answers

1. **Q: Compare and contrast Lipofection and Calcium Phosphate precipitation.**
   A: Both are chemical methods. Lipofection uses cationic lipids to form liposomes that encapsulate DNA; it is very efficient and works on primary cells but is more expensive. Calcium Phosphate involves forming an insoluble salt-DNA complex. It is the cheapest method and works well for large-scale transient transfection in robust cell lines like HEK293, but it is sensitive to small changes in pH and is less effective on delicate cells.
2. **Q: Explain the mechanism and application of Electroporation in animal biotechnology.**
   A: Electroporation uses an electrical field to induce a voltage across the cell membrane. When this voltage reaches a "critical threshold," the membrane undergoes a dielectric breakdown, forming temporary pores. DNA moves through these pores driven by electrophoresis and

diffusion. It is widely used for cells that are "hard to transfect," such as stem cells and lymphoid cells.

3. **Q: Describe the use of Retroviral vectors in animal cell transformation.**
   A: Retroviruses (like HIV-based Lentiviruses) are engineered to be replication-deficient. They carry the gene of interest as RNA. Once they enter the cell, the RNA is reverse-transcribed into DNA and then integrated into the host cell's genome by the enzyme Integrase. This ensures that the gene is passed on to all daughter cells, making it the preferred method for creating stable cell lines and gene therapy.
4. **Q: Why is Microinjection considered the "Gold Standard" for embryos?**
   A: Microinjection is the only method where the scientist has 100% control over the dose and the exact location (the nucleus) of the DNA delivery. Because it bypasses the cytoplasm and membrane barriers physically, the success rate per injected cell is extremely high compared to other methods. It is the primary technique for producing transgenic animals
5. **Q: What are the primary factors that influence the choice of a transformation strategy?**
   A: The choice depends on: (1) Cell type (primary cells vs. established lines), (2) Duration (transient vs. stable), (3) Efficiency required, (4) Toxicity (some chemicals or electricity kill cells), and (5) Budget (Microinjection requires expensive equipment, while Calcium Phosphate is nearly free).

---

15 Multiple Choice Questions (MCQs)

1. The introduction of DNA into animal cells using viral vectors is called:

   a) Transfection

   b) Transduction ✓

   c) Translation

   d) Transcription

2. Which chemical method uses a positively charged polymer?

   a) Lipofection

   b) Calcium Phosphate

   c) DEAE-Dextran ✓

   d) Microinjection

3. Liposomes are made of:

   a) Proteins

   b) Sugars

c) Lipids ✓

d) DNA

4. In Electroporation, DNA enters the cell through:

a) Receptors

b) Nanopores ✓

c) Ion channels

d) Vesicles

5. Which method is most labor-intensive?

a) Microinjection ✓

b) Lipofection

c) Electroporation

d) Calcium Phosphate

6. Calcium Phosphate precipitation depends heavily on the _____ of the buffer.

a) Temperature

b) pH ✓

c) Color

d) Volume

7. Lentiviruses are a type of:

a) Retrovirus ✓

b) Adenovirus

c) Bacteria

d) Plant virus

8. Cationic lipids are _____ charged.

a) Positively ✓

b) Negatively

c) Neutrally

d) Not

9. Stable transformation requires the DNA to:

a) Stay in the cytoplasm

b) Integrate into the genome ✓

c) Be destroyed

d) Stay outside the cell

10. Biolistics in animals is often used for _____ delivery.

a) In vivo (into living tissue) ✓

b) Liquid culture

c) Bacterial

d) Dead cell

11. Adenoviral vectors are typically used for:

a) Stable integration

b) Transient expression ✓

c) Killing cells

d) Making yogurt

12. The "Male Pronucleus" is a target for:

a) Microinjection ✓

b) Lipofection

c) Electroporation

d) DEAE-Dextran

13. Which method is the most cost-effective for large-scale work?

a) Lipofection

b) Calcium Phosphate ✓

c) Microinjection

d) Gene Gun

14. Endocytosis is the primary uptake mechanism for:

a) Calcium Phosphate precipitates ✓

b) Electroporation

c) Microinjection

d) Biolistics

15. Transfection efficiency refers to the:

    a) Speed of DNA replication

    b) Percentage of cells that take up the DNA ✓

    c) Amount of DNA used

    d) Size of the cell

Topic 146 – Chemical Transfection Techniques

1. Principles of Chemical Transfection

Animal cell membranes and DNA are both **negatively charged**. Since like charges repel, DNA cannot naturally cross the lipid bilayer. Chemical transfection methods work by:

1. Neutralizing the negative charge of DNA.
2. Forming a complex that the cell can internalize via **endocytosis** or **membrane fusion**.

2. Calcium Phosphate (CaPO_4) Precipitation

This is one of the oldest and most cost-effective methods.

- **The Process:** DNA is mixed with Calcium Chloride (CaCl_2) in a Phosphate buffer. A fine precipitate of Calcium Phosphate-DNA forms.
- **Mechanism:** The insoluble particles settle onto the cell surface and are taken up by the cell through **phagocytosis** or **endocytosis**.
- **Pros/Cons:** Very cheap; works well for robust cells like HEK293. However, it is highly sensitive to pH changes (optimal pH is 6.95–7.05) and can be toxic to delicate cells.

3. Lipofection (Liposome-Mediated Transfection)

This is the "Gold Standard" for modern chemical transfection.

- **The Process:** DNA is mixed with **Cationic Lipids** (positively charged lipids). These lipids spontaneously form a sphere called a **Liposome** around the DNA.
- **Mechanism:** The positively charged liposome is attracted to the negative cell membrane. It then fuses with the membrane or is taken up via endocytosis, releasing DNA into the cytoplasm.
- **Pros/Cons:** High efficiency and low toxicity; works on a wide range of cell types. The main drawback is the high cost of commercial lipid reagents.

4. DEAE-Dextran Method

- **The Process:** Uses a large, positively charged carbohydrate polymer called Diethylaminoethyl-dextran.

- **Mechanism:** The polymer binds to the DNA, giving it an overall positive charge, which helps it stick to the cell membrane for uptake.
- **Pros/Cons:** Simple and repeatable, but primarily used for **transient transfection** (temporary expression) as it is not efficient for creating stable cell lines.

---

10 Short Questions & Answers

1. **Q: Why do we need chemicals for transfection? A:** To neutralize the negative charge of DNA so it can pass through the negative cell membrane.
2. **Q: What is the ideal pH for Calcium Phosphate transfection? A:** It is very strict, usually between **6.95 and 7.05**.
3. **Q: What is a Liposome? A:** A small artificial vesicle made of a lipid bilayer used to deliver DNA or drugs into cells.
4. **Q: Define "Endocytosis." A:** The cellular process where the membrane folds inward to bring external materials into the cell.
5. **Q: Which chemical method is the most expensive? A:** Lipofection (Liposome-mediated).
6. **Q: What is the main use of DEAE-Dextran? A:** It is used mainly for transient expression in certain animal cell lines.
7. **Q: Name the two ions that react to form the precipitate in the CaPO_4 method. A:** Calcium ions (Ca^{2+}) and Phosphate ions (PO_4^{3-}).
8. **Q: What are "Cationic Lipids"? A:** Lipids with a positive charge that interact with the negatively charged backbone of DNA.
9. **Q: Why is Calcium Phosphate transfection often called "Inexpensive"? A:** It uses basic lab chemicals (CaCl_2 and HEPES buffer) rather than patented commercial kits.
10. **Q: What is the fate of DNA in "Transient Transfection"? A:** It remains in the nucleus as an episome and is eventually lost as the cell divides.

---

5 Long Questions & Answers

1. **Q: Compare Calcium Phosphate and Lipofection in terms of mechanism and efficiency.**
   A: Calcium Phosphate relies on the formation of a physical precipitate that cells "eat" via endocytosis; its efficiency is moderate and depends heavily on environmental factors like pH. Lipofection uses chemical affinity; cationic lipids form vesicles that encapsulate DNA and can fuse directly with the cell membrane. Lipofection is generally much more efficient and less stressful for the cells, making it suitable for a wider variety of cell types, including primary cells.
2. **Q: Explain the importance of "Cationic Lipids" in modern biotechnology.**
   A: Cationic lipids are essential because they solve the problem of charge repulsion. They have a hydrophilic head with a positive charge and a hydrophobic tail. The head binds to the DNA, while the tails form a protective bilayer (liposome). This protects the DNA from being degraded by enzymes (nucleases) and provides a "greasy" exterior that can easily slide through the cell's lipid-based membrane.

3. **Q: Describe the step-by-step procedure of Calcium Phosphate transfection.**
   A: (1) The DNA is dissolved in a sterile water/calcium chloride solution. (2) This mixture is added drop-wise to a buffered saline solution containing phosphate (HEPES-buffered saline). (3) The mixture is incubated for 20 minutes to allow the fine precipitate to form. (4) The precipitate is added to the cell culture medium. (5) After several hours, the cells are washed to remove excess precipitate to prevent toxicity, and the medium is replaced.
4. **Q: What are the limitations of using DEAE-Dextran for animal cell transformation?**
   A: While DEAE-Dextran is simple, it has two major limitations: Toxicity and Integration. At high concentrations, the polymer can damage the cell membrane. Furthermore, it is very poor at facilitating the integration of the foreign DNA into the host genome. Therefore, if a scientist needs a "Stable" cell line that expresses a gene forever, DEAE-Dextran is rarely used.
5. **Q: Why is the choice of chemical method critical for industrial protein production?**
   A: In industry (like making vaccines or antibodies), efficiency and cost are balanced. For massive 100-liter bioreactors, lipofection might be too expensive. Companies often optimize Calcium Phosphate or use polymers like Polyethylenimine (PEI) because they are scalable. The method must ensure high protein yield without killing the cells that are the "factories" for the product.

---

15 Multiple Choice Questions (MCQs)

1. Animal cell membranes are primarily composed of:

   a) Cellulose

   b) Phospholipids ✓

   c) Chitin

   d) Peptidoglycan

2. DNA carries a _____ charge due to its phosphate backbone.

   a) Positive

   b) Negative ✓

   c) Neutral

   d) Variable

3. The "Cationic" part of a lipid refers to its:

   a) Positive charge ✓

   b) Negative charge

   c) Water-fearing nature

   d) Ability to glow

4. Which method involves the formation of an insoluble salt?
    a) Calcium Phosphate ✓
    b) Lipofection
    c) DEAE-Dextran
    d) Electroporation
5. DEAE-Dextran is a positively charged:
    a) Lipid
    b) Protein
    c) Carbohydrate polymer ✓
    d) Virus
6. Lipofection involves DNA being encapsulated in a:
    a) Nucleus
    b) Liposome ✓
    c) Cell wall
    d) Protein coat
7. Phagocytosis is a type of:
    a) Endocytosis ✓
    b) Exocytosis
    c) Diffusion
    d) Active transport
8. Calcium Phosphate transfection is highly sensitive to:
    a) Light
    b) pH ✓
    c) Sound
    d) Oxygen levels
9. The term "Transfection" usually refers to gene transfer in:
    a) Bacteria
    b) Plants

c) Animal cells ✓

d) Fungi

10. A "Stable" transfection means the DNA has:

a) Integrated into the host genome ✓

b) Dissolved in the cytoplasm

c) Been rejected by the cell

d) Turned into a virus

11. Which method is considered most cost-effective for large-scale work?

a) Calcium Phosphate ✓

b) Lipofection

c) Microinjection

d) Gene Gun

12. Commercial reagents like "Lipofectamine" are used for:

a) CaPO_4 method

b) Lipofection ✓

c) DEAE-Dextran

d) SCNT

13. Chemical methods primarily rely on _____ to enter the cell.

a) Physical holes

b) The endocytic pathway ✓

c) Viral infection

d) Magnetic force

14. In the CaPO_4 method, the DNA is found:

a) Inside the Calcium atoms

b) Within the precipitate matrix ✓

c) Attached to a virus

d) In the lipid bilayer

15. The primary disadvantage of DEAE-Dextran is its:

a) High cost

b) Use of heavy metals

c) Inability to produce stable cell lines ✓

d) Need for expensive lasers

Topic 147 – Transfection with Polyplexes

1. What are Polyplexes?

A **polyplex** is a complex formed by the electrostatic interaction between negatively charged DNA and positively charged polymers (cationic polymers). Unlike lipoplexes (which use lipids), polyplexes do not contain a lipid component.

2. Common Cationic Polymers

Several polymers are used to create polyplexes, with the most common being:

- **Polyethylenimine (PEI):** The "Gold Standard" for polyplexes. It has a high "cationic charge density."
- **Poly-L-Lysine (PLL):** One of the first polymers used, though it can be somewhat toxic to cells.
- **Chitosan:** A natural, biocompatible polymer derived from shellfish shells, often used for non-toxic delivery.
- **Dendrimers:** Highly branched, tree-like macromolecules that provide a very precise structure for DNA binding.

3. The "Proton Sponge Effect"

This is the most critical concept for Topic 147. Polyplexes often enter the cell via endocytosis and get trapped in **endosomes**.

1. As the endosome tries to become acidic by pumping in protons (H+), the polymer (especially PEI) acts like a sponge and absorbs them.
2. To maintain electrical balance, Chloride ions (Cl^-) and water follow the protons into the endosome.
3. The endosome swells and eventually **bursts**, releasing the DNA into the cytoplasm before it can be destroyed by lysosomal enzymes.

4. Advantages of Polyplexes

- **Stability:** Polyplexes are generally more stable than lipoplexes.
- **High DNA Loading:** They can condense large amounts of DNA into very small particles.

- **Protection:** The polymer "wraps" the DNA, protecting it from degradation by extracellular nucleases.

---

10 Short Questions & Answers

1. **Q: Define a Polyplex. A:** A complex formed by the interaction of negatively charged DNA and positively charged (cationic) polymers.
2. **Q: Which polymer is considered the "Gold Standard" for polyplex transfection? A:** Polyethylenimine (PEI).
3. **Q: What is the primary charge of the polymers used in this method? A:** Positive (Cationic).
4. **Q: How do polyplexes enter the animal cell? A:** Primarily through endocytosis.
5. **Q: What does PEI stand for? A:** Polyethylenimine.
6. **Q: What is a Dendrimer? A:** A highly branched, symmetric polymer used to form very stable polyplexes.
7. **Q: Why is the "Proton Sponge Effect" important? A:** It allows the DNA to escape the endosome before it is degraded by the cell's "waste disposal" system (lysosomes).
8. **Q: Name a natural polymer used for polyplexes. A:** Chitosan.
9. **Q: What type of bond holds the DNA and polymer together? A:** Electrostatic (ionic) bonds between positive and negative charges.
10. **Q: Are polyplexes more or less stable than liposomes? A:** Generally more stable.

---

5 Long Questions & Answers

1. **Q: Describe the step-by-step mechanism of the "Proton Sponge Effect."**
   A: After the polyplex enters the cell via an endosome, the cell attempts to acidify the endosome using proton pumps. Cationic polymers like PEI have many amino groups that can be protonated; they soak up these protons like a sponge. This prevents the pH from dropping but causes an influx of Cl^- ions to maintain charge neutrality. This leads to high osmotic pressure, causing water to rush into the endosome. The resulting osmotic swelling causes the endosome to rupture, releasing the polyplex into the cytoplasm.
2. **Q: Compare Polyplexes and Lipoplexes (Lipofection).**
   A: Lipoplexes use cationic lipids and rely heavily on membrane fusion or endocytosis; they are very efficient but can be unstable in the presence of serum. Polyplexes use cationic polymers; they are chemically more stable and can condense DNA more tightly. While polyplexes (like PEI) often have better endosomal escape due to the proton sponge effect, they can sometimes be more toxic to cells than modern lipids.
3. **Q: What are the advantages of using Chitosan in gene delivery?**
   A: Chitosan is a natural polysaccharide. Its primary advantage is biocompatibility and low toxicity. Unlike synthetic polymers like PEI, which can damage cell membranes, Chitosan is biodegradable and well-tolerated by living tissues. This makes it an excellent candidate for in vivo (inside the body) gene therapy and mucosal vaccine delivery.

4. **Q: Explain the structural importance of Dendrimers in transfection.**
   A: Dendrimers are unique because they are synthesized with a highly controlled, tree-like branching structure. This "globular" shape allows for a very high density of positive charges on the surface. This structure allows them to condense DNA into very compact, uniform nanoparticles, which improves the efficiency of cellular uptake and protects the DNA more effectively than linear polymers.
5. **Q: Why is "Endosomal Escape" the biggest challenge in chemical transfection?**
   A: When a cell takes in DNA via endocytosis, it treats it like food or a pathogen, moving it into an endosome which eventually fuses with a lysosome. Lysosomes contain acidic enzymes that shred DNA. If the DNA cannot "escape" into the cytoplasm and move to the nucleus quickly, the transfection fails. Polyplexes are specifically engineered to solve this "bottleneck" through mechanisms like the proton sponge effect.

---

15 Multiple Choice Questions (MCQs)

1. A polyplex is a combination of DNA and:

   a) Lipids

   b) Cationic polymers ✓

   c) Gold particles

   d) Sugars

2. Which polymer acts as a "Proton Sponge"?

   a) Polyethylenimine (PEI) ✓

   b) Agarose

   c) Cellulose

   d) Starch

3. Polyplexes enter the cell through:

   a) Direct injection

   b) Osmosis

   c) Endocytosis ✓

   d) Pores in the cell wall

4. The charge of DNA is _____, which allows it to bind to cationic polymers.

   a) Positive

   b) Negative ✓

c) Neutral

d) Changing

5. What follows protons into the endosome during the sponge effect?

a) Chloride ions and Water ✓

b) DNA and RNA

c) Oxygen and Nitrogen

d) Lipids

6. Which polymer is derived from natural sources?

a) PEI

b) PLL

c) Chitosan ✓

d) Dendrimers

7. The "Proton Sponge Effect" leads to endosomal:

a) Shrinkage

b) Rupture/Bursting ✓

c) Fusion

d) Growth

8. Dendrimers are characterized by their:

a) Linear structure

b) Highly branched structure ✓

c) Lack of charge

d) High fat content

9. Poly-L-Lysine (PLL) is a type of:

a) Polypeptide (Polymer of amino acids) ✓

b) Fatty acid

c) Vitamin

d) Mineral

10. The main advantage of polyplexes over lipoplexes is their:

a) Lower cost

b) Increased physical stability ✓

c) Larger size

d) Bright color

11. Endosomal escape must occur before the endosome fuses with a:

a) Nucleus

b) Lysosome ✓

c) Ribosome

d) Mitochondria

12. Cationic polymers bind to DNA via _____ interactions.

a) Electrostatic ✓

b) Covalent

c) Metallic

d) Disulfide

13. Which of the following is NOT a polymer used in polyplexes?

a) PEI

b) Chitosan

c) Dendrimers

d) Cholesterol ✓

14. High osmotic pressure in the endosome is caused by an influx of:

a) Protons only

b) Water ✓

c) DNA

d) Amino acids

15. Transfection with polyplexes is a type of _____ gene delivery.

a) Non-viral ✓

b) Biological

c) Physical

d) Spontaneous

Topic 148 – Liposomes and Lipoplexes

1. Understanding the Components

- **Liposomes:** These are artificial, spherical vesicles composed of one or more lipid bilayers. They have a hollow core that can carry aqueous solutions (DNA/RNA).
- **Cationic Lipids:** These are specialized synthetic lipids with a **positively charged** head group and a hydrophobic tail.
- **Lipoplexes:** When cationic lipids are mixed with negatively charged DNA, they spontaneously organize into a stable complex called a **Lipoplex**.

2. The Mechanism of Lipofection

The process follows four main steps:

1. **Complex Formation:** The positive heads of the lipids bind to the negative phosphate backbone of the DNA, "packaging" the DNA.
2. **Cell Attachment:** The resulting lipoplex has an overall positive charge, which allows it to stick to the negatively charged surface of the animal cell membrane.
3. **Uptake:** The cell takes in the complex primarily through **Endocytosis** or by direct **Membrane Fusion** (since the lipoplex is made of the same material as the cell membrane).
4. **Release:** The DNA is released from the endosome into the cytoplasm and eventually moves into the nucleus.

3. Advantages of Lipofection

- **High Efficiency:** It is much more effective than Calcium Phosphate for many cell types.
- **Low Toxicity:** Modern lipid formulations are designed to be "gentle" on cells.
- **Versatility:** Can deliver DNA of almost any size and can also deliver RNA or proteins.
- **Ease of Use:** Requires simple mixing of reagents without complex equipment like a Gene Gun.

4. Limitations

- **Cost:** Commercial lipofection reagents (like Lipofectamine) are expensive.
- **Serum Interference:** Proteins found in blood serum can sometimes coat the lipoplex and prevent it from entering the cell.

---

10 Short Questions & Answers

1. **Q: What is Lipofection? A:** A transfection method using cationic lipids to deliver DNA into cells.
2. **Q: Why are "Cationic" lipids used? A:** Because their positive charge allows them to bind to negatively charged DNA.
3. **Q: What is a Lipoplex? A:** A complex formed between cationic lipids and nucleic acids.

4. **Q: Name the two ways a lipoplex enters a cell. A:** Endocytosis and direct membrane fusion.
5. **Q: Are liposomes natural or artificial? A:** They are artificial vesicles.
6. **Q: What is the charge of an animal cell membrane? A:** Negatively charged.
7. **Q: Why is lipofection better than chemical precipitation? A:** It generally offers higher efficiency and lower cell death.
8. **Q: Can lipofection be used for RNA delivery? A:** Yes, it is commonly used for delivering mRNA and siRNA.
9. **Q: What is the main barrier to DNA after it enters the cell? A:** Escaping the endosome before it is degraded by lysosomes.
10. **Q: Define "Vesicle." A:** A small fluid-filled bladder or cyst within the body or a cell.

---

5 Long Questions & Answers

1. **Q: Explain the structural difference between a simple liposome and a lipoplex. A:** A **liposome** is a simple lipid bilayer sphere surrounding an aqueous core. It is often used to carry drugs. A **lipoplex** is a more complex arrangement. When cationic lipids meet DNA, the DNA acts as a scaffold. The lipids wrap around the DNA strands, often forming a "sandwich" structure (lamellar) or a "honeycomb" structure (hexagonal). This complex protects the DNA from being "chewed up" by enzymes outside the cell.
2. **Q: Discuss the importance of the charge-to-charge ratio in lipofection experiments. A:** For successful transfection, the ratio of positive charges (from the lipid) to negative charges (from the DNA) must be carefully balanced. Usually, a slight **excess of positive charge** is desired. This ensures that all the DNA is completely "hidden" within the lipid and that the resulting lipoplex has a net positive charge to attract it to the negative cell surface. If there is too much DNA, the complex will be negative and will be repelled by the cell.
3. **Q: Describe the process of Endosomal Escape in lipofection. A:** Once the lipoplex is taken into the cell via endocytosis, it is trapped in a bubble called an endosome. To be successful, the DNA must get out. Cationic lipids are thought to interact with the endosomal membrane, causing it to "flip" or destabilize. This allows the DNA to "leak" into the cytoplasm. If the DNA fails to escape, the endosome will fuse with a lysosome, and the DNA will be destroyed by acid and enzymes.
4. **Q: Compare Lipofection with Transfection using Polyplexes (Topic 147). A: Lipofection** uses lipids and relies heavily on membrane fusion and lipid-membrane interactions. It is generally very efficient for a wide range of cells. **Polyplexes** use cationic polymers (like PEI) and rely on the "Proton Sponge Effect" to escape endosomes. While polyplexes are often cheaper and more stable, lipofection is often preferred for sensitive primary cells because it is less toxic to the cell's internal environment.
5. **Q: Why is Lipofection a preferred method in pharmaceutical research? A:** In drug development, scientists need to test genes in many different cell types. Lipofection is highly **reproducible** and **scalable**. It doesn't require the specialized training of microinjection or the high-voltage risks of electroporation. Furthermore, since it can deliver mRNA, it is a key technology in the development of modern mRNA vaccines and gene therapies.

15 Multiple Choice Questions (MCQs)

1. The term "Lipofection" was coined to describe:
   a) Protein synthesis
   b) Lipid-mediated DNA delivery ✓
   c) Cell division
   d) Viral infection
2. Cationic lipids have a head group that is:
   a) Positively charged ✓
   b) Negatively charged
   c) Neutrally charged
   d) Hydrophobic
3. A lipoplex is formed via _____ interactions.
   a) Covalent
   b) Electrostatic ✓
   c) Metallic
   d) Disulfide
4. Which of the following is an artificial lipid vesicle?
   a) Liposome ✓
   b) Nucleus
   c) Ribosome
   d) Cell wall
5. Lipoplexes stick to cell membranes because membranes are:
   a) Positively charged
   b) Negatively charged ✓
   c) Made of wood
   d) Very hot
6. The most common pathway for lipoplex entry is:
   a) Simple diffusion

b) Endocytosis ✓

c) Active transport through pumps

d) Osmosis

7. "Lipofectamine" is a well-known:

a) Antibiotic

b) Commercial transfection reagent ✓

c) Type of sugar

d) Laboratory glass

8. DNA is protected within a lipoplex from:

a) Nucleases (degrading enzymes) ✓

b) Sunlight

c) Gravity

d) Water

9. Fusion refers to the lipoplex merging with the:

a) Nucleus

b) Plasma membrane ✓

c) Mitochondria

d) Vacuole

10. Lipofection is generally _____ toxic than Calcium Phosphate.

a) More

b) Less ✓

c) Equally

d) Extremely

11. Excess _____ charge is usually required for the lipoplex to bind to cells.

a) Positive ✓

b) Negative

c) Neutral

d) Nuclear

12. Serum interference can _____ transfection efficiency.
    a) Increase
    b) Decrease ✓
    c) Stabilize
    d) Not change
13. Which part of the lipid interacts with the DNA?
    a) Hydrophilic head ✓
    b) Hydrophobic tail
    c) The entire molecule
    d) None
14. Lipofection can be used to deliver:
    a) DNA
    b) RNA
    c) Proteins
    d) All of the above ✓
15. Endosomal escape allows DNA to enter the:
    a) Cytoplasm ✓
    b) Extracellular space
    c) Cell wall
    d) Bloodstream

Topic 149 – Physical Transfection Techniques

1. Introduction to Physical Methods

Physical transfection involves the use of external physical force to create temporary openings in the plasma membrane or to directly "inject" DNA into the cell. These methods are generally highly efficient but often require specialized, expensive equipment.

2. Electroporation

This is the most widely used physical method for animal cells.

- **Mechanism:** Cells are subjected to a high-voltage electrical pulse. This creates temporary "nanopores" (dielectric breakdown) in the lipid bilayer.
- **DNA Entry:** The negatively charged DNA moves through these pores driven by both diffusion and electrophoresis.
- **Optimization:** Success depends on the **pulse duration** and **field strength**. If the pulse is too strong, the cells die; if too weak, no pores form.

3. Microinjection

The most precise and direct method of gene delivery.

- **Mechanism:** A microscopic glass needle (micropipette) is used to physically pierce the cell and nuclear membranes.
- **Application:** It is the "Gold Standard" for producing transgenic animals. DNA is usually injected into the **male pronucleus** of a fertilized zygote.
- **Efficiency:** Nearly 100% of the cells successfully injected will express the gene, but it can only be done one cell at a time.

4. Biolistics (Gene Gun)

Originally developed for plants, but adapted for animal tissues.

- **Mechanism:** Heavy metal microparticles (gold or tungsten) are coated with DNA and accelerated to high speeds using compressed helium.
- **Application:** Useful for transforming skin cells, muscle tissue, or organs *in vivo* (in a living organism).

5. Sonoporation and Laserfection

- **Sonoporation:** Uses ultrasound waves to create pores (cavitation) in the cell membrane.
- **Laserfection (Optoporation):** Uses a highly focused laser beam to "drill" a tiny hole in the cell membrane for a fraction of a second.

---

10 Short Questions & Answers

1. **Q: What is the primary advantage of physical transfection? A:** It can transform cells that are resistant to chemical methods (like primary cells or stem cells).
2. **Q: What are "nanopores" in electroporation? A:** Temporary, microscopic holes in the cell membrane created by an electric field.
3. **Q: Which physical method is best for transforming a single embryo? A:** Microinjection.
4. **Q: What gas is usually used to power the Gene Gun? A:** Helium.
5. **Q: Define "Dielectric Breakdown." A:** The process where an insulator (like the cell membrane) becomes conductive due to a high electric field, leading to pore formation.
6. **Q: Why is Gold preferred over Tungsten in Biolistics? A:** Gold is chemically inert (non-toxic) and has a uniform spherical shape.

7. **Q: What is the main drawback of Electroporation? A:** High cell mortality; many cells do not survive the electric shock.
8. **Q: What is "In vivo" transfection? A:** Introducing DNA into the cells of a living animal rather than in a lab dish.
9. **Q: What does "Sonoporation" use to create pores? A:** Ultrasound waves.
10. **Q: In microinjection, where is the DNA typically delivered? A:** Directly into the nucleus (or male pronucleus).

---

5 Long Questions & Answers

1. **Q: Explain the factors that must be balanced to achieve successful Electroporation.**
   A: The two most critical parameters are Field Strength (voltage) and Pulse Length (time). Field strength determines the size and number of pores, while pulse length determines how long they stay open. There is an inverse relationship between efficiency and cell viability. If the voltage is too high or the pulse too long, the pores become permanent, and the cell's contents leak out, leading to death. The "optimal" settings are those that allow maximum DNA entry while keeping at least 50-70% of the cells alive.
2. **Q: Why is Microinjection the preferred method for creating transgenic animals?**
   A: While chemical methods or electroporation are used for populations of cells, microinjection is a "surgery" on a single cell. Because the scientist manually places the DNA into the male pronucleus of a zygote, they ensure that the gene is present at the very first stage of life. This ensures that as the zygote divides, every single cell in the resulting animal (including the sperm or eggs) will contain the transgene, which is essential for "germline" transmission.
3. **Q: Describe the mechanism and limitations of Biolistics in animal biotechnology.**
   A: Biolistics involves coating heavy metal micro-carriers with DNA and "firing" them into target tissue. It is excellent for "shallow" tissues like skin or exposed organs during surgery. However, its limitations include: (1) Low penetration depth, as the particles cannot travel far through dense tissue. (2) Physical damage to the cells at the center of the "blast" zone. (3) The specialized equipment and consumables (gold/helium) are very expensive.
4. **Q: Contrast Sonoporation and Laserfection.**
   A: Sonoporation uses ultrasound to create "cavitation bubbles." When these bubbles collapse near a cell, the energy ruptures the membrane briefly. It is non-invasive and can be used for deep tissue delivery. Laserfection uses a precise laser (femtosecond laser) to create a single, specific hole in the membrane. Laserfection is much more precise and has a higher cell survival rate, but it is much slower as it targets cells individually, similar to microinjection.
5. **Q: Evaluate the pros and cons of Physical methods versus Chemical methods.**
   A: Pros of Physical: They are "universal" (not dependent on cell receptors or charge); they are generally more efficient for hard-to-transfect cells; and they don't involve potentially toxic chemicals. Cons of Physical: They require expensive machinery (Electroporators, Gene Guns, Micromanipulators); they often cause more physical damage to the cells; and they are often lower "throughput" (harder to do millions of cells quickly compared to adding a chemical to a flask).

---

15 Multiple Choice Questions (MCQs)

1. Electroporation creates temporary holes in the:
   a) Nucleus
   b) Plasma membrane ✓
   c) Mitochondria
   d) Ribosome
2. The microscopic needle used in microinjection is made of:
   a) Glass ✓
   b) Steel
   c) Plastic
   d) Gold
3. Which method is used to deliver DNA to skin or muscle tissues in vivo?
   a) Calcium Phosphate
   b) DEAE-Dextran
   c) Biolistics (Gene Gun) ✓
   d) Lipofection
4. In Electroporation, DNA moves into the cell partly because it is:
   a) Positively charged
   b) Negatively charged ✓
   c) Neutral
   d) Hydrophobic
5. Cavitation is the process associated with:
   a) Electroporation
   b) Sonoporation ✓
   c) Laserfection
   d) Microinjection
6. The most inert and common micro-carrier for Biolistics is:
   a) Gold ✓
   b) Iron

c) Aluminum

d) Lead

7. Microinjection is primarily used for the production of:

a) Bacteria

b) Plant seeds

c) Transgenic animals ✓

d) Fungi

8. Which physical method has the highest cell-by-cell precision?

a) Microinjection ✓

b) Electroporation

c) Sonoporation

d) Calcium Phosphate

9. High-voltage pulses in electroporation lead to:

a) Cell growth

b) Mutation

c) Dielectric breakdown ✓

d) DNA replication

10. A femtosecond laser is used in:

a) Sonoporation

b) Laserfection ✓

c) Biolistics

d) Electroporation

11. Physical methods are often called "Mechanical" because they:

a) Use force to bypass the membrane ✓

b) Use robots

c) Only work on machines

d) Use cold temperatures

12. The biggest disadvantage of Electroporation is:

a) It's too slow

b) It's too cheap

c) High cell mortality ✓

d) It only works in plants

13. Compressed _____ gas is the power source for a Gene Gun.

a) Oxygen

b) Carbon Dioxide

c) Helium ✓

d) Nitrogen

14. In microinjection, a "Holding Pipette" is used to:

a) Keep the cell stationary ✓

b) Suck out the DNA

c) Heat the cell

d) Kill the cell

15. Which of the following is NOT a physical transfection method?

a) Electroporation

b) Microinjection

c) Lipofection ✓

d) Biolistics

Topic 150 – Microinjection

1. Definition and Concept

Microinjection is a mechanical method where a microscopic glass needle (micropipette) is used to physically penetrate the cell membrane and nuclear envelope to deliver genetic material (DNA, RNA) or proteins directly into a specific compartment of a single cell.

2. The Apparatus (The Micromanipulator)

Since human hands are too "shaky" for microscopic work, a **Micromanipulator** is used. It converts large movements of the hand into microscopic movements of the needle.

- **Holding Pipette:** A blunt-ended pipette that uses gentle suction to hold the target cell (like an egg) in place.

- **Injection Needle:** A needle with a tip diameter of roughly 0.5 to 5.0 µ.
- **Inverted Microscope:** Used to visualize the cells from below, providing space for the needles to move above.

3. The Procedure for Transgenic Animals

1. **Preparation:** A female animal is superovulated to produce multiple eggs, which are then fertilized.
2. **Targeting:** The fertilized egg (zygote) is held by the holding pipette.
3. **The Male Pronucleus:** Scientists target the **male pronucleus** (the nucleus from the sperm) because it is larger and more visible than the female one.
4. **Injection:** The needle pierces the pronucleus, and a specific volume of DNA (usually a few picoliters) is injected.
5. **Implantation:** The surviving embryos are transferred into a pseudo-pregnant surrogate mother.

4. Advantages and Limitations

- **Advantages:** * No need for a viral vector or chemical carrier.
    - High efficiency per cell (nearly 100% delivery).
    - Direct delivery into the nucleus ensures better integration.
- **Limitations:**
    - **Low Throughput:** Only one cell can be treated at a time.
    - **Technical Skill:** Requires highly trained personnel.
    - **Cost:** The equipment is extremely expensive.
    - **Physical Damage:** The puncture can sometimes kill the cell.

---

10 Short Questions & Answers

1. **Q: What is a micromanipulator? A:** A device that allows for precise, microscopic movement of needles during the injection process.
2. **Q: Why is the male pronucleus preferred for injection? A:** It is physically larger than the female pronucleus, making it an easier target.
3. **Q: What is the average tip diameter of an injection needle? A:** Usually between 0.5 and 1.0 µ.
4. **Q: What is the role of the holding pipette? A:** It uses suction to keep the cell stationary so it does not move when the needle strikes.
5. **Q: Define "Superovulation." A:** Using hormones to make a female produce more eggs than she normally would in a single cycle.
6. **Q: What volume of DNA is typically injected? A:** Very small amounts, measured in **picoliters** (10^-12 liters).
7. **Q: Can microinjection be used for things other than DNA? A:** Yes, it can deliver RNA, proteins, or even organelles into a cell.
8. **Q: What type of microscope is used? A:** An **inverted microscope**.

9. **Q: What is a "pseudo-pregnant" mother? A:** A female animal that has been mated with a sterile male to prepare her uterus for embryo implantation.
10. **Q: What is the main success indicator for a successful injection? A:** A slight "swelling" of the pronucleus upon fluid entry.

---

5 Long Questions & Answers

1. **Q: Explain why microinjection is used for germline transformation rather than simple cell culture.**
   A: In cell culture, we only need a few cells to take up the DNA. However, to make a transgenic animal, the DNA must be in every cell of the body. By injecting a zygote (a single-cell embryo), the foreign DNA is replicated along with the animal's own DNA during every subsequent cell division. This ensures the trait is in the "germline" (sperm/eggs), allowing the trait to be passed to future generations.
2. **Q: Describe the physical challenges and risks associated with the microinjection proces**s.
   A: The primary challenge is the fragility of the cell. Piercing the plasma membrane and the nuclear envelope is a violent act at the cellular level. If the needle is too thick, or if the pressure of the injection is too high, the cell will lyse (burst). Additionally, the DNA must integrate into the host genome; if it simply floats in the nucleus, it will be lost. This leads to a low overall survival rate (often only 10–20% of injected eggs become live transgenic animals).
3. **Q: Compare Microinjection with Electroporation**
   A: Electroporation is a "bulk" method; it can treat millions of cells at once by using electricity to open pores. It is faster but has a higher cell death rate and lower precision. Microinjection is a "bespoke" method; it targets individual cells with 100% delivery success to that specific cell. While microinjection is much slower and requires more expensive gear, it is the only reliable way to ensure a specific embryo receives a specific gene.
4. **Q: Discuss the importance of the Micromanipulator in biotechnology.**
   A: The micromanipulator acts as the interface between the macro-world and the micro-world. It uses mechanical or hydraulic linkages to scale down hand movements. Without this, the fine control required to hit a pronucleus (which is only a few microns wide) would be impossible. Modern manipulators often include "joy-stick" controls and automated pressure systems to ensure that the exact picoliter volume is delivered every time.
5. **Q: What are the requirements for the DNA used in microinjection?**
   A: The DNA must be extremely pure (free of salts and phenol) to avoid poisoning the embryo. It must be "linearized" (cut out of the circular plasmid) because linear DNA fragments integrate into the host genome much more efficiently than circular ones. Usually, the DNA is dissolved in a specific "Injection Buffer" (like TE buffer) that is compatible with the internal environment of the nucleus.

---

15 Multiple Choice Questions (MCQs)

1. Microinjection is a _____ method of gene transfer.
   a) Chemical
   b) Physical/Mechanical ✓
   c) Biological
   d) Spontaneous
2. The DNA is injected into the _____ of the zygote.
   a) Cytoplasm
   b) Male Pronucleus ✓
   c) Mitochondria
   d) Vacuole
3. Which device is used to hold the egg stationary?
   a) Centrifuge
   b) Holding Pipette ✓
   c) Electroporator
   d) Vortex
4. The volume of liquid injected into a cell is measured in:
   a) Milliliters
   b) Microliters
   c) Nanoliters
   d) Picoliters ✓
5. The "Inverted" microscope is used because:
   a) It provides space for the micromanipulator needles ✓
   b) It makes the DNA look bigger
   c) It uses less light
   d) It is cheaper
6. Success in microinjection is visually confirmed by:
   a) The cell turning blue
   b) The pronucleus swelling ✓

c) The cell dividing immediately

d) Bubbles appearing

7. The main advantage of microinjection is:

a) It is very fast

b) It is very cheap

c) High delivery efficiency to the target cell ✓

d) It works on dead cells

8. Superovulation is induced by using:

a) Antibiotics

b) Hormones ✓

c) Sugars

d) Electricity

9. The needle used for injection is usually made of:

a) Borosilicate Glass ✓

b) Stainless Steel

c) Plastic

d) Copper

10. A "Surrogate" mother is used to:

a) Provide the eggs

b) Carry the injected embryo to term ✓

c) Provide the sperm

d) Provide the DNA

11. DNA used for microinjection should be:

a) Circular

b) Linearized and pure ✓

c) In the form of a virus

d) Mixed with lipids

12. The "Male" pronucleus is targeted because it is:

a) Stronger

b) Larger and more visible ✓

c) More receptive to DNA

d) Darker in color

13. Which of the following is NOT a part of a micromanipulation system?

a) Injection needle

b) Holding pipette

c) Gene Gun ✓

d) Inverted microscope

14. The term "Micron" (μ) represents:

a) 10^-3 meters

b) 10^-6 meters ✓

c) 10^-9 meters

d) 10^-12 meters

15. Microinjection is the "Gold Standard" for producing:

a) Recombinant bacteria

b) GM yeast

c) Transgenic animals ✓

d) Vaccines

Topic 151 – Selectable Markers

1. Why are Selectable Markers Necessary?

Even with advanced methods like lipofection or electroporation, only a small percentage (often 1-10%) of animal cells successfully integrate foreign DNA. Selectable markers are genes co-transfected with the gene of interest that allow the researcher to kill off the non-transformed cells, leaving only a pure population of transgenic cells.

2. Types of Selection

- **Positive Selection:** The marker gene provides a new ability to the cell (like antibiotic resistance) that allows it to survive in a medium that kills normal cells.
- **Negative Selection:** The marker gene produces a protein that is toxic to the cell under certain conditions (often used to eliminate cells that have integrated DNA incorrectly).

3. Common Dominant Selectable Markers (Antibiotics)

Since most animal cells are naturally sensitive to certain toxins, providing a resistance gene is the most common strategy:

- **Neomycin Phosphotransferase (*neo*):** Provides resistance to **G418** (Geneticin). This is the most widely used marker in animal biotechnology.
- **Hygromycin B Phosphotransferase (*hph*):** Provides resistance to the antibiotic **Hygromycin B**.
- **Puromycin N-acetyltransferase (*pac*):** Provides resistance to **Puromycin**, which acts very quickly (2-3 days).

4. Metabolic/Amplifiable Markers

These markers use specific enzyme deficiencies or "overdrive" mechanisms:

- **Dihydrofolate Reductase (DHFR):** Cells are grown in a medium lacking thymidine. If the *DHFR* gene is present, they survive.
- **Methotrexate (MTX) Amplification:** By increasing the concentration of the drug MTX, the cell is forced to create multiple copies (amplification) of the *DHFR* gene and the linked "gene of interest," leading to massive protein production.

5. Screenable Markers (Reporters)

Unlike selectable markers that kill cells, screenable markers make the transformed cells look different:

- **Green Fluorescent Protein (GFP):** Cells glow green under UV light.
- **Luciferase:** Cells produce light through a chemical reaction.

---

10 Short Questions & Answers

1. **Q: What is a selectable marker? A:** A gene introduced into a cell that confers a trait suitable for artificial selection.
2. **Q: What antibiotic does the *neo* gene provide resistance to? A:** G418 (also known as Geneticin).
3. **Q: How does Puromycin kill cells? A:** It inhibits protein synthesis by causing premature chain termination.
4. **Q: What is "Gene Amplification"? A:** The process of increasing the copy number of a specific gene within a cell to increase protein yield.
5. **Q: Why is GFP considered a "Screenable" rather than "Selectable" marker? A:** Because it allows you to see transformed cells but doesn't kill the non-transformed ones.
6. **Q: What is the drug used to select for the *DHFR* gene? A:** Methotrexate (MTX).
7. **Q: Define "Co-transfection." A:** The simultaneous introduction of two or more genes (e.g., a gene of interest and a marker gene) into the same cell.
8. **Q: What is a "Dominant" selectable marker? A:** A marker that works in wild-type cells without needing a specific pre-existing mutation.

9. **Q: How long does G418 selection typically take? A:** Usually 10 to 14 days.
10. **Q: Name a marker used for negative selection. A:** Thymidine Kinase (*tk*) from the Herpes Simplex Virus.

---

5 Long Questions & Answers

1. **Q: Compare and contrast Selectable Markers and Reporter Genes.**
   A: Selectable markers (like neo or dhfr) are used to physically isolate transformed cells by killing those that lack the gene. They are essential for creating stable cell lines. Reporter genes (like GFP or LacZ) are used to "report" where and when a gene is active. They do not kill cells but provide a visual or measurable signal (fluorescence or color change). Reporters are often used for transient assays to check if a promoter is working.
2. **Q: Explain the mechanism of DHFR selection and its role in gene amplification.**
   A: DHFR is an enzyme essential for DNA synthesis. In cells lacking DHFR, they cannot grow unless provided with thymidine. When a DHFR gene is transfected into these cells, they can grow in thymidine-free medium. Furthermore, if the drug Methotrexate (MTX) is added, it inhibits DHFR. Only cells that "amplify" (copy many times) the DHFR gene can survive the drug. Because the gene of interest is linked to DHFR, it gets amplified too, leading to very high protein production.
3. **Q: Why is G418 (Geneticin) the most common selection agent in animal labs?**
   A: G418 is an aminoglycoside antibiotic similar to neomycin, but unlike neomycin, it is highly effective at killing eukaryotic (animal) cells by interfering with the 80S ribosome. The neo gene produces an enzyme that phosphorylates G418, making it harmless. It is popular because it is reliable, the selection pressure is strong, and the neo gene is small and easy to incorporate into many different expression vectors.
4. **Q: Describe the process of creating a "Stable Cell Line" using selectable markers.**
   A: First, the gene of interest and a selectable marker (e.g., puromycin resistance) are transfected into a population of cells. For the first 24-48 hours, the cells express the genes "transiently." Then, the selection drug is added to the medium. Cells that did not take up the DNA or failed to integrate it into their chromosomes die. Over 1-2 weeks, the surviving cells grow into "colonies." These colonies are then expanded, resulting in a stable cell line where every cell contains the new gene.
5. **Q: What is the "Dual Marker" strategy in gene targeting?**
   A: In advanced experiments like Gene Knockouts, scientists use a positive marker (like neo) inside the target gene and a negative marker (like HSV-tk) at the end of the DNA fragment. If the DNA integrates correctly (homologous recombination), the tk gene is lost, and the cell lives in G418. If it integrates randomly, the tk gene is kept, and the cell dies when a specific drug (Ganciclovir) is added. This ensures only the "perfect" genetic edits survive.

---

15 Multiple Choice Questions (MCQs)

1. Selectable markers are used to identify:
   a) All cells in a dish
   b) Only cells that have taken up the transgene ✓
   c) Only dead cells
   d) Viral particles
2. The neo gene provides resistance to which drug?
   a) Penicillin
   b) Ampicillin
   c) G418 (Geneticin) ✓
   d) Methotrexate
3. Which marker causes cells to glow green?
   a) GFP ✓
   b) DHFR
   c) PAC
   d) Neo
4. Methotrexate (MTX) is an inhibitor of:
   a) Protein synthesis
   b) RNA polymerase
   c) Dihydrofolate Reductase (DHFR) ✓
   d) Cell wall synthesis
5. Selection with Puromycin is usually faster than G418, taking about:
   a) 2-3 days ✓
   b) 2-3 months
   c) 24 hours
   d) 4 weeks
6. "Negative Selection" aims to:
   a) Keep all cells alive
   b) Kill cells containing a specific gene ✓

c) Make cells grow faster

d) Change the cell color

7. Aminoglycoside phosphotransferase is the enzyme produced by the _____ gene.

a) Neomycin (neo) ✓

b) Hygromycin (hph)

c) Luciferase

d) DHFR

8. Gene amplification results in:

a) Smaller cells

b) Multiple copies of the transgene ✓

c) Faster cell death

d) Loss of DNA

9. Which of these is a metabolic marker?

a) Puromycin

b) GFP

c) DHFR ✓

d) Geneticin

10. A "Reporter Gene" is primarily used to study:

a) Gene expression and localization ✓

b) Antibiotic resistance

c) Cell motility

d) DNA replication speed

11. G418 interferes with the function of:

a) The nucleus

b) The 80S ribosome ✓

c) The mitochondria

d) The cell membrane

12. The hph gene provides resistance to:

a) Puromycin

b) Hygromycin B ✓

c) Kanamycin

d) Tetracycline

13. In co-transfection, the ratio of the gene of interest to the marker gene is often:

a) 10:1 ✓

b) 1:100

c) 0:1

d) 1:1

14. Luciferase requires a substrate called _____ to produce light.

a) Glucose

b) Luciferin ✓

c) ATP

d) Methotrexate

15. The process of introducing two genes at once is called:

a) Bi-transformation

b) Co-transfection ✓

c) Dual-injection

d) Re-integration

Topic 152 – Bactofection

1. What is Bactofection?

Bactofection is a method of gene transfer where attenuated (weakened) bacteria are used to deliver a functional gene into a target animal cell. Unlike viral vectors, which use a protein shell, bactofection uses an entire living organism to bypass cellular barriers.

2. Commonly Used Bacterial Vectors

The bacteria used are usually **intracellular pathogens** that have been genetically modified to be harmless to the host:

- ***Listeria monocytogenes***
- ***Salmonella typhi***
- ***Shigella flexneri***
- ***Escherichia coli*** **(modified)**

3. The Mechanism of Gene Delivery

The process of bactofection generally follows these steps:

1. **Entry:** The bacteria enter the animal cell through **phagocytosis** or **endocytosis**.
2. **Endosomal Escape:** Once inside the "phagosome" (a bubble inside the cell), the bacteria must escape or be released.
3. **Lysis:** The bacteria are engineered to undergo "self-destruction" (lysis) once they reach the cytoplasm of the host cell.
4. **DNA Release:** Upon lysis, the plasmid DNA (carrying the gene of interest) is released.
5. **Nuclear Entry:** The plasmid moves into the host nucleus, where the animal cell's machinery begins expressing the gene.

4. Advantages of Bacterial Vectors

- **Large Cargo Capacity:** Bacteria can carry very large segments of DNA (much larger than viruses).
- **Targeting:** Some bacteria naturally prefer specific tissues (e.g., *Salmonella* often targets tumor cells).
- **Cost-Effective:** Bacteria are much cheaper and easier to grow in large quantities compared to viral vectors.
- **Safety:** They can be controlled using antibiotics if the infection spreads too far.

5. Limitations and Risks

- **Immunogenicity:** The human immune system may recognize and attack the bacteria before they can deliver the DNA.
- **Toxicity:** Even weakened bacteria can sometimes cause an inflammatory response.

---

10 Short Questions & Answers

1. **Q: Define Bactofection. A:** The use of bacteria to deliver genetic material into eukaryotic (animal) cells.
2. **Q: Name one bacterium commonly used for bactofection. A:** *Salmonella typhi* or *Listeria monocytogenes*.
3. **Q: What is an "Attenuated" bacterium? A:** A bacterium that has been weakened or modified so it cannot cause disease.
4. **Q: How do the bacteria release the DNA inside the cell? A:** Through programmed lysis (self-destruction) of the bacterial cell wall.

5. **Q: What is the primary method of bacterial entry into host cells? A:** Phagocytosis or endocytosis.
6. **Q: Why is *Salmonella* used in cancer research? A:** Because it has a natural tendency to accumulate inside solid tumors.
7. **Q: What is a "Phagosome"? A:** The vesicle formed around a bacterium as it is taken into the host cell.
8. **Q: Can bacteria deliver RNA? A:** Yes, they can be used to deliver DNA, shRNA, or even proteins.
9. **Q: What happens to the bacteria after DNA delivery? A:** They are cleared by the host's immune system or destroyed during the lysis process.
10. **Q: Is bactofection a viral or non-viral method? A:** It is considered a **biological non-viral** method.

---

5 Long Questions & Answers

1. **Q: Describe the molecular modifications required to make a bacterium a safe vector.**
   A: To ensure safety, scientists perform "gene knockouts" on the bacteria. For example, genes responsible for virulence (making the host sick) are removed. Additionally, "auxotrophic" mutations are often introduced—this means the bacteria can only survive if a specific nutrient is provided in the lab. Once inside the animal body, they cannot find that nutrient and will eventually die or self-lyse, preventing a runaway infection.
2. **Q: Explain the significance of "Endosomal Escape" in bactofection.**
   A: When an animal cell "swallows" a bacterium, it traps it in a phagosome/endosome. If the bacterium stays there, it will be destroyed by lysosomes. Vectors like Listeria produce a protein called Listeriolysin O (LLO), which punches holes in the endosome membrane. This allows the bacterium to escape into the cytoplasm, where it can safely release the plasmid DNA for transport to the nucleus.
3. **Q: Compare Bacterial Vectors and Viral Vectors.**
   A: Viral vectors (like Adenovirus) are very efficient but have a small "carrying capacity" for DNA and can be difficult to produce. Bacterial vectors can carry significantly larger plasmids (up to 100 kb or more) and are very inexpensive to produce in fermenters. However, bacteria trigger a much stronger immune response (inflammation) than most modern viral vectors, which limits their use in sensitive tissues.
4. **Q: Discuss the application of bactofection in vaccine development.**
   A: Bacteria are excellent for "DNA Vaccines." By delivering a gene that codes for a viral or bacterial antigen directly into a host's immune cells (like macrophages), the host cell starts producing the antigen. This "tears" the immune system to recognize the pathogen without ever exposing the person to the actual virus. This method provides both humoral and cellular immunity.
5. **Q: What is "Tumor-Targeted Bactofection"?**
   A: Certain anaerobic bacteria like Clostridium or Salmonella thrive in the "hypoxic" (low oxygen) environment found inside solid tumors. By using these as vectors, scientists can deliver "suicide genes" or "anti-angiogenic genes" directly to the heart of a tumor while leaving healthy,

oxygen-rich tissues untouched. This reduces the side effects associated with traditional chemotherapy.

---

15 Multiple Choice Questions (MCQs)

1. Bactofection is a method used to transform:

   a) Bacteria

   b) Animal cells ✓

   c) Plant seeds

   d) Fungi

2. Which of these is a Gram-negative bacterium used as a vector?

   a) Listeria

   b) Salmonella ✓

   c) Bacillus

   d) Staphylococcus

3. The release of DNA from the bacteria occurs due to:

   a) Bacterial Lysis ✓

   b) Bacterial reproduction

   c) Host cell death

   d) Sunlight

4. Attenuated means the bacterium is:

   a) Extra strong

   b) Weakened and non-pathogenic ✓

   c) Dead

   d) Radioactive

5. Bacterial vectors are particularly useful because they can carry:

   a) Large DNA sequences ✓

   b) Only small RNA

   c) No DNA

   d) Only proteins

6. Listeriolysin O (LLO) is used by bacteria to:
    a) Kill the host
    b) Escape the phagosome ✓
    c) Reproduce faster
    d) Attach to the cell wall
7. Bactofection can be controlled using:
    a) Fertilizers
    b) Antibiotics ✓
    c) UV light
    d) High salt
8. The DNA delivered by bacteria is usually in the form of a:
    a) Linear fragment
    b) Circular Plasmid ✓
    c) Single nucleotide
    d) Protein
9. Which organelle is the final destination for the delivered DNA?
    a) Nucleus ✓
    b) Mitochondria
    c) Golgi apparatus
    d) Lysosome
10. Auxotrophic bacteria are those that:
    a) Require a specific nutrient to survive ✓
    b) Can live anywhere
    c) Do not have DNA
    d) Are resistant to all drugs
11. Bactofection is a _____ method of gene delivery.
    a) Physical
    b) Chemical

c) Biological ✓

d) Spontaneous

12. The use of bacteria to treat cancer is effective because tumors are:

a) Very cold

b) Hypoxic (low oxygen) ✓

c) Full of sugar

d) Very small

13. Which bacterium is Gram-positive and escapes the endosome easily?

a) Listeria ✓

b) E. coli

c) Shigella

d) Salmonella

14. Phagocytosis is the process of:

a) Cell eating ✓

b) Cell drinking

c) Cell breathing

d) Cell division

15. A major disadvantage of bacterial vectors is:

a) They are expensive

b) High immunogenicity ✓

c) They are too small

d) They cannot be grown in labs

Topic 153 – Viral Vectors

1. What is a Viral Vector?

A viral vector is a virus that has been genetically engineered to remove its disease-causing (pathogenic) genes and replace them with a therapeutic or experimental "gene of interest." The process of using a virus to deliver DNA into a cell is called **Transduction**.

2. Essential Modifications for Safety

To make a virus a safe vector, scientists ensure it is:

- **Replication-Deficient:** The genes required for the virus to make copies of itself and burst out of the cell are removed.
- **Non-Pathogenic:** The genes that cause sickness are deleted.
- **Helper Systems:** Since the vector cannot replicate on its own, it is produced in "Packaging Cell Lines" that provide the missing viral proteins.

3. Key Types of Viral Vectors

Vector Type	Genetic Material	Integration	Application
Retrovirus	RNA	Yes (Random)	Stable gene expression; dividing cells only.
Lentivirus	RNA	Yes	Subclass of retrovirus; can infect non-dividing cells.
Adenovirus	dsDNA	No (Episomal)	Vaccines (e.g., COVID-19); transient expression.
AAV	ssDNA	Minimal	Gene therapy; low immune response.

4. The Transduction Process

1. **Adsorption:** The virus binds to specific receptors on the animal cell surface.
2. **Entry:** The virus enters via endocytosis or membrane fusion.
3. **Uncoating:** The viral protein coat (capsid) breaks down, releasing the genetic cargo.
4. **Expression:** The DNA travels to the nucleus. If it is an integrating virus (like Lentivirus), it joins the host genome. If not (like Adenovirus), it remains as an **episome**.

5. Advantages and Disadvantages

- **Advantages:** Extremely high efficiency (nearly 100% of cells can be infected); natural ability to target specific cell types (**tropism**).
- **Disadvantages:** Limited cargo capacity (small genes only); risk of **Immune Response** (inflammation); risk of **Insertional Mutagenesis** (disrupting a host gene during integration).

---

10 Short Questions & Answers

1. **Q: Define Transduction. A:** The process of transferring genetic material into a cell using a viral vector.
2. **Q: What does "Replication-Deficient" mean? A:** The virus can enter a cell and deliver DNA but cannot produce new viral particles to infect other cells.
3. **Q: Which viral vector is commonly used for vaccines? A:** Adenovirus.
4. **Q: What is "Viral Tropism"? A:** The specific preference of a virus for infecting a particular cell type or tissue (e.g., liver vs. lung).
5. **Q: Name a virus that can infect non-dividing cells. A:** Lentivirus or Adeno-associated virus (AAV).
6. **Q: What is an "Episome"? A:** DNA that remains in the nucleus but does not integrate into the host's chromosomes.
7. **Q: Why are Retroviruses risky? A:** They integrate randomly into the genome, which might accidentally turn on a cancer-causing gene (oncogene).
8. **Q: What are "Packaging Cells"? A:** Specialized lab cells that provide the proteins needed to assemble a replication-deficient virus.
9. **Q: Which vector is used for stable, long-term gene expression? A:** Lentivirus or Retrovirus.
10. **Q: What is the genetic material of a Lentivirus? A:** RNA (which is reverse-transcribed into DNA after entry).

---

5 Long Questions & Answers

1. **Q: Explain the mechanism of Retroviral integration into the host genome.**
   A: After entering the cell, the retroviral RNA is converted into double-stranded DNA by the enzyme Reverse Transcriptase. This DNA is then transported into the nucleus. An enzyme called Integrase makes a "nick" in the host cell's DNA and inserts the viral DNA. This ensures that the transgene is permanently part of the cell's genetic blueprint and will be passed on to all future daughter cells.
2. **Q: Compare Adenoviruses and Lentiviruses in terms of clinical application.**
   A: Adenoviruses do not integrate into the DNA, meaning the gene expression is temporary (transient). They are excellent for vaccines because they trigger a strong immune response. Lentiviruses integrate into the DNA, providing permanent (stable) expression. They are preferred for treating genetic diseases (like Sickle Cell Anemia) where the patient needs a working gene for their entire life.
3. **Q: What is "Insertional Mutagenesis" and how can it be prevented?**
   A: This occurs when a viral vector integrates its DNA in the middle of a vital host gene,

potentially causing mutations or cancer (leukemia). To prevent this, scientists are developing "Self-Inactivating" (SIN) vectors or using AAVs, which tend to integrate at specific, safe "harbor" sites in the genome, or CRISPR-Cas9 systems to guide the virus to a safe location.

4. **Q: Describe the structural components of a typical viral vector system.**
   A: A viral vector system usually consists of two parts: (1) The Vector Plasmid, which contains the gene of interest and the "packaging signal" (φ); and (2) The Packaging/Helper Plasmids, which contain the genes for the capsid proteins and enzymes but lack the packaging signal. When these are put into a cell together, the cell produces "empty-shell" viruses that only grab the vector plasmid containing the gene of interest.
5. **Q: Discuss the challenges of using viruses for in vivo gene therapy.**
   A: The primary challenge is the Immune System. The body often recognizes the viral capsid as a foreign invader and destroys the vector before it reaches the target cells. This can cause severe inflammation (cytokine storm). Another challenge is Delivery: it is difficult to get a virus to reach every cell in a large organ (like the brain or heart) simply through an injection into the blood.

---

15 Multiple Choice Questions (MCQs)

1. The transfer of DNA into animal cells by a virus is called:

   a) Transformation

   b) Conjugation

   c) Transduction ✓

   d) Electroporation

2. Which enzyme converts retroviral RNA into DNA?

   a) DNA Polymerase

   b) Integrase

   c) Reverse Transcriptase ✓

   d) RNA Ligase

3. Adenoviral vectors are characterized by being:

   a) Non-integrating ✓

   b) Smallest in size

   c) Derived from plants

   d) Only capable of infecting bacteria

4. Lentiviruses are a sub-group of:

   a) Bacteriophages

b) Retroviruses ✓

c) Adenoviruses

d) Mimiviruses

5. The term "Tropism" refers to the virus's:

a) Cell-type specificity ✓

b) Speed of replication

c) Size of DNA

d) Color under a microscope

6. A major disadvantage of Retroviruses is that they:

a) Are too large

b) Only infect dividing cells ✓

c) Do not have a capsid

d) Are made of protein only

7. Episomal DNA is found in the:

a) Nucleus (but not in chromosomes) ✓

b) Mitochondria

c) Cytoplasm

d) Cell membrane

8. Which vector has the lowest immunogenicity (immune response)?

a) Adenovirus

b) Bacteria

c) AAV (Adeno-associated virus) ✓

d) HIV

9. The "Packaging Signal" (φ) ensures that:

a) Only the gene of interest is put into the virus ✓

b) The virus kills the cell

c) The virus reproduces indefinitely

d) The DNA stays outside the cell

10. Insertional mutagenesis is a risk associated with:
    a) Integrating vectors (Retrovirus) ✓
    b) Non-integrating vectors (Adenovirus)
    c) Chemical methods
    d) Microinjection
11. Helper plasmids are used to provide:
    a) Food for the virus
    b) Viral structural proteins ✓
    c) Antibiotic resistance
    d) Heat for the experiment
12. Which of these is NOT a viral vector?
    a) Lentivirus
    b) Herpes Simplex Virus
    c) Agrobacterium ✓
    d) Vaccinia Virus
13. Stable transformation is achieved when DNA:
    a) Integrates into the host genome ✓
    b) Remains in the cytoplasm
    c) Is degraded by lysosomes
    d) Is turned into a protein
14. The genetic material of Adenovirus is:
    a) Double-stranded DNA ✓
    b) Single-stranded RNA
    c) Double-stranded RNA
    d) Single-stranded DNA
15. A "Self-Inactivating" (SIN) vector is designed to improve:
    a) Speed
    b) Cost

c) Safety ✓

d) Color

Topic 154 – Transgenic Mice Production

1. Overview of Transgenesis

A transgenic mouse is a "Genetically Modified" (GM) mouse that carries a foreign gene (transgene) in all of its cells, including its germline (sperm/eggs), allowing the trait to be inherited by its offspring.

2. The DNA Microinjection Method

This is the most common technique for adding a new gene.

1. **Superovulation:** Female donor mice are injected with hormones (FSH and LH) to produce up to 40 eggs instead of the usual 5-10.
2. **Fertilization:** These females are mated with fertile males to produce zygotes.
3. **Microinjection:** The transgene is injected into the **male pronucleus** (the sperm nucleus within the egg) before it fuses with the egg nucleus.
4. **Implantation:** The surviving embryos are surgically transferred into the oviduct of a **pseudo-pregnant** surrogate mother.

3. The Embryonic Stem (ES) Cell Method

This method is used for "Gene Targeting" or "Knockouts."

1. **ES Cell Culture:** Stem cells are harvested from the inner cell mass of a mouse blastocyst.
2. **Transfection:** Foreign DNA is introduced into these cells via electroporation.
3. **Selection:** Cells that successfully integrated the DNA are selected using antibiotics.
4. **Injection into Blastocyst:** These modified ES cells are injected into a fresh blastocyst from a different colored mouse.
5. **Chimeras:** The resulting offspring are "Chimeras"—animals with some cells from the host blastocyst and some from the modified ES cells.

4. Verification and Breeding

- **Genotyping:** Once pups are born, DNA is extracted from their tails and analyzed using **PCR** (Polymerase Chain Reaction) to confirm the presence of the transgene.
- **Founder Mice:** The first confirmed transgenic mice are called "Founders." They are bred to create a stable, homozygous transgenic line.

10 Short Questions & Answers

1. **Q: Why are mice preferred for transgenesis? A:** They have short gestation periods (20 days), produce large litters, and are genetically similar to humans.
2. **Q: What is a "Founder Mouse"? A:** The original transgenic animal that first successfully integrated the foreign gene.
3. **Q: Define "Superovulation." A:** The use of hormones to stimulate a female to release a higher-than-normal number of eggs.
4. **Q: What is a Chimera? A:** An animal that contains two genetically distinct types of cells (e.g., modified ES cells and host blastocyst cells).
5. **Q: Why target the male pronucleus? A:** It is physically larger and more visible under a microscope than the female pronucleus.
6. **Q: What is the role of a surrogate mother? A:** To provide a womb for the development of the injected embryos.
7. **Q: How is a surrogate mother prepared? A:** By mating with a sterile (vasectomized) male to induce hormonal changes.
8. **Q: What does "Germline Transmission" mean? A:** The transgene is present in the sperm or eggs, allowing it to be passed to offspring.
9. **Q: What is the "Inner Cell Mass"? A:** The part of the blastocyst from which Embryonic Stem cells are derived.
10. **Q: Name a common method to confirm transgenesis in pups. A:** PCR (Polymerase Chain Reaction).

5 Long Questions & Answers

1. **Q: Compare DNA Microinjection and the ES Cell method.**
   A: DNA Microinjection is simpler and faster; it involves injecting DNA directly into a zygote. However, the DNA integrates randomly into the genome, which can cause unpredictable results. The ES Cell method is more complex but allows for "Gene Targeting." Because researchers can select specific cells in a dish before putting them back into an embryo, they can precisely delete (Knockout) or replace (Knock-in) specific genes.
2. **Q: Describe the hormonal steps involved in Superovulation.**
   A: To maximize egg yield, a female mouse is first injected with Pregnant Mare Serum Gonadotropin (PMSG), which mimics Follicle Stimulating Hormone (FSH) to promote egg maturation. Approximately 48 hours later, she is injected with Human Chorionic Gonadotropin (hCG), which mimics Luteinizing Hormone (LH) to trigger ovulation. She is then immediately mated with a male to ensure fertilization.
3. **Q: Explain the significance of "Chimeric Mice" in the production of transgenic lines.**
   A: Chimeras are not the final product but a "middle step." In ES cell technology, the pups born are usually mixtures (e.g., black and white fur). The goal is to breed these chimeras with wild-type mice. If the modified ES cells successfully formed the germline (sperm/eggs), the resulting

pups will be 100% transgenic. If the pups don't show the trait, the chimera's germline was not modified.

4. **Q: What are the primary applications of Transgenic Mice in biotechnology?**
   A: (1) Disease Models: Creating mice with human-like cancer or Alzheimer's to test drugs. (2) Gene Function: Studying what a specific gene does by knocking it out. (3) Pharming: Using mice as small-scale models to see if a human protein can be produced in milk before moving to larger animals like goats. (4) Toxicology: Testing the safety of new chemicals or cosmetics.
5. **Q: Discuss the limitations and efficiency of transgenic mouse production.**
   A: The process is inefficient. In microinjection, many embryos die from the physical trauma of the needle. Even if they survive, the transgene often fails to integrate. Typically, only 1-5% of injected eggs result in a live transgenic mouse. Additionally, random integration can lead to "Position Effect," where the gene is integrated into a silent part of the chromosome and is never expressed.

---

15 Multiple Choice Questions (MCQs)

1. Superovulation is induced using:

   a) Antibiotics

   b) FSH and LH (or PMSG and hCG) ✓

   c) Saline

   d) Glucose

2. In DNA microinjection, the transgene is delivered into the:

   a) Cytoplasm

   b) Male Pronucleus ✓

   c) Mitochondria

   d) Ribosome

3. Embryonic Stem (ES) cells are derived from the:

   a) Inner Cell Mass of the blastocyst ✓

   b) Tail of the adult mouse

   c) Liver

   d) Brain

4. A pup with patches of two different colors is called a:

   a) Clone

   b) Mutant

c) Chimera ✓

d) Hybrid

5. Which drug is often used to select for transformed ES cells?

a) Neomycin (G418) ✓

b) Penicillin

c) Aspirin

d) Insulin

6. The gestation period of a mouse is approximately:

a) 9 months

b) 19-21 days ✓

c) 5 days

d) 60 days

7. A surrogate mother must be _____ to accept the embryos.

a) Very old

b) Pseudo-pregnant ✓

c) Sick

d) Genetically modified

8. The enzyme used in PCR to confirm transgenesis is:

a) Taq Polymerase ✓

b) Ligase

c) Helicase

d) Amylase

9. Random integration of a gene is a hallmark of:

a) DNA Microinjection ✓

b) ES cell targeting

c) Natural mating

d) Cloning

10. A "Knockout" mouse is one where a gene has been:

a) Overexpressed

b) Inactivated or deleted ✓

c) Made to glow

d) Doubled

11. The "Founder" generation is denoted as:

a) F0 ✓

b) F1

c) P1

d) G1

12. ES cells are "Pluripotent," meaning they can:

a) Only become skin

b) Become any cell type in the body ✓

c) Only live in a lab

d) Not divide

13. Electroporation is used in the ES cell method to:

a) Introduce DNA into the stem cells ✓

b) Kill the blastocyst

c) Freeze the embryo

d) Magnify the cells

14. Confirming that a transgene is passed to offspring is proof of:

a) Mutation

b) Toxicity

c) Germline transmission ✓

d) Random luck

15. Which gas is used to anesthetize mice during surgery?

a) Isoflurane or CO2 ✓

b) Oxygen

c) Nitrogen

d) Hydrogen

Topic 155 – Pronuclear Microinjection

1. The Biological Window

When a sperm enters an egg, it does not immediately merge its DNA with the egg's DNA. For several hours, the two sets of chromosomes exist in separate structures called **pronuclei**.

- **Male Pronucleus:** Usually larger and more distinct; the preferred target.
- Female Pronucleus: Smaller and often less accessible.
  The goal of this technique is to inject the transgene into one of these pronuclei before syngamy (fusion) occurs.

2. Technical Setup

The procedure requires a high-end microscope equipped with **micromanipulators**.

- **Holding Pipette:** A wide, blunt glass tube that uses gentle suction to secure the zygote.
- **Injection Pipette:** A needle with an incredibly sharp tip (approx. 0.5 - 1.0 μ).

3. The Step-by-Step Process

1. **Zygote Recovery:** Fertilized eggs are harvested from the oviduct of a superovulated female.
2. **Positioning:** The zygote is held by the holding pipette so the male pronucleus is in the field of view.
3. **Injection:** The needle pierces the zona pellucida (outer layer), the plasma membrane, and finally the pronuclear membrane.
4. **Verification:** A small volume of DNA solution is expelled. A successful injection is confirmed by the **visible swelling** of the pronucleus.
5. **Re-implantation:** Injected zygotes are cultured for a few hours. Healthy-looking ones are transferred into the oviduct of a **pseudo-pregnant** female.

4. Fate of the Injected DNA

The injected DNA is usually linear. Once inside the nucleus, the cell's natural repair enzymes often integrate the foreign DNA into a random location in the host's genome. If this happens at the 1-cell stage, the transgene will be present in every cell of the developing animal.

---

10 Short Questions & Answers

1. **Q: What is a "Zygote"? A:** A fertilized egg before it begins to divide into multiple cells.
2. **Q: At what stage is pronuclear microinjection performed? A:** The 1-cell stage.
3. **Q: Why target the male pronucleus over the female one? A:** It is physically larger and more visible under differential interference contrast (DIC) microscopy.

4. **Q: What is syngamy? A:** The fusion of the male and female pronuclei to form a single diploid nucleus.
5. **Q: What indicates a successful injection? A:** The pronucleus visibly swells or expands.
6. **Q: What is the "Zona Pellucida"? A:** The thick, transparent outer membrane surrounding the mammalian egg.
7. **Q: How many copies of the gene are usually injected? A:** Hundreds to thousands of copies in a few picoliters of buffer.
8. **Q: What type of DNA integrates most efficiently? A:** Linear DNA.
9. **Q: What is a "Pseudo-pregnant" female? A:** A female whose body is hormonally prepared for pregnancy after mating with a sterile male.
10. **Q: What is the main limitation of this method? A:** Random integration, which can lead to "position effects" or gene disruption.

---

5 Long Questions & Answers

1. **Q: Describe the significance of timing in Pronuclear Microinjection.**
   A: Timing is the most critical factor. The injection must occur after fertilization but before the two pronuclei fuse (syngamy). If the injection is too late and the cell has already divided into two, the resulting animal will be a mosaic (only some cells carry the gene). By injecting at the 1-cell stage, researchers ensure that the transgene is copied during every subsequent round of mitosis, leading to a fully transgenic "Founder" animal.
2. A: A micromanipulator is a mechanical device that translates l**Q: Explain the role of the micromanipulator system in this technique.**
   arge movements of the researcher's hand into microscopic movements of the pipettes. Because the pronucleus is only about 10-15 μ in diameter, any manual tremor would destroy the cell. The system allows for precise X, Y, and Z axis control, enabling the needle to pierce the membrane without causing lethal trauma to the zygote.
3. Q: Discuss the possible outcomes of DNA integration after injection.
   A: Integration is a random process governed by the cell's DNA repair machinery. There are three main outcomes: (1) No integration: The DNA is degraded and lost. (2) Successful integration: The DNA inserts into a single site (often in a head-to-tail concatemer). (3) Multiple site integration: The DNA inserts into different chromosomes, which can complicate breeding later on.
4. Q: What are the primary equipment requirements for a microinjection lab?
   A: A specialized lab requires: (1) An Inverted Microscope with high-quality optics (Nomarski or DIC). (2) Two Micromanipulators (one for holding, one for injecting). (3) A Micro-injector pump to control the pressure and volume of the DNA. (4) A Pipette Puller to create glass needles from capillaries using heat and tension.
5. Q: Evaluate the efficiency and survival rates of this procedure.
   A: Pronuclear microinjection is quite inefficient. Roughly 50-80% of zygotes survive the physical trauma of injection. Of those that survive and are implanted, only 10-25% result in live births. Finally, only about 1-5% of the total eggs injected actually result in a "Founder" transgenic mouse. This necessitates the use of many donor animals to produce a single transgenic line.

15 Multiple Choice Questions (MCQs)

1. Pronuclear microinjection is performed at the:
   a) 1-cell stage ✓
   b) 8-cell stage
   c) Blastocyst stage
   d) Adult stage
2. The male pronucleus is derived from the:
   a) Egg
   b) Sperm ✓
   c) Polar body
   d) Somatic cell
3. The visible expansion of the pronucleus during injection is called:
   a) Lysis
   b) Syngamy
   c) Swelling ✓
   d) Mutation
4. The DNA used for injection is typically dissolved in:
   a) A specialized buffer (e.g., TE buffer) ✓
   b) Pure alcohol
   c) Tap water
   d) Blood
5. Syngamy refers to the:
   a) Injection of DNA
   b) Fusion of pronuclei ✓
   c) Division of the cell
   d) Death of the zygote
6. Which microscope is most commonly used for this procedure?
   a) Electron microscope

b) Scanning probe microscope

c) Inverted microscope with DIC optics ✓

d) Dissecting microscope

7. The "Holding Pipette" works based on:

a) High heat

b) Gentle suction (vacuum) ✓

c) Magnetic force

d) Glue

8. The diameter of a mammalian zygote is approximately:

a) 70-100 µ ✓

b) 1-2 mm

c) 10 nm

d) 1 cm

9. Integration of the transgene in this method is:

a) Site-specific

b) Random ✓

c) Guided by RNA

d) Impossible

10. Pseudo-pregnancy in mice lasts approximately:

a) 2 days

b) 9 months

c) 19-21 days (same as regular pregnancy) ✓

d) 1 year

11. A "Mosaic" animal is one where:

a) The gene is in every cell

b) The gene is only in some cells ✓

c) The animal is multi-colored

d) The DNA is circular

12. Which gas is used to pull the glass pipettes into fine needles?
    a) Oxygen
    b) Helium
    c) No gas (uses heat and gravity/tension) ✓
    d) Nitrogen
13. The picoliter (pL) is equal to:
    a) 10^-6 Liters
    b) 10^-9 Liters
    c) 10^-12 Liters ✓
    d) 10^-15 Liters
14. Superovulation helps the process by:
    a) Providing more zygotes for injection ✓
    b) Making the DNA stay linear
    c) Speeding up the microscope
    d) Making the needles sharper
15. The first transgenic mouse was created in the:
    a) 1950s
    b) 1980s ✓
    c) 2010s
    d) 1920s

Topic 156 – Recombinant Retroviruses

1. Nature of Retroviruses

Retroviruses are a family of enveloped viruses that have an **RNA genome**. Their most unique feature is the ability to convert their RNA into DNA and integrate it permanently into the host cell's chromosomes.

2. Structure of a Recombinant Retroviral Vector

To use a retrovirus as a safe tool, its natural genes—**gag** (structural proteins), **pol** (enzymes like reverse transcriptase), and **env** (envelope proteins)—are removed. These are replaced by:

- **The Gene of Interest (GOI):** The trait we want to introduce.
- **Selectable Marker:** To identify infected cells.
- **LTRs (Long Terminal Repeats):** Sequences at the ends required for integration and expression.
- **\psi (Psi) Sequence:** The "packaging signal" that tells the virus to wrap the RNA into a capsid.

3. The Life Cycle & Transduction

1. **Entry:** The virus binds to receptors and enters via membrane fusion.
2. **Reverse Transcription:** The viral enzyme **Reverse Transcriptase** converts RNA into dsDNA.
3. **Integration:** The enzyme **Integrase** inserts the viral DNA into a random site in the host genome.
4. **Expression:** The host cell now treats the viral DNA as its own, producing the protein encoded by the gene of interest.

4. Advantages and Limitations

- **Advantages:**
  - **High Efficiency:** Can infect nearly 100% of a cell population.
  - **Stable Expression:** The gene is passed to all daughter cells.
- **Limitations:**
  - **Dividing Cells Only:** Standard retroviruses (like MLV) cannot enter the nucleus unless the nuclear envelope breaks down during cell division.
  - **Size Limit:** They can only carry about 8 \text kb of foreign DNA.
  - **Insertional Mutagenesis:** Random integration might disrupt a host tumor-suppressor gene.

---

10 Short Questions & Answers

1. **Q: What is a Recombinant Retrovirus? A:** A retrovirus modified to carry a foreign gene instead of its own viral replication genes.
2. **Q: Define "Reverse Transcription." A:** The process of synthesizing DNA from an RNA template using the enzyme Reverse Transcriptase.
3. **Q: What is the role of the \psi (Psi) sequence? A:** It acts as a packaging signal to ensure the viral RNA is encapsulated into the virus particle.
4. **Q: What is the function of the "env" gene? A:** It encodes the envelope proteins that determine which cell types the virus can infect (tropism).
5. **Q: Why are retroviruses used for "Stable" transformation? A:** Because they integrate their genetic material directly into the host cell's genome.
6. **Q: Name the enzyme responsible for inserting viral DNA into host DNA. A:** Integrase.
7. **Q: What is a "Packaging Cell Line"? A:** A specialized cell line that provides the *gag*, *pol*, and *env* proteins in trans to assemble the virus.
8. **Q: What are LTRs? A:** Long Terminal Repeats; they are necessary for the integration and transcription of the viral genome.

9. **Q: What is the main safety risk of retroviral vectors? A:** Insertional mutagenesis (potential to cause cancer by disrupting host genes).
10. **Q: Can standard retroviruses infect brain neurons? A:** No, because neurons are non-dividing cells, and standard retroviruses require cell division to enter the nucleus.

---

5 Long Questions & Answers

1. **Q: Explain how a replication-deficient retrovirus is produced in the lab.**
   A: To ensure safety, the vector carrying the Gene of Interest lacks the genes needed to replicate (gag, pol, env). To "package" this vector into a viral particle, it is transfected into a Packaging Cell Line. These cells have the gag, pol, and env genes integrated into their own chromosomes but lack the \psi (packaging signal). Consequently, the cell produces all the "parts" of a virus, but only the vector RNA (which has the \psi signal) is stuffed inside the shells. The resulting viruses can infect a target cell once, but cannot produce new viruses.
2. **Q: Discuss the importance of LTRs (Long Terminal Repeats) in retroviral vectors.**
   A: LTRs are found at both ends of the retroviral genome and are divided into U3, R, and U5 regions. They serve two vital roles: (1) Integration: They are recognized by the Integrase enzyme to facilitate the insertion of DNA into the host chromosome. (2) Expression: The 5' LTR acts as a powerful promoter/enhancer, recruiting the host's RNA Polymerase to transcribe the gene of interest.
3. **Q: What is "Insertional Mutagenesis" and why is it a concern in gene therapy?**
   A: Retroviruses integrate into the host genome randomly. If the virus happens to land inside a gene that regulates cell growth (a tumor-suppressor gene) or near an oncogene, it can cause the cell to divide uncontrollably, leading to leukemia or other cancers. This was famously observed in early gene therapy trials for X-SCID, where some patients developed T-cell leukemia due to the site of viral integration.
4. **Q: Compare Retroviruses and Lentiviruses.**
   A: Both are retroviruses that integrate into the genome. However, simple Retroviruses (like Moloney Murine Leukemia Virus) can only infect cells that are actively dividing because they wait for the nuclear membrane to dissolve during mitosis. Lentiviruses (like HIV-based vectors) have specialized proteins that allow the viral DNA to be actively transported through the intact nuclear pores, enabling them to infect non-dividing cells like neurons and muscle cells.
5. **Q: Describe the "Transduction" process step-by-step.**
   A: (1) Adsorption: The virus binds to a receptor on the host cell. (2) Fusion: The viral envelope fuses with the host membrane. (3) Uncoating: The RNA core is released into the cytoplasm. (4) Reverse Transcription: RNA is converted to dsDNA. (5) Nuclear Entry: The DNA enters the nucleus. (6) Integration: Integrase inserts the DNA into the host chromosome. (7) Translation: The host cell produces the protein from the new gene.

---

15 Multiple Choice Questions (MCQs)

1. The genetic material of a retrovirus is:
   a) Double-stranded DNA
   b) Single-stranded DNA
   c) Single-stranded RNA ✓
   d) Proteins only
2. Which enzyme is unique to retroviruses?
   a) DNA Ligase
   b) RNA Polymerase
   c) Reverse Transcriptase ✓
   d) Helicase
3. The "gag" gene in a retrovirus encodes for:
   a) Structural proteins (capsid) ✓
   b) Envelope proteins
   c) Enzymes
   d) The gene of interest
4. Transduction efficiency with retroviruses is generally:
   a) Very low
   b) 10%
   c) Very high (up to 100% in some cultures) ✓
   d) Zero
5. Retroviruses integrate into the host genome using the enzyme:
   a) Protease
   b) Integrase ✓
   c) Amylase
   d) Catalase
6. A "Replication-Deficient" vector cannot:
   a) Make new viral particles after the first infection ✓
   b) Enter the cell

c) Integrate into DNA

d) Produce protein

7. The packaging signal is known as the:

a) LTR

b) Pol gene

c) \psi (Psi) sequence ✓

d) TATA box

8. The 5' LTR primarily functions as a:

a) Tail

b) Promoter ✓

c) Protein

d) Membrane

9. Standard retroviral vectors can only infect:

a) Dead cells

b) Dividing cells ✓

c) Neurons

d) Plant cells

10. The maximum DNA cargo capacity for a retrovirus is approximately:

a) 8 \text kb ✓

b) 100 \text kb

c) 1 \text kb

d) 50 \text kb

11. Which of these is a safety feature in modern retroviral vectors?

a) Adding more viral genes

b) Deleting viral replication genes ✓

c) Using unfiltered viruses

d) Increasing the dose

12. Reverse transcription occurs in the:

a) Cytoplasm ✓

b) Nucleus

c) Ribosome

d) Cell wall

13. "Tropism" refers to:

a) The color of the virus

b) The range of host cells a virus can infect ✓

c) The speed of the virus

d) The age of the virus

14. Insertional mutagenesis can potentially lead to:

a) Faster cell growth

b) Fluorescent cells

c) Cancer ✓

d) Smaller cells

15. Integration of the viral DNA into the host chromosome is:

a) Random ✓

b) Site-specific

c) Temporary

d) Only in the mitochondria

Topic 157 – Transfection of ES Cells

1. What are Embryonic Stem (ES) Cells?

ES cells are **pluripotent** cells derived from the **inner cell mass (ICM)** of a blastocyst-stage embryo. Pluripotency means they have the potential to differentiate into any cell type in the body (skin, muscle, bone, etc.), including the germline.

2. The Workflow of ES Cell Transfection

Unlike microinjection, which is done on a single zygote, ES cell transfection is done on millions of cells in a culture dish.

- **Cell Culture:** ES cells are grown on "feeder layers" (usually inactivated fibroblasts) to prevent them from differentiating prematurely.
- **Transfection:** The transgene is introduced into the ES cells. Because ES cells are delicate, **Electroporation** is the most common method used.
- **Selection:** Since the efficiency of "Homologous Recombination" (precise gene targeting) is very low, a dual-selection strategy is used:
    - **Positive Selection:** Using the *neo* gene to keep cells that took up the DNA.
    - **Negative Selection:** Using the *HSV-tk* gene to kill cells that integrated the DNA randomly.

3. Generating the Transgenic Animal

1. **Blastocyst Injection:** The successfully modified (and verified) ES cells are injected into the cavity of a fresh blastocyst.
2. **Implantation:** This blastocyst is transferred into a surrogate mother.
3. **Chimera Formation:** The resulting pup is a **Chimera**—its body is made of two types of cells (original blastocyst cells and the modified ES cells).
4. **Germline Transmission:** The chimera is bred with a wild-type mouse. If the modified ES cells formed the sperm/eggs, the offspring will be 100% transgenic.

4. Advantages of ES Cell Transfection

- **Precision:** Allows for "Knockout" (deleting a gene) or "Knock-in" (replacing a gene).
- **Verification:** You can verify the genetic edit in the lab dish *before* ever creating an animal.

---

10 Short Questions & Answers

1. **Q: From which part of the embryo are ES cells derived? A:** The inner cell mass (ICM) of the blastocyst.
2. **Q: Define Pluripotency. A:** The ability of a cell to develop into all types of cells in an adult organism.
3. **Q: Why is electroporation used for ES cells? A:** It is efficient for delivering DNA into large populations of cells simultaneously.
4. **Q: What is a "Knockout" mouse? A:** A mouse in which a specific gene has been intentionally inactivated using ES cell technology.
5. **Q: What is a "Feeder Layer"? A:** A layer of inactivated cells that provide nutrients and signals to keep ES cells in an undifferentiated state.
6. **Q: How do you identify a chimeric mouse visually? A:** Usually through coat color; the chimera will have patches of fur from both the host blastocyst and the donor ES cells.
7. **Q: What is the role of *G418* in this process? A:** It acts as the selection agent for cells containing the *neo* (neomycin resistance) gene.

8. **Q: Why is negative selection important? A:** It eliminates cells where the DNA integrated randomly instead of at the specific target site.
9. **Q: What is "Homologous Recombination"? A:** The process where DNA sequences exchange between two similar or identical molecules, used for precise gene targeting.
10. **Q: Can ES cells contribute to the germline? A:** Yes, that is their most critical feature for creating stable transgenic lines.

---

5 Long Questions & Answers

1. **Q: Describe the "Positive-Negative Selection" strategy in ES cell gene targeting.**
   A: To ensure precise integration, two markers are used. The Positive marker (neo) is placed inside the target gene sequence; it allows cells to survive G418 treatment. The Negative marker (HSV-tk) is placed at the end of the DNA fragment. If homologous recombination occurs, only the neo gene is integrated, and the cell lives. If random integration occurs, the whole fragment (including tk) integrates. When the drug Ganciclovir is added, it reacts with tk to kill the cell. This leaves only the precisely modified cells.
2. **Q: Why is the ES cell method preferred over microinjection for functional genomics?**
   A: Microinjection results in random integration, meaning you can add a gene but you can't control where it goes or "turn off" an existing gene. ES cell transfection allows for site-specific changes. This is essential for studying the function of a specific gene by removing it (Knockout) or creating exact models of human genetic diseases.
3. **Q: Explain the process of creating a 100% transgenic mouse from a chimera.**
   A: A chimera is a mosaic. To get a "pure" transgenic line, the chimera must be mated with a wild-type mouse. If the modified ES cells successfully populated the gonads (testes/ovaries) of the chimera, some of its gametes will carry the transgene. When these "transgenic gametes" fertilize, the resulting F1 generation pups will carry the transgene in every single cell of their bodies.
4. **Q: What are the challenges in maintaining ES cell cultures?**
   A: ES cells are extremely sensitive. They must be kept in a "pluripotent" state. If the culture conditions are slightly off, they will start to differentiate (turn into specialized cells like skin or nerve), at which point they lose the ability to create a whole animal. This requires specialized media, specific growth factors (like LIF - Leukemia Inhibitory Factor), and often a feeder layer of cells.
5. **Q: What is the "Knock-in" technique and how does it differ from a "Knockout"?**
   A: A Knockout involves disrupting a gene so it no longer functions. A Knock-in involves replacing a gene with a different version (e.g., replacing a mouse gene with a human version) or inserting a reporter gene (like GFP) exactly under the control of a specific native promoter. Both rely on homologous recombination in ES cells.

---

15 Multiple Choice Questions (MCQs)

1. Embryonic Stem cells are obtained from the:
   a) Zygote
   b) Blastocyst ✓
   c) Gastrula
   d) Fetus
2. The ability of ES cells to become any tissue is called:
   a) Pluripotency ✓
   b) Multipotency
   c) Totipotency
   d) Unipotency
3. Precise gene editing in ES cells relies on:
   a) Random integration
   b) Homologous recombination ✓
   c) Viral infection
   d) Heat shock
4. The negative selection marker HSV-tk makes cells sensitive to:
   a) Penicillin
   b) Ganciclovir ✓
   c) Neomycin
   d) Water
5. Chimeric mice are characterized by having:
   a) Extra legs
   b) Only one type of DNA
   c) Two genetically distinct cell populations ✓
   d) Viral DNA in their cytoplasm
6. Which method is most commonly used to transfect ES cells?
   a) Electroporation ✓
   b) Microinjection

c) Gene gun

d) Calcium phosphate

7. In positive-negative selection, the neo gene is used for:

a) Positive selection ✓

b) Negative selection

c) Screening only

d) Cell lysis

8. The offspring of a chimera that carry the transgene in every cell are the:

a) F0 generation

b) F1 generation ✓

c) Founders

d) Clones

9. LIF (Leukemia Inhibitory Factor) is used in ES cell culture to:

a) Kill the cells

b) Speed up division

c) Inhibit differentiation ✓

d) Induce mutations

10. Gene targeting allows for:

a) Only adding genes

b) Only removing genes

c) Specific site-directed changes ✓

d) Killing all bacteria

11. The "Inner Cell Mass" eventually develops into the:

a) Embryo proper ✓

b) Placenta

c) Yolk sac

d) Egg shell

12. Which coat color is often used to track ES cell contribution in chimeras?

a) Only white

b) Only black

c) Contrasting colors (e.g., Agouti and Black) ✓

d) Green

13. "Targeting Vectors" must have sequences _____ to the target gene.

a) Homologous ✓

b) Identical to viruses

c) Made of protein

d) Extremely short

14. Stable germline transmission is confirmed when:

a) Offspring of the chimera carry the transgene ✓

b) The chimera survives

c) The ES cells glow green

d) The surrogate mother gives birth

15. The primary reason for using ES cells is to create _____ mice.

a) Larger

b) Faster

c) Knockout/Knock-in ✓

d) Multi-colored

Topic 158 – Disease Models

1. Why Use Animal Models?

Human diseases are often too complex to study in a petri dish (cell culture). Transgenic animals provide a "whole-body" system where researchers can see how a disease affects various organs, the immune system, and behavior.

2. Methods of Creating Disease Models

- **Gene Knockouts:** Deleting a specific tumor-suppressor gene to study cancer or deleting metabolic genes to study obesity/diabetes.

- **Gene Knock-ins:** Replacing a mouse gene with a human gene that carries a specific mutation (e.g., the mutated *HTT* gene for Huntington's Disease).
- **Transgenic Overexpression:** Adding multiple copies of a human disease-linked gene (e.g., *APP* for Alzheimer's) to see the effects of protein buildup.

3. Key Examples of Transgenic Models

- **Alzheimer's Disease:** Mice engineered to express human **Amyloid Precursor Protein (APP)**. These mice develop plaques in the brain and memory loss similar to human patients.
- **OncoMouse:** The first patented transgenic mouse, carrying the *v-ras* oncogene, making it highly susceptible to breast cancer. It revolutionized cancer drug testing.
- **Cystic Fibrosis (CF):** Mice created with a disrupted **CFTR** gene. These models help in testing gene therapy treatments for the lungs and digestive tract.
- **Huntington's Disease:** Mice carrying the human expanded polyglutamine (CAG) repeats, showing neurodegeneration and motor defects.

4. Advantages and Ethical Considerations

- **Advantages:** Short generation time of mice allows for rapid study of life-long diseases; high genetic similarity (85%) to humans.
- **Ethical Constraints:** The **3Rs** principle (Replacement, Reduction, and Refinement) is strictly followed to ensure animal welfare while advancing medical science.

---

10 Short Questions & Answers

1. **Q: What is a "Disease Model"? A:** A laboratory animal with a specific condition that mimics a human disease.
2. **Q: Name the first patented transgenic animal used for cancer research. A:** The OncoMouse.
3. **Q: Which human gene is often mutated in Alzheimer's mouse models? A:** *APP* (Amyloid Precursor Protein) or Presenilin.
4. **Q: What is the goal of a "Knockout" in disease research? A:** To determine if the loss of a specific gene causes a disease.
5. **Q: Why are mice the preferred models for human disease? A:** Due to their genetic similarity to humans and their high reproductive rate.
6. **Q: Define "Plaques" in the context of Alzheimer's models. A:** Abnormal clusters of protein fragments (Amyloid-beta) that build up between nerve cells.
7. **Q: What is the *CFTR* gene related to? A:** Cystic Fibrosis.
8. **Q: Can transgenic models be used to test drug toxicity? A:** Yes, they are a requirement before human clinical trials.
9. **Q: What are the "3Rs" in animal research? A:** Replacement, Reduction, and Refinement.
10. **Q: What is "Phenotyping"? A:** The process of observing the physical and behavioral characteristics of the transgenic animal to see if it matches the human disease.

5 Long Questions & Answers

1. **Q: Describe the significance of the OncoMouse in the history of biotechnology.**
   A: Created at Harvard in the 1980s, the OncoMouse was engineered to carry an activated human oncogene (ras). This caused the mice to develop consistent breast cancer. Its significance is twofold: (1) It provided a reliable platform to test chemotherapy drugs. (2) It sparked a massive legal and ethical debate over whether a living organism could be patented, eventually leading to the commercialization of transgenic research.
2. **Q: How are transgenic mice used to study neurodegenerative diseases?**
   A: Diseases like Parkinson’s or Alzheimer’s involve the buildup of toxic proteins in the brain. Scientists inject human genes that produce these proteins into mouse embryos. As the mice age, researchers can use MRI scans and behavioral tests (like the Morris Water Maze) to track memory loss and brain decay. This allows for the testing of "neuroprotective" drugs that cannot be safely tested on humans in early stages.
3. **Q: Explain the role of "Knock-in" technology in studying rare genetic disorders.**
   A: Some diseases are caused by a single "typo" in the DNA. Simple overexpression isn't enough to study this. In a "Knock-in," the mouse's version of the gene is replaced by the human version containing the exact mutation. This is used for diseases like Sickle Cell Anemia or Hemophilia, ensuring that the animal produces the human version of the protein, which is essential for testing specific gene therapies.
4. **Q: What are the limitations of using mice as models for human diseases?**
   A: Despite the similarities, mice are not "small humans." Some human diseases do not manifest the same way in mice. For example, early Cystic Fibrosis mouse models showed intestinal issues but not the lung infections seen in humans. Additionally, the lifespan of a mouse (2-3 years) is too short to fully replicate the decades-long progression of some human age-related diseases.
5. **Q: Discuss the ethical framework governing the creation of disease models.**
   A: The creation of animals that are "born to be sick" is ethically sensitive. Research is governed by Institutional Animal Care and Use Committees (IACUC). The 3Rs are applied: Replace animal models with computer simulations where possible; Reduce the number of animals used to the statistical minimum; and Refine procedures to minimize pain (using anesthesia and humane endpoints).

15 Multiple Choice Questions (MCQs)

1. The first transgenic animal model for cancer was the:

   a) CF-Mouse

   b) OncoMouse ✓

   c) Alzheimer's Rat

   d) Obesity Mouse

2. Which human condition is modeled by disrupting the CFTR gene?
   a) Diabetes
   b) Cancer
   c) Cystic Fibrosis ✓
   d) Hemophilia
3. In Alzheimer's models, "Amyloid-beta" refers to:
   a) Toxic protein fragments ✓
   b) A type of nerve cell
   c) A memory-enhancing drug
   d) A viral vector
4. A "Knock-in" mouse involves:
   a) Deleting a gene
   b) Replacing a gene with a specific version ✓
   c) Doubling the size of the mouse
   d) Randomly adding bacteria
5. The study of the physical characteristics of a disease model is called:
   a) Genotyping
   b) Phenotyping ✓
   c) Sequencing
   d) Cloning
6. Mice share approximately _____ genetic similarity with humans.
   a) 50%
   b) 70%
   c) 85% ✓
   d) 100%
7. The Morris Water Maze is a behavioral test used to measure:
   a) Strength
   b) Memory and learning ✓

c) Sight

d) Hunger

8. The v-ras gene used in the OncoMouse is an example of an:

a) Oncogene ✓

b) Antibiotic marker

c) Enzyme

d) Hormone

9. Which organ is primarily studied in the APP transgenic mouse?

a) Heart

b) Liver

c) Brain ✓

d) Kidney

10. The "Refinement" part of the 3Rs refers to:

a) Using fewer animals

b) Minimizing pain and distress ✓

c) Using computers instead of animals

d) Cleaning the cages

11. Transgenic models for Huntington’s Disease focus on:

a) CAG repeat expansions ✓

b) Insulin production

c) Bone density

d) Lung capacity

12. A "Reporter Gene" in a disease model might be used to:

a) Kill the animal

b) Track where the disease gene is active ✓

c) Stop the disease

d) Make the animal grow

13. Testing a drug in animals before human trials is part of _____ studies.

a) Pre-clinical ✓

b) Clinical Phase III

c) Post-market

d) Historical

14. Why are non-human primates (monkeys) used less often than mice?

a) They are genetically identical to humans

b) They are smaller

c) High cost and major ethical concerns ✓

d) They don't get human diseases

15. The term "Founder" in disease models refers to:

a) The first animal to show the stable transgene ✓

b) The scientist who discovered the gene

c) The drug company

d) The laboratory building

Topic 159 – Isolation of Genomic DNA

1. Introduction to Genomic DNA (gDNA)

Genomic DNA represents the total genetic blueprint of an organism. Unlike plasmid DNA, which is small and circular, genomic DNA is extremely long and coiled within the nucleus of eukaryotic cells or the cytoplasm of prokaryotes.

2. The General Principles of Isolation

To isolate pure DNA, four major barriers must be overcome:

1. **Cell Lysis:** Breaking open the cell wall (if present) and the plasma membrane.
2. **Inactivation of Nucleases:** Stopping enzymes (DNases) that want to shred the DNA.
3. **Separation from Proteins/Lipids:** Removing cellular "junk."
4. **Purification:** Precipitating the DNA and washing away salts.

3. The Step-by-Step Procedure

Step A: Cell Harvesting and Lysis

Cells are collected via centrifugation. The cell pellet is treated with a **Lysis Buffer**.

- **Detergents (e.g., SDS):** These break down the lipid bilayer of membranes.

- **Enzymes: Lysozyme** is used for bacteria (to digest peptidoglycan); **Proteinase K** is used to digest proteins associated with DNA (histones).

Step B: Deproteinization (Removal of Proteins)

Phenol and Chloroform are added to the lysate.

- When centrifuged, the mixture separates into three layers:
    1. **Top Aqueous Phase:** Contains the DNA and RNA.
    2. **Middle Interphase:** Contains denatured proteins.
    3. **Bottom Organic Phase:** Contains lipids and hydrophobic debris.

Step C: RNA Removal

RNase A is added to digest any contaminating RNA, leaving only the DNA in the aqueous solution.

Step D: DNA Precipitation

DNA is insoluble in alcohols. **Ice-cold Ethanol** or **Isopropanol** is added in the presence of salts (Na^+ ions). The DNA will appear as white, stringy fibers (spooling).

Step E: Washing and Resuspension

The DNA pellet is washed with **70% Ethanol** to remove residual salts and then dissolved in **TE Buffer** (Tris-EDTA) or sterile water for storage.

---

10 Short Questions & Answers

1. **Q: What is the primary purpose of SDS in lysis buffer? A:** It is a detergent that disrupts the cell membrane by dissolving lipids and denaturing proteins.
2. **Q: Why is Proteinase K added during isolation? A:** To digest proteins, especially histones that bind to DNA, and to inactivate DNases.
3. **Q: What happens if the pH is not controlled during isolation? A:** Extreme pH can cause DNA denaturation or degradation.
4. **Q: Why must DNA isolation be done on ice or at 4°C? A:** To slow down the activity of any residual DNase enzymes that could destroy the DNA.
5. **Q: What is the role of EDTA in DNA storage buffers? A:** EDTA chelates (binds) divalent metal ions like Mg^2+, which are required cofactors for DNase enzymes.
6. **Q: How can you check the purity of isolated DNA? A:** By calculating the ratio of absorbance at 260nm and 280nm (A_260)/(A_280). A ratio of ~1.8 is pure.
7. **Q: What is "Spooling"? A:** The process of collecting long-chain genomic DNA onto a glass rod after alcohol precipitation.
8. **Q: Name the two organic solvents used in extraction. A:** Phenol and Chloroform.
9. **Q: Why is Genomic DNA more difficult to isolate than plasmid DNA? A:** Because it is much larger and more susceptible to "shearing" (physical breakage).

10. **Q: What is the aqueous phase? A:** The top, water-based layer in an extraction that contains the polar DNA molecules.

---

5 Long Questions & Answers

1. **Q: Explain the chemical basis of DNA precipitation using Ethanol.**
   A: DNA is a highly polar molecule due to its phosphate backbone. In water, H_2O molecules surround the DNA, keeping it dissolved. When ethanol is added, it reduces the "dielectric constant" of the solution. This allows positive ions (Na^+ from added salt) to get close enough to the negative phosphate groups to neutralize their charge. Once neutral, the DNA molecules clump together and fall out of the solution (precipitate).
2. **Q: Compare and contrast Phenol-Chloroform extraction with modern Column-based purification.**
   A: Phenol-Chloroform extraction is a liquid-liquid method; it is very effective for high yields but involves toxic chemicals and is time-consuming. Column-based purification (Spin columns) uses a silica membrane. DNA binds to the silica in the presence of "chaotropic salts," while contaminants are washed through. It is faster, safer, and easier to automate, though the yield can sometimes be lower than the organic method.
3. **Q: What is "Shearing" of DNA and how can it be prevented during isolation?**
   A: Shearing is the physical breaking of long DNA strands into smaller fragments due to mechanical stress. It is caused by vigorous vortexing, fast pipetting, or using small-bore needles. To prevent it, genomic DNA should be handled gently: using wide-bore pipette tips, "inverting" tubes slowly instead of vortexing, and avoiding unnecessary agitation.
4. **Q: Describe the role of different enzymes used in isolating DNA from different sources (Bacteria, Plant, Animal).**
   A: For Animal cells, only Proteases (Proteinase K) and detergents are needed. For Bacteria, Lysozyme is required to break the peptidoglycan cell wall. For Plants, Cellulase or physical grinding in liquid nitrogen is needed to break the tough cellulose cell wall. Additionally, plants require CTAB (a specialized detergent) to remove complex carbohydrates and polyphenols.
5. **Q: How do we quantify the concentration and quality of isolated genomic DNA?**
   A: We use a Spectrophotometer. DNA absorbs UV light at 260nm. Using the formula 1 OD_260 = 50µg/mL, we can calculate the concentration. Quality is checked by the (A_260)/(A_280) ratio. If the ratio is <1.7, there is protein contamination; if it is >2.0, there is likely RNA contamination. Quality can also be checked by running the DNA on an Agarose Gel; genomic DNA should appear as a single, high-molecular-weight band near the top.

---

15 Multiple Choice Questions (MCQs)

1. The primary genetic material of a cell is called:

   a) Plasmid DNA

b) Genomic DNA ✓

c) Mitochondrial DNA

d) RNA

2. Which enzyme is used to digest the bacterial cell wall?

a) Lysozyme ✓

b) Lipase

c) Amylase

d) Helicase

3. DNA is insoluble in:

a) Water

b) Buffer

c) Cold Ethanol ✓

d) Saline

4. In a phenol-chloroform extraction, DNA is found in the:

a) Aqueous phase ✓

b) Organic phase

c) Interphase

d) Pellet

5. The A_260/A_280 ratio for pure DNA is approximately:

a) 1.0

b) 1.5

c) 1.8 ✓

d) 2.5

6. What is the purpose of RNase?

a) To make DNA

b) To degrade RNA contamination ✓

c) To break cell membranes

d) To dye the DNA

7. Which chemical is used to denature and remove proteins?
    a) Phenol ✓
    b) Water
    c) Glucose
    d) Agarose
8. EDTA in TE buffer prevents DNA degradation by:
    a) Boiling the DNA
    b) Chelating Mg^2+ ions ✓
    c) Adding oxygen
    d) Changing the color
9. Vigorous shaking or vortexing during gDNA isolation causes:
    a) Better purity
    b) DNA shearing ✓
    c) RNA synthesis
    d) Protein activation
10. The "Interphase" in organic extraction contains:
    a) Pure DNA
    b) Lipids
    c) Denatured proteins ✓
    d) Alcohol
11. DNA precipitates when ethanol is added because it:
    a) Decreases the solubility of DNA ✓
    b) Increases the temperature
    c) Destroys the DNA
    d) Dissolves the phosphate
12. What is the common concentration of Ethanol used for washing the DNA pellet?
    a) 100%
    b) 70% ✓

c) 10%

d) 50%

13. SDS is a type of:

a) Anionic detergent ✓

b) Enzyme

c) Sugar

d) Vitamin

14. Genomic DNA is usually visualized on a/an:

a) Microscope

b) Agarose Gel ✓

c) Staining plate

d) pH paper

15. Which of the following requires liquid nitrogen for DNA isolation?

a) Human blood

b) E. coli

c) Plant tissue ✓

d) Virus particles

Topic 160 – Agarose Gel Electrophoresis

1. Principle of Electrophoresis

Electrophoresis is the movement of charged particles through a matrix under the influence of an electric field.

- **Charge:** DNA and RNA are **negatively charged** (due to their phosphate backbone). Therefore, they migrate toward the **positive electrode (Anode)**.
- **Matrix:** Agarose (a polysaccharide from seaweed) forms a porous 3D mesh.
- **Separation:** Smaller DNA fragments move through the pores faster than larger ones. Thus, molecules are separated by **size**.

2. Components of the System

- **Agarose Gel:** Usually 0.7% to 2.0%. Low concentrations (0.7%) are better for large DNA, while high concentrations (2.0%) are better for small DNA.
- **Buffer (TAE or TBE):** Conducts electricity and maintains a stable pH.
- **Loading Dye:** Contains glycerol (to make the DNA sink into the well) and tracking dyes (to visualize progress).
- **Ethidium Bromide (EtBr) or SYBR Safe:** Fluorescent dyes that bind to DNA so it can be seen under **UV light**.
- **DNA Ladder (Marker):** A mixture of DNA fragments of known sizes used as a ruler for comparison.

3. The Procedure

1. **Gel Casting:** Melted agarose with buffer is poured into a tray with a "comb" to create wells.
2. **Loading:** DNA samples mixed with loading dye are placed into the wells.
3. **Running:** An electric current is applied (50\text--150\text V).
4. **Visualization:** The gel is placed on a UV transilluminator to see the "bands" of DNA.

---

10 Short Questions & Answers

1. **Q: Why does DNA move toward the positive electrode? A:** Because DNA has a negative charge due to its phosphate groups.
2. **Q: What is the source of Agarose? A:** It is a linear polymer extracted from seaweed (red algae).
3. **Q: What is a "DNA Ladder"? A:** A solution of DNA molecules of known lengths used to estimate the size of unknown DNA fragments.
4. **Q: What is the role of Glycerol in the loading dye? A:** It adds density to the sample, ensuring the DNA sinks to the bottom of the well.
5. **Q: Name a common buffer used in electrophoresis. A:** TAE (Tris-Acetate-EDTA) or TBE (Tris-Borate-EDTA).
6. **Q: How does the concentration of agarose affect separation? A:** Higher concentration creates smaller pores, which is better for resolving small DNA fragments.
7. **Q: Why is Ethidium Bromide considered dangerous? A:** It is a potent mutagen because it intercalates (slides) between DNA base pairs.
8. **Q: What color does DNA appear under UV light when stained with EtBr? A:** Orange/Pink.
9. **Q: What are "Bands"? A:** A group of DNA molecules of the same size that have migrated to the same position in the gel.
10. **Q: Define "Pore Size." A:** The size of the gaps in the agarose matrix that determine how easily a molecule can pass through.

---

5 Long Questions & Answers

1. Q: Describe the relationship between DNA molecular weight and migration rate.
   A: There is an inverse relationship between the size of a DNA fragment and its migration rate.

The agarose matrix acts as a molecular sieve. Small, compact DNA molecules move easily through the pores, while large, bulky strands are slowed down by the mesh. Mathematically, the distance migrated (d) is inversely proportional to the log_10 of the molecular weight (MW).

2. Q: Compare TAE and TBE buffers.
   A: TAE (Tris-Acetate-EDTA) is better for recovering DNA from the gel (gel extraction) and is cheaper, but it exhausts faster during long runs. TBE (Tris-Borate-EDTA) has a higher buffering capacity and provides better resolution for very small DNA fragments, but the borate can interfere with some downstream enzymatic reactions.
3. Q: Explain the visualization of DNA using intercalating agents.
   A: DNA itself is invisible to the naked eye. We use intercalating agents like Ethidium Bromide. These molecules slide into the spaces between the rungs of the DNA ladder (base pairs). When exposed to Ultraviolet (UV) light, the EtBr absorbs the UV energy and re-emits it as visible orange light, allowing us to see exactly where the DNA "bands" are located.
4. Q: What are the factors that affect the mobility of DNA in a gel?
   A: (1) Size of DNA: Smaller is faster. (2) Agarose Concentration: Higher concentration slows everything down. (3) DNA Conformation: Supercoiled DNA moves faster than linear DNA of the same size. (4) Voltage: Higher voltage increases speed but can cause the gel to melt. (5) Buffer: Ionic strength affects the current and heat.
5. Q: What is "Gel Extraction" and why is it performed?
   A: Gel extraction (or gel purification) is the process of cutting a specific DNA band out of the agarose gel with a blade and then dissolving the gel to recover the DNA. This is performed when a researcher needs a specific piece of DNA (e.g., a PCR product or a restricted gene) for further experiments like cloning or sequencing, free from other contaminating fragments.

---

15 Multiple Choice Questions (MCQs)

1. DNA fragments are separated in agarose gel based on:

   a) Color

   b) Size and charge ✓

   c) Temperature

   d) Volume

2. The charge of DNA is _____, so it moves toward the _____.

   a) Positive, Cathode

   b) Negative, Anode ✓

   c) Neutral, Anode

   d) Negative, Cathode

3. Which of the following concentrations is best for resolving very large DNA?
   a) 0.7% ✓
   b) 2.0%
   c) 5.0%
   d) 10%
4. The tracking dye in electrophoresis allows the researcher to:
   a) See the DNA directly
   b) Stop the DNA from moving
   c) Monitor the progress of the run ✓
   d) Change the DNA sequence
5. Which chemical is used to visualize DNA under UV light?
   a) Ethidium Bromide ✓
   b) Bromophenol Blue
   c) Coomassie Blue
   d) Methylene Blue
6. "Comb" is used in gel electrophoresis to:
   a) Clean the gel
   b) Create wells for sample loading ✓
   c) Heat the agarose
   d) Measure the voltage
7. The unit of measurement for DNA size on a gel is:
   a) Millimeters
   b) Base pairs (bp) or Kilobases (kb) ✓
   c) Grams
   d) Volts
8. If the power supply is accidentally reversed, the DNA will:
   a) Run out of the top of the gel into the buffer ✓
   b) Move faster toward the bottom

c) Turn green

d) Explode

9. EDTA in electrophoresis buffers serves to:

a) Provide color

b) Inactivate DNases by chelating metal ions ✓

c) Melt the agarose

d) Speed up migration

10. Which DNA shape moves the fastest through a gel?

a) Supercoiled ✓

b) Linear

c) Nicked circular

d) Relaxed circular

11. The movement of the "front" of the run is usually marked by:

a) DNA Ladder

b) Bromophenol Blue ✓

c) UV light

d) Agarose

12. High voltage during electrophoresis can cause:

a) DNA to disappear

b) The gel to melt due to heat ✓

c) DNA to become positive

d) Pores to get smaller

13. SYBR Safe is often used as an alternative to EtBr because it is:

a) Cheaper

b) Less toxic/mutagenic ✓

c) Brighter

d) Smells better

14. Agarose is a _____ of agar.

a) Purified component ✓

b) Synthetic version

c) Liquid form

d) Protein

15. A DNA band that is higher up on the gel (closer to the well) is:

a) Larger in size ✓

b) Smaller in size

c) More negative

d) More positive

Topic 161 – Nucleic Acid Blotting

1. What is Blotting?

Blotting refers to the transfer of biological macromolecules (DNA, RNA, or proteins) from a gel matrix onto the surface of a solid supporting membrane (usually nitrocellulose or nylon). Once on the membrane, the molecules are "immobilized," meaning they are stuck in place and can be analyzed.

2. The Main Types of Blotting

There are three primary blotting techniques, named based on the molecule being studied:

- **Southern Blotting:** Used for the detection of specific **DNA** sequences. (Named after Edwin Southern).
- **Northern Blotting:** Used for the detection of specific **RNA** sequences (to study gene expression).
- **Western Blotting:** Used for the detection of **Proteins**.

3. The Southern Blotting Procedure

1. **Digestion:** Genomic DNA is cut into fragments using **Restriction Enzymes**.
2. **Electrophoresis:** Fragments are separated by size on an agarose gel.
3. **Denaturation:** The gel is treated with an alkali (NaOH) to turn double-stranded DNA into **single-stranded DNA**.
4. **Transfer (The Blot):** Through capillary action or vacuum, the DNA moves from the gel to a **nylon/nitrocellulose membrane**.
5. **Hybridization:** The membrane is incubated with a **Probe** (a labeled, single-stranded DNA fragment) that is complementary to the target sequence.

6. **Detection:** The membrane is washed and exposed to X-ray film (**Autoradiography**) or a digital scanner to see where the probe bound.

4. Why Use Blotting?

- **Specific Detection:** It can find one specific gene among millions of other fragments.
- **Quantification:** Northern blots show how much of a certain mRNA is being produced (gene expression level).
- **Medical Diagnosis:** Used to identify genetic diseases, viral infections, or for DNA fingerprinting in forensics.

---

10 Short Questions & Answers

1. **Q: Who developed Southern Blotting? A:** Edwin Southern in 1975.
2. **Q: What is the purpose of Northern Blotting? A:** To detect specific RNA sequences and measure gene expression.
3. **Q: Why must DNA be denatured before blotting? A:** So it becomes single-stranded, allowing the probe to bind (hybridize) to it.
4. **Q: Name two types of membranes used in blotting. A:** Nitrocellulose and Nylon.
5. **Q: What is a "Probe"? A:** A short, single-stranded piece of DNA/RNA labeled with radioactivity or a fluorescent dye.
6. **Q: How does DNA move from the gel to the membrane? A:** Primarily through **capillary action** using a stack of paper towels and a buffer.
7. **Q: What is the difference between Southern and Northern blotting? A:** Southern detects DNA; Northern detects RNA.
8. **Q: What chemical is used to denature DNA in the gel? A:** Sodium Hydroxide (NaOH).
9. **Q: What is Hybridization? A:** The process where the labeled probe binds to its complementary sequence on the membrane.
10. **Q: How are results visualized in radioactive blotting? A:** Via **Autoradiography** (exposure to X-ray film).

---

5 Long Questions & Answers

1. **Q: Explain the role and importance of the "Probe" in blotting experiments.**
   A: The probe is the most critical part of a blotting experiment because it provides the specificity. Without a probe, the membrane would just be a smudge of millions of DNA fragments. The probe is a pre-synthesized, single-stranded sequence that is perfectly complementary to the gene of interest. Because it is labeled (with a radioactive isotope like ^{32}P or a fluorescent tag), it acts like a "homing beacon" that only sticks to its match. This allows scientists to "see" a specific gene even if it represents only 0.0001% of the total DNA.
2. **Q: Describe the "Capillary Transfer" setup in Southern Blotting.**
   A: The transfer setup acts like a "sandwich." At the bottom is a reservoir of transfer buffer.

Above this is a "wick" (filter paper), then the gel containing the DNA. The membrane is placed directly on top of the gel. Finally, a thick stack of dry paper towels is placed on top. As the dry towels soak up the buffer from the reservoir, the liquid is pulled upward through the gel and the membrane. This upward flow carries the DNA fragments out of the gel and onto the membrane, where they become permanently trapped.

3. **Q: Why is Northern Blotting technically more difficult than Southern Blotting?**
   A: Northern blotting deals with RNA, which is much more fragile than DNA. The environment is full of enzymes called RNases (found on human skin and in the air) that quickly destroy RNA. Researchers must use specialized chemicals like DEPC to treat all water and equipment. Furthermore, RNA has significant secondary structures (folding), so the gel must contain "denaturing agents" like formaldehyde to keep the RNA linear during electrophoresis.
4. **Q: What are the clinical and forensic applications of Southern Blotting?**
   A: (1) DNA Fingerprinting: Comparing RFLP (Restriction Fragment Length Polymorphism) patterns between a crime scene sample and a suspect. (2) Genetic Screening: Identifying carriers of diseases like Sickle Cell Anemia or Cystic Fibrosis. (3) Infectious Disease: Detecting the presence of viral DNA (like HIV or Hepatitis) in a patient's blood sample before the virus has even caused symptoms.
5. **Q: Compare and contrast Blotting with PCR (Polymerase Chain Reaction).**
   A: PCR is much faster and more sensitive than blotting (it can amplify a single molecule of DNA). However, Blotting provides information that PCR cannot, such as the physical size of a gene fragment in the genome and the quantity of intact mRNA (in Northern blots). While PCR is better for quick detection, Blotting is often used as a "confirmatory test" because it provides a visual map of the genetic material.

---

15 Multiple Choice Questions (MCQs)

1. Southern blotting is used for the detection of:

   a) DNA ✓

   b) RNA

   c) Proteins

   d) Lipids

2. Northern blotting is used for the detection of:

   a) DNA

   b) RNA ✓

   c) Proteins

   d) Carbohydrates

3. The transfer of molecules from gel to membrane is called:
    a) Electrophoresis
    b) Blotting ✓
    c) Translation
    d) Transcription
4. Which of the following is NOT a step in Southern Blotting?
    a) Electrophoresis
    b) Hybridization
    c) Protein synthesis ✓
    d) Denaturation
5. In blotting, the "Probe" must be:
    a) Double-stranded
    b) Single-stranded and labeled ✓
    c) A protein
    d) Circular
6. The chemical used to denature DNA in Southern blotting is:
    a) Formaldehyde
    b) NaOH ✓
    c) HCl
    d) Alcohol
7. Autoradiography is used to detect probes labeled with:
    a) Radioactive isotopes ✓
    b) Fluorescent dyes
    c) Heavy metals
    d) Sugar
8. Western blotting is used for the study of:
    a) DNA
    b) RNA

c) Proteins ✓

d) Bacteria

9. The membrane used in blotting is typically made of:

a) Plastic

b) Paper

c) Nitrocellulose or Nylon ✓

d) Glass

10. Northern blotting uses _____ to keep RNA denatured.

a) Formaldehyde ✓

b) Salt

c) Water

d) Oxygen

11. DNA fragments for Southern blotting are usually generated using:

a) Heat

b) Restriction enzymes ✓

c) Magnets

d) Centrifugation

12. What force moves DNA during a capillary transfer?

a) Gravity

b) Magnetism

c) Suction/Wicking action ✓

d) Electricity

13. Hybridization refers to the binding of:

a) Probe to target ✓

b) Gel to membrane

c) DNA to RNA

d) Protein to lipid

14. Blotting is a _____ technique.

a) Synthetic

b) Analytical ✓

c) Destructive

d) Computational

15. RNase-free conditions are most critical for:

a) Southern Blotting

b) Northern Blotting ✓

c) Western Blotting

d) PCR

Topic 162 – Autoradiography

1. Definition

Autoradiography is a technique that uses **X-ray film** or digital imaging sensors to detect the location of radioactive isotopes. The "auto" prefix indicates that the sample itself provides the radiation source to create its own image.

2. Principle of Operation

The process is very similar to traditional photography, but instead of using visible light, it uses **ionizing radiation** (alpha, beta, or gamma particles).

- **Isotopes used:** Common isotopes in biotechnology include ^32P (for DNA/RNA), ^35S (for proteins), and ^3H (Tritium for cell localization).
- **Silver Halide Crystals:** The X-ray film contains silver halide crystals. When radioactive particles strike these crystals, they "excite" the silver ions.
- **Development:** When the film is chemically developed, the excited crystals are converted into **black metallic silver**, creating an image (spots or bands).

3. The Procedure

1. **Labeling:** The biological sample (DNA, protein, or tissue) is labeled with a radioactive isotope.
2. **Exposure:** The labeled sample is placed in direct contact with a sheet of X-ray film inside a light-tight cassette.
3. **Incubation:** The cassette is often stored at -70^°C. The low temperature stabilizes the chemicals and improves the sharpness of the image.
4. **Development:** The film is processed using a developer and a fixer (similar to darkroom photography).

5. **Analysis:** The black spots or bands on the film correspond to the exact location of the radioactive molecules in the original sample.

4. Applications

- **DNA Sequencing:** Visualizing the results of Sanger sequencing.
- **Blotting:** Detecting specific bands in Southern or Northern blotting.
- **Metabolic Tracking:** Following a "labeled" nutrient through a plant or animal's circulatory system.
- **In situ Hybridization:** Locating specific genes directly on a chromosome.

---

10 Short Questions & Answers

1. **Q: What does the prefix "Auto" signify in Autoradiography? A:** It means the sample is the source of its own radiation.
2. **Q: Which isotope is most commonly used for labeling DNA? A:** Phosphorus-32 (^32P).
3. **Q: What is the function of silver halide in X-ray film? A:** It acts as the radiation-sensitive component that records the image.
4. **Q: Why is the film kept in a "Cassette"? A:** To protect the film from visible light, which would ruin the image.
5. **Q: What is the result of the chemical "development" of the film? A:** The silver ions turn into black metallic silver, visible to the eye.
6. **Q: For what purpose is Sulfur-35 (^35S) used? A:** It is commonly used to label proteins (since sulfur is in amino acids like methionine).
7. **Q: Why is exposure often done at -70^°C? A:** To increase sensitivity and prevent the "fuzziness" caused by the movement of molecules.
8. **Q: What is "In situ" autoradiography? A:** Detecting radioactivity directly within a tissue or on a chromosome slide.
9. **Q: What is the main safety concern with this technique? A:** Exposure to ionizing radiation requires specialized shielding (Lead or Plexiglass).
10. **Q: Can autoradiography be used for non-radioactive samples? A:** No, it strictly requires radioactive isotopes or chemiluminescent markers.

---

5 Long Questions & Answers

1. **Q: Explain the chemical process of image formation on X-ray film.**
   A: The X-ray film consists of a plastic base coated with an emulsion of silver halide (AgBr) crystals. When a radioactive particle (like a beta-electron from ^32P) passes through the emulsion, it hits a silver ion, providing energy to knock an electron loose. This electron gets trapped in a "sensitivity speck" within the crystal, attracting a silver ion to form a single atom of metallic silver. This is called the Latent Image. During chemical development, this single atom acts as a catalyst to turn the entire crystal into black metallic silver, making the spot visible.

2. **Q: Discuss the advantages of using ^32P versus ^3H (Tritium) in** autoradiography.
   A: ^32P is a "hard" beta-emitter with high energy. It creates images very quickly (hours to days), but the image is less precise because the particles travel further through the film. ^3H is a "soft" beta-emitter with low energy. It takes much longer (weeks) to create an image, but because the particles don't travel far, the resolution is extremely high, allowing scientists to see exactly which organelle inside a cell contains the label.
3. **Q: How has the digital age modified the traditional autoradiography technique?**
   A: Traditional X-ray film is being replaced by Phosphor Imager screens. Instead of silver halide, these screens contain "storage phosphors." When hit by radiation, they store the energy. The screen is then scanned by a laser, which releases the stored energy as light. This light is captured digitally. Advantages include a wider "dynamic range" (it can see very faint and very bright spots at once) and the fact that the screens can be "erased" and reused thousands of times.
4. **Q: Describe the setup for a Southern Blot autoradiography experiment.**
   A: After the DNA has been transferred to a membrane and hybridized with a radioactive probe, the membrane is dried. In a darkroom, a sheet of X-ray film is placed directly on top of the membrane. This "sandwich" is placed inside an intensifying screen (which converts radiation to light to speed up the process) and locked in a light-tight cassette. After 24–48 hours in a freezer, the film is developed to reveal the specific DNA bands.
5. **Q: What safety protocols are essential when performing autoradiography?**
   A: Since isotopes like ^32P emit high-energy beta radiation, researchers must use Plexiglass (Lucite) shields (lead is actually not recommended for beta-emitters as it produces X-rays). Personal protective equipment (PPE) including double gloves and lab coats is required. All waste must be stored in specialized containers until the radioactivity decays (usually 10 half-lives). Researchers must also wear Dosimeter badges to track their total radiation exposure.

---

15 Multiple Choice Questions (MCQs)

1. Autoradiography is used to detect:

   a) Fluorescent light

   b) Magnetism

   c) Ionizing radiation ✓

   d) Sound waves

2. The most common isotope for DNA research in autoradiography is:

   a) ^32P ✓

   b) ^14C

   c) ^12C

   d) ^16O

3. X-ray film is sensitive to radiation because of:
    a) Glucose
    b) Silver halide crystals ✓
    c) Gold dust
    d) Lead paint
4. The black color on a developed autoradiograph represents:
    a) Metallic silver ✓
    b) Carbon deposits
    c) Bacteria
    d) Ink
5. Which isotope is best for high-resolution cell localization?
    a) ^32P
    b) ^3H (Tritium) ✓
    c) ^131I
    d) ^60Co
6. A "Cassette" in this context is:
    a) A light-tight container for film ✓
    b) A type of music tape
    c) A radioactive probe
    d) A centrifuge tube
7. Isotopes that emit _____ particles are most common in this technique.
    a) Delta
    b) Beta ✓
    c) Gravity
    d) Sonic
8. Storing the film at -70^°C helps to:
    a) Kill the DNA
    b) Improve image clarity and sensitivity ✓

c) Freeze the bacteria

d) Melt the silver

9. The "Latent Image" is:

a) The final developed film

b) The invisible image on the film before development ✓

c) The radioactive probe

d) The DNA gel

10. The digital alternative to X-ray film is:

a) Phosphor Imager ✓

b) Polaroid camera

c) Electron microscope

d) Spectrophotometer

11. ^35S is typically used to label:

a) Carbohydrates

b) Proteins ✓

c) Lipids

d) Water

12. What is the typical thickness of a beta-shield for ^32P?

a) 1 cm Plexiglass ✓

b) 1 mm Paper

c) 1 meter Concrete

d) 10 cm Lead

13. In Sanger sequencing, autoradiography is used to visualize:

a) The color of the DNA

b) The bands representing the DNA sequence ✓

c) The temperature of the reaction

d) The weight of the gel

14. Autoradiography results are often expressed in:

a) Volts

b) Grams

c) Optical Density (OD) ✓

d) Liters

15. If a film is completely black after development, it means:

a) The experiment was perfect

b) It was exposed to visible light accidentally ✓

c) No radiation was present

d) The freezer was too cold

Topic 163 – Southern Blotting

1. Definition and Purpose

Southern Blotting is a laboratory method used to detect the presence of a specific DNA fragment in a sample. It combines **agarose gel electrophoresis** for size separation with **biochemical probes** for sequence identification.

2. The Five Main Stages

The process is methodical and requires several days to complete:

Stage 1: DNA Digestion

High-molecular-weight genomic DNA is too large to enter a gel. It is treated with **Restriction Endonucleases** (Restriction Enzymes) that cut the DNA at specific sequences, creating thousands of smaller fragments of varying lengths.

Stage 2: Electrophoresis

The DNA fragments are loaded onto an agarose gel. An electric current separates them by size. Larger fragments stay near the top (wells), while smaller fragments move toward the bottom.

Stage 3: Denaturation

Because the DNA is double-stranded, it must be converted to single-stranded DNA (ssDNA) so the probe can bind to it. The gel is soaked in an **alkali solution (NaOH)**, which breaks the hydrogen bonds between base pairs.

Stage 4: The Transfer (Blotting)

DNA is transferred from the fragile gel to a sturdy **nitrocellulose or nylon membrane**. This is usually done via **capillary action**: buffer moves from a reservoir, through the gel, and into a stack of paper towels, carrying the DNA onto the membrane where it sticks permanently.

Stage 5: Hybridization and Detection

The membrane is exposed to a **labeled probe** (a small piece of ssDNA complementary to the target gene). The probe "finds" and binds to its match on the membrane. The location of the probe is then visualized using **autoradiography** or fluorescence.

3. Applications of Southern Blotting

- **Gene Discovery:** Locating a specific gene in a new species.
- **DNA Fingerprinting:** Used in forensics and paternity testing (RFLP analysis).
- **Medical Diagnosis:** Identifying genetic mutations or viral infections (e.g., HIV, Sickle Cell).
- **Transgenic Confirmation:** Verifying if a foreign gene has successfully integrated into an animal's genome.

---

10 Short Questions & Answers

1. **Q: Who invented the Southern Blot? A:** Edwin Southern (1975).
2. **Q: What is the primary molecule detected? A:** DNA.
3. **Q: Why is NaOH used in the process? A:** To denature double-stranded DNA into single-stranded DNA.
4. **Q: What is a "Restriction Digest"? A:** The process of using enzymes to cut genomic DNA into manageable fragments.
5. **Q: Why is a nylon membrane preferred over nitrocellulose? A:** Nylon is more durable and can be "stripped" and re-probed multiple times.
6. **Q: What is "Hybridization"? A:** The process of a single-stranded probe binding to a complementary DNA strand on the membrane.
7. **Q: What provides the specificity in a Southern Blot? A:** The DNA probe.
8. **Q: How is DNA fixed to the membrane? A:** Usually by baking the membrane at 80^\circ C or using UV cross-linking.
9. **Q: What is a "depurination" step? A:** A brief soak in HCl to break very large DNA fragments into even smaller ones to aid transfer.
10. **Q: Can Southern blotting detect RNA? A:** No; the detection of RNA is called Northern Blotting.

---

5 Long Questions & Answers

1. **Q: Explain the detailed mechanism of Capillary Transfer.**
   A: Capillary transfer uses the flow of high-salt buffer to move DNA. A "sandwich" is built: a buffer reservoir at the bottom, a filter paper wick, the gel, the membrane, and a stack of dry paper

towels on top. As the dry towels absorb liquid, the buffer is pulled upward through the gel. The buffer carries the DNA out of the gel matrix. Because the membrane has a high affinity for nucleic acids (positive charge), the DNA fragments are trapped on its surface in the exact same pattern they held in the gel.

2. **Q: Why is the "Denaturation" step critical for the hybridization phase?**
   A: DNA naturally exists as a double helix held together by hydrogen bonds. Probes work by following the rules of base pairing (A-T, G-C). If the target DNA remains double-stranded, the probe cannot access the bases to form a bond. Treatment with NaOH disrupts these hydrogen bonds, leaving the DNA "unzipped." This ensures that when the probe is added, it can successfully hybridize to its complementary sequence.
3. **Q: Compare Southern Blotting with PCR in terms of diagnostic utility.**
   A: PCR is faster, requires less DNA, and is highly sensitive. However, Southern Blotting is superior for determining the copy number of a gene and analyzing large structural rearrangements or "Restriction Fragment Length Polymorphisms" (RFLP). While PCR can tell you if a sequence is present, Southern Blotting can tell you how many copies exist and where they sit relative to other DNA markers.
4. **Q: What is a DNA probe and how is it prepared for detection?**
   A: A probe is a short (100-1000 bp) sequence of single-stranded DNA. It is prepared by labeling it with either a radioactive isotope (^32P) or a non-radioactive tag (like Biotin or Digoxigenin). During the "Labeling" reaction, modified nucleotides are incorporated into the probe sequence. This label allows the researcher to see the final results on X-ray film or through a color change reaction.
5. Q: Describe the steps taken to minimize "background noise" in a Southern Blot.
   A: "Background noise" occurs when the probe sticks non-specifically to the membrane rather than the target DNA. To prevent this, researchers use a Pre-hybridization (Blocking) step. The membrane is soaked in a solution containing "nonspecific" DNA (like salmon sperm DNA) or proteins (like BSA). These molecules "plug up" the empty spaces on the membrane so that the probe only binds to the actual target DNA sequences.

---

15 Multiple Choice Questions (MCQs)

1. Southern Blotting is specifically used to identify:

   a) Proteins

   b) mRNA

   c) Specific DNA sequences ✓

   d) Lipids

2. DNA fragments move through the gel based on:

   a) Color

   b) Size ✓

c) Temperature

d) Age

3. The enzyme used to cut DNA before electrophoresis is:

   a) Restriction Endonuclease ✓

   b) DNA Ligase

   c) RNA Polymerase

   d) Helicase

4. Transfer of DNA from gel to membrane is driven by:

   a) Gravity

   b) Magnetism

   c) Capillary action ✓

   d) Centrifugation

5. In Southern Blotting, the gel is treated with NaOH to _____ the DNA.

   a) Denature ✓

   b) Staining

   c) Fragment

   d) Freeze

6. A radioactive probe is typically labeled with:

   a) ^12C

   b) ^32P ✓

   c) ^14N

   d) ^16O

7. DNA is fixed to a nylon membrane using:

   a) Glue

   b) Water

   c) UV Light or Heat ✓

   d) Alcohol

8. The "Pre-hybridization" step is used to:

   a) Block non-specific binding sites ✓

b) Cut the DNA again

c) Melt the gel

d) Add the probe

9. Detection of radioactive probes is done via:

a) Microscope

b) PCR

c) Autoradiography ✓

d) Centrifuge

10. The correct order of steps is:

a) Digestion --> Electrophoresis --> Blotting --> Hybridization ✓

b) Hybridization --> Blotting --> Digestion

c) Blotting --> Digestion --> Electrophoresis

d) Electrophoresis --> Hybridization --> Digestion

11. DNA Fingerprinting (RFLP) was traditionally based on:

a) Northern Blotting

b) Southern Blotting ✓

c) Western Blotting

d) Eastern Blotting

12. The "wick" in a capillary transfer is made of:

a) Filter paper ✓

b) Plastic wrap

c) Metal foil

d) Agarose

13. Nylon membranes are superior to nitrocellulose because they are:

a) Cheaper

b) More physically robust ✓

c) Transparent

d) Made of sugar

14. Which isotope is a "soft" beta emitter often used in high-resolution blotting?
    a) ^32P
    b) ^35S ✓
    c) ^60Co
    d) ^238U
15. If the NaOH step is skipped, the probe will:
    a) Bind too strongly
    b) Break the membrane
    c) Not bind at all ✓
    d) Change color

Topic 164 – Northern Blotting

1. Definition and Purpose

Northern Blotting is a technique used to study gene expression by observing the abundance and size of specific RNA strands. It allows researchers to determine if a specific gene is "turned on" in a particular tissue (e.g., liver vs. brain) or under specific conditions (e.g., before and after drug treatment).

2. The Procedure: Key Differences from Southern Blotting

The workflow is similar to Southern Blotting but adapted for the unique chemistry of RNA:

- **RNA Isolation:** Total RNA is extracted from cells. Unlike DNA, RNA does not need to be cut with restriction enzymes because mRNA molecules are already relatively small.
- **Denaturation:** RNA is highly prone to forming secondary structures (folding into loops). To keep it linear for size separation, the gel must contain **Formaldehyde** or another denaturant.
- **Electrophoresis:** RNA is separated by size on an agarose gel.
- **Transfer:** The RNA is transferred to a positively charged nylon or nitrocellulose membrane.
- **Hybridization:** A labeled DNA or RNA probe is added. It binds to the complementary target RNA sequence.
- **Detection:** Visualized via autoradiography or chemiluminescence.

3. Challenges: The RNA Problem

- **Fragility:** RNA is much less stable than DNA.

- **RNases:** The environment is full of **RNases** (enzymes that destroy RNA). These are found on human skin and are very hard to kill. All equipment must be treated with **DEPC** (Diethyl pyrocarbonate) to ensure it is "RNase-free."

4. Applications

- **Tissue-Specific Expression:** Finding out which organs express a certain gene.
- **Developmental Biology:** Tracking gene activity during different stages of embryo growth.
- **Disease Diagnosis:** Detecting overexpressed genes in cancer cells.
- **RNA Splicing:** Identifying different sizes of mRNA produced from the same gene (alternative splicing).

---

10 Short Questions & Answers

1. **Q: What is the main molecule detected in Northern Blotting? A:** mRNA (Messenger RNA).
2. **Q: Why is it called "Northern" blotting? A:** It was named as a play on the "Southern" blot (the DNA version).
3. **Q: What is the primary use of Northern blotting? A:** To study gene expression patterns.
4. **Q: Why is formaldehyde added to the gel? A:** To denature the RNA and prevent it from forming secondary structures (folding).
5. **Q: What are RNases? A:** Enzymes that degrade RNA; they are the biggest obstacle in Northern blotting.
6. **Q: What is the role of DEPC? A:** It is a chemical used to inactivate RNase enzymes in water and on lab equipment.
7. **Q: How do you determine the size of an RNA band? A:** By comparing it to an **RNA Ladder** (standard markers of known size).
8. **Q: Does Northern blotting require restriction enzyme digestion? A:** No, because RNA molecules are already small enough to be separated on a gel.
9. **Q: What is a "Loading Control" in Northern blotting? A:** A probe for a "housekeeping gene" (like Actin) used to prove that equal amounts of RNA were loaded in each lane.
10. **Q: How is RNA fixed to the membrane? A:** Through UV cross-linking or baking at 80^\circ C.

---

5 Long Questions & Answers

1. **Q: Compare and contrast Southern and Northern Blotting.**
   A: * Molecule: Southern detects DNA; Northern detects RNA.
   - **Preparation:** Southern requires DNA digestion with restriction enzymes; Northern uses intact RNA.
   - **Denaturation:** Southern uses NaOH to unzip double-stranded DNA; Northern uses formaldehyde to prevent single-stranded RNA from folding.
   - **Information Provided:** Southern tells us about the **presence/structure** of a gene; Northern tells us about the **activity/expression** of a gene.

2. **Q: Why is maintaining an "RNase-free" environment critical?**
   A: RNA is naturally unstable due to its single-stranded nature and the 2'-OH group on its ribose sugar. RNase enzymes are extremely robust, require no co-factors, and can even survive being autoclaved. A single fingerprint can introduce enough RNase to destroy an entire RNA sample. Therefore, researchers use DEPC-treated water, dedicated "RNA-only" pipettes, and wear gloves at all times.
3. **Q: Explain how Northern blotting can be used to study "Alternative Splicing."**
   A: A single gene can produce multiple different mRNA transcripts by including or excluding certain exons. Because Northern blotting separates RNA by size, these different transcripts will appear as separate bands at different heights on the membrane. By looking at a Northern blot, a scientist can see if a cell is producing a "short" or "long" version of a protein's instructions.
4. A: To compare gene expression between two samples (e.g., healthy vs. cancerous), you must ensure you started with the same amount of total RNA**Q: What is the significance of the "Loading Control" in a Northern blot?**
   . Scientists probe for a housekeeping gene (like GAPDH or rRNA) that is expressed at the same level in all cells. If the housekeeping bands look equal but the target gene bands are different, the researcher knows the change in expression is real and not just a pipetting error.
5. **Q: Discuss the modern alternatives to Northern Blotting.**
   A: While Northern blotting is accurate, it is slow and requires large amounts of RNA. Modern alternatives include RT-qPCR (Reverse Transcription Quantitative PCR), which is much faster and more sensitive, and RNA-Seq (Next-Generation Sequencing), which allows scientists to look at the expression of every gene in a cell at once, rather than just one gene at a time.

---

15 Multiple Choice Questions (MCQs)

1. Northern blotting is primarily used to measure:

   a) DNA sequence

   b) Gene expression (mRNA levels) ✓

   c) Protein folding

   d) Lipid concentration

2. Which chemical is used to inactivate RNases?

   a) DEPC ✓

   b) SDS

   c) Agarose

   d) Ethidium Bromide

3. RNA is separated in the gel based on:

   a) Charge only

b) Size/Molecular weight ✓

c) Color

d) Radioactivity

4. Formaldehyde is used in Northern blotting to:

a) Dye the RNA

b) Keep the RNA denatured/linear ✓

c) Glue the RNA to the gel

d) Kill bacteria

5. A probe used in Northern blotting is typically made of:

a) Labeled DNA or RNA ✓

b) Pure protein

c) Carbohydrates

d) Salt

6. Northern blotting does NOT require _____, which is essential for Southern blotting.

a) Electrophoresis

b) A membrane

c) Restriction enzyme digestion ✓

d) A probe

7. Ethidium Bromide can be used to visualize _____ in a Northern blot gel.

a) DNA

b) Ribosomal RNA (rRNA) bands ✓

c) Proteins

d) Probes

8. The transfer of RNA to the membrane is usually done via:

a) Capillary action ✓

b) Simple gravity

c) Heat

d) Stirring

9. If you see no bands on a Northern blot, the most likely reason is:
    a) The DNA was too long
    b) RNase contamination degraded the sample ✓
    c) The gel was too hot
    d) The room was too bright
10. A "Housekeeping Gene" is used as a:
    a) Loading control ✓
    b) Negative control
    c) Restriction enzyme
    d) Membrane cleaner
11. RNA moves toward the _____ electrode during electrophoresis.
    a) Negative (Cathode)
    b) Positive (Anode) ✓
    c) Neutral
    d) Ground
12. UV cross-linking is used to:
    a) See the DNA
    b) Attach RNA permanently to the membrane ✓
    c) Cut the RNA
    d) Denature the probe
13. Detection using a radioactive probe is called:
    a) Autoradiography ✓
    b) Photography
    c) Chromatography
    d) Spectroscopy
14. Northern blotting can identify:
    a) Gene mutations in DNA
    b) mRNA transcript size ✓

c) Amino acid sequences

d) Bacterial cell wall thickness

15. Which of the following is a common housekeeping gene?

a) \beta-Actin ✓

b) Insulin

c) Hemoglobin

d) Adrenaline

Topic 165 – Western Blotting

1. Definition and Purpose

Western Blotting is used to identify a specific protein, determine its relative amount (quantification), and check its molecular weight. It is widely used in research to see if a gene has been successfully translated into a protein and in clinical settings to diagnose diseases like HIV or Lyme disease.

2. The Core Principle: Antibody-Antigen Interaction

While Southern and Northern blots use nucleic acid probes, Western blots use **Antibodies**. Antibodies are highly specific proteins that bind only to their target "antigen" (the protein of interest).

3. The Procedure Step-by-Step

Step 1: Sample Preparation (Lysis)

Cells or tissues are broken open using a buffer containing detergents (like **SDS**) and protease inhibitors to prevent the protein from being degraded.

Step 2: Gel Electrophoresis (SDS-PAGE)

Proteins are separated by size using **SDS-PAGE** (Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis).

- **SDS** coats proteins with a negative charge and unfolds them (denatures), ensuring they are separated strictly by molecular weight.
- **Polyacrylamide** is used instead of agarose because it has much smaller pores, which is necessary for the smaller size of protein molecules.

Step 3: Transfer (Electroblotting)

Proteins are transferred from the gel to a **PVDF** or **Nitrocellulose** membrane. Because proteins are coated in negative SDS, they are "pulled" onto the membrane using an electric current (not capillary action).

Step 4: Blocking

The membrane is soaked in a protein solution (like **non-fat dry milk** or BSA). This prevents the antibodies from sticking to the empty spaces on the membrane, which would cause "noise."

Step 5: Antibody Incubation

1. **Primary Antibody:** Binds specifically to the target protein.
2. **Secondary Antibody:** Binds to the primary antibody. This secondary antibody is usually linked to an enzyme (like HRP) that produces light or color.

Step 6: Detection

The membrane is exposed to a chemical substrate. If the target protein is present, a reaction occurs, producing a visible band on a digital imager or X-ray film.

---

10 Short Questions & Answers

1. **Q: What is the primary molecule detected in Western Blotting? A:** Protein.
2. **Q: What does SDS stand for? A:** Sodium Dodecyl Sulfate.
3. **Q: Why is Polyacrylamide used instead of Agarose? A:** It provides better resolution for small protein molecules due to its smaller pore size.
4. **Q: What is the role of the "Secondary Antibody"? A:** It carries the detection enzyme and amplifies the signal by binding to the primary antibody.
5. **Q: What is "Blocking"? A:** Using a protein like milk to cover non-specific binding sites on the membrane.
6. **Q: Name a common membrane used in Western Blotting. A:** PVDF (Polyvinylidene difluoride) or Nitrocellulose.
7. **Q: What is the charge of a protein after treatment with SDS? A:** Negative.
8. **Q: What is a "Molecular Weight Marker" (Ladder)? A:** A mixture of pre-stained proteins of known weights used for comparison.
9. **Q: How are proteins transferred to the membrane? A:** Using an electric field (Electroblotting).
10. **Q: What is "Denaturation" in this context? A:** Unfolding the 3D structure of a protein into a linear chain using heat and SDS.

---

5 Long Questions & Answers

1. **Q: Explain the role of SDS and Beta-mercaptoethanol in sample preparation.**
   A: Proteins have complex 3D shapes and varying charges. To separate them by size alone, we must neutralize these factors. SDS is a detergent that denatures the protein and coats it with a

uniform negative charge. Beta-mercaptoethanol (or DTT) is a reducing agent that breaks disulfide bonds (S-S bonds) that hold different parts of the protein together. Together, they ensure the protein becomes a linear, negatively charged string that migrates through the gel purely based on its length.

2. **Q: Compare Indirect Detection (Primary + Secondary Antibody) with Direct Detection.**
   A: In Direct Detection, the primary antibody itself is labeled with an enzyme. This is faster but less sensitive. In Indirect Detection, a labeled secondary antibody is used to find the primary. This is preferred because multiple secondary antibodies can bind to a single primary antibody, greatly amplifying the signal (making faint bands visible). It is also more cost-effective as one secondary antibody can be used for many different primary antibodies.
3. **Q: Describe the "Electroblotting" transfer setup.**
   A: The "transfer sandwich" consists of a sponge, filter paper, the gel, the membrane, more filter paper, and another sponge. This sandwich is clamped and placed in a buffer tank. The membrane is placed on the side of the positive electrode (Anode). Since the proteins are negatively charged (thanks to SDS), the electric current drives them out of the gel and onto the surface of the membrane, where they bind via hydrophobic and electrostatic interactions.
4. **Q: Why is Western Blotting often used as a confirmatory test for HIV?**
   A: Rapid tests like ELISA are sensitive but can sometimes give "false positives." Western Blotting is more specific. It allows doctors to see exactly which viral proteins the patient's antibodies are attacking (e.g., gp120, p24). A positive diagnosis is only confirmed if bands appear at multiple specific viral protein weights, making it the "gold standard" for verification.
5. **Q: Discuss the importance of "Loading Controls" in Western Blotting.**
   A: To prove that a change in protein levels is due to an experimental treatment and not just because one lane had more sample than another, we use a loading control. We probe for a housekeeping protein like Actin or Tubulin, which is produced at constant levels in all cells. If the Actin bands are the same across all lanes, we can confidently say that any differences seen in our target protein bands are scientifically valid.

---

15 Multiple Choice Questions (MCQs)

1. Western blotting is used for the detection of:

   a) DNA

   b) RNA

   c) Proteins ✓

   d) Glucose

2. The gel matrix used for protein separation is:

   a) Agarose

   b) Polyacrylamide ✓

c) Silica

d) Gelatin

3. SDS gives proteins a uniform _____ charge.

a) Negative ✓

b) Positive

c) Neutral

d) Variable

4. Which chemical breaks disulfide bonds in proteins?

a) SDS

b) Beta-mercaptoethanol ✓

c) Milk

d) Water

5. The transfer of proteins to a membrane is called:

a) Hybridization

b) Electroblotting ✓

c) Transformation

d) Translation

6. Blocking the membrane is usually done using:

a) Pure water

b) Alcohol

c) Non-fat dry milk or BSA ✓

d) DNA

7. The "Primary Antibody" binds to the:

a) Target Protein ✓

b) Secondary Antibody

c) Membrane

d) Gel

8. HRP (Horseradish Peroxidase) is a common _____ used in detection.

a) Protein

b) Enzyme ✓

c) Membrane

d) Detergent

9. Proteins migrate through the gel toward the:

a) Positive Anode ✓

b) Negative Cathode

c) Center

d) Top

10. A Western blot band's "thickness" indicates the:

a) Age of the protein

b) Quantity of the protein ✓

c) Color of the protein

d) Shape of the protein

11. Which of the following is a common housekeeping protein?

a) \beta-Actin ✓

b) Insulin

c) Estrogen

d) Keratin

12. Chemiluminescence is the production of _____ from a chemical reaction.

a) Heat

b) Light ✓

c) Electricity

d) Sound

13. The pore size of a polyacrylamide gel is determined by the concentration of:

a) Acrylamide ✓

b) SDS

c) Protein

d) Buffer

14. Western blotting is also known as:
    a) Southern Blot
    b) Northern Blot
    c) Immunoblot ✓
    d) RNA-Seq
15. If the primary antibody is made in a rabbit, the secondary antibody must be:
    a) Anti-rabbit ✓
    b) Anti-human
    c) Anti-mouse
    d) Pure rabbit protein

Topic 166 – STS in Physical Mapping

1. What is a Sequence Tagged Site (STS)?

An **STS** is a short DNA sequence (200 \text to 500 base pairs) that has a unique occurrence in the genome and whose exact location and base sequence are known. Because an STS appears only **once** in the entire genome, it serves as a highly specific landmark.

2. The Role of STS in Physical Mapping

Physical mapping aims to determine the actual number of nucleotides between genes. STS markers allow scientists to:

- **Connect Overlapping Clones:** In a "Contig" map, if two different DNA clones both contain the same unique STS, those clones must overlap at that specific site.
- **Universal Standard:** Unlike other markers, STS can be shared between labs as simple digital sequence data, allowing anyone to synthesize the necessary primers for PCR.

3. Generating and Detecting STS

- **Generation:** STS are often derived from expressed sequence tags (ESTs), random genomic sequences, or the ends of cloned DNA fragments.
- **Detection via PCR:** This is the most critical feature. Since the sequence is known, specific **primers** are designed to flank the STS. If the STS is present in a DNA sample, a PCR reaction will produce a visible band on an agarose gel.

4. Types of STS Markers

1. **ESTs (Expressed Sequence Tags):** STS markers derived from cDNAs (representing expressed genes).
2. **SSRs (Simple Sequence Repeats):** Highly polymorphic STS used in DNA profiling.

5. Importance in the Human Genome Project

STS markers provided the "scaffolding" for the Human Genome Project. By creating a high-density STS map, researchers could organize thousands of small, sequenced fragments into their correct positions along the 23 human chromosomes.

---

10 Short Questions & Answers

1. **Q: Define STS. A:** A short, unique DNA sequence used as a landmark in physical mapping.
2. **Q: What is the typical length of an STS? A:** Usually between 200 and 500 base pairs.
3. **Q: Why must an STS be unique? A:** If it appeared in multiple places, it couldn't be used to determine a specific physical location on a chromosome.
4. **Q: What is the primary method used to detect an STS? A:** Polymerase Chain Reaction (PCR).
5. **Q: What is a "Contig"? A:** A set of overlapping DNA segments that together represent a continuous region of a chromosome.
6. **Q: How does STS help in contig assembly? A:** It acts as a bridge; if two clones share the same STS, they are known to overlap.
7. **Q: What is the advantage of STS over RFLP? A:** STS is PCR-based, making it faster, easier to automate, and easier to share digitally.
8. **Q: What are ESTs? A:** Expressed Sequence Tags; they are STS markers derived from the coding regions of the genome.
9. **Q: What does "Physical Mapping" measure? A:** The actual distance in base pairs (bp), kilobases (kb), or megabases (Mb).
10. **Q: Can an STS be a gene? A:** Yes, any unique sequence, whether it codes for a protein or not, can serve as an STS.

---

5 Long Questions & Answers

1. **Q: Explain how STS markers are used to create a physical map.**
   A: To create a physical map, a genome is broken into many overlapping clones (e.g., in BACs or YACs). Each clone is tested via PCR for the presence of various STS markers. By recording which clones contain which markers, researchers create a matrix. Clones that share markers are grouped together. Eventually, this allows the "anchoring" of the clones to specific physical coordinates on the chromosome, creating a continuous map of landmarks.
2. **Q: Why is the PCR-based nature of STS considered a "revolution" in mapping?**
   A: Before STS, mapping required the physical exchange of biological materials (probes or clones) between labs, which was slow and prone to error. With STS, a lab in one country can

simply email a DNA sequence to a lab in another. The second lab can then synthesize the primers and perform PCR to find that exact landmark. This digitized the mapping process and allowed for global collaboration.

3. **Q: Compare Genetic Maps and Physical Maps.**
   A: Genetic Maps are based on recombination frequencies and measured in centiMorgans (cM). They show the relative order of genes. Physical Maps (using STS) show the actual physical distance in base pairs. While the order of markers is usually the same in both, the distances can vary because recombination is not uniform across chromosomes (some areas are "hotspots" and others are "coldspots").
4. **Q: What is the relationship between ESTs and STS?**
   A: An EST is a specific type of STS. It is generated by sequencing a small part of a cDNA molecule. Because cDNAs are made from mRNA, an EST represents a gene that is actively being expressed in a cell. Using ESTs as STS markers allows researchers to map the "functional" part of the genome (the genes) directly onto the physical map.
5. **Q: Describe the "Sequence-Tagged Connector" (STC) strategy.**
   A: In large-scale sequencing projects, the ends of BAC (Bacterial Artificial Chromosome) clones are sequenced to create STS markers. These end-sequences are called "connectors." By matching the end-sequence of one BAC to the middle sequence of another, researchers can rapidly "walk" along the chromosome, connecting the pieces like a giant jigsaw puzzle.

---

15 Multiple Choice Questions (MCQs)

1. STS stands for:

   a) Sequence Total System

   b) Sequence Tagged Site ✓

   c) Systemic Tagged Sequence

   d) Single Targeted Site

2. The uniqueness of an STS means it occurs _____ in the genome.

   a) Exactly once ✓

   b) Thousands of times

   c) Only in bacteria

   d) Twice per chromosome

3. The detection of STS markers primarily relies on:

   a) Southern Blotting

   b) PCR ✓

c) Microscopy

d) Centrifugation

4. Physical maps measure distance in:

a) CentiMorgans

b) Base pairs ✓

c) Liters

d) Grams

5. Which of these is a type of STS derived from cDNA?

a) EST ✓

b) RFLP

c) SNP

d) Plasmid

6. A "Contig" is a set of:

a) Broken genes

b) Overlapping DNA clones ✓

c) Radioactive probes

d) Different species

7. To visualize the results of an STS-PCR, one uses:

a) Agarose Gel Electrophoresis ✓

b) Western Blotting

c) Spectrophotometry

d) Autoclaving

8. STS markers are useful because they are:

a) Very long

b) Made of protein

c) Easy to automate and share digitally ✓

d) Found only in plants

9. If Clone A and Clone B share the same STS, they are said to:

a) Overlap ✓

b) Be identical

c) Be from different chromosomes

d) Repel each other

10. Mapping that uses STS landmarks is called:

a) Genetic mapping

b) Physical mapping ✓

c) Cytogenetic mapping

d) Protein mapping

11. The primers for STS-PCR are designed based on:

a) The known sequence of the STS ✓

b) The weight of the DNA

c) The color of the cell

d) The length of the chromosome

12. High-density STS maps were crucial for:

a) Inventing the microscope

b) The Human Genome Project ✓

c) Discovering penicillin

d) Creating insulin

13. Which marker is most likely to be found in an exon?

a) EST ✓

b) Repetitive DNA

c) Junk DNA

d) Intron

14. The "Contig map" is a type of:

a) Physical map ✓

b) Genetic map

c) Climate map

d) Population map

15. An STS sequence length is typically:

    a) 1-10 bp

    b) 200-500 bp ✓

    c) 50,000 bp

    d) 1 million bp

Topic 167 – SNPs as Physical Markers

1. What is an SNP?

A **Single Nucleotide Polymorphism (SNP)**, pronounced "snip," is a variation at a single position in a DNA sequence among individuals. For a variation to be considered an SNP, it must occur in at least 1% of the population.

- **Example:** In Individual 1, the sequence is AAGG**C**TTA; in Individual 2, it is AAGG**G**TTA. The C/G variation is the SNP.

2. Abundance and Distribution

SNPs are the most common type of genetic variation. In the human genome, there is an SNP approximately every 300 to 1,000 base pairs. This means there are roughly **10 million SNPs** in the human genome.

- **Coding Regions:** SNPs can change amino acids (nonsynonymous) or have no effect (synonymous).
- **Non-coding Regions:** Most SNPs occur here but can still affect gene splicing, transcription factor binding, or non-coding RNA sequence.

3. SNPs as Physical Markers

SNPs are superior physical markers for several reasons:

1. **High Density:** Because they occur so frequently, they provide a much higher resolution map than STS or RFLP markers.
2. **Stability:** SNPs have a very low mutation rate, making them reliable for tracking inheritance over many generations.
3. **Biallelic Nature:** Most SNPs have only two possible alleles (e.g., C or T), which simplifies automated computational analysis.

4. Detection Technologies

Unlike older markers that require gel electrophoresis, SNPs are analyzed using high-throughput digital technologies:

- **DNA Microarrays (SNP Chips):** Can screen millions of SNPs simultaneously.
- **Next-Generation Sequencing (NGS):** Directly identifies SNPs by reading the code.
- **Mass Spectrometry:** Measures the mass of DNA fragments to identify the single base change.

---

10 Short Questions & Answers

1. **Q: Define an SNP. A:** A single base pair site in the genome where different individuals have different nucleotides.
2. **Q: What is the frequency threshold for an SNP? A:** The variation must be present in at least 1% of the population.
3. **Q: Approximately how many SNPs are in the human genome? A:** About 10 million.
4. **Q: What is the difference between an SNP and a mutation? A:** A mutation is usually rare (<1%), while an SNP is a common population variant (>1%).
5. **Q: What is a "Nonsynonymous" SNP? A:** An SNP in a coding region that changes the amino acid sequence of a protein.
6. **Q: Define a "Haplotype." A:** A set of SNPs on a single chromosome that are inherited together.
7. **Q: Why are SNPs biallelic? A:** Because it is statistically very rare for a single position to mutate into three or four different bases across a population.
8. **Q: What is the "HapMap Project"? A:** An international effort to identify and catalog genetic similarities and differences in human beings.
9. **Q: Can SNPs occur in non-coding DNA? A:** Yes, the majority of SNPs are located in non-coding regions.
10. **Q: How are SNPs used in personalized medicine? A:** They can predict an individual's response to certain drugs or their risk of developing specific diseases.

---

5 Long Questions & Answers

1. **Q: Discuss the significance of SNPs in Genome-Wide Association Studies (GWAS).**
   A: GWAS uses SNPs as markers to find genetic variations associated with complex diseases like diabetes or heart disease. Researchers compare the SNPs of thousands of people with a disease to those without it. If certain SNPs are found significantly more often in people with the disease, those SNPs are said to be "associated" with it. This helps point scientists toward the specific regions of the genome where the disease-causing genes may be located.
2. **Q: Compare SNPs with Microsatellites (SSRs) as genetic markers.**
   A: Microsatellites are highly polymorphic (many alleles) and very informative, but they are less abundant and difficult to analyze with high-throughput automation. SNPs are less informative individually (usually only two alleles), but their sheer density (millions vs. thousands) and ease of automated digital analysis make them the preferred choice for modern large-scale genomic mapping and clinical diagnostics.

3. **Q: Explain the potential impact of an SNP located in a "Promoter" region.**
   A: Even if an SNP does not change a protein's amino acid sequence, it can still have a massive biological effect if it is in a promoter. An SNP in a promoter can change the "binding affinity" of transcription factors. This might cause a gene to be expressed too much (overexpression) or too little (underexpression), which can lead to disease or altered physiological traits.
4. **Q: How does SNP analysis contribute to Pharmacogenomics?**
   A: Pharmacogenomics is the study of how genes affect a person's response to drugs. Many enzymes that break down medications (like the Cytochrome P450 family) have SNPs. If an individual has a specific SNP that makes an enzyme slow, a standard dose of medicine could become toxic to them. SNP profiling allows doctors to prescribe the "right drug at the right dose" based on a patient's genetic makeup.
5. **Q: Describe the structure and function of an SNP Microarray.**
   A: An SNP microarray is a small glass or silicon chip with millions of microscopic spots. Each spot contains a unique DNA probe designed to match a specific SNP. When a patient's fluorescently labeled DNA is washed over the chip, it binds (hybridizes) to the matching probes. A laser scanner then reads the fluorescence to determine which allele (e.g., A or G) the patient carries at every single one of those millions of sites simultaneously.

---

15 Multiple Choice Questions (MCQs)

1. SNP stands for:

   a) Single Nucleotide Protein

   b) Single Nucleotide Polymorphism ✓

   c) Systemic Nucleotide Process

   d) Single Node Polyploidy

2. To be an SNP, a variation must occur in at least _____ of the population.

   a) 1% ✓

   b) 10%

   c) 50%

   d) 0.1%

3. Most SNPs are:

   a) Triallelic

   b) Biallelic ✓

   c) Tetra-allelic

   d) Non-allelic

4. How often do SNPs occur in the human genome on average?
   a) Every 10 bp
   b) Every 300-1000 bp ✓
   c) Every 1,000,000 bp
   d) Only once per chromosome
5. An SNP that does not change the amino acid is called:
   a) Synonymous ✓
   b) Nonsynonymous
   c) Nonsense
   d) Frameshift
6. A group of SNPs inherited together is known as a:
   a) Genotype
   b) Phenotype
   c) Haplotype ✓
   d) Isotype
7. GWAS stands for:
   a) Genome-Wide Association Study ✓
   b) General Wide Allele Survey
   c) Genomic Western Analysis System
   d) Gene-Wide Assay Study
8. The primary advantage of SNPs over RFLPs is:
   a) They are larger
   b) They require gels
   c) Their high density and ease of automation ✓
   d) They only occur in plants
9. Detection of millions of SNPs at once is done using:
   a) DNA Microarrays ✓
   b) Southern Blotting

c) SDS-PAGE

d) Centrifugation

10. The study of drug response based on SNPs is called:

a) Proteomics

b) Pharmacogenomics ✓

c) Metabolomics

d) Bioinformatics

11. Nonsynonymous SNPs are found in:

a) Coding regions (exons) ✓

b) Introns only

c) Centromeres only

d) Telomeres only

12. SNP variations typically involve:

a) Large deletions

b) A single base substitution ✓

c) Chromosome doubling

d) Viral integration

13. Which project aimed to map the human haplotypes?

a) The Manhattan Project

b) The HapMap Project ✓

c) The Proteome Project

d) The Blue Brain Project

14. In SNP analysis, the "A" in A/G variation is called an:

a) Allele ✓

b) Enzyme

c) Locus

d) Primer

15. Which technology uses a laser to detect SNP hybridization?

a) Microarray Scanner ✓

b) Autoclave

c) Electron Microscope

d) PCR Machine

Topic 168 – Detection Without Sequence Info

1. The Core Challenge

To perform standard PCR or SNP analysis, you must know the DNA sequence to design specific primers. When sequence information is **unavailable**, scientists use "Universal" or "Arbitrary" primers that bind to many locations in the genome by chance.

2. RAPD (Random Amplified Polymorphic DNA)

RAPD is the most common technique for this purpose. It uses a single, short (10-nucleotide) primer with an arbitrary sequence.

- **Mechanism:** The primer scans the genome and binds wherever it find a match. If two primers bind close enough to each other on opposite strands, the DNA between them is amplified.
- **Polymorphism:** If a mutation occurs at a binding site in one individual but not another, a band will appear in one and be absent in the other.

3. AFLP (Amplified Fragment Length Polymorphism)

AFLP is more robust than RAPD. It combines restriction digestion with PCR.

1. **Digestion:** DNA is cut with two restriction enzymes.
2. **Ligation:** "Adapters" (known DNA sequences) are glued to the cut ends.
3. Amplification: Primers matching the adapters are used to amplify the fragments.
   Because the adapters provide a known "handle," we can amplify the DNA even though we don't know what is in the middle of the fragments.

4. DAF (DNA Amplification Fingerprinting)

Similar to RAPD but uses much shorter primers (5–8 nucleotides) and higher concentrations, resulting in very complex banding patterns (fingerprints) visualized on polyacrylamide gels.

5. Applications

- **Genetic Diversity:** Comparing different varieties of plants.
- **Phylogenetics:** Determining how closely related two unknown species are.
- **Mapping:** Finding markers linked to specific traits in unsequenced organisms.

---

10 Short Questions & Answers

1. **Q: What is "Arbitrary Primer"? A:** A primer with a random sequence used to amplify DNA without prior knowledge of the target sequence.
2. **Q: How long is a typical RAPD primer? A:** Usually 10 nucleotides.
3. **Q: What does "Polymorphic" mean in this context? A:** Differences in DNA banding patterns between individuals of the same species.
4. **Q: Why is RAPD considered "Dominant"? A:** Because it can only tell if a band is present or absent; it cannot distinguish between homozygous and heterozygous states.
5. **Q: What is the main advantage of AFLP over RAPD? A:** AFLP is much more reproducible and reliable.
6. **Q: What are "Adapters" in AFLP? A:** Known DNA sequences ligated to restriction fragments to provide a binding site for primers.
7. **Q: Does RAPD require a high amount of DNA? A:** No, very small amounts are sufficient due to PCR amplification.
8. **Q: How are the results of these techniques visualized? A:** Via Gel Electrophoresis (Agarose for RAPD, Polyacrylamide for AFLP/DAF).
9. **Q: What is a "Null Allele" in RAPD? A:** The absence of a band due to a mutation at the primer binding site.
10. **Q: Why is sequence information unnecessary for these methods? A:** Because the primers bind randomly or to synthetic adapters rather than specific known genes.

---

5 Long Questions & Answers

1. **Q: Describe the step-by-step mechanism of RAPD and how it identifies polymorphisms.**
   A: In RAPD, a single short primer is added to a PCR mix. During the annealing step, the primer binds to multiple random sites on the genome. If two primer molecules bind in an inverted orientation (facing each other) within 200-3000 base pairs, the DNA between them is amplified. Polymorphism occurs if: (a) a base change happens at the binding site, preventing the primer from sticking, or (b) an insertion/deletion occurs between the sites, changing the band's size.
2. **Q: Explain the AFLP procedure and why it is more "stringent" than RAPD.**
   A: AFLP involves cutting genomic DNA with a frequent cutter (MseI) and a rare cutter (EcoRI). Known adapters are ligated to the ends. PCR is then performed in two stages (pre-selective and selective) using primers that match the adapters plus a few extra "selective" nucleotides. This makes it more stringent because the conditions for binding are much tougher, leading to highly reproducible "fingerprints" that are consistent across different laboratories.
3. **Q: Discuss the limitations and drawbacks of using RAPD in biotechnology.**
   A: The main drawback of RAPD is reproducibility. Because the primers are short and the

annealing temperature is low, slight changes in salt concentration, DNA quality, or the brand of thermal cycler can produce different results. Additionally, RAPD markers are "dominant," meaning a single band doesn't tell you if the organism has one or two copies of that allele, which limits its use in sophisticated genetic calculations.

4. **Q: How does DAF (DNA Amplification Fingerprinting) differ from RAPD?**
   A: While both use arbitrary primers, DAF uses shorter primers (5-8 bp) and much higher concentrations. This leads to the amplification of many more fragments, creating a very dense and complex "fingerprint" of the genome. Because the fragments are often very similar in size, DAF requires high-resolution Polyacrylamide Gel Electrophoresis (PAGE) and silver staining to visualize the numerous bands.
5. **Q: Why are these techniques vital for "Orphan Crops" and Biodiversity studies?**
   A: Sequencing a full genome is expensive. For many local crops in developing nations or rare endangered plants, there is no funding for full sequencing. These "sequence-independent" techniques allow researchers to perform "Genetic Mapping" and "Diversity Assessment" immediately. They can identify which plants are most different from each other to ensure a wide genetic base for conservation or breeding.

---

15 Multiple Choice Questions (MCQs)

1. RAPD uses _____ primer(s) per reaction.

   a) One ✓

   b) Two

   c) Four

   d) Zero

2. The length of a RAPD primer is typically:

   a) 50 bp

   b) 100 bp

   c) 10 bp ✓

   d) 1000 bp

3. In AFLP, DNA is first treated with:

   a) Heat

   b) Restriction enzymes ✓

   c) Alcohol

   d) RNA

4. Which technique is the most reproducible?
    a) RAPD
    b) AFLP ✓
    c) DAF
    d) All are equal
5. AFLP uses _____ to provide known sequences for primer binding.
    a) Adapters ✓
    b) Plasmids
    c) Viruses
    d) Probes
6. RAPD markers are considered _____ markers.
    a) Co-dominant
    b) Dominant ✓
    c) Recessive
    d) Sex-linked
7. DAF stands for:
    a) DNA Amplification Fingerprinting ✓
    b) Digital Allele Frequency
    c) DNA Analysis Factor
    d) Diagnostic Antigen Filter
8. Short primers in DAF are usually _____ nucleotides long.
    a) 20-30
    b) 5-8 ✓
    c) 100
    d) 2
9. Detection of polymorphisms without sequence info is ideal for:
    a) Humans
    b) E. coli

c) Unsequenced species ✓

d) Viruses

10. A mutation at a primer binding site in RAPD leads to:

a) Loss of a band ✓

b) A change in band color

c) Duplication of the genome

d) Cell death

11. Which gel is used for high-resolution AFLP visualization?

a) Agarose

b) Polyacrylamide ✓

c) Gelatin

d) Silica

12. The "selective nucleotides" in AFLP primers help to:

a) Dye the DNA

b) Reduce the number of amplified bands ✓

c) Speed up the PCR

d) Protect the DNA

13. Which of the following does NOT require DNA ligase?

a) RAPD ✓

b) AFLP

c) Cloning

d) Recombinant DNA

14. Arbitrary primers work by:

a) Seeking specific genes

b) Binding to many random locations ✓

c) Destroying the DNA

d) Making RNA

15. Genetic distance in unsequenced species is calculated by:

a) Counting the number of cells

b) Comparing shared vs. unique bands ✓

c) Measuring the weight of the animal

d) Sequencing the whole genome

Topic 169 – AFLP Detection

1. What is AFLP?

AFLP is a PCR-based tool used to detect DNA polymorphisms. It does not require any prior knowledge of the target DNA sequence. It works by "selective amplification" of a subset of restriction fragments from a complex mixture of genomic DNA.

2. The Four Major Steps

Step A: Digestion and Ligation

The genomic DNA is cut with two different restriction enzymes:

1. **Rare Cutter:** Usually *EcoRI* (6-bp recognition site).
2. Frequent Cutter: Usually MseI (4-bp recognition site).
   Immediately after cutting, double-stranded Adapters (known DNA sequences) are ligated to the sticky ends of the fragments.

Step B: Pre-selective Amplification

PCR is performed using primers that match the Adapter sequences plus **one** extra nucleotide at the 3' end. This reduces the number of fragments amplified by 16-fold, making the mixture less "crowded."

Step C: Selective Amplification

A second PCR is performed using primers with **three** extra "selective" nucleotides. Only fragments that perfectly match these three bases will be amplified. This ensures that only 50–100 clear bands appear on the gel rather than thousands.

Step D: Gel Analysis

The resulting fragments are separated using **Polyacrylamide Gel Electrophoresis (PAGE)**. Because the primers are often fluorescently or radioactively labeled, the resulting "fingerprint" is highly detailed.

3. Advantages of AFLP

- **High Reproducibility:** Much more stable than RAPD.
- **Genome Coverage:** Samples the entire genome simultaneously.
- **No Sequence Info Needed:** You only need to know the sequence of the synthetic adapters.

10 Short Questions & Answers

1. **Q: What does AFLP stand for? A:** Amplified Fragment Length Polymorphism.
2. **Q: Which two enzymes are commonly used in AFLP? A:** *EcoRI* and *MseI*.
3. **Q: What is an "Adapter" in the AFLP context? A:** A short, known DNA sequence ligated to the ends of restriction fragments.
4. **Q: Why are two different enzymes used? A:** To create a variety of fragments and ensure that most fragments have two different ends for selective primer binding.
5. **Q: What is the purpose of "Selective Nucleotides"? A:** To reduce the number of amplified fragments so the bands can be clearly separated on a gel.
6. **Q: Is AFLP a dominant or co-dominant marker? A:** It is typically a dominant marker (detects presence/absence of a band).
7. **Q: What is the primary method of visualization for AFLP? A:** Polyacrylamide Gel Electrophoresis (PAGE) followed by silver staining or fluorescence.
8. **Q: Does AFLP require prior sequence information of the organism? A:** No.
9. **Q: What is "Selective PCR"? A:** The second round of PCR that uses primers with 2-3 additional bases to narrow down the pool of amplified DNA.
10. **Q: How many bands are typically generated in one AFLP reaction? A:** Between 50 and 100 bands.

5 Long Questions & Answers

1. **Q: Explain the chemical necessity of the Ligation step in AFLP.**
   A: In AFLP, we do not know the sequence of the genomic DNA. Therefore, we cannot design primers to stick to it. By ligating Adapters of a known sequence to the ends of the DNA fragments produced by restriction enzymes, we provide a "universal docking station." The PCR primers are then designed to match these adapters. This allows us to amplify unknown DNA using known primer sequences.
2. **Q: Describe how AFLP reduces a complex genome into a readable "fingerprint."**
   A: A total restriction digest of a genome produces millions of fragments, which would appear as a smear on a gel. AFLP uses Selective Amplification. In the first PCR, primers have +1 extra base, allowing only 1/16th of fragments to grow. In the second PCR, primers have +3 extra bases. Statistically, only a tiny fraction of the original millions of fragments will match these specific 3-base extensions. This results in a manageable number of distinct, sharp bands.
3. **Q: Compare the reliability of AFLP to RAPD markers.**
   A: RAPD uses very short (10-bp) primers at low temperatures, making it highly sensitive to small changes in lab conditions (reproducibility is poor). AFLP uses longer primers (matching adapters) and higher "annealing" temperatures (stringent conditions). Because the primers are longer and the enzymes provide specific starting points, AFLP results are highly consistent between different labs and different days.
4. **Q: What is the role of the "Rare Cutter" and "Frequent Cutter" in the digest?**
   A: The Frequent Cutter (MseI) cuts the DNA into many small fragments (averaging 256 bp),

which are ideal for PCR and gel separation. The Rare Cutter (EcoRI) cuts less often. Most fragments analyzed in AFLP are those that have an EcoRI cut at one end and an MseI cut at the other. This dual-enzyme system ensures that the fragments generated are within the optimal size range for high-resolution imaging.

5. **Q: Discuss the applications of AFLP in plant breeding and biodiversity.**
   A: AFLP is used to assess the "Genetic Diversity" of a population. In plant breeding, it helps identify markers linked to traits like drought resistance in crops where the genome isn't sequenced. In biodiversity, it helps distinguish between nearly identical species or varieties. Because it scans the whole genome, it is excellent for creating high-density genetic maps.

---

15 Multiple Choice Questions (MCQs)

1. AFLP was originally developed by:

   a) Edwin Southern

   b) Zabeau and Vos ✓

   c) Kary Mullis

   d) Gregor Mendel

2. The rare cutter enzyme EcoRI recognizes a _____ sequence.

   a) 6-bp ✓

   b) 4-bp

   c) 10-bp

   d) 2-bp

3. The frequent cutter enzyme MseI recognizes a _____ sequence.

   a) 8-bp

   b) 4-bp ✓

   c) 20-bp

   d) 6-bp

4. Adapters are joined to DNA fragments using the enzyme:

   a) Polymerase

   b) Ligase ✓

   c) Helicase

   d) Nuclease

5. Selective nucleotides are added to the _____ end of the primer.
   a) 5' end
   b) 3' end ✓
   c) Middle
   d) Both ends
6. A PCR primer with "+3" selective nucleotides will amplify _____ fragments than a "+1" primer.
   a) Fewer ✓
   b) More
   c) The same number of
   d) Zero
7. The secondary structure of the AFLP primer matches the:
   a) Adapter and Restriction site ✓
   b) Only the gene
   c) Only the centromere
   d) The cell wall
8. AFLP fragments are typically visualized using:
   a) Agarose gel
   b) Polyacrylamide gel (PAGE) ✓
   c) Centrifugation
   d) Spectrophotometry
9. The main advantage of AFLP is its:
   a) Low cost
   b) High reproducibility and resolution ✓
   c) Ability to detect proteins
   d) Speed (takes 5 minutes)
10. AFLP is a "Sequence Independent" technique because it uses:
    a) Universal Adapters ✓
    b) Human blood

c) Only RNA

d) Radioactive water

11. In the pre-selective amplification, primers usually have _____ extra base(s).

a) 1 ✓

b) 5

c) 10

d) 0

12. The use of two different enzymes in AFLP prevents:

a) Self-ligation of fragments ✓

b) DNA from melting

c) PCR from starting

d) RNA contamination

13. Which of the following is a common label for AFLP primers?

a) Glucose

b) Fluorescent dye or ^32P ✓

c) Salt

d) Iron

14. AFLP analysis is widely used for:

a) DNA Fingerprinting ✓

b) Measuring protein weight

c) Counting bacteria

d) Determining cell pH

15. If the selective nucleotides don't match the DNA, the result is:

a) A larger band

b) No amplification (no band) ✓

c) A mutation

d) The DNA glows green

Topic 170 – The Suitability of Bioremediation

1. What is Bioremediation?

Bioremediation is a biotechnology process that uses microorganisms (bacteria, fungi) or plants to degrade, transform, or remove contaminants from soil and water.

2. The "Suitability" Criteria

Not every polluted site is a candidate for bioremediation. The suitability depends on three main factors:

A. The Nature of the Pollutant (Chemical Factor)

- **Biodegradability:** The contaminant must be "biologically available." Simple hydrocarbons (oil, gasoline) are highly suitable.
- **Recalcitrance:** Some synthetic chemicals (like certain plastics or DDT) are "recalcitrant," meaning microbes cannot easily break them down.
- **Toxicity:** If the concentration of the pollutant is so high that it kills the microbes themselves, bioremediation is not suitable.

B. Environmental Conditions (Physical Factor)

- **Nutrients:** Microbes need Nitrogen, Phosphorus, and Carbon to grow.
- **Environmental Parameters:** pH (usually 6.5–8.5), temperature (15°C–35°C), and moisture must be optimal.
- **Oxygen:** Aerobic bioremediation requires oxygen. In deep soil or groundwater, oxygen may be a limiting factor.

C. Site Characteristics (Geological Factor)

- **Soil Permeability:** If the soil is too dense (like heavy clay), nutrients and oxygen cannot reach the microbes.
- **Location:** *In-situ* (at the site) is cheaper but harder to control; *Ex-situ* (removing soil to another location) is faster but more expensive.

3. Advantages vs. Disadvantages

Advantages	Disadvantages
Environmentally friendly (Natural)	Slower than mechanical methods

Lower cost than incineration	Not effective for heavy metals (Lead/Mercury)
Can be done *in-situ* (no transport needed)	Biological by-products might be toxic

[Image comparing In-situ and Ex-situ bioremediation techniques]

---

10 Short Questions & Answers

1. **Q: Define Bioremediation. A:** The use of biological systems to reduce or eliminate environmental pollutants.
2. **Q: What are "Recalcitrant" compounds? A:** Chemicals that resist biodegradation and persist in the environment for a long time.
3. **Q: Why is moisture important for bioremediation? A:** Microbes require a water film for metabolic activity and the transport of nutrients.
4. **Q: What is the ideal pH range for most bioremediation microbes? A:** Between 6.5 and 8.5.
5. **Q: What is "Bioaugmentation"? A:** Adding specialized, high-performance microbes to a contaminated site to speed up the process.
6. **Q: What is "Biostimulation"? A:** Adding nutrients (N and P) to stimulate the growth of indigenous (local) microbes already present at the site.
7. **Q: Name one limitation of bioremediation. A:** It takes a significantly longer time (months to years) compared to chemical or physical removal.
8. **Q: Can heavy metals be "degraded" by microbes? A:** No; metals can only be transformed into less toxic forms or accumulated, but they cannot be destroyed.
9. **Q: What is "Phytoremediation"? A:** The use of plants to clean up contaminated soil or water.
10. **Q: Define *In-situ* bioremediation. A:** Treating the contaminated material directly at the site without moving it.

---

5 Long Questions & Answers

1. **Q: Discuss the importance of the "Microbial Population" in determining suitability.**
   A: For bioremediation to work, the site must contain microbes capable of degrading the specific pollutant. If the indigenous population is too small or lacks the necessary enzymes, the site is unsuitable unless Bioaugmentation is performed. Furthermore, these microbes must be in an active growth phase; if the environment contains inhibitors that prevent microbial metabolism, bioremediation will fail regardless of how many microbes are added.
2. **Q: Explain how soil texture and permeability influence the success of in-situ bioremediation.**
   A: Soil permeability determines how easily water, oxygen, and nutrients can move through the ground. Sandy soils have high permeability and are very suitable for bioremediation. In contrast,

clay soils have very small pores that restrict the flow of air and nutrients. If the microbes in the center of a clay patch cannot receive oxygen, they will die or become dormant, leaving the pollution untreated.

3. **Q: Compare Biostimulation and Bioaugmentation.**
   A: Biostimulation assumes that the right microbes are already there but are "hungry" or "suffocating"; it involves adding nutrients or pumping in oxygen. Bioaugmentation assumes the right microbes are missing; it involves adding lab-grown "super-bugs" to the site. Biostimulation is generally preferred because local microbes are already adapted to the site's specific climate and conditions.
4. **Q: What are the risks of "Incomplete Degradation"?**
   A: Sometimes, microbes start breaking down a pollutant but stop halfway. The resulting "intermediate" chemicals can sometimes be more toxic or more mobile than the original pollutant. For example, the breakdown of Trichloroethylene (TCE) can produce Vinyl Chloride, which is a much more potent carcinogen. Therefore, suitability studies must include an assessment of the potential metabolic pathways.
5. **Q: Why is bioremediation considered a "Cost-Effective" technology?**
   A: Traditional methods like "Dig and Dump" require heavy machinery, transportation to a landfill, and the cost of the landfill space. Incineration requires massive amounts of energy. Bioremediation, especially in-situ, uses natural processes and requires minimal equipment. The primary costs are simply the nutrients (fertilizers) and monitoring, making it significantly cheaper for large-scale spills.

---

15 Multiple Choice Questions (MCQs)

1. Bioremediation is most suitable for which type of pollutant?

   a) Heavy metals

   b) Organic hydrocarbons (oil/gas) ✓

   c) Radioactive waste

   d) Plastic bottles

2. Compounds that are resistant to microbial degradation are called:

   a) Biodegradable

   b) Labile

   c) Recalcitrant ✓

   d) Aerobic

3. Adding nitrogen and phosphorus to a site is called:

   a) Biostimulation ✓

b) Bioaugmentation

c) Biomagnification

d) Biosynthesis

4. In-situ bioremediation refers to treatment:

a) At the site of contamination ✓

b) In a laboratory

c) In a bioreactor

d) After moving soil to a landfill

5. Which soil type is MOST suitable for bioremediation?

a) Sand ✓

b) Clay

c) Silt

d) Rock

6. Aerobic bioremediation requires the presence of:

a) Methane

b) Oxygen ✓

c) Carbon dioxide

d) Darkness

7. The use of plants for remediation is known as:

a) Mycoremediation

b) Phytoremediation ✓

c) Bacterial remediation

d) Zooremediation

8. Microbial activity typically _____ as temperature increases (up to 35°C).

a) Increases ✓

b) Decreases

c) Stops

d) Fluctuates

9. A major disadvantage of bioremediation is that it is:
    a) Expensive
    b) Unnatural
    c) Time-consuming ✓
    d) Dangerous to humans
10. Bioaugmentation involves the addition of:
    a) Nutrients
    b) Oxygen
    c) Exogenous microorganisms ✓
    d) Water
11. The process where microbes turn contaminants into less toxic forms is:
    a) Biotransformation ✓
    b) Digestion
    c) Combustion
    d) Evaporation
12. Which of the following is a heavy metal NOT easily removed by bioremediation?
    a) Benzene
    b) Lead ✓
    c) Crude oil
    d) Phenol
13. Bioremediation is considered "Green Technology" because it:
    a) Uses natural processes ✓
    b) Always uses green plants
    c) Produces electricity
    d) Only works in forests
14. If a pollutant concentration is too high, it may become _____ to the microbes.
    a) A nutrient
    b) Toxic ✓

c) Invisible

d) Heavier

15. Which gas is often pumped into the ground to aid aerobic bioremediation?

a) Nitrogen

b) Air/Oxygen ✓

c) Helium

d) Argon