PREPARED BY VU SQUAD 0313-7248280
LMS AND PROJECTS HANDLER EXPERT

MATH-201 FINAL TERM QUIZES BY VU SQUAD

LMS HANDLING

ASSIGNMENT

QUIZ

GDB

LECTURE WATCH

CISCO ASSIGNMENT

PROJECTS

CS-519,619 MATH-600,620 MGT MKT EDU

ALL INTERSHIP AND THESIS

PREPARED BY VU SQUAD 0313-7248280
LMS AND PROJECTS HANDLER EXPERT

CONTACT ON WP / CALL 0313-7248280

SUBSCRIBE MY YOUTUBE CHANNEL VU SQUAD

https://www.youtube.com/@vusquad9329

Question:

\int\limits_0^{\frac{\pi }{2}} {\int\limits_0^1 {r\,\,dr\,\,d\theta } } \,\,\, = \,\,\, - - - - - -

Options:

A. \frac{\pi }{6}

B. \frac{\pi }{4}

C. \frac{\pi }{3}

D. \frac{\pi }{2}

Question:

{\text{The differential equation }}\,\,dz\, = \,8y\,dx\,\, + \,\,8x\,dy\,\,\,\,\,{\text{is}}\,{\text{an exact differential equation}}{\text{.}}

Options:

A. {\text{False}}

B. {\text{True}}

Question:

PREPARED BY VU SQUAD 0313-7248280
LMS AND PROJECTS HANDLER EXPERT

{\text{One of the line integral properties is}} \int\limits_{AB} {Fds = } - \int\limits_{BA} {Fds}

Options:

A. False

B. True

Question:

{\text{For a vector valued function }}\vec r(t){\text{ }} = {\text{ }}3{\text{ }}i + 4j - 0k{\text{ then lenght of }}\vec r(t){\text{ }}

Options:

A. \left\| r \right\| = 9

B. \left\| r \right\| = 5

C. \left\| r \right\| = 25

D. \left\| r \right\| = 4

Question:

Natural domain of a vector valued function is --------

Options:

A. intersection of the natural domains of its components

B. union of the natural domains of its components

C. not dependent on domains of its components

D. none of these

Question:

A vector-valued function is continuous at some point t0 if ----------

PREPARED BY VU SQUAD 0313-7248280
LMS AND PROJECTS HANDLER EXPERT

Options:

A. each of its component is differentiable at that point.

B. each of its component is continuous at that point.

C. some of its component is continuous at that point.

D. all of its component should be continuous every where in the domain.

Question:

{\text{If the path of integration c joining A and B is divided into two parts AK and KB, then}}

Options:

A. {I_c} = {I_{AK}} - {I_{KB}}

B. - {I_c} = {I_{AK}} + {I_{KB}}

C. {I_c} = - {I_{AK}} - {I_{KB}}

D. {I_c} = {I_{AK}} + {I_{KB}}

Question:

\int\limits_0^1 {\int\limits_0^1 {\int\limits_0^1 {xyz} } } \,dx\,\,dy\,\,dz = \,\,\, -

- - - - - -

Options:

A. \frac{1}{4}

B. \frac{1}{8}

C. \frac{1}{{10}}

D. \frac{1}{2}

PREPARED BY VU SQUAD 0313-7248280
LMS AND PROJECTS HANDLER EXPERT

Question:

{\text{If}}\,\,\vec r(t) = {t^2}\,\hat i + t\,\hat j + 3\,t\,\hat k,\,\,{\text{then}}\,\,\,\frac{d}{{dt}}\,\left[{\vec r(t)} \right] = _____.

Options:

A. \,2\,t\,\hat i + \hat j + 3\,\hat k

B. \,\,\hat i + t\,\hat j + 3\,\hat k

C. \,2\,t\,\hat i + t\,\hat j + 3\,\hat k

D. \,\,\hat i + \hat j + \,\hat k

Question:

{\text{For a vector valued function }}\vec r(t){\text{ }} = {\text{ }}\sqrt {2t} {\text{ }}i + (t + 1){\text{ }}j{\text{ then lenght of }}\vec r(t){\text{ }}

Options:

A. \left\| {r(t)} \right\| = \sqrt {4{t^2} + {{(t - 1)}^2}}

B. \left\| {r(t)} \right\| = \sqrt {2t + {{(t - 1)}^2}}

C. \left\| {r(t)} \right\| = \sqrt {2{t^2} + {{(t - 1)}^2}}

D. \left\| {r(t)} \right\| = \sqrt {4t + {{(t - 1)}^2}}