4/3/2023.

- — is a polymerase chain reaction that actually takes place inside cell on slide
  In situ
- In southern blotting fragments detec
  DNA
- Northern blotting is used for
  Gene expression
- Enzyme responsible for joining DNA fragments
  DNA ligase
- Sucide PCR is typically used in
  Paleogenetics
- Mutation in DNA can be identified by
  Southern blot
- RFLP: **Restriction fragment length poly**
- The type of PCR which defines bacterial yeast clone.
  Colony PCR
- WCA PCR stands for:-
- 10% DMSO work well as-
  Freezing medium
- — reduces nonspecific amplification during initial set up stages of PCR
  Hot start PCR.

- haplotype wla meqs (def thi).
- ___ determine specificity of PCR reaction
- **Primers**
- ___ reduce double stranded to single stranded.
- **Denaturation**
- ___ which reverse transcribes RNA into DNA
- **Reverse transcription**
- **Mono primer** — 16s RNA gene
- PCR has facilitated research in detecting of pathogens in - **plants, animals & environment**
- R-M system types
  4
- Radioactive probe
  P-32
- **SNP**.— Single nucleotide polymorphism

Date / /20

Subunits, two modification subunits and one recognition subunit? Type I

15 → Enzyme that catalyzes the template dependent synthesis of DNA is known as DNA polymerase

16 → Assembly PCR is used to assemble two or more pieces of DNA into one piece.

17 → Which is true for nonradioactive probe Biotin

18 → Transfer of denatured DNA fragments into a solid support Southern Blotting

19 → DNA ligase has been used to joint blunt ended DNA molecules T4

Date

- Palaeontology refers to the study of ancient human, animal and plant remains.
- WGA is stands for Whole genome amplification
- ALFP is the abbreviation of Amplified fragment length polymorphism
- How many Screening Methods SNPs have 2
- Immunosorbent Electron Microscopy contains antibodies (both (monoclonal & polyclonal))
- How many % attribute make a person unique 0.1%.
- Which of the following type of enzyme consists of two restriction

Date: / /20

Sequence genomes of a large number of people SNP Maps

Distance b/w the primer binding sites will determin the Size of PCR Product

Wet transfer is recommended for large proteins >100KD

During northen blot, Separation of RNA according to size through a Agarose gel

Second PCR Product is Shorter than first one Nested PCR

Fluorophore-containing DNA probes such as to measure the amount of amplified product in real time Taq Man

Date 1/120

1. What happens in the denature Step of PCR? Double-stranded DNA into Single Strand of DNA
2. Allele Specific PCR uses primers whose 3 ends encompass the SNP.
3. Suicide PCR is used in Paleogenetics
4. In situ PCR is a polymerase chain reaction that actually takes place inside the cell on a slide.
5. A heat stable enzyme called Taq Polymerase is Thermus aquaticus
6. Colony PCR is used for the screening of bacterial or yeast clones for correct ligation or plasmid products.
7. Minf PCR is used to amplify 16S rRNA gene.

Date:

26 → This diagnostic or cloning technique is used to identify or utilize single nucleotide polymorphisms Allele Specific PCR

27 → Applications of PCR: All

28 → RAPD is stands for Random Amplified polymorphic DNA

29 → Can repair the nicks produced after the association of cohesive ends of DNA Stands DNA ligase

Four base recognition site occurs ever 4^4 (256) bp