بِسۡمِ ٱللَّهِ ٱلرَّحۡمَٰنِ ٱلرَّحِيمِ

In the name of God, Most Gracious, Most Merciful



.....STUDY CHANELL LINK.....

join our WhatsApp channel

https://whatsapp.com/channel/0029VaODheTDp2Q8YVcoON2p

BIO 302 MIDTERM PART

OBJECTIVE

1. The term molecular biology was first used in **1945** by William Astbury.
2. The study of the chemical and physical structure of biological macromolecules is referred to as **molecular biology**.
3. By the time the term molecular biology was introduced, biochemists had already discovered many fundamental **intracellular** reactions.
4. The most advantageous approaches in molecular biology involve studying **bacteria** and bacteriophages.
5. Molecular biology became rapidly developed due to the belief in the basic **uniformity** of life processes.
6. Francis Crick and James Watson proposed the double helical model of **DNA** in 1953.
7. The double helix structure of DNA is tied together by **hydrogen** bonds between purines and pyrimidines.
8. The experiments by Griffith in 1928 showed the phenomenon of **transformation** in bacteria.
9. The DNA nature of the transforming principle was recognized in 1944 by Avery, MacLeod, and **McCarthy**.
10. Chargaff demonstrated that the number of adenine and thymine groups in DNA are **equal**.
11. The experiment by Hershey and Chase in 1952 showed that only the **DNA** of T2 bacteriophage enters the host.
12. The model for DNA proposed by Watson and Crick in 1953 consists of two **helically intertwined** chains.
13. The 1953 discovery of DNA structure was followed by investigations into the molecular nature of living cell reactions and the realization that all living organisms have **remarkable** similarities.
14. RNA molecules like ribosomal RNA, messenger RNA, and transfer RNA play a key role in **transcription**.
15. The process by which genetic information in RNA is used to synthesize a polypeptide is called **translation**.
16. In 1941, G.W. Beadle and E.L. Tatum studied the biochemical genetics of **Neurospora**.

17. The discovery of DNA's role in heredity was confirmed by Avery, MacLeod, and **McCarthy** in 1944.
18. In 1957, G. Beadle and E. Tatum received the **Nobel** Prize for their work on biochemical genetics.
19. In 1961, Nirenberg and Matthaei cracked the **messenger** RNA code.
20. The concept of the operon, crucial to gene regulation, was introduced by **Jacob** and Monod in 1961.
21. In 1968, Watson and Crick received the **Nobel** Prize for the discovery of the molecular nature of DNA.
22. The first recombinant DNA molecules were produced in 1972 by Mertz and **Davis**.
23. In 1973, Cohen and others successfully cloned DNA using **plasmids** as a vector.
24. In 1977, Sharp and Roberts discovered the phenomenon of **split** genes in adenoviruses.
25. Khorana, in 1979, completed the total synthesis of a biologically **functional** gene.
26. The technique for transferring mRNA bands from a gel to a membrane is called **Northern** blotting.
27. In 1983, Kary Mullis discovered the **polymerase chain** reaction (PCR), revolutionizing gene cloning.
28. In 1984, Alec Jeffreys discovered the technique of **DNA** fingerprinting.
29. The discovery of the role of RNA molecules as enzymes (ribozymes) won the Nobel Prize in 1989 for **Cech** and Altman.
30. The technique of **DNA** was developed by S. Tonegawa in 1987 to study DNA sequence rearrangements in immunoglobulin genes.
31. In 1987, researchers discovered the technique for high-speed introduction of foreign DNA into plant cells, known as the **particle** bombardment gun.
32. The sequencing of a 5400 nucleotide DNA sequence from the φ × 174 bacteriophage was accomplished by **Fredrick Sanger**.
33. The Nobel Prize for providing the three-dimensional structure of tRNA in 1982 went to **A. Klug**.
34. In 1984, Robert Tijan identified the DNA-binding protein called **SP1** involved in gene regulation.
35. The scanning hypothesis for eukaryotic translation initiation was proposed by **Marilyn Kozak** in 1983.
36. The western blotting technique, used to detect proteins, was developed by **Towbin** in 1979.
37. The Nobel Prize in 1975 for the discovery of reverse transcriptase was awarded to **Howard Temin** and Baltimore.

38. In 1972, **Southern** demonstrated the technique of Southern blotting for analyzing related genes in DNA.
39. The Nobel Prize in 1982 for the creation of transgenic mice was awarded to **Palmiter** and Brinster.
40. The transformation of yeast cells using plasmids was first described by **Hinnen** in 1978.
41. In 1978, the term **exon** was coined by W. Gilbert for genes that are interrupted by non-coding sequences.
42. The discovery of split genes in adenoviruses was made by Sharp and **Roberts** in 1977.
43. The method developed by Marilyn Kozak in 1983 is known as the **scanning** hypothesis for translation initiation.
44. The molecule that catalyzes the polymerization of DNA in the PCR process is **Taq** polymerase.
45. The discovery of the chemical structure of antibodies led to the Nobel Prize in Physiology in 1972 for **Porter** and Edelman.

SUBJECTIVE

---

1. What is the significance of epigenetics in gene regulation, and how has it evolved in recent years?

Epigenetics plays a crucial role in gene regulation by mediating changes in gene activity without altering the DNA sequence. Over the past decade, it has become clear that chromatin serves as a platform for post-translational modifications of DNA and histones, which act as signaling mechanisms for various cellular processes. These modifications can influence gene expression, development, and disease states. Advances in this field have revealed the dynamic nature of epigenetic markers and their potential for reversibility, making them promising targets for therapeutic interventions. The discoveries surrounding non-coding RNAs and their regulatory roles have further enriched our understanding of

epigenetics, paving the way for groundbreaking research into long non-coding RNAs and their functions.

---

2. How did Shinya Yamanaka and Kazutoshi Takahashi revolutionize the field of stem cell research?

Shinya Yamanaka and Kazutoshi Takahashi made a paradigm-shifting discovery in stem cell research by demonstrating that a combination of four transcription factors—OCT4, SOX2, MYC, and KLF4—could reprogram differentiated cells, such as mouse skin fibroblasts, into induced pluripotent stem (iPS) cells. This breakthrough showed that mature cells could revert to a pluripotent state similar to embryonic stem cells, offering immense potential for regenerative medicine. This technique enables the creation of patient-specific iPS cells, which can be used to model diseases, test drugs, and potentially develop personalized therapies. Their work marked a significant milestone in understanding cellular plasticity and has since stimulated extensive research in stem cell biology.

---

3. What is the reaction-diffusion mechanism, and how has it contributed to our understanding of cellular processes?

The reaction-diffusion mechanism is a system in which signaling molecules create gradients that organize spatial information within cells. This concept, initially proposed by Eric Karsenti in the 1980s, gained empirical support with the development of FRET-based biosensors in the past decade. Gradients of molecules like RAN•GTP and Aurora B have been shown to play crucial roles in mitotic chromatin signaling and cytokinesis. The mechanism has expanded our understanding of intracellular spatial organization, illustrating how cells utilize chemical gradients to coordinate complex processes such as division and development. It has also bridged developmental biology and cellular dynamics by highlighting the universality of gradient-based signaling.

---

4. How has the study of metabolism reshaped our understanding of growth control and disease?

Recent research has highlighted the intricate link between metabolism and growth control, particularly in the context of diseases like cancer. Autophagy, a conserved

cellular process for recycling cellular components, has emerged as a key mechanism that is often deregulated in human diseases. Studies have unveiled the roles of central regulators like mTOR, sirtuins, and the Hippo pathway in maintaining cellular homeostasis. These findings have transformed our understanding of how metabolic pathways influence cell proliferation, survival, and death, providing insights into potential therapeutic targets for metabolic and cancer-related disorders.

---

5. What are some key technical advancements that have revolutionized molecular biology in recent years?

Several technical advancements have reshaped molecular biology, including high-throughput sequencing, RNA interference (RNAi), and super-resolution microscopy techniques like PALM, STORM, and STED. These technologies allow for precise manipulation, imaging, and analysis of cellular and genetic components. Additionally, advances in bioinformatics and computational modeling have enabled scientists to process and interpret large datasets, facilitating genome-wide studies. Techniques like ChIP–seq and fluorescence biosensors have further deepened our understanding of gene regulation and protein interactions, marking a new era of integrative and interdisciplinary research.

---

6. How has the genomic revolution influenced biological research and its applications?

The genomic revolution, driven by advancements in sequencing technologies, has transformed biological research by enabling rapid and cost-effective sequencing of entire genomes. This has facilitated comparative genomics, identification of genetic mutations, and the study of coding and non-coding RNAs. The insights gained have advanced fields like personalized medicine, evolutionary biology, and microbial ecology. For example, studies on the human microbiome have revealed the profound impact of microbial communities on human health and disease. The ability to analyze genome-wide data has also opened avenues for developing targeted therapies and understanding complex biological systems.

---

7. What role do non-coding RNAs play in cellular processes, and why are they significant?

Non-coding RNAs, including microRNAs (miRNAs) and long non-coding RNAs (lncRNAs), play critical roles in regulating gene expression and various cellular processes. miRNAs typically function by targeting messenger RNAs for degradation or translation inhibition, while lncRNAs can modulate chromatin structure, transcription, and post-transcriptional processes. These molecules have been implicated in diverse biological phenomena, from development and differentiation to stress responses and disease mechanisms. The discovery of their regulatory functions has unveiled new layers of complexity in gene expression and highlighted their potential as diagnostic markers and therapeutic targets.

---

8. How have interdisciplinary collaborations enhanced molecular biology research?

Interdisciplinary collaborations have become essential in modern molecular biology, bringing together expertise from fields like bioinformatics, physics, chemistry, and engineering. For instance, bioinformaticians help process and analyze large genomic datasets, while physicists and chemists contribute to developing advanced imaging and analytical tools. Collaborative efforts have led to breakthroughs like super-resolution microscopy, high-throughput sequencing, and RNAi technology. By integrating diverse perspectives and methodologies, these partnerships have accelerated discoveries, enhanced our understanding of biological systems, and driven innovations in areas like drug development and regenerative medicine.

---

9. What is the significance of RNAi technology, and how has it advanced biological research?

RNA interference (RNAi) technology has revolutionized gene function studies by enabling targeted knockdown of specific genes. This technique allows researchers to silence genes systematically, providing insights into their roles in cellular processes and disease mechanisms. RNAi has been instrumental in functional genomics, drug discovery, and therapeutic development. It has also been used to identify and validate molecular targets for treating diseases such as cancer and viral infections. The precision and versatility of RNAi continue to make it a cornerstone of molecular biology research.

10. How has the study of chromatin structure contributed to our understanding of genome organization and gene expression?

The study of chromatin structure has provided foundational insights into genome organization and gene expression. Chromatin, composed of DNA and histone proteins, undergoes dynamic modifications that regulate access to genetic information. Epigenetic markers, such as DNA methylation and histone modifications, influence chromatin compaction and gene activity. Advances in chromatin analysis have elucidated how these modifications govern cell identity, differentiation, and response to environmental cues. Genome-wide studies have further linked chromatin alterations to various diseases, highlighting their significance in both normal development and pathological conditions.

11. Explain the importance of nucleic acids in biological systems.

Answer:
Nucleic acids, including DNA and RNA, are critical biopolymers that store, transmit, and execute genetic information. DNA encodes hereditary information in the form of genes, which direct the synthesis of proteins vital for cellular functions. RNA, on the other hand, plays a pivotal role in converting DNA-encoded information into proteins. It serves as a messenger (mRNA), a catalytic component of ribosomes (rRNA), and a transporter of amino acids (tRNA). Without nucleic acids, the processes of heredity, evolution, and cellular function would cease, making them indispensable for life.

12. Describe the structural differences between DNA and RNA.

Answer:
The primary structural differences between DNA and RNA include:

1. **Sugar Molecule:** DNA contains deoxyribose, while RNA contains ribose. The absence of an oxygen atom on the 2' carbon of deoxyribose distinguishes DNA.
2. **Strand Configuration:** DNA is typically double-stranded, forming a double helix, whereas RNA is usually single-stranded.

3. **Nitrogenous Bases:** DNA uses thymine (T) as one of its bases, while RNA replaces thymine with uracil (U).
4. **Function:** DNA serves as the genetic blueprint, while RNA functions in protein synthesis and other cellular roles.

---

13. What is the significance of the double-helix structure of DNA?

Answer:
The double-helix structure of DNA, discovered by Watson and Crick, is critical for its function. Its antiparallel strands allow for complementary base pairing (A-T, C-G), enabling accurate replication and transcription. The helical structure provides stability while compactly storing genetic information within chromosomes. Additionally, the accessibility of the bases allows for efficient transcription and repair mechanisms, ensuring genetic fidelity.

---

14. How does DNA replication ensure genetic consistency during cell division?

Answer:
DNA replication is a semi-conservative process where each new DNA molecule contains one original and one newly synthesized strand. Enzymes like DNA helicase unwind the double helix, and DNA polymerase synthesizes new strands by complementary base pairing. Proofreading mechanisms correct errors, ensuring genetic information is accurately copied. This process guarantees that each daughter cell inherits an identical set of DNA.

---

15. Discuss the types and functions of RNA in protein synthesis.

Answer:
There are three primary types of RNA involved in protein synthesis:

1. **Messenger RNA (mRNA):** Carries genetic instructions from DNA to ribosomes.
2. **Ribosomal RNA (rRNA):** Forms the structural and catalytic core of ribosomes, facilitating peptide bond formation.

3. **Transfer RNA (tRNA):** Delivers specific amino acids to the ribosome based on codons in the mRNA.
   Together, these RNAs ensure the accurate translation of genetic information into proteins.

---

16. Explain the chemical composition of a nucleotide.

Answer:
A nucleotide comprises three components:

1. **Nitrogenous Base:** Can be a purine (adenine or guanine) or a pyrimidine (cytosine, thymine in DNA, or uracil in RNA).
2. **Sugar Molecule:** Ribose in RNA or deoxyribose in DNA.
3. **Phosphate Group:** One to three phosphate groups provide the nucleotide with energy and allow polymerization.
   These components link to form DNA or RNA, with sequences encoding genetic information.

---

17. How are nucleosides different from nucleotides?

Answer:
A nucleoside consists of a nitrogenous base covalently bonded to a sugar (ribose or deoxyribose) but lacks a phosphate group. A nucleotide, however, includes a nitrogenous base, sugar, and one to three phosphate groups. Nucleotides play essential roles in DNA and RNA synthesis and serve as energy carriers (e.g., ATP).

---

18. Describe the phosphodiester bond formation in DNA.

Answer:
Phosphodiester bonds link nucleotides to form the DNA backbone. During DNA synthesis, the phosphate group of one nucleotide's 5' carbon reacts with the hydroxyl group on the 3' carbon of the previous nucleotide, releasing a water molecule. This bond creates a sugar-phosphate backbone, ensuring the structural integrity and directionality (5' to 3') of DNA strands.

---

19. What are artificial nucleic acid analogs, and how are they used?

Answer:
Artificial nucleic acid analogs, such as peptide nucleic acids and locked nucleic acids, are synthetic molecules that mimic DNA or RNA. They differ in their backbone structures, enhancing stability or specificity. These analogs are used in genetic research, therapeutic applications, and diagnostics, such as targeting specific genes in gene silencing or designing antisense therapies.

---

20. Summarize the process of transcription in DNA.

Answer:
Transcription is the process where DNA is transcribed into RNA. It involves:

1. **Initiation:** RNA polymerase binds to the promoter region of a gene.
2. **Elongation:** RNA polymerase reads the DNA template strand and synthesizes a complementary RNA strand by adding ribonucleotides.
3. **Termination:** Transcription ends when RNA polymerase reaches a termination signal.
   The resulting mRNA is processed (e.g., splicing, capping) before translation into proteins.

---

21. What are the essential characteristics of genetic material, and why is DNA considered the hereditary material?

Answer:
Genetic material must fulfill several essential criteria:

1. **Complex Information**: It should carry detailed instructions for the entire structure and function of an organism. DNA's sequence of four nucleotides (adenine, thymine, cytosine, and guanine) provides this complexity.
2. **Faithful Replication**: It must replicate accurately to ensure that genetic information is passed down without errors during cell division. DNA replication uses complementary base pairing, ensuring fidelity.

3. **Phenotype Encoding**: It must direct the synthesis of proteins, which determine the traits of an organism. DNA achieves this through the transcription of mRNA and subsequent protein synthesis.

Experiments by Griffith, Avery, and Hershey-Chase demonstrated that DNA, not proteins, is the genetic material, as DNA could transform bacteria, retain its activity after protein digestion, and enter cells during bacteriophage infection.

---

22. Explain Griffith's experiment and its significance in understanding genetic material.

Answer:
Griffith's 1928 experiment demonstrated the phenomenon of transformation. He used two strains of *Streptococcus pneumoniae*:

- **Type III-S (smooth)**: Virulent, protected by a polysaccharide capsule.
- **Type II-R (rough)**: Non-virulent, lacking the capsule.

Griffith heat-killed the III-S strain and mixed its remains with live II-R strain bacteria. Injecting this combination into mice resulted in death, with live III-S bacteria recovered from the mice.

Conclusion:
Griffith concluded that a "transforming principle" from the dead III-S bacteria enabled the II-R bacteria to become virulent. This laid the foundation for identifying DNA as the genetic material.

---

23. Discuss Avery's experiments and how they proved DNA is the transforming principle.

Answer:
In 1944, Avery, MacLeod, and McCarty purified the transforming substance from Griffith's experiment. They demonstrated that:

1. The substance matched DNA in its chemical properties.
2. Protein-digesting enzymes and ribonuclease (RNA-digesting enzyme) had no effect on transformation.

3. DNA-digesting enzymes destroyed the transforming ability.

Their findings strongly suggested that DNA is the hereditary material, not proteins or other molecules, marking a pivotal shift in molecular biology.

---

24. How did the Hershey-Chase experiment confirm DNA as the genetic material?

Answer:
Hershey and Chase conducted experiments in 1952 using bacteriophages (viruses that infect bacteria).

- They labeled DNA with radioactive phosphorus (^32P) and protein with radioactive sulfur (^35S).
- Infected bacteria retained ^32P (DNA) but not ^35S (protein).
- Newly formed phages also contained radioactive DNA but not protein.

Conclusion:
These results confirmed that DNA, not protein, is transferred to bacterial cells during infection and is responsible for genetic inheritance.

---

25. What is the role of proteins in genetic material and why was it initially assumed to be the genetic material?

Answer:
Proteins were initially considered genetic material due to their structural complexity and diverse functionality. They are made of 20 amino acids, providing a variety of combinations that could hypothetically encode genetic information.

However, experiments like those of Avery, Griffith, and Hershey-Chase demonstrated that DNA, despite being simpler in composition, is the true carrier of hereditary information. Proteins play critical roles as enzymes and structural components but do not store genetic information.

---

26. Discuss the chemical composition of proteins and their primary structure.

Answer:
Proteins are polymers made of amino acids linked by peptide bonds. Their **chemical composition** includes:

- **Amino Acids**: Building blocks, each with an amino group ($-NH_2$), a carboxyl group (-COOH), and a unique side chain (R group).
- **Peptide Bond**: A covalent bond formed between the carboxyl group of one amino acid and the amino group of another.

Primary Structure:
This refers to the unique sequence of amino acids in a polypeptide chain, determining the protein's overall shape and function. Even slight changes in the sequence can alter the protein's properties.

---

27. Define secondary structure of proteins and describe the α-helix.

Answer:
Secondary Structure: It refers to the local three-dimensional arrangement of protein segments, primarily stabilized by hydrogen bonds. Two common structures are α-helix and β-sheets.

α-Helix:

- A right-handed coil stabilized by hydrogen bonds between the carbonyl oxygen of one amino acid and the amide hydrogen of another four residues ahead.
- Each turn of the helix has about 3.6 amino acids.
- The α-helix is common in structural proteins like keratin and contributes to their elasticity.

---

28. What are the β-sheets in protein secondary structures, and how are they stabilized?

Answer:
β-Sheets:

- These are formed by extended polypeptide chains lying side-by-side, connected by hydrogen bonds between carbonyl and amine groups of the backbone.
- Chains can be arranged in **parallel** (same direction) or **antiparallel** (opposite directions) configurations.

Stabilization:

- Hydrogen bonds provide structural stability.
- The planarity of peptide bonds and the flexibility of the polypeptide backbone contribute to forming β-sheets.

They are common in fibrous proteins like silk fibroin.

---

29. What is the significance of peptide bonds in protein structure?

Answer:
Peptide bonds are the covalent links between amino acids in proteins. **Key features**:

1. **Planarity**: The peptide bond is rigid and planar due to its partial double-bond character.
2. **Stability**: It resists hydrolysis under physiological conditions, ensuring protein integrity.
3. **Orientation**: Peptide bonds have a specific orientation, contributing to the protein's primary structure.

Peptide bonds are crucial for maintaining the protein's structure and enabling folding into higher-order structures.

---

30. How does protein folding depend on primary and secondary structures?

Answer:
Protein folding is a hierarchical process:

1. **Primary Structure**: The amino acid sequence determines how the protein will fold. Specific sequences favor particular secondary and tertiary structures.
2. **Secondary Structure**: Local structures like α-helices and β-sheets form first, stabilized by hydrogen bonds.
3. **Tertiary Structure**: Interactions between side chains (R-groups) fold the protein into its final three-dimensional shape.

Misfolding can result in nonfunctional or toxic proteins, often linked to diseases like Alzheimer's and Parkinson's.

---

31. How is the genetic material organized in eukaryotic cells?

Answer:
In eukaryotic cells, the genetic material is segmented into linear chromosomes, unlike the circular DNA in bacteria. A eukaryotic genome comprises all the unique chromosomes in a haploid set, while diploids and polyploids have multiple copies. The DNA is wrapped around histone proteins, forming nucleosomes, which are further supercoiled and compacted during cell division. Proteins such as condensin and cohesin play roles in chromosomal structure and segregation during mitosis and meiosis. Regulatory sequences along the chromosomes control gene expression, and the chromosomes are divided into euchromatin (active) and heterochromatin (inactive) regions based on their compaction levels.

---

32. What are histone proteins, and what roles do they play in DNA organization?

Answer:
Histones are alkaline proteins abundant in eukaryotic cell nuclei, crucial for DNA packaging. They act as spools around which DNA winds, forming nucleosomes. The five classes of histones—H1, H2A, H2B, H3, and H4—facilitate chromatin structure and regulation of gene expression. The positively charged amino acids, lysine and arginine, bind to the negatively charged DNA, stabilizing the chromatin. Without histones, the DNA's extensive length would make nuclear packaging unmanageable. For example, 1.8 meters of human DNA is compacted into about 90 micrometers of chromatin.

---

33. Describe the structure and significance of nucleosomes.

Answer:
Nucleosomes are the basic units of DNA packaging in eukaryotic cells. They consist of DNA (~146 bp) wrapped around an octamer of histones (two each of H2A, H2B, H3, and H4). Linker DNA connects adjacent nucleosomes, and the histone H1 helps secure the structure. Nucleosomes compact the DNA and regulate access for processes like transcription and replication. They also carry epigenetic information through histone modifications, influencing gene expression. Observed as "beads on a string" under an electron microscope, nucleosomes enable efficient nuclear packing of the genome while maintaining functionality.

---

34. What are the 30-nm fibers, and what models describe their structure?

Answer:
The 30-nm fiber is a higher-order chromatin structure formed by folding chains of nucleosomes. Two major models explain its formation:

1. **Zigzag Model:** Linker DNA forms a straight path between nucleosomes, arranging them on opposite sides of the fiber.
2. **Solenoid Model:** The nucleosome chain forms a helical structure, with about six nucleosomes per turn, and bent linker DNA connects neighboring nucleosomes.

Both models highlight how chromatin achieves compaction beyond nucleosomes, although their biological context remains under study.

---

35. How does DNA replication occur, and what is the semiconservative model?

Answer:
DNA replication involves copying the DNA to produce two identical molecules. According to the semiconservative model, each parental DNA strand serves as a template for a new complementary strand. The daughter DNA molecules consist of one parental and one newly synthesized strand, ensuring genetic consistency across generations. This model, validated by the Meselson-Stahl experiment, contrasts with alternative conservative and dispersive models, which were disproven.

36. Describe the Meselson-Stahl experiment and its significance.

Answer:
The Meselson-Stahl experiment demonstrated that DNA replication is semiconservative. They labeled E. coli DNA with heavy nitrogen (^15N) and then transferred the cells to a medium with light nitrogen (^14N). Density gradient centrifugation revealed intermediate and light-density DNA after replication cycles, consistent with one parental and one new strand per molecule. This elegant experiment confirmed the Watson-Crick model of DNA replication.

37. What are the two key components required for DNA synthesis, and what are their roles?

Answer:

1. **Deoxynucleoside triphosphates (dNTPs):** These serve as the building blocks for DNA synthesis, providing the nucleotide units added to the growing DNA strand.
2. **Primer-template junction:** This provides a starting point for DNA polymerase to add dNTPs, where the primer offers a free 3' hydroxyl group required for chain elongation.
   Together, these components ensure accurate replication of the DNA template.

38. How is DNA packaged in eukaryotic cells, and how does this differ from bacterial DNA packaging?

Answer:
Eukaryotic DNA is packaged around histone proteins into nucleosomes, which fold into higher-order structures like the 30-nm fiber. These structures compact the DNA and regulate gene accessibility. Unlike eukaryotes, bacterial DNA lacks histones and is organized into a single circular chromosome with supercoiled loops anchored to a protein matrix. While both systems achieve DNA compaction, the mechanisms differ significantly due to structural and functional variances.

BIO 302 FINALTERM PART

OBJECTIVE

1. The **Central Dogma** refers to the process by which the instructions in ___ are converted into a functional product.
   Answer: DNA
2. The Central Dogma was first proposed in 1958 by ___, one of the discoverers of DNA's structure.
   Answer: Francis Crick
3. The flow of genetic information follows the sequence: DNA → ___ → Protein.
   Answer: RNA
4. RNA acts as a ___ that carries genetic information to the ribosomes.
   Answer: messenger
5. Ribosomes serve as ___ where information is translated into functional products.
   Answer: factories
6. The process of converting DNA instructions into functional products is called ___.
   Answer: gene expression
7. The two key stages of gene expression are ___ and ___.
   Answer: transcription, translation
8. During transcription, information in ___ is converted into RNA messages.
   Answer: DNA
9. Translation occurs in the ___, where mRNA messages are read to make proteins.
   Answer: ribosomes

10. DNA replication involves the copying of DNA to make new ___.
    Answer: DNA
11. Reverse transcription occurs when information is transferred from ___ to ___.
    Answer: RNA, DNA
12. Reverse transcription is a key process in retroviruses, such as ___.
    Answer: HIV
13. The Central Dogma suggests that ___ contains all the information needed to make proteins.
    Answer: DNA
14. The enzyme responsible for transcription is ___.
    Answer: RNA polymerase
15. Transcription produces ___, which must be processed before translation.
    Answer: pre-mRNA
16. The 5' cap and ___ tail are added to pre-mRNA during processing.
    Answer: poly-A
17. The mature RNA product of transcription is called ___.
    Answer: mRNA
18. Translation begins with the ___ codon, which is typically AUG.
    Answer: start
19. The amino acid associated with the AUG codon is ___.
    Answer: methionine
20. During translation, ___ molecules carry amino acids to the ribosome.
    Answer: tRNA
21. The anticodon on tRNA matches the ___ on mRNA.
    Answer: codon
22. Translation ends with a ___ codon, such as UAA, UAG, or UGA.
    Answer: stop
23. Proteins often require additional ___ to become functional after translation.
    Answer: processing
24. Proteins that assist in folding other proteins are called ___.
    Answer: chaperones
25. The enzymes involved in DNA replication include ___, which unwinds the DNA helix.
    Answer: helicase
26. The enzyme that synthesizes new DNA strands is called ___.
    Answer: DNA polymerase
27. The small RNA fragments required to initiate DNA replication are added by ___.
    Answer: RNA primase

28. Okazaki fragments are joined together by ___.
    Answer: DNA ligase
29. RNA-dependent RNA polymerase is involved in ___ replication.
    Answer: RNA
30. Special transfers include RNA to DNA and direct translation from ___ to protein.
    Answer: DNA
31. The Central Dogma states that information cannot flow from ___ to RNA or DNA.
    Answer: protein
32. In prokaryotes, transcription and translation occur ___, without clear separation.
    Answer: simultaneously
33. In eukaryotes, transcription occurs in the ___, and translation occurs in the ___.
    Answer: nucleus, cytoplasm
34. The genetic code is read in groups of three bases, called ___.
    Answer: codons
35. The Central Dogma describes three major biopolymers: DNA, RNA, and ___.
    Answer: protein
36. Alternative ___ increases the diversity of proteins produced from a single gene.
    Answer: splicing
37. The enzyme telomerase uses ___ as a template for reverse transcription.
    Answer: RNA
38. Transfer RNA (tRNA) is part of the ___ family.
    Answer: ribonucleic acid
39. RNA editing involves altering an RNA sequence with the help of guide RNA and ___.
    Answer: proteins
40. The ribosome reads mRNA codons from the ___ to the ___ direction.
    Answer: 5', 3'
41. The functional product of translation is a ___.
    Answer: protein
42. tRNA molecules bring ___ to the ribosome during translation.
    Answer: amino acids
43. A nascent polypeptide chain must fold into its correct ___ to become functional.
    Answer: conformation

44. DNA replication typically occurs during the ___ phase of the cell cycle.
Answer: S
45. A polypeptide chain that undergoes processing to remove internal segments releases ___.
Answer: inteins

SUBJECTIVE

1. **What is the primary function of tRNA in the cell?**

The primary function of transfer RNA (tRNA) is to translate the genetic code carried by messenger RNA (mRNA) into a sequence of amino acids to form proteins. During translation, tRNA reads the mRNA codons, each consisting of three nucleotides, and brings the corresponding amino acid to the ribosome. Each tRNA molecule has a specific anticodon that pairs with a complementary mRNA codon, ensuring the correct amino acid is added to the growing polypeptide chain.

2. **How does the redundancy in the genetic code affect tRNA function?**

The genetic code's redundancy means that multiple codons can code for the same amino acid. For example, the amino acid valine is coded by GUU, GUC, GUA, and GUG. Despite this redundancy, each codon has a specific tRNA that carries the appropriate amino acid, ensuring no ambiguity in

protein synthesis. This characteristic helps maintain precision in translation while allowing for genetic variations without altering the resulting protein.

3. **What distinguishes bacterial transcription from eukaryotic transcription?**

Bacterial transcription occurs in the cytoplasm and can happen simultaneously with translation, as there is no nuclear membrane to separate these processes. In contrast, eukaryotic transcription occurs in the nucleus, and translation takes place in the cytoplasm. Additionally, eukaryotic transcription involves three types of RNA polymerases with distinct functions, whereas bacteria have a single RNA polymerase for all transcription processes.

4. **What roles do the three eukaryotic RNA polymerases play?**

- **RNA Polymerase I (Pol I):** Transcribes ribosomal RNA (rRNA) genes, producing 28S, 18S, and 5.8S rRNAs, which are critical components of ribosomes.
- **RNA Polymerase II (Pol II):** Transcribes messenger RNA (mRNA) and other small RNAs like snRNAs, siRNAs, and miRNAs. These RNAs play roles in protein synthesis and regulation.
- **RNA Polymerase III (Pol III):** Produces small RNAs, including tRNAs, 5S rRNA, and other non-coding RNAs essential for various cellular functions.

5. **How does the initiation phase of transcription differ between bacteria and eukaryotes?**

In bacteria, RNA polymerase binds to promoter regions with the help of a

sigma factor to form a holoenzyme. This recognizes specific sequences like the -10 and -35 regions. In eukaryotes, RNA polymerase II requires several general transcription factors to assemble on the promoter and form the preinitiation complex. Eukaryotic promoters often include a TATA box and other regulatory elements.

6. **What is the function of the carboxyl terminal domain (CTD) of RNA Polymerase II?**

The CTD of RNA Polymerase II plays a crucial role in coordinating transcription with RNA processing. Its heptapeptide repeats (YSPTSPS) undergo phosphorylation and other post-translational modifications during transcription. These modifications regulate the transitions between initiation, elongation, and termination phases and facilitate the recruitment of factors necessary for RNA splicing, capping, and polyadenylation.

7. **What is intrinsic termination in bacterial transcription?**

Intrinsic termination, also known as Rho-independent termination, relies on specific sequences in the RNA transcript. These sequences form a stem-loop hairpin structure followed by a string of uracil residues. The hairpin causes the RNA polymerase to pause and destabilize, leading to the dissociation of the RNA transcript and polymerase from the DNA template.

8. **How do transcription factors influence eukaryotic transcription?**

Transcription factors bind to specific DNA sequences near or within promoter regions and either activate or repress transcription. They assist RNA polymerase in recognizing promoters, stabilize the transcriptional machinery, and modulate gene expression in response to cellular signals. Some transcription factors also interact with enhancers or silencers to regulate transcription over long distances.

9. **Why is simultaneous transcription and translation possible in bacteria but not in eukaryotes?**

In bacteria, the absence of a nuclear membrane allows ribosomes to attach to mRNA transcripts as they are being synthesized. This enables transcription and translation to occur concurrently. In eukaryotes, transcription occurs in the nucleus, while translation takes place in the cytoplasm. The separation of these processes is necessary to allow for RNA processing events like splicing, capping, and polyadenylation.

10. **What are the main stages of transcription, and what happens during each?**

- **Initiation:** RNA polymerase binds to the promoter, and the DNA strands are separated to allow the enzyme to access the template strand.
- **Elongation:** RNA polymerase synthesizes the RNA transcript by adding nucleotides complementary to the DNA template strand.
- **Termination:** Transcription ends when RNA polymerase encounters a terminator sequence. The RNA transcript is released, and the polymerase detaches from the DNA template.

11. Explain the process of gene-specific repression in eukaryotic transcription.

Gene-specific repression is a mechanism by which eukaryotic transcription is inhibited. Eukaryotic transcription repressors can act through several strategies. One way is by binding to DNA sites that overlap with activator binding sites, preventing the activator from binding. More frequently, repressors inhibit the function of activators by masking their activating domains, preventing their nuclear localization, promoting degradation, or chemically modifying them. Additionally, repressors can directly interfere with transcription initiation by binding to upstream promoter regions and interacting with the transcriptional machinery. They can also recruit histone modifiers (such as deacetylases and methylases) or nucleosome remodeling enzymes, which decrease DNA accessibility, thereby repressing

transcription. This gene silencing can spread along the chromatin, affecting multiple genes simultaneously.

12. What are the mechanisms behind elongation and termination control in transcription?

Elongation and termination control are vital for regulating transcription. Elongation starts after the transcription complex is assembled and continues until a termination sequence is encountered. Some genes, especially in eukaryotes, experience promoter-proximal pausing, where RNA polymerase pauses shortly after initiation. This pause, influenced by factors like DSIF and NELF, affects chromatin structure and allows for rapid or coordinated transcription in response to activation signals. Release of this pause is controlled by the recruitment of P-TEFb kinase. Transcription termination is also tightly regulated, involving efficient recycling of RNA polymerase and gene looping, which ensures proper termination and re-initiation. Termination factors can influence gene looping, which in turn determines how effectively transcription can restart.

13. Describe the process of transcription-coupled DNA repair.

Transcription-coupled DNA repair (TC-NER) is a mechanism that repairs DNA lesions that interfere with transcription. When RNA polymerase encounters a DNA lesion, it stalls, which signals the recruitment of DNA repair proteins. Central to this process is the general transcription factor TFIIH, which has ATPase activity. TFIIH causes a conformational change in the RNA polymerase, opening up the transcription bubble to allow repair enzymes to access the damaged DNA strand. This repair mechanism ensures that actively transcribed genes are protected from mutations, making RNA polymerase itself a damage-sensing protein that directs repair enzymes to the site of transcription.

14. Compare prokaryotic and eukaryotic transcription in terms of complexity and regulation.

Eukaryotic transcription is far more complex than prokaryotic transcription. In eukaryotes, DNA is confined to the nucleus, separated from the cytoplasm where translation occurs. This separation allows for temporal regulation of gene expression and selective transport of mRNAs to the cytoplasm. In contrast, bacteria do not have a nuclear membrane, so transcription and translation occur simultaneously, providing an efficient mechanism for gene regulation. In terms of transcription initiation, prokaryotic RNA polymerase binds strongly to promoters

and initiates transcription with a high basal rate, while in eukaryotes, transcription initiation is more complex due to chromatin structure, requiring a multi-protein complex and ATP hydrolysis to begin transcription. As a result, eukaryotic transcription initiation tends to occur at a lower basal rate.

15. What is RNA splicing, and why is it necessary in eukaryotic cells?

RNA splicing is the process of removing introns (non-coding sequences) from pre-mRNA and joining the exons (coding sequences) together. This process is crucial for the maturation of mRNA in eukaryotic cells, as it ensures that only the coding sequences are retained for translation into proteins. Since most eukaryotic genes contain introns, their removal is essential for the proper functioning of the gene. In addition, splicing allows for alternative splicing, where different combinations of exons can be joined, leading to multiple protein variants from a single gene.

16. What is the role of the spliceosome in pre-mRNA splicing?

The spliceosome is a large and complex molecular machine that performs the splicing of pre-mRNA in eukaryotic cells. It is composed of small nuclear RNAs (snRNAs) and numerous protein factors. The spliceosome recognizes the splice sites at the junctions of introns and exons and catalyzes the removal of introns. The snRNAs (U1, U2, U4, U5, and U6) are essential components of the spliceosome, interacting with each other and with the pre-mRNA to facilitate the splicing process. The spliceosome's activity results in the precise removal of introns and the ligation of exons, producing a mature mRNA molecule.

17. Describe the mechanisms involved in RNA splicing by the spliceosome.

RNA splicing begins with the recognition of splice sites at the 5' and 3' ends of the introns. The U1 snRNA binds to the 5' splice site, and the branch-point-binding protein (BBP) interacts with the branch point sequence. The U2 auxiliary factors, U2AF35 and U2AF65, bind to the 3' splice site and polypyrimidine tract. Once these components are in place, the spliceosome assembles, and U1 and U4 snRNPs are released. The spliceosome then catalyzes the first transesterification reaction, forming a lariat structure by connecting the 5' end of the intron to the branch point adenosine. The second transesterification reaction results in the ligation of the exons and the removal of the intron in the form of a lariat.

18. What is alternative splicing, and why is it important for genetic diversity?

Alternative splicing is a process by which different combinations of exons from the same pre-mRNA are joined together to produce multiple mRNA isoforms. This process is a major source of genetic diversity in eukaryotes, allowing a single gene to encode multiple protein variants with potentially different functions. For example, the Drosophila DSCAM gene can be spliced into over 38,000 different mRNA variants. This diversity plays a crucial role in cellular functions and adaptation, particularly in complex organisms like humans, where alternative splicing increases the functional potential of a limited number of genes.

19. What are the key components of the spliceosome and their roles in RNA splicing?

The spliceosome is composed of five small nuclear RNAs (snRNAs) — U1, U2, U4, U5, and U6 — which form complexes known as snRNPs (small nuclear ribonucleoproteins). These snRNAs and their associated proteins recognize and interact with the splice sites and other key sequences in the pre-mRNA. U1 snRNA binds to the 5' splice site, U2 snRNA binds to the branch point sequence, and U4/U6 snRNAs form a complex that is involved in catalyzing the splicing reactions. The U2AF complex helps in recognizing the 3' splice site, while SR proteins and other auxiliary factors facilitate spliceosome assembly and function.

20. How does the minor spliceosome differ from the major spliceosome?

The minor spliceosome is responsible for splicing a rare class of pre-mRNA introns that differ from the typical introns spliced by the major spliceosome. While the major spliceosome is composed of U1, U2, U4, U5, and U6 snRNAs, the minor spliceosome includes U11, U12, U4atac, and U6atac snRNAs, along with U5. The minor spliceosome splices U12-type introns, which are found in certain genes in metazoans. Unlike the major spliceosome, the minor spliceosome is less abundant and is generally active in specific cells or conditions, such as anucleate platelets or specialized neurons.

21. **What is RNA editing, and how does it differ from other RNA processing events like splicing or capping?**

RNA editing is a molecular process by which cells make discrete changes to specific nucleotide sequences in an RNA molecule after it has been synthesized by RNA polymerase. This differs from other RNA processing events such as splicing,

5’-capping, and 3’-polyadenylation, which are typically involved in the modification of RNA without altering its nucleotide sequence. RNA editing can involve insertion, deletion, or base substitution within the RNA molecule, which can ultimately result in a change to the encoded protein, unlike splicing which removes introns and joins exons.

22. **What are the different types of RNA editing mechanisms, and how do they function?**

RNA editing mechanisms are diverse and include processes like insertion or deletion of nucleotides, and base modifications such as C-to-U and A-to-I editing. In insertion/deletion editing, uracil nucleotides are added or deleted in a gene sequence, affecting the protein's sequence and function. C-to-U editing involves the deamination of cytidine to uridine, while A-to-I editing involves the conversion of adenosine to inosine. These changes can alter the protein's amino acid sequence or modify RNA function, such as miRNA target recognition.

23. **How does pan-editing work in the mitochondria of organisms like *Trypanosoma brucei*?**

Pan-editing involves widespread changes to RNA sequences, typically through the addition of uracil nucleotides in mitochondria of organisms such as *Trypanosoma brucei*. This process begins with the alignment of an unedited RNA transcript with a guide RNA (gRNA). The gRNA has complementary sequences to regions needing editing, and the editosome, a protein complex, catalyzes the insertion of uracils. This results in frameshift mutations, creating proteins with sequences differing from those predicted by the genomic DNA. This extensive RNA editing plays a crucial role in maintaining proper mitochondrial function.

24. **Describe the C-to-U editing process in RNA and its significance using the example of the human ApoB gene.**

C-to-U editing involves the deamination of cytidine to uridine within RNA molecules. In humans, one key example is the apolipoprotein B (ApoB) gene. In the liver, the full-length ApoB100 protein is produced, while in the intestines, the mRNA undergoes C-to-U editing that converts a CAA codon into a UAA stop codon, leading to the production of the shorter ApoB48 protein. This selective editing is crucial for tissue-specific protein synthesis, highlighting the regulatory power of RNA editing in gene expression.

25. **What is A-to-I RNA editing, and what role does it play in mammals?**

A-to-I RNA editing is a common form of RNA editing in mammals, where adenosine is converted to inosine by the enzyme ADAR (adenosine deaminases acting on RNA). Inosine behaves similarly to guanosine in both translation and secondary structure formation, thereby altering the protein product encoded by the edited mRNA. This editing can affect coding capacity, alter miRNA or siRNA target populations, and impact splicing and other RNA-related processes. It plays an essential role in regulating gene expression, immune responses, and various other cellular activities.

26. **How does RNA editing in plant mitochondria and plastids differ from RNA editing in other organisms?**

RNA editing in plant mitochondria and plastids typically involves the conversion of cytidine (C) to uridine (U), with very rare instances of U-to-C editing. This process is essential for correcting RNA sequences to maintain functional tRNAs and ensure proper translation and respiration within these organelles. The editing mechanisms in plants are carried out by proteins from the PPR (pentatricopeptide repeat) and MORF (multiple organellar RNA editing factor) families, which help guide the editing process. This system is critical for maintaining organellar functions and producing proteins vital for respiration and other metabolic pathways.

27. **What are the roles of the PPR and MORF proteins in RNA editing in plants?**

PPR and MORF proteins are key components of the RNA editing machinery in plants. PPR proteins act as trans-acting factors that recognize specific RNA sequences and guide the editing process, while MORF proteins are involved in the assembly of the RNA editing complex known as the editosome. These proteins ensure that the RNA editing process in mitochondria and plastids is accurate and effective, facilitating proper translation, tRNA functionality, and protein synthesis crucial for organellar functions.

28. **What is the significance of A-to-I editing in altering miRNA or siRNA target populations?**

A-to-I editing has a significant role in altering the target specificity of miRNAs and siRNAs. In this process, the conversion of adenosine to inosine in the RNA molecule can change the base-pairing properties of the RNA, which can affect the binding of miRNAs or siRNAs. As a result, A-to-I editing can lead to changes in gene silencing or regulation, impacting processes like gene expression, cell

differentiation, and immune responses. This type of editing provides a layer of regulation that can adapt to environmental or physiological changes.

29. **How does RNA editing influence protein diversity in organisms?**

RNA editing contributes to protein diversity by allowing cells to produce multiple protein isoforms from a single gene. By altering the nucleotide sequence of mRNA after transcription, RNA editing can change the amino acid sequence of the encoded protein, leading to functional differences. This mechanism is particularly important in processes like tissue-specific expression and adaptive responses to environmental changes. In some cases, RNA editing can even generate proteins with distinct functional properties, increasing the organism's ability to respond to various cellular needs.

30. **What potential therapeutic applications could arise from understanding RNA editing processes?**

Understanding RNA editing holds significant therapeutic potential, particularly in the context of genetic diseases, cancer, and viral infections. By manipulating RNA editing, it may be possible to correct genetic mutations at the RNA level, effectively bypassing the need for DNA-based treatments. For example, in diseases caused by point mutations, RNA editing could potentially restore normal protein function by editing the RNA transcript. Additionally, understanding RNA editing in viruses could lead to the development of antiviral strategies that target the RNA editing machinery to inhibit viral replication or modify viral gene expression.

31. What are the key features of small regulatory RNAs (sRNAs) in bacteria, and how have their discovery and roles evolved over time?

Small RNAs (sRNAs) in bacteria have been recognized for their role as regulators of gene expression, a function initially identified in the early 1980s. These RNA molecules, typically under 200 nucleotides, function mainly in the regulation of transcription, translation, and mRNA stability. The discovery of these regulatory RNAs began with the identification of the RNA I in *Escherichia coli* in 1981, which was involved in the regulation of plasmid replication, followed by the discovery of the MicF RNA in 1984, which modulates translation by inhibiting the expression of the outer membrane porin OmpF. However, it was not until the early 2000s that the prevalence and biological importance of sRNAs were truly appreciated. Advances in computational techniques, deep sequencing, and microarray analysis facilitated the identification of hundreds of new sRNAs in

bacteria. The current understanding of sRNAs has expanded significantly, and they are now recognized as essential components of bacterial gene regulatory networks.

32. What are riboswitches, and how do they regulate gene expression in bacteria?

Riboswitches are conserved RNA elements found typically in the 5' untranslated region (5' UTR) of bacterial mRNAs that act as sensors for small molecules. These RNA structures can change conformation in response to the binding of specific metabolites, which then modulate gene expression. The binding of the metabolite can lead to the formation of terminator or antiterminator structures, influencing transcription termination or elongation. Riboswitches can also impact translation by occluding or exposing the ribosome-binding site. There are many types of riboswitches that regulate the expression of genes involved in metabolite synthesis, transport, or usage. For example, the B. subtilis riboswitch can bind flavin mononucleotide (FMN) or thiamin pyrophosphate to regulate genes responsible for vitamin uptake. Riboswitches are classified based on their ability to bind specific metabolites, and they represent a direct and efficient means of regulating gene expression in response to environmental changes.

33. Explain the function of trans-encoded base pairing sRNAs and their regulatory mechanisms in bacterial cells.

Trans-encoded base pairing sRNAs are a class of regulatory RNAs that interact with their target mRNAs, typically in the 5' untranslated region (UTR). These sRNAs usually exhibit limited complementarity with their target mRNAs, and their regulation generally leads to translational repression or mRNA degradation. The sRNAs bind to their target mRNAs and, through base pairing, either block the ribosome-binding site or recruit ribonucleases, leading to mRNA degradation. One of the best-known examples is the GcvB sRNA, which regulates the expression of genes involved in glycine utilization. The interaction between trans-encoded sRNAs and their target mRNAs is often facilitated by an RNA chaperone protein called Hfq. This protein assists in the formation of RNA-RNA complexes by increasing the local concentration of sRNAs and mRNAs, and it also helps protect sRNAs from degradation. In this way, Hfq plays a crucial role in maintaining the stability of sRNAs and ensuring the efficiency of regulatory interactions.

34. How do cis-encoded base pairing sRNAs function in bacteria, and what are their roles in gene regulation?

Cis-encoded base pairing sRNAs are transcribed from the opposite DNA strand of their target mRNAs and typically exhibit a high degree of complementarity with their target RNA. These sRNAs are often involved in regulating gene expression within the same genomic region. In bacteria, cis-encoded antisense sRNAs can repress translation, promote mRNA degradation, or prevent transcription termination. For example, in *Escherichia coli*, the sRNA IstR acts as an antitoxin by base pairing with the mRNA encoding a toxic protein, thus preventing the translation of the toxin. These sRNAs are typically expressed constitutively, and their expression can be influenced by environmental stress or other signals. They are essential in maintaining the balance between gene expression and cellular function, especially under stressful conditions, where they may help regulate potentially harmful genes or proteins. The roles of cis-encoded sRNAs are still being explored, and they represent an important class of regulatory elements in bacterial genomes.

35. What is the relationship between small RNAs and protein-binding in bacteria, and how do they modulate protein activity?

In bacteria, some small RNAs (sRNAs) regulate protein activity by binding to specific proteins. These protein-binding sRNAs can either enhance or inhibit the activity of their target proteins. For instance, the CsrB and CsrC sRNAs in *Escherichia coli* regulate the CsrA protein, which controls carbon metabolism and motility. The CsrB and CsrC sRNAs contain multiple binding sites for CsrA and sequester the protein, preventing it from binding to its target mRNAs and thus modulating gene expression. Similarly, the 6S RNA in *E. coli* binds to the RNA polymerase holoenzyme containing the sigma factor σ^70, inhibiting transcription from σ^70-dependent promoters during the stationary phase. Another example is the GlmY RNA, which modulates the activity of the GlmZ sRNA involved in the regulation of glucosamine-6-phosphate synthesis. These protein-binding sRNAs provide a dynamic mechanism for regulating protein activity and are integral components of bacterial cellular processes.

****
****
****

ZB

Advice

MA apsy bohat zada payar krta ho ap janta b nahe phr b just for sake of Allah

Meri ap sbky leay duay ha Allah ap sbko hamesha kamyab kryn khush rakhyn Zindagi k har moor pe kamyab kryn ameeeen mri jaaaan k tooto

Study zada handouts se kea kryn zada depend handouts pe kea kryn

Or Vu k groups ko avoid kea kryn yaha bohat zada choty moty keery makory hn jo students ki help krny ke bahany unki life tk tbah kr dety hn ap bs apni Handouts se parha kro bs

Dunya m koi ksi ke Sath sincere nahe har insan matalbi ha agr koi apky Sath acha b ha to tab tk jab tk usy apsy koi mtlb Hoga so payar bs khud se kro ksi or se kbi b nahe or na ksi se umeed rakho

Ye dunya takleef k elawa ksi ko Kuch nahe de saqti

Be brave be strong my family 

Be kind with Everyone for no reason because kindness charges nothing.

Hmaesha relax ho ke parha kryn panic mat hua kryn apky parents ky bohat khawab hoty hn wo pura krny ki puri koshish kea kryn kbi b ksi pe depend na kea kryn apko bs ak insan success kr saqta ha or wo insan ap khud hn apky elawa koi nahe ap bs koshish kryn Allah pak pe strong yakeen rakha kryn or mehnat krke sb Allah ke hawaly kr dea kryn everything is possible be brave be strong stay blessed

Hum insan hamesha moat se darty or bhagty hn or moat se bachny ki koshish krty hn jabke moat ny ana hi ana ha hum moat se nahe bach saqty humy jahanam se bachny ki koshish krni chaheay hum jahanam se bach saqty hn

Insan ko 3 chezo se dar lagta ha Moat Risk/Dolat Ezat Shohrat Fame

Moat

Humy pta hona chaheay Moat tab ani ha jab Allah chahy us se pehly puri dunya bi ak tararf ho ke apko marna chahy to apka kuch nahe beggar saqti

Risk/Dolat

Risk Dolat sb Allah pak ke hath m ha puri dunya ak taraf b ho jay na apsy apka risk cheen saqti ha na de saqti ha Ye sirf ALLAh pak ky hath m ha

Izat Zilat Shohrat Fame

Izat Zilat Sirf Allah de saqty hn puri dunya mel ke b apko zra brabr b demage nahe kr saqti Agr ALLAH Apko izat dena chahay to puri dunya ak tararf ho ke b apko 1% b nuksan nahe pohancha saqti

Phr Dar ks bat ka ??????????????

Be brave be strong Just put your trust To ALLAH Hamesha Confident or himat se raho kbi ksi k samne mat jhuko puri dunya mel k b apka kuch b nahe beggar saqti

VU Examination Software Demo:

This Online Demo allows students to get themselves familiar with VU Examination Software and it's available features. This will help students to attempt their paper conveniently under real time environment. Click this link:

https://qb.vu.edu.pk/Vutes_demo/default.aspx Virtual University Official website:

Click this link:

https://www.vu.edu.pk/

Software / Tools:

Is drive m vu k related sb k sb software add hain jin ko kio bhi software ki zaroorat hai tu wo is link ko click kr k download kr sakty hain.

Mega drive:

https://mega.nz/folder/tA5x1LYa#9QpZ11pGc9jERh16nJduKg Note:

Is file m kisi bhi kisam ka kio bhi virus nahi hai sb safe hai aur sb software/tools sirf PC/laptop k liye hain.

Join our WhatsApp channel for more

https://whatsapp.com/channel/0029VaODheTDp2Q8YVcoON2p

second WhatsApp channel

https://whatsapp.com/channel/0029VafvOcy90x2tPbUaX02I

ZB

MY REQUEST FROM ALL OF YOU

ZB Request from All of you My family

Mjy ap sbki help or support ki zarort ha or wo ye k hum sb Mel kr Allah ka Quran ki Urdu translation logo tak pohanchy or Quran ko samjna asaaan kryn dosro k leay sb tak Allah ka Quran pohanchy it's my compaign hum log 70+ age k ho jaty hn phr b hmy namaz tak ki translation nahe pta hoti k hum Allah pak se Kya Dua kr rhy hn so hum youngster's ko Mel kr puri takat or energy ke sath Allah ka Quran spread krna ha

Hum puri Koshish kr rhy hn k hum sb Mel k Quran ki translation or most important topics ko maximum share kry with translation hmara maksad Quran ko spread krna ha be a part of us

m apni pocket se ye sb kr rha ho Allah ka Quran spread krny k leay apne or mene sbny Marna ha ak din to q na Allah k leay Kuch kryn apni energies apni power ko bajay negative use krny k Allah k leay invest kryn khud ko be a part of us

Agr m 23 ki age m apny sare sources use kr rha ho pocket se heavy amounts give away kr rha ho Quran spread krny k leay to ap just share to kr saqty hn itna e krdyn

Contact ZB +92 300 5427579

Agr ap mujsy contact na kr sky too b ap jo b hn jaha bi hn waha Allah ka quran pheelay zada se zada with urdu translation jitni himat ha utna share kryn

ZB +92 300 5427579

WhatsApp:

https://whatsapp.com/channel/0029VaODheTDp2Q8YVcoON2p

Second WhatsApp channel

channel on WhatsApp:

https://whatsapp.com/channel/0029VafvOcy90x2tPbUaX02I

YouTube:

https://youtu.be/-AMj4iNz0UE?si=hg29pJZ1kujzg57J