MTH202 Mcq's Final Term By Vu Topper RM

Discreate mathematics (Virtual University of Pakistan)



Scan to open on Studocu

Studocu is not sponsored or endorsed by any college or university
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Mth-202 Important Mid Term Mcq's Solution 100% Correct : Solve By Vu-Topper RM!!



This document is available on studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:1 (Marks:1) Vu-Topper RM

Range of function /[f(x)=(e^x)/] is _____

➢**Set of positive real numbers**

Question No:2 (Marks:1) Vu-Topper RM

Composite relation symbolically written as _____

➢**SoR={(a,c)aeA, ceC, 3eB, (a,b)eR and (b,c)eS}**

Question No:3 (Marks:1) Vu-Topper RM

If x=17(mod 5) which of the following integers are valid solution for x ?

➢**12**

Question No:4 (Marks:1) Vu-Topper RM

Range of the relation {(0,1),(3,22),(90,34)}

➢**{1,22,34}**

Question No:5 (Marks:1) Vu-Topper RM

Let A= {0,1,2} and R = {(0,2),(1,1),(2,0)} be a relation on A. The which of the following ordered pairs are needed to make it transitive?

➢ **(0,0) and (2,2)**

Question No:6 (Marks:1) Vu-Topper RM

Operation of subtraction is a binary operation on the set of _____

➢**Integers**

Question No:7 (Marks:1) Vu-Topper RM

Let S=R and define the 'square' relation R= $\{(x,y)Ix^2=y^2\}$. The square relation is an _____ relation

➢**Equivalence relation**

Question No:8 (Marks:1) Vu-Topper RM

The logic gate NOT is a uniary operation on {0,1}.

➢**True**

Question No:9 (Marks:1) Vu-Topper RM

Let A= {1,2}, then P(A)=_____

➢**{{},{1},{2},{1,2}}**

Question No:10 (Marks:1) Vu-Topper RM

If a relation R is reflexive, anti symmetric and transitive then which of the following is not true for the inverse relation.

➢**Inverse relation will be irreflexive.**

Question No:11 (Marks:1) Vu-Topper RM

Let R be a binary relation on a set A,R is anti-symmetric iff _____

➢**a,beA if (a,b)eR and (b,a)eR then a=b**

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:12 (Marks:1) Vu-Topper RM

"-" is a binary operation on the set of integers Z.

➢**True**

Question No:13 (Marks:1) Vu-Topper RM

The inverse relation R^(-1) from B to A is defined as _____

➢**R^(-1) = {(b,a) e B*A I (a,b)eR}**

Question No:14 (Marks:1) Vu-Topper RM

Which of the following is always true for the matrix representation of a symmetric relation?

➢**Matrix is equal to its transpose**

Question No:15 (Marks:1) Vu-Topper RM

Let A= {1,2,3,4} and let R and S be transitive binary relations on A defined as; R= {(1,2), (1,3), (2,2), (3,3), (4,2), (4,3) and S={(2,1), (2,4), (3,3)} then RuS= {(1,2), (1,3), (2,1), (2,2), (2,4), (3,3), (4,2), (4,3)}

➢**R union S is transitive**

Question No:16 (Marks:1) Vu-Topper RM

Let S=R and define the ' square' relation R = $\{(x,y) I x^2=y^2\}$. The square is an _____ relation.

➢**Equivalance relation**

Question No:17 (Marks:1) Vu-Topper RM

If x=-10(mod 15). Which of the following integers are valid solution for x?

➢**5**

Question No:18 (Marks:1) Vu-Topper RM

Let R be a binary relation on a set A. If R is anti symmetric then _____

➢**Inverse of R is anti symmetric**

Question No:19 (Marks:1) Vu-Topper RM

If A={1,2,3} is a set and R = {(1,2),(2,2),(2,1)} is a relation on A, R is

➢**Symmetric**

Question No:20 (Marks:1) Vu-Topper RM

Let A={0,1} and B=(1). Let R and S be two binary relations on Cartesian product of A and B such that R={(0,1)} and S= {(1,1)}. Then R intersection S=_____

➢**Empty**

Question No:21 (Marks:1) Vu-Topper RM

A relation R is said to be _____ iff it is reflexive, antisymmetric and transitive.

➢**Partial order Relation**

Question No:22 (Marks:1) Vu-Topper RM

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Let X={1,2,3} and Y={7,8,9} and let f be function defined from X to Y such that f is onto then which of the following statement about f is true?

➢**Co-domain of f must contain 1 element**

Question No:23 (Marks:1) Vu-Topper RM

The function fog and gof are always equal

➢**False**

Question No:24 (Marks:1) Vu-Topper RM

If a relation R={(1,2),(2,3),(3,4)(4,1)(2,2) is given then which of the following is true about this relation.

➢**R is reflexive**

Question No:25 (Marks:1) Vu-Topper RM

A set is called countable if , and only if, it is_____

➢**finite**

Question No:26 (Marks:1) Vu-Topper RM

Let $f(x) = x^2-1$ define function f from R to R and c=2 be any scalar, then c,f(x) is _____

➢**$2x^2+2$**

Question No:27 (Marks:1) Vu-Topper RM

The set Z of all integers is _____

➢**Countable**

Question No:28 (Marks:1) Vu-Topper RM

Let R be a binary relation on a set A. If R is anti symmetric then _____

➢**Inverse of R is symmetric**

Question No:29 (Marks:1) Vu-Topper RM

For (2x-3, 4y+2) = (1,10). What will be the value of x and y ?

➢ **(2,2)**

Question No:30 (Marks:1) Vu-Topper RM

Let f and g be the two functions from R to R defined by f(x) = IxI and g(x)= square root of x^2 for all xeR. Then_____

➢**F(x) is not equal to g(x)**

Question No:31 (Marks:1) Vu-Topper RM

If a set A has 15 elements then P(A) (power set of A) has _____ elements.

➢**2^15**

Question No:32 (Marks:1) Vu-Topper RM

For the relation below to be a function, x cannot be what values {(12,14),(13,5),(-2,7),(x,13)}?

➢**X cannot be 12, 13, or -2**

Question No:33 (Marks:1) Vu-Topper RM

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Let the set A = {1,2,3,4}. Then the relation {(2,4),(4,2)} is _____

➢**Symmetric**

Question No:34 (Marks:1) Vu-Topper RM

For the following relation to be a function, x can not be what values? R={(2,4),(x,1), (4,2),(5,6)}.

➢**x cannot be 2,4 and 5**

Question No:35 (Marks:1) Vu-Topper RM

Vertical line test is used to determine that whether the graph of a relation is a function or not.

➢**True**

Question No:36 (Marks:1) Vu-Topper RM

The properties of being symmetric and being anti symmetric are _____

➢**Not negative of each other**

Question No:37 (Marks:1) Vu-Topper RM

The number of elements in AxB are _____ if A is a set with '5' elements and B is a set with '4' elements.

➢**20**

Question No:38 (Marks:1) Vu-Topper RM

R={(a,1)(b,2)(c,3)(d,4)} then the inverse of this relation is _____

➢**{(1,a)(2,b)(3,c)(4,d)}**

Question No:39 (Marks:1) Vu-Topper RM

Logic gate NOT does not define a binary operation on (0,1) because _____

➢**It takes a single input and gives a single output**

Question No:40 (Marks:1) Vu-Topper RM

How many real numbers exist between 1 and 5

➢**3**

Question No:41 (Marks:1) Vu-Topper RM

The number pi is

➢**Irrational**

Question No:42 (Marks:1) Vu-Topper RM

The number square root 2 is

➢**Irrational**

Question No:43 (Marks:1) Vu-Topper RM

Range the relation {(0,1)(3,22),(90,34)}

➢**{0,3,90}**

Question No:44 (Marks:1) Vu-Topper RM

Supherical coordinate 0 is related to the cylindrical coordinate _____

Free document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

➢@=@

Question No:45 (Marks:1) Vu-Topper RM

Operation of subtraction is a binary operation on the set_____

➢**Integers**

Question No:46 (Marks:1) Vu-Topper RM

Let A {1,2,3,4} and R={(1,2),(2,3),(3,3),(3,4)} be a relation on A. Then which one of the following ordered pair has made R not an irreflexive relation?

➢ **(3,3)**

Question No:47 (Marks:1) Vu-Topper RM

Input values of the function are called the _____

➢**Domain**

Question No:48 (Marks:1) Vu-Topper RM

Range of function f(x)=IxI will be

➢**Set of positive real numbers**

Question No:49 (Marks:1) Vu-Topper RM

Which of the following is not a binary operation on the set of integers?

➢**Division**

Question No:50 (Marks:1) Vu-Topper RM

In the matrix representation of an irreflexive relation all the entries in the main diagonal are _____

➢**0**

Question No:51 (Marks:1) Vu-Topper RM

If the partition set of A is $\{A_1, A_2\}$ then

➢**$A_1 n A_2$= not empty set**

Question No:52 (Marks:1) Vu-Topper RM

Let A= {a,b} then P(A) =

➢**{Non empty set, {a},{b},{a,b}}**

Question No:53 (Marks:1) Vu-Topper RM

Which relations below are not functions?

➢**{(13,14),(13,5),(16,7),(18,13)}**

Question No:54 (Marks:1) Vu-Topper RM

In the directed graph of an antisymmetric relation there is _____ pair of arrows between two distinct elements of the set.

➢**No**

Question No:55 (Marks:1) Vu-Topper RM

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

If a relation R= (1,1),(2,1)(2,2) is given then which of the following is not true about this relation

➢**R is irreflexive**

Question No:56 (Marks:1) Vu-Topper RM

Let R and S be transitive relations on a set A then _____

➢**Neither R union S is transitive nor R intersection S is transitive**

Question No:57 (Marks:1) Vu-Topper RM

Let R={(1,2)(3,4)(5,6)(7,8)}. Domain of the inverse of the relation is _____

➢**{2,4,6,8}**

Question No:58 (Marks:1) Vu-Topper RM

Let A={1,2,3,4,5} and B={4,9,,16,17,25}. Then the relation R={(2,4),(3,9),(4,16),93,17)} The inverse of R is'

➢**{(4,2),(9,3),(16,4),(17,3)}**

Question No:59 (Marks:1) Vu-Topper RM

Let R be a relation on a set A. If R is symmetric then its compliment is

_____

➢**Irreflexive**

Question No:60 (Marks:1) Vu-Topper RM

Which is not a binary operation on the set of natural numbers N?

➢**Subtraction**

Question No:61 (Marks:1) Vu-Topper RM

If a relation R={(1,1)(2,1)(1,2)(2,2)} is given then which of the following is not true about this relation.

➢**R is irreflexive**

Question No:62 (Marks:1) Vu-Topper RM

R={(a,1)(b,2)(c,3)(d,4)} then the inverse of this relation is _____

➢**{(1,a)(2,b)(3,c)(4,d)}**

Question No:63 (Marks:1) Vu-Topper RM

For any set A, the Cartesian product of A and A is known as _____

➢**Universal relation**

Question No:64 (Marks:1) Vu-Topper RM

Let A={p,q,r,s} and define a relation R on A by R={(p,p),(p,r),(q,r),(q,s),(r,s)} Then which one of the following is the correct statement about R:

➢**R is not reflexive**

Question No:65 (Marks:1) Vu-Topper RM

A={1,2} B={3,4}, R={(1,3)(2,4)}. Then the complement of R is _____.

➢**{(1,4)(2,3)}**

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:66 (Marks:1) Vu-Topper RM

Domain of a relation symbolically written as_____.

➢Dom(R)={aeAI(a,b)eR}

Question No:67 (Marks:1) Vu-Topper RM

Let X={2,4,5} and Y={1,2,4} and R be a relation from X to Y defined by R={(2,4)(4,1)(a,2)}. For what value of 'a' the relation R is a function?

➢5

Question No:68 (Marks:1) Vu-Topper RM

Let A={1,2,3,4,5,6,7,8,9}, then which of the following sets represent the partition of the set A?

➢A={1,3,5,7,9}, B={2,4,6}, C={8}

Question No:69 (Marks:1) Vu-Topper RM

Let A={1,2,3} and B={2,4} then number of binary relations from A to B are _____.

➢64

Question No:70 (Marks:1) Vu-Topper RM

A relation R is said to be _____iff it is reflexive, antisymmetric and transitivde.

➢Partial order Relation

Question No:71 (Marks:1) Vu-Topper RM

Let f be a function from X={2,4,5} to Y={1,2,4,6} defined as:f={(2,6),(4,2),(5,1)} . The range of f is _____

➢{1,2,6}

Question No:72 (Marks:1) Vu-Topper RM

Let A={0,1,2} and R={(0,2),(1,1),(2,0)} be a relation on A. Then which of the following statement about R is true?

➢R is symmetric

Question No:73 (Marks:1) Vu-Topper RM

Let A={2,3,4} and B={2,6,8} and let R be the "divides" relation from A to B i.e for all (a,b) belong to (Cartesian product of A and B), a,R b iff a I b (a divides b). Then

➢R={(2,2),(2,6),(2,8),(3,6),(4,8)}

Question No:74 (Marks:1) Vu-Topper RM

Let A={1,2,3,…,50} and B={2,4,6,8,10}. Then the Cartesian product of A and B has _____ elements.

➢250

Question No:75 (Marks:1) Vu-Topper RM

In the matrix representation of an reflexive relation all the entries in the main diagonal are _____

➢1

Question No:76 (Marks:1) Vu-Topper RM

Which of the following is not a type of a relation?

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

➢**Permutation**

Question No:77 (Marks:1) Vu-Topper RM

Let X={2,4,5} and Y={1,2,4} and R be a relation from X to Y defined by R={(2,4),(4,1),(a,2)}. For what value of 'a' the relation R is a function ?

➢5

Question No:78 (Marks:1) Vu-Topper RM

Which of the following is not a representation of a relation?

➢**Venn diagram**

Question No:79 (Marks:1) Vu-Topper RM

Let A={1,2,3,4} and define the following relations on A. Then R={(1,3),(1,4),(2,3),(2,4),(3,1),(3,4)} is _____

➢**R is irreflexive**

Question No:80 (Marks:1) Vu-Topper RM

The range of f:X-> Y is also called the image of

➢**True**

Question No:81 (Marks:1) Vu-Topper RM

Which of the following logical connective is not a binary operation ?

➢**Implication**

Question No:82 (Marks:1) Vu-Topper RM

A set may be dividend up into its disjoint subsets, such division is called_____

➢**Partition**

Question No:83 (Marks:1) Vu-Topper RM

If A=(1,2,3)&B=(4,5,6) and R ={(1,4)(2,5)(3,6)(3,4)} The complementary relation is _____

➢**A*B(difference or –) R**

Question No:84 (Marks:1) Vu-Topper RM

In matrix representation of a_____ relation, the diagonal entries are always 1.

➢**Reflexive**

Question No:85 (Marks:1) Vu-Topper RM

R is not symmetric iff there are elements a and b in A such that _____

➢ **(a,b) belongs to R but (b,a) does not belong to R**

Question No:86 (Marks:1) Vu-Topper RM

Which relations below are functions?
R1={(3,4),(4,5),(6,7),(8,9)}
R2={(3,4),(4,5),(6,7),(3,9)}
R3={(-3,4),(4,-5),(0,0),(8,9)}
R4={(8,11),(34,5),(6,17),(8,19)}

➢**R1 and R3 are functions**

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:87 (Marks:1) Vu-Topper RM

The logic gate OR and AND are uniary operation on {0,1}

➢**False**

Question No:88 (Marks:1) Vu-Topper RM

There is atleast one loop in the graph of an irreflexive relation

➢**False**

Question No:89 (Marks:1) Vu-Topper RM

There is atleast one loop in the graph of an reflexive relation

➢**True**

Question No:90 (Marks:1) Vu-Topper RM

A contains 3 elements and B contains 2 elements, then number of subsets of A*B are _____

➢**64**

Question No:91 (Marks:1) Vu-Topper RM

Let A={1,2,3} and B={0,1,2} and C={a,b} R={(1,0),(1,2),(3,1),(3,2)} S={(0,b),(1,a),(2,b)} composite of R and S=_____

➢{(1,b),(1,a),(3,a),(3,b)}

Question No:92 (Marks:1) Vu-Topper RM

If R is transitive then the inverse relation will be transitive

True

Question No:93 (Marks:1) Vu-Topper RM

The number of elements in the power set of P(not empty set) denoted by P(P(not empty set) is

➢**2**

Question No:94 (Marks:1) Vu-Topper RM

The function defined from Z to Z as f(x)= 1/(x+2)(x-2) is not well defined because _____

➢**Function is not defined at x=2 and x=-2**

Question No:95 (Marks:1) Vu-Topper RM

Rang of relation {(0,1),(3,22),(90,34)} is _____

➢**{1, 22, 34}**

Question No:96 (Marks:1) Vu-Topper RM

The number of elements in the power set of P(not empty set) denoted by P(not empty set) is

➢**1**

Question No:97 (Marks:1) Vu-Topper RM

Let R={(1,2),(3,4),(5,6),(7,8)}. Domain of inverse of the relation is _____

➢**{2,4,6,8}**

Question No:98 (Marks:1) Vu-Topper RM

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

The relation ‘divides’ on the set of integers is _____

➢**A symmetric relation**

Question No:99 (Marks:1) Vu-Topper RM

Operation of subtraction is a binary operation on the set of_____

➢**Integers**

Question No:100 (Marks:1) Vu-Topper RM

If R is transitive then the inverse relation will be transitive.

➢**True**

Question No:101 (Marks:1) Vu-Topper RM

Let A={1,2,3,4} and define the relation R on A by R={(1,2),(2,3),(3,3),(3,4)}.
Then_____

➢**R is both reflexive and irreflexive**

Question No:102 (Marks:1) Vu-Topper RM

A set may be divided up into its disjoints subsets,such division is called

➢**Partition**

Question No:103 (Marks:1) Vu-Topper RM

If a set A contains is elements then the number of elements in its power set P(A) is _____

➢**2^n**

Question No:104 (Marks:1) Vu-Topper RM

Range of a relation symbolically written as _____.

➢**Ran R = {b ∈ B I (a, b) ∈ R }**

Question No:105 (Marks:1) Vu-Topper RM

Let R be a binary relation on a set A. IF R is anti symmetric then _____.

➢**Inverse of R is anti symmetric**

Question No:106 (Marks:1) Vu-Topper RM

Let A be a set with m elements and B be a set with n elements then the number of elements in A*B are _____

➢**m.n**

Question No:107 (Marks:1) Vu-Topper RM

Let A{1,2,3,4} and define the following relation on A. Then R={(1,3),(2,2),(2,4),(3,1),(4,2)} Is _____

➢**R is symmetric**

Question No:108 (Marks:1) Vu-Topper RM

If A={1,2,3}& B={4,5,6} and R={1,4},{2,5},{3,6},{3,4}. The complementary relation is _____

➢**A*B (difference or -) R**

Question No:109 (Marks:1) Vu-Topper RM

This document is available on VU-Studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Let R be a relation on a set A. If R is reflexive then its compliment is _____

➢**Irreflexive**

Question No:110 (Marks:1) Vu-Topper RM

25=1(mod 3) means that 3 divides _____

➢**25-1**

Question No:111 (Marks:1) Vu-Topper RM

Let R be the universal relation on a set A then which one of the following statement about R is true ?

➢**R is reflexive, symmetric and transitive**

Question No:112 (Marks:1) Vu-Topper RM

Domain of a relation symbolically written as _____

➢**Dom(R)= {aeR I (a,b) e R}**

Question No:113 (Marks:1) Vu-Topper RM

Let R be a relation on a set A. R is transitive if and only if for all a,b,ceA then

➢ **(a,b)eR and (b,c)eR then (a,c)eR**

Question No:114 (Marks:1) Vu-Topper RM

Let A be a non-empty set and P(A) the power set of A Deifne the ‘ subset’ relation ,c as follows for all X,Y e P(A), XcY<-> for all x, iffxeX then xeY . Then c is _____

➢**c is partial order relation**

Question No:115 (Marks:1) Vu-Topper RM

Define a relation R={(1,1),(2,2),(3,3),(1,3) the relation is

➢**R is reflexive and transitive**

Question No:116 (Marks:1) Vu-Topper RM

Let R be a relation on a set A. R is transitive if and only if for all a,b,ceA then.

➢ **(a,b)eR and (b,c)eR then (a,c)eR**

Question No:117 (Marks:1) Vu-Topper RM

Let R and S be reflexive relations on a set A then R intersection S is reflexive

➢**True**

Question No:118 (Marks:1) Vu-Topper RM

A function whose range consists of only one element is called_____

➢**One to one function**

Question No:119 (Marks:1) Vu-Topper RM

The set Z of all integers is _____

➢**Countable**

Question No:120 (Marks:1) Vu-Topper RM

Let X={1,5,9} and Y={3,4,7}, Define a function f from X to Y such that f(1)=7,f(5)=3,f(9)=_____. Which is true f(9) to make it a one-to-one

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

(injective) function?
➢**4**

Question No:121 (Marks:1) Vu-Topper RM
What will be the fourth term of the following sequence 1/2,2/3,3/4 _____?
➢**4/5**

Question No:122 (Marks:1) Vu-Topper RM
The value of 6!=
➢**720**

Question No:123 (Marks:1) Vu-Topper RM
A constant function is one to one iffits _____ is a singleton.
➢**Domain**

Question No:124 (Marks:1) Vu-Topper RM
A constant function is onto iffits _____ is a singleton.
➢**Co-domain**

Question No:125 (Marks:1) Vu-Topper RM
Number of one to one functions from X={a,b} to Y={u,v} are equal to _____-
➢**2**

Question No:126 (Marks:1) Vu-Topper RM
If f is defined recursively by f(0) = -1 and f(n+1)=f(n)+3, then f(2)=_____.
➢**5**

Question No:127 (Marks:1) Vu-Topper RM
A function whose inverse function exists is called a/an_____
➢**Invertible function**

Question No:128 (Marks:1) Vu-Topper RM
Let f(2)=3, g(2)=3, f(4)=1 and g(4)=2 then the value of fog(4) is……
➢**3**

Question No:129 (Marks:1) Vu-Topper RM
Let A={1,2,3,4} and B={7} then the constant function from A to B is _____.
➢**Onto**

Question No:130 (Marks:1) Vu-Topper RM
Composition of a function is a commutative operation.
➢**False**

Question No:131 (Marks:1) Vu-Topper RM
Composition of a function is not a commutative operation
➢**True**

This document is available on studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:132 (Marks:1) Vu-Topper RM

The sum of first five whole number is _____.

➢**10**

Question No:133 (Marks:1) Vu-Topper RM

If f and g are two one-to-one functions, then their composition that is gof is one-to-one.

➢**True**

Question No:134 (Marks:1) Vu-Topper RM

Inverse of a surjective function is always a function.

➢**False**

Question No:135 (Marks:1) Vu-Topper RM

Inverse of a surjective function may not be a function

➢**True**

Question No:136 (Marks:1) Vu-Topper RM

Let X={1,2,3,4} and Y={7,8,9} and let f be function defined from X to Y such that f is onto then which of the following statement about f is true?

➢**Co-domain of f must contain 3 elements**

Question No:137 (Marks:1) Vu-Topper RM

If f; X->Y and g; Y->Z are both onto functions. Then gof; X->Z is _____

➢onto

Question No:138 (Marks:1) Vu-Topper RM

If f and g are two one-to-one functions, then their composition gof is_____

➢**One-to-one**

Question No:139 (Marks:1) Vu-Topper RM

If f: W →X, g:X →Y, and h:Y →Z are functions, then_____

➢ **(hog)of = ho(gof)**

Question No:140 (Marks:1) Vu-Topper RM

Cardinality of positive prime numbers less than 20 is _____.

➢**8**

Question No:141 (Marks:1) Vu-Topper RM

IF $f(x)=\sin^{-1}(x)$ and g(x)=sin x then gof(x) is _____.

➢X

Question No:142 (Marks:1) Vu-Topper RM

0!=_____.

➢**1**

Question No:143 (Marks:1) Vu-Topper RM

An important data type in computer programming consists of _____.

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

➢**Finite sequences**

Question No:144 (Marks:1) Vu-Topper RM

Let f(x)=2x and g(x)=x+2 define functions f and g from R to R, then (f-g)(x) is _____.

➢**x-2**

Question No:145 (Marks:1) Vu-Topper RM

The total number of terms in an arithmetic series 0+5+10+15+….50 are _____.

➢**11**

Question No:146 (Marks:1) Vu-Topper RM

9!/6!=_____

➢**504**

Question No:147 (Marks:1) Vu-Topper RM

Let f(x)=3x and g(x)=3x-2 define functions f and g from R to R, THEN (F+G)(X) is _____

➢**6x-2**

Question No:148 (Marks:1) Vu-Topper RM

If f is a bijective function then $(f^{-1}f(x))$ is equal to

➢**X**

Question No:149 (Marks:1) Vu-Topper RM

A sequence whose terms alternate in sign is called an_____

➢**Alternating sequence**

Question No:150 (Marks:1) Vu-Topper RM

Common ration in the sequence “4, 16, 64, 256,…” is…..12.

➢**4**

Question No:151 (Marks:1) Vu-Topper RM

0.8181818181 is a infinite geometric series.

➢**True**

Question No:152 (Marks:1) Vu-Topper RM

The word ‘algorithm’ refers to a step-by-step method for performing some action.

➢**True**

Question No:153 (Marks:1) Vu-Topper RM

A predicate become _____ when its variables are given specific values

➢**Sentence**

Question No:154 (Marks:1) Vu-Topper RM

The sum of two irrational numbers must be an irrational number.

➢**False**

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:155 (Marks:1) Vu-Topper RM

The sum of two irrational numbers need not be irrational number

➢**True**

Question No:156 (Marks:1) Vu-Topper RM

The division by zero is allowed in mathematics.

➢**Fasle**

Question No:157 (Marks:1) Vu-Topper RM

The product of any two consecutive positive integers is divisible by 2

➢**True**

Question No:158 (Marks:1) Vu-Topper RM

If 'n' is an odd integer then n^3+n is _____.

➢**Even**

Question No:159 (Marks:1) Vu-Topper RM

For integers a,b,c, If divides b and a divides c, then a divides (a+b).

➢**False**

Question No:160 (Marks:1) Vu-Topper RM

Quotient remainder theorem states that for any positive integer d, there exist unique integer q and r such that _____ and 0<r<d

➢**N=d.q+r**

Question No:161 (Marks:1) Vu-Topper RM

A rule that assigns a numerical value to each outcome in a simple space is called

➢**Random variable**

Question No:162 (Marks:1) Vu-Topper RM

If A and B are two disjoint (mutually exclusive) events then P(AB)=

➢**P(A) + P(B)**

Question No:163 (Marks:1) Vu-Topper RM

How many ways are there to select five players from a 10 member tennis team to make a trip to a match to another school?

➢**C(10,5)**

Question No:164 (Marks:1) Vu-Topper RM

The expectation of x is equal to

➢**Sum xf(x)**

Question No:165 (Marks:1) Vu-Topper RM

If P(A intersection B) = P(A) P(B) THEN THE events A and B are called

➢**Independent**

Question No:166 (Marks:1) Vu-Topper RM

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

A walk that starts and ends at the same vertex is called.
➢**None optins**

Question No:167 (Marks:1) Vu-Topper RM
How many integers from 1 through 1000 are neither multiple of 3 nor multiple of 5
➢**497**

Question No:168 (Marks:1) Vu-Topper RM
What is the probability of getting a number greater than 4 when a die is thrown?
➢**3/5**

Question No:169 (Marks:1) Vu-Topper RM
Eater formula for graphs is_____.
➢**F=e-v+2**

Question No:170 (Marks:1) Vu-Topper RM
X+a,x+3a,x+5a…..is an_____
➢**Arithmetic sequence**

Question No:171 (Marks:1) Vu-Topper RM
Composition of a function is a commutative operation.
➢**False**

Question No:172 (Marks:1) Vu-Topper RM
Real valued function is a function that assigns _____ to each member of its domain.
➢**Only a real number**

Question No:173 (Marks:1) Vu-Topper RM
Let X={1,2,3,4} and a function ‘f’ defined on X f(1)=1,f(2)=2,f(3)=3,f(4)=4 then _____
➢**F is an identity function**

Question No:174 (Marks:1) Vu-Topper RM
A constant function is surjective if and only if _____
➢**The co-domain consists of a single element**

Question No:175 (Marks:1) Vu-Topper RM
Cardinality means the total number of elements in a set.
➢**True**

Question No:176 (Marks:1) Vu-Topper RM
If f(x)=2x and g(x)=x then g(f(x)) is _____.
➢**$2x^2$**

Question No:177 (Marks:1) Vu-Topper RM
Let f:R->R is one to one function then c,f, c is not equal 0 is also one to one function.
➢**True**

This document is available on studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:178 (Marks:1) Vu-Topper RM

Let X={1,2,3,4} and a function 'f' defined from X to X by f(1)=1, f(2)=1, f(3)=1, f(4)=1 then which of the following is true?

➢**F is a constant function**

Question No:179 (Marks:1) Vu-Topper RM

If f and g are two one-to-one functions, then their composition that is gof is one-to-one.

➢**True**

Question No:180 (Marks:1) Vu-Topper RM

Which of the following is not correct for a 'sequence'?

➢**A sequence is a relation whose domain is the set of natural numbers**

Question No:181 (Marks:1) Vu-Topper RM

$F(x)=x^2$ is not one to one function from R to R^+

➢**True**

Question No:182 (Marks:1) Vu-Topper RM

Let f: R->R is one to one function then c,f c is not equal to is also one to one function.

➢**True**

Question No:183 (Marks:1) Vu-Topper RM

Let f(x)=x+2 then $f^{-1}(x)$ is_____

➢**x-2**

Question No:184 (Marks:1) Vu-Topper RM

Let $f(x)=x^2+1$ define functions f from R to R and c=2 be any scalar, then c,f(x) is _____.

➢**$2x^2-1$**

Question No:185 (Marks:1) Vu-Topper RM

A function F: R-> R defined by f(x) = square root x is a real valued function.

➢**False**

Question No:186 (Marks:1) Vu-Topper RM

If g:R->R defined by $g(x)=e^2$ is a real valued function of a real variable.

➢**True**

Question No:187 (Marks:1) Vu-Topper RM

A function F:R-> R defined by f(x) = log x is a real valued function

➢**True**

Question No:188 (Marks:1) Vu-Topper RM

½, then 3rd term of sequence is _____.

➢**1/2**

Question No:189 (Marks:1) Vu-Topper RM

The process of defining an object in terms of smaller versions of itself is called recursion.

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

➢**True**

Question No:190 (Marks:1) Vu-Topper RM

Which of the following is not correct for a 'sequence'?

➢**A sequence is a relation whose domain is the set of natural numbers**

Question No:191 (Marks:1) Vu-Topper RM

A set is called countable if, and only if, it is _____.

➢**Finite and countable infinite…..both**

Question No:192 (Marks:1) Vu-Topper RM

A set that is not countable is called _____.

➢**Uncountable**

Question No:193 (Marks:1) Vu-Topper RM

A sequence whose terms alternate in sign is called an _____.

➢**Alternating sequence**

Question No:194 (Marks:1) Vu-Topper RM

Let f: R->R is one to one function then c,f is also one to one function for _____.

➢**C is not equal 0**

Question No:195 (Marks:1) Vu-Topper RM

Let f(x) = x+3 then $f^{-1}(x)$ is _____-

➢**x-3**

Question No:196 (Marks:1) Vu-Topper RM

Let f and g be two functions defined by f(x) = x+2 and g(x)= 2x+1. Then the composition of f and g is_____.

➢**2x+3**

Question No:197 (Marks:1) Vu-Topper RM

Number of one to one functions form X={a,b} to Y={u,v} are equal to _____

➢**2**

Question No:198 (Marks:1) Vu-Topper RM

The flbonacci sequence is deined as $F_0=1, F_1=1, F_k=F_{k-1}+F_{k-2}$ for all integers $k \geq 2$ then which of the following is true for F_2

➢**$F_2-F_1= 2+1=3$**

Question No:199 (Marks:1) Vu-Topper RM

x+a, x+3a, x+5a,…… is a/an_____.

➢**Arithmetic sequence**

Question No:200 (Marks:1) Vu-Topper RM

Inverse of a function may not be a function.

➢**True**

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:201 (Marks:1) Vu-Topper RM

In the following sequence $a_k=K/(k+1)$, for k=1, a_1 will be _____.

➢**½**

Question No:202 (Marks:1) Vu-Topper RM

If f(x)=x and g(x)= -x are both one to one function then (f+g)(x) is also one to one function.

➢**False**

Question No:203 (Marks:1) Vu-Topper RM

If f(x)=x and g(x)= -x are both one to one function then (f+g)(x) is not one to one function.

➢**True**

Question No:204 (Marks:1) Vu-Topper RM

The function 'f' and 'g' are inverse of each other if and only if their composition gives_____.

➢**Identity function**

Question No:205 (Marks:1) Vu-Topper RM

Which of the following set is the domain of a sequence?

➢**Set of real numbers**

Question No:206 (Marks:1) Vu-Topper RM

Let C is defined as the set of all countries in the world then C is a _____.

➢**Finite set**

Question No:207 (Marks:1) Vu-Topper RM

A constant function is surjective if and only if_____.

➢**The co domain consists of a single element**

Question No:208 (Marks:1) Vu-Topper RM

Let X={1,2,3,4} and a function 'f' defined from X to X by f(1)=1, f(2)=1, f(3)=1, f(4)=1 then which of the following is true?

➢**F is a constant function**

Question No:209 (Marks:1) Vu-Topper RM

The composition of function is always

➢**Associative**

Question No:210 (Marks:1) Vu-Topper RM

A set is countably infinite if, and only if, it has the same cardinality as the set of

➢**Positive integers**

Question No:211 (Marks:1) Vu-Topper RM

Two functions 'f' and 'g' from 'X' to 'Y' are said to be equal if and only if _____.

➢**F(x)=g(x) for all 'x' belongs to X**

Question No:212 (Marks:1) Vu-Topper RM

$Y=x^3$ is a graph of bijective function from R to R.

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

➢**True**

Question No:213 (Marks:1) Vu-Topper RM

Domain and range are same for_____.

➢**Identity function**

Question No:214 (Marks:1) Vu-Topper RM

Let X={1,2,3,4} and a function 'f' defined on X by f(1)=1,f(2)=2,f(3)=3

➢**F is an identity function**

Question No:215 (Marks:1) Vu-Topper RM

Composition of a function is a commutative operation.

➢**True**

Question No:216 (Marks:1) Vu-Topper RM

Inverse of a surjective function is always a function.

➢**False**

Question No:217 (Marks:1) Vu-Topper RM

A function whose inverse function exists is called a/an_____.

➢**Invertible**

Question No:218 (Marks:1) Vu-Topper RM

Given a set X define a function I from X to X by i(x)=x from all x belonging to X . Then_____.

➢**I is both injective and surjective**

Question No:219 (Marks:1) Vu-Topper RM

Let f: R->R is a one to one function then c,f is also one to one function for.

➢**C is not equal 0**

Question No:220 (Marks:1) Vu-Topper RM

Let X={1,5,9} and Y={3,4,7}. Define a function f from X to Y such that f(1)=7, f(5)=3, f(9)=_____. Which is true for f(9) to make it a one-to-one (injective) function?

➢**4**

Question No:221 (Marks:1) Vu-Topper RM

Which of the following is not a predecessors of ak?

➢**A_k+1**

Question No:222 (Marks:1) Vu-Topper RM

Two functions 'f' and 'g' from X to Y are said to be equal if and only if_____.

➢**F(x)=g(x) for all 'x' belongs to x**

Question No:223 (Marks:1) Vu-Topper RM

The two functions 'f' and 'g' are equal if _____.

➢**$F(x) =3x$ and $g(x)= 6x^2+3x/2x^2+1$ for all xeR**

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:224 (Marks:1) Vu-Topper RM

If first term of a geometric sequence is 2 and common ratio is ½, then 3rd ter, of sequence.

➢¼

Question No:225 (Marks:1) Vu-Topper RM

Y= squre root x is an _____function form R^+ to R

➢**One to one function**

Question No:226 (Marks:1) Vu-Topper RM

Y= x^2 is an _____function form R to R^+

➢**NOT ONE TO ONE FUNTION**

Question No:227 (Marks:1) Vu-Topper RM

If a function (gof)(x) : X->Z is defined as (gof)(x)=g(f(x)) for all xeX, Then the function_____—.

➢ **(gof)**

Question No:228 (Marks:1) Vu-Topper RM

If 0 is the first term and -2 be the common difference of an arithmetic series, then the sum of first five terms of series is _____.

➢**-20**

Question No:229 (Marks:1) Vu-Topper RM

If $f(x)=\sin^{-1}(x)$ and g(x) = sin x then gof(x) is _____.

➢**X**

Question No:230 (Marks:1) Vu-Topper RM

F: X->Y that is both one to one and onto is called a _____.

➢**Bijective function**

Question No:231 (Marks:1) Vu-Topper RM

What does 'y' denotes in a geometric sequence?

➢**Common ratio**

Question No:232 (Marks:1) Vu-Topper RM

Let g be a function defined by g(x)=x+1. Then the composition of (gog).

➢**X+2**

Question No:233 (Marks:1) Vu-Topper RM

A graph of a function f is one to one iff every horizontal line intersects the graph in at most one point.

➢**True**

Question No:234 (Marks:1) Vu-Topper RM

Which of the following is true for the following sequence?

➢**If n is even, then Cn=2 and if n is odd, then Cn = 0**

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:232 (Marks:1) Vu-Topper RM

The function 'f' and 'g' are inverse of each other if and only if their composition gives_____.

➢**Identity function**

Question No:233 (Marks:1) Vu-Topper RM

N! is defined to be _____.

➢**The product of the integers from 1 to n**

Question No:234 (Marks:1) Vu-Topper RM

Let A = {1,2,3,4} and B={7} then the constant function from A to B

➢**Both one to one and onto**

Question No:235 (Marks:1) Vu-Topper RM

A set is called countable if, and only if, it is _____.

➢**Countably infinite and finite**

Question No:236 (Marks:1) Vu-Topper RM

If f(x) = x and g(x) = -x are both one to one functions then (f+g)(x) is also one to one function.

➢**False**

Question No:237 (Marks:1) Vu-Topper RM

If a is the 1st term and d be the common difference of an arithmetic sequence then the sequence is a, a+d, a+2d, a+3d….

➢**True**

Question No:238 (Marks:1) Vu-Topper RM

x+a, x+3a, x+5…… is a/an_____.

➢**Arithmetic sequence**

Question No:239 (Marks:1) Vu-Topper RM

inverse of an injective function may not be a function.

➢**True**

Question No:240 (Marks:1) Vu-Topper RM

$y=x^3$ is a graph of bijective function form R to R

➢**False**

Question No:241 (Marks:1) Vu-Topper RM

Two functions 'f' and 'g' from x to y are said to be equal if and only if _____.

➢**F(x)=g(x) for all 'x' belongs to x**

Question No:242 (Marks:1) Vu-Topper RM

Common ration in sequence '36, 12, 4 , 4/3, …..' is ……

➢**1/3**

Question No:243 (Marks:1) Vu-Topper RM

This document is available on studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

A set is called finite if, and only if, is the _____ or there is _____.

➢**Empty set or one-to-one**

Question No:244 (Marks:1) Vu-Topper RM

Let $f(X)=x^2-1$ and $g(x)=x+1$ define functions f and g from R to R, then (f/g)x

➢**x-1**

Question No:245 (Marks:1) Vu-Topper RM

A graph of a function f is one to one iff every horizontal line intersects the graph in at most one point.

➢**True**

Question No:246 (Marks:1) Vu-Topper RM

Let g be a function defined by g(x) = x+1. Then the composition of (gog).

➢**X+2**

Question No:247 (Marks:1) Vu-Topper RM

F:X->Y that is both one to one and onto is called a _____.

➢**Bijective function**

Question No:248 (Marks:1) Vu-Topper RM

What does 'I' denotes in a geometric sequence?

➢**Common ratio**

Question No:249 (Marks:1) Vu-Topper RM

Let A={1,2,3,4} and B={7} then the constant function from A to B is _____.

➢**Both one to one and onto**

Question No:250 (Marks:1) Vu-Topper RM

N! is defined to be _____.

➢**The product of the integers from 1 to n**

Question No:251 (Marks:1) Vu-Topper RM

A set is called countable if, and only if, it is _____.

➢**Both b and c**

Question No:252 (Marks:1) Vu-Topper RM

Which of the following is the example of an alternating sequence?

➢**$C_n=n/n+1$ for $n \geq 0$**

Question No:253 (Marks:1) Vu-Topper RM

X+a, x+3a, x+5a….. is an_____

➢**Arithmetic sequence**

Question No:254 (Marks:1) Vu-Topper RM

0, -5, -10, -15, … is an _____.

➢**Arithemetic sequence**

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:255 (Marks:1) Vu-Topper RM

5, 9, 13, 17, … is an _____.

➢**Arithemetic sequence**

Question No:256 (Marks:1) Vu-Topper RM

An important data type in computer programming consists of _____.

➢**Finite sequence**

Question No:257 (Marks:1) Vu-Topper RM

One-to-one correspondence means the condition of _____.

➢**One-one and onto ….both**

Question No:258 (Marks:1) Vu-Topper RM

Cardinality means the total number of elements in a set.

➢**True**

Question No:259 (Marks:1) Vu-Topper RM

Inverse of a function may not be a function.

➢**True**

Question No:260 (Marks:1) Vu-Topper RM

The functions 'f' and 'g' are inverse of each other if and only if their composition gives _____.

➢**Identity function**

Question No:261 (Marks:1) Vu-Topper RM

A set is called finite if , and only if, it is the _____ or there is _____.

➢**Empty set or one-to-one**

Question No:262 (Marks:1) Vu-Topper RM

Let f and g be the two functions from R to R defined by f(x) = IXI and g(x) = square root x^2 for all xeR, then _____.

➢**F(x) is not equal to g(x)**

Question No:263 (Marks:1) Vu-Topper RM

If $f^{-1}(x) = 6-x/2$ then $f^{-1}(2)$ is _____.

➢**2**

Question No:264 (Marks:1) Vu-Topper RM

Let f(2)=3, g(2)=3, f(4)=1 and g(4)=2 then the value of fog(4) is _____.

➢**3**

Question No:265 (Marks:1) Vu-Topper RM

An important date type in computer programming consists of _——.

➢**Finite sequence**

Question No:266 (Marks:1) Vu-Topper RM

This document is available on studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Let f(x)=x+3 then $f^{-1}(x)$ is _____.

➢**X-3**

Question No:267 (Marks:1) Vu-Topper RM

If f:X->Y amdg:Y->Z are both onto function. Then gof : X->Z is _____.

➢**One-to-one function**

Question No:268 (Marks:1) Vu-Topper RM

The functions 'f' and 'g' are inverse of each other if and only if their composition gives

➢**Identity function**

Question No:269 (Marks:1) Vu-Topper RM

The two function 'f' and 'g' are equal if _____.

➢**F(x) = 3x and g(x) = $6x^2$+3x/2x+1 for all xeR**

Question No:270 (Marks:1) Vu-Topper RM

Two functions 'f' and 'g' from X To Y are said to be equal if and only if——.

➢**F(x) and g(x) for all 'x' belongs to X**

Question No:271 (Marks:1) Vu-Topper RM

Which of the following set is the domain of a sequence?

➢**Set of natural numbers**

Question No:272 (Marks:1) Vu-Topper RM

If 1st term of a geometric sequence is 2 and common ratio is 1/2 , then 3rd term of sequence is _____.

➢**1/2**

Question No:273 (Marks:1) Vu-Topper RM

Composition of a function is a commutative operation.

➢**True**

Question No:274 (Marks:1) Vu-Topper RM

If a function (gof)(x): X-> Z is defined as (gof)(x)=g(f(x)) for all xeX. Then the function _____ is known as composition of f and g.

➢ **(g o f)**

Question No:275 (Marks:1) Vu-Topper RM

A set is countably infinite if and only if and only if, it has the same cardinality as the set of _____.

➢**Positive integers**

Question No:276 (Marks:1) Vu-Topper RM

The 3rd term of the sequence $b_n=5^n$ is _____.

➢**125**

Question No:277 (Marks:1) Vu-Topper RM

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

If f is a bijective function then $(f^{-1}(f(x))$ Is equal to _____.

➢**X**

Question No:278 (Marks:1) Vu-Topper RM

Let f(x) = x and g(x) = -x for all xeR, then (f+g)(x) is ____.

➢ **0**

Question No:279 (Marks:1) Vu-Topper RM

An infinite sequence may have only a finite number of values.

➢**True**

Question No:280 (Marks:1) Vu-Topper RM

The functions fog and gof are always equal .

➢**False**

Question No:281 (Marks:1) Vu-Topper RM

If f and g are two one-to-one functions, then their composition that is gof is one-to-one.

➢**True**

Question No:282 (Marks:1) Vu-Topper RM

A function whose range consists of only one element is called _____.

➢**Constant function**

Question No:283 (Marks:1) Vu-Topper RM

Let X ={1,5,9} and Y={3,4,7}. Define a function f from X to Y such that f(1)=7, f(5)=3, f(9)=4 then which of the following statement about 'f' is true?

➢**F is both one-to-one and onto**

Question No:284 (Marks:1) Vu-Topper RM

$Y=x^3$ is a graph of bijective function from R to R.

➢**True**

Question No:285 (Marks:1) Vu-Topper RM

A function whose inverse function exists is called a/an_____.

➢**Invertible**

Question No:286 (Marks:1) Vu-Topper RM

Let F and g be two functions defined by f(x)= x+2 and g(x)= 2x+1. Then the composition of f and g is _____.

➢**2x + 5**

Question No:287 (Marks:1) Vu-Topper RM

If r is a positive real number, then the value of r in 3, r,r=-27r is

➢**-9**

Question No:288 (Marks:1) Vu-Topper RM

The _____ of the terms of a sequence forms a series.

This document is available on studocu

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

➢**Sum**

Question No:289 (Marks:1) Vu-Topper RM

The sum of first five whole number is _____.

➢**10**

Question No:290 (Marks:1) Vu-Topper RM

If $f_k=f_k-1+f_k-2$then $f_0=1$, $f_1=2$, then f_2= _____.

➢**3**

Question No:291 (Marks:1) Vu-Topper RM

Let f(x) = 3x and g(x) = x + 2 define functions f and g from R to R, then (f.g)(x) is _____.

➢**$3x^2 + 6x$**

Question No:292 (Marks:1) Vu-Topper RM

Let R be a relation on a set A. If R is reflexive then its compliment is _____ .

➢**Irreflexive**

Question No:293 (Marks:1) Vu-Topper RM

If A = Set of students of virtual university then A has been written in the _____.

➢**Descriptive form**

Question No:294 (Marks:1) Vu-Topper RM

If a function (g o f)(x):X→Z is defined as (g o f)(x) = g(f(x)) for all x ∈ X. Then the function _____ is known as composition of f and g.

➢ **(g o f)**

Question No:295 (Marks:1) Vu-Topper RM

If X and Y are independent random variables and a and b are constants, then Var(aX + bY)is equal to

➢**aVar(X) + bVar(Y)**

Question No:296 (Marks:1) Vu-Topper RM

Let A = {1, 2, 3} and B = {2, 4} then number of functions from A to B are _____.

➢**8**

Question No:297 (Marks:1) Vu-Topper RM

p is equivalent to q' means _____.

➢**p is necessary and sufficient for q.**

Question No:298 (Marks:1) Vu-Topper RM

Let A and B be subsets of U with n(A) = 12, n(B) = 15, n(A') = 17, and n(A intersection B) = 8, then n(U)=_____ .

➢**29**

Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)

Question No:299 (Marks:1) Vu-Topper RM

For the following relation to be a function, x can not be what values?
R = {(2,4), (x,1), (4,2), (5,6)}
➢x cannot be 2, 4 or 5

Question No:300 (Marks:1) Vu-Topper RM

Find the number of the word that can be formed of the letters of the word "ELEVEN".
➢120

This document is available on studocu
Downloaded by Shahzeb Mughal (shahzebvip01@gmail.com)