ZOO-202 ... Chordates .

MID+FINAL TERM... Spring 2022

シ Effort By SAFDAR RIND シ

OBJ ECTIVE

Mid Term:

1 The organisms belonging to the Protochordata are generally known as the *lower chordates.*

2 All blood flowing anteriorly passes into a series of blood

sinuses, called the *glomerulus* , at the base of the proboscis.

3 *cillia and mucus* helps in acorn worm in feeding

4 The earliest tetrapods (having four legs) are the *amphibians*

5 which one is not present in tunicate

A.. pharynx B... atrium

C endostyl D. glomerulus

6 Tunicate blood plasma is *colorless*

7 in cephalochordates mucus secreted by the *endostyle* .

8 *Bony or cartilaginous* vertebrae that completely or partially replace the

notochord characterizes the vertebrates.

9 A group of ancient eel-like animals, the

conodonts , are known from fossils that date back about 510 million years.

10 *Ostracoderms* are extinct agnathans that belonged to several classes.

11 The larval stage of the primitive lamprey, known as an *ammocoetes larva*

12 Salamander belonging to order *Caudata*

13 Salamander *30 to 90%* exchange through skin.

14 blood secretes gases (various mixtures of nitrogen and oxygen) into the swim bladder using a countercurrent exchange mechanism in a vascular network called *"miraculous* net"

14 Larva of frog is called *tadpole*

15 Metamorphosis is under the control of *neurosecretions* of the hypothalamus.

16 All reptiles periodically shed the outer, epidermal layers of the skin in a

process called *ecdysis*

17 largest group of living vertebrate is called cartilaginous fishes*

18 The male Brazilian catfish (Loricaria typhys) broods embryos in an enlarged *lower lip*.

19 *Semicircular canals* detect rotational movement in vertebrates.

20 In fishes *muscles* are in zigzag. species) and Asymmetron (about six species).

FINAL TERM

1 Cilia And Mucus assist acorn worms in feeding.

2 The ASCIDIANS comprise the largest class of tunicates.

3 Members of the subphylum cephalochordata are called LANCELETS.

mucus secreted by the endostyle in cephalochordates.

4 WHALES don't have hair on the adult stage.

5 The male Brazilian catfish (Loricaria typhys) broods embryos in an enlarged LOWER LIP.

6 The PHOTOPERIOD is an important migratory cue for many birds, particularly for birds in temperate zones.

7 Larval stage of the frog is TADPOLE.

8 •A coat of hair, called PELAGE, usually consists of two kinds of hair.

9•Hair color depends on the amount of pigment...(MELANIN).

10Most mammals have two sets of teeth during their lives.

11Even the giraffe and the whale have SEVEN neck vertebrae, which are greatly elongate or

compressed, respectively.

12Chiroptera include BATS.

13•The middle ear contains **THREE** ear ossicles that conduct vibrations to the inner ear.

14 Cats, however, are really staking claim to their territory, using odors from **FACIAL SCENT GLANDS.**

15 In Mammals, Fertilization-usually occurs in the upper third of the **OVIDUCT** within hours of copulation.

16 The only venomous lizards are the gila MONSTER (Heloderma suspectum) and the **MEXICAN BEADED LIZARD.**

17 The venoms of vipers are primarily **HEMOTOXINS**.

18 In snakes and some lizards, the upper and lower eyelids fuse in the embryo to form a protective window of clear skin, called the **SPECTACLE**.

19 Sperm may be stored for up to **FOUR** years in some turtles, and up to six years in some snakes.

20 Birds also have a reduced number of skull bones, and a lighter, keratinized sheath called a **BILL** replaces the teeth.

21 The **CROP** of pigeons produces “pigeon’s milk.

22 A special voice box, called the **SYRINX**.

23 Birds maintain **38 TO 45** body temperature.

24.Cones are especially **CONCENTRATED (1,000,000/MM2)** at a focal point called the fovea.

25 Like amphibians, reptiles possess **TWO atria** that are completely separated in the adult and have veins from the body and lungs emptying into them.

26. • A kidney with many more blood-filtering units, called **NEPHRONS**.

27 Eoalulavis, was found in 115- million-year-old deposits and had a wingspan of 17 cm to (roughly the same as a goldfinch). This fossil showed that Eoalulavis had a wing structure called an **ALULA**.